

## COMMENT

10.1002/2014JF003211

This article is a comment on *Bolla Pittaluga and Imran* [2014] doi:10.1002/2013JF002812.

## Correspondence to:

J. Peakall,  
j.peakall@leeds.ac.uk

## Citation:

Peakall, J., S. E. Darby, R. M. Dorrell, D. R. Parsons, E. J. Sumner, and R. B. Wynn (2014), Comment on "A simple model for vertical profiles of velocity and suspended sediment concentration in straight and curved submarine channels" by M. Bolla Pittaluga and J. Imran, *J. Geophys. Res. Earth Surf.*, 119, 2070–2073, doi:10.1002/2014JF003211.

Received 14 MAY 2014

Accepted 26 AUG 2014

Accepted article online 30 AUG 2014

Published online 29 SEP 2014

# Comment on "A simple model for vertical profiles of velocity and suspended sediment concentration in straight and curved submarine channels" by M. Bolla Pittaluga and J. Imran

J. Peakall<sup>1</sup>, S. E. Darby<sup>2</sup>, R. M. Dorrell<sup>1</sup>, D. R. Parsons<sup>3</sup>, E. J. Sumner<sup>4,5</sup>, and R. B. Wynn<sup>6</sup>

<sup>1</sup>School of Earth and Environment, University of Leeds, Leeds, UK, <sup>2</sup>Geography and Environment, University of Southampton, Southampton, UK, <sup>3</sup>Department of Geography, Environment and Earth Sciences, University of Hull, Hull, UK, <sup>4</sup>Ocean and Earth Science, University of Southampton, Southampton, UK, <sup>5</sup>Monterey Bay Aquarium Research Institute, Moss Landing, California, USA, <sup>6</sup>National Oceanography Centre, Southampton, UK

## 1. Introduction

*Bolla Pittaluga and Imran* [2014] present a model of flow in straight and curved submarine channels which discusses the influence of flow stratification on flow field dynamics. Here we examine why this elegant model for submarine channels breaks down in the case of bend flow, highlighting that it does not incorporate some key physics. We also show how more complex modeling is required to produce realistic secondary flows in submarine channels. The associated model for submarine channel bend sedimentation is shown here to conflict with observations from physical modeling, field data, theory, and other numerical modeling. We discuss sedimentation in submarine channel bends and demonstrate that this is a function of the three-dimensional helical flow field.

## 2. Controls on Secondary Flows in Submarine Channels

Secondary flow at submarine channel bend apices can crucially be either the same as rivers (normal) with inward directed basal flows or opposite to rivers (reversed) with outward directed basal flows, as first shown by *Kassem and Imran* [2004] and *Keevil et al.* [2006], respectively. *Bolla Pittaluga and Imran* [2014] provide a two-dimensional secondary flow model that incorporates stratification. This produces the key result that increasing stratification increases the probability of river-like secondary flow, with the absence of stratification favoring reversed secondary flow conditions. Such a result appears paradoxical, since a key control on secondary flow orientation, assuming other parameters such as channel planform and cross section are constant, is the position of the downstream velocity maximum,  $U_{\max}$ , with low  $U_{\max}$  positions enhancing the probability of flow reversal [*Corney et al.*, 2008; *Giorgio Serchi et al.*, 2011]. Given that stratification and downstream velocity are coupled, increasing stratification would be expected to lead to lower values of  $U_{\max}$  and therefore increased likelihood of secondary flow reversal [*Parsons et al.*, 2010; *Giorgio Serchi et al.*, 2011]. The same paradoxical relationship observed by *Bolla Pittaluga and Imran* [2014] between stratification and basal secondary flow orientations was also produced in the two-dimensional closure models of *Dorrell et al.* [2013]; for example, see their Figure 7. These model results stem from the requirement that the net lateral fluid and mass transport (material) fluxes vanish for secondary flows constrained within a two-dimensional plane [*Dorrell et al.*, 2013; *Bolla Pittaluga and Imran*, 2014]. While the net lateral material fluxes are constrained to be zero, the sum lateral flux in the near-bed region must exactly oppose the sum lateral flux between the near-bed region and the flow interface. This in turn leads to reversed secondary flows being favored by limited stratification and normally oriented secondary flows becoming more likely as a function of increasing stratification [*Dorrell et al.*, 2013; *Bolla Pittaluga and Imran*, 2014].

Submarine channel bend flows do not exhibit zero material fluxes around a bend, instead they exhibit prominent flow superelevation at bend apices [*Imran et al.*, 1999] and, therefore, positive radial material fluxes upstream of the bend apex and negative fluxes downstream. The magnitude of superelevation in submarine channel bends is around 2 orders higher than in river bends, for a given width, reflecting the differences in density between the channelized flow and the surrounding ambient fluid; typical transverse water slopes at the apex are order  $10^{-2}$  in submarine bends [*Komar*, 1969; *Pirmez and Imran*, 2003], versus order  $10^{-4}$  in rivers [*Leopold*, 1982]. Such high superelevation in submarine channel bends is also reflected by

outer channel bend levees being consistently higher than inner channel bend levees [e.g., Pirmez and Imran, 2003]. In addition, radial material fluxes will be significantly enhanced by any flow overspill at bends, which is thought to occur frequently in submarine channels [Peakall *et al.*, 2000; Mohrig and Buttles, 2007; Dorrell *et al.*, 2014], and from any variations from a uniform channel bathymetry [Dorrell *et al.*, 2013; Sumner *et al.*, 2014]. As a consequence of these factors, a three-dimensional framework with a nonzero flux condition at bend apices is required for realistic modeling. Dorrell *et al.* [2013] identified the importance of this three-dimensional framework and implemented a closure of the secondary flow dynamics that incorporated downstream convective radial transport. They validated the model against fully three-dimensional laboratory data and numerical models [Corney *et al.*, 2006; Abad *et al.*, 2011] and demonstrated that radial material fluxes are the crucial control on the vertical structure of secondary flow.

A key outcome of the three-dimensional modeling of Dorrell *et al.* [2013] is that stratified flows with nonnegligible material fluxes oriented toward the outer bank (i.e., as superelevation is increasing around a bend) will dominantly exhibit basal flows that are reversed relative to rivers. This finding of Dorrell *et al.* [2013] is further supported by the submarine channel bend measurements of Sumner *et al.* [2014] and past work on highly stratified flows in curved estuaries [Chant and Wilson, 1997; Seim and Gregg, 1997; Lacy and Monismith, 2001; Nidzieko *et al.*, 2009], all of which exhibit reversed secondary circulation relative to rivers. We note that this result is the opposite of that predicted by the simple two-dimensional closure model of Bolla Pittaluga and Imran [2014] and the two-dimensional model implemented by Dorrell *et al.* [2013], which both predict that stratified flows are more likely to exhibit river-like secondary flows.

### 3. HelicalFlow-Driven Sedimentation in Submarine Channels

Bolla Pittaluga and Imran [2014] replicate the argument of Abad *et al.* [2011] that the implication of reversed secondary flows is to create deposition on the outer channel bank, in contrast to the inner bend accumulations (point bars) associated with normal river-dominated secondary flows. In so doing, they overlook the direct evidence for traction-dominated inner bend sediment accumulation during reversed secondary flow (in the absence of significant Coriolis forcing) that is derived from physical experiments [Peakall *et al.*, 2007; Amos *et al.*, 2010; Cossu and Wells, 2013; Wells and Cossu, 2013], numerical modeling [Darby and Peakall, 2012], and field outcrops [Pyles *et al.*, 2012].

The proposed model of Bolla Pittaluga and Imran [2014] is based on a two-dimensional consideration of channel bends, where in the case of normal river secondary circulation, “sediment is eroded from the outer bank to be deposited in the inner bank” (p. 500). However, fluvial workers have long recognized that sediment is not moved directly across the channel but is instead eroded from upstream concave banks into the downstream convex bar; reflecting the dominance of along-stream sediment transport [Friedkin, 1945; Nelson and Smith, 1989; Bridge, 1992]. As such, sediment accumulation is a three-dimensional process that is in turn linked to the three-dimensional flow field. Furthermore, deposition dominantly occurs where there is a convergence of streamlines and therefore sediment flux [Nelson and Smith, 1989]. In river-like secondary flows, this streamline convergence occurs prior to the bend apex producing sedimentation around the bend apex [Nelson and Smith, 1989]. In contrast, flow is still diverging (outwardly directed) at bend apices under reversed secondary circulation conditions, such that convergence is delayed to farther around the bend [Keevil *et al.*, 2006, Figure 6; Amos *et al.*, 2010, Figure 5]. This spatial lag in the convergence of sediment flux therefore leads to inner bank deposition being located farther downstream, past the bend apex [Keevil *et al.*, 2006; Peakall *et al.*, 2007; Amos *et al.*, 2010; Darby and Peakall, 2012].

Bolla Pittaluga and Imran [2014] support their two-dimensional outer bank model of channel sedimentation by comparison with the work of Janocko *et al.* [2013], who across their series of experiments observed deposition at all points along both inner and outer banks. As such, only a small part of the experimental data set of Janocko *et al.* [2013] fits the Bolla Pittaluga and Imran [2014] model. In so doing, Bolla Pittaluga and Imran [2014] chose an example in which additional processes were operating. The work of Janocko *et al.* [2013] includes the following: (i) both traction-dominated sedimentation as occurs in point bars and the 2-D Bolla Pittaluga and Imran [2014] model, and large-scale deposition from suspension as flows collapse; (ii) flow separation in the lee of sharp bends which leads to deposition in these zones [Straub *et al.*, 2008, 2011]; (iii) runup and collapse, and deposition from flows against outer channel banks [Straub *et al.*, 2008, 2011]; and (iv) the interaction of overbank and intrachannel flow, most notably where overbank flow reenters the channel

[Amos *et al.*, 2010; Ezz and Imran, 2014]. This combination of processes explains the presence of deposition in such a wide range of positions in the Janocko *et al.* [2013] experiments. In particular, in suspension-dominated flows the orientation of the basal secondary flow will have little effect on sediment position, and deposits preferentially occur where flows interact with outer banks producing outer bank bars [Nakajima *et al.*, 2009; Huang *et al.*, 2012; Ezz *et al.*, 2013; Janocko *et al.*, 2013]. In contrast to the range of processes observed in the Janocko *et al.* [2013] experiments, the works of Peakall *et al.* [2007], Amos *et al.* [2010] (excluding a very high sinuosity channel which exhibited overbank flow reentering the channel), and Darby and Peakall [2012], consist of purely tractional transport and do not exhibit flow separation zones, runup and collapse, or significant overbank—in-channel interaction. As such, these studies are directly comparable to the conditions postulated in the Bolla Pittaluga and Imran [2014] model, while nevertheless producing results that conflict, demonstrating inner bend accumulation of sediment in reversed secondary flows.

#### 4. Conclusions

The elegant two-dimensional model of Bolla Pittaluga and Imran [2014] for velocity and density distributions in straight submarine channels breaks down for curved flows, since it does not incorporate the critical three-dimensional advective terms which have been recognized in submarine channel bend flows, highly stratified curved estuarine flows, and the numerical modeling of submarine channels of Dorrell *et al.* [2013]. The latter study was validated against both physical modeling and three-dimensional numerical modeling data sets [Dorrell *et al.*, 2013]. The key conclusion of Bolla Pittaluga and Imran [2014], that stratification enhances river-like secondary flow in submarine channel bends, is shown to be incorrect. Rather, increasing stratification leads to dominantly reversed secondary circulation in submarine channel bends, as shown by submarine channel bend data, three-dimensional numerical modeling, and analogous studies of highly stratified curved estuaries. Similarly, it is shown that three-dimensional models must be considered for submarine channel bend sedimentation. The two-dimensional model of Bolla Pittaluga and Imran [2014] predicts that outer bank sedimentation will occur from traction-dominated flows with reversed secondary circulation, yet physical and numerical modeling, and field data from traction-dominated reversed secondary flows, have demonstrated that sedimentation occurs instead at the inner bank, albeit with the locus of sedimentation translated farther around the bend.

#### Acknowledgments

This work was supported by Natural Environment Research Council grants NE/F020511/1, NE/F020120/1, and NE/F020279/1.

#### References

- Abad, J. D., O. E. Sequeiros, B. Spinewine, C. Pirmez, M. H. Garcia, and G. Parker (2011), Secondary current of saline underflow in a highly meandering channel: Experiments and theory, *J. Sediment. Res.*, *81*, 787–813.
- Amos, K. J., J. Peakall, P. W. Bradbury, M. Roberts, G. Keevil, and S. Gupta (2010), The influence of bend amplitude and planform morphology on flow and sedimentation in submarine channels, *Mar. Pet. Geol.*, *27*(7), 1431–1447.
- Bolla Pittaluga, M., and J. Imran (2014), A simple model for vertical profiles of velocity and suspended sediment concentration in straight and curved submarine channels, *J. Geophys. Res. Earth Surf.*, *119*, 483–503, doi:10.1002/2013JF002812.
- Bridge, J. S. (1992), A revised model for water-flow, sediment transport, bed topography and grain-size sorting in natural river bends, *Sedimentology*, *40*(4), 801–810.
- Chant, R. J., and R. E. Wilson (1997), Secondary circulation in a highly stratified estuary, *J. Geophys. Res.*, *102*, 23,207–23,215, doi:10.1029/97JC00685.
- Corney, R. K. T., J. Peakall, D. R. Parsons, L. Elliott, K. J. Amos, J. L. Best, G. Keevil, and D. B. Ingham (2006), The orientation of helical flow in curved channels, *Sedimentology*, *53*(2), 249–257.
- Corney, R. K. T., J. Peakall, D. R. Parsons, L. Elliott, J. L. Best, R. E. Thomas, G. Keevil, D. B. Ingham, and K. J. Amos (2008), Reply to Discussion of Imran *et al.* on “The orientation of helical flow in curved channels” by Corney *et al.*, *Sedimentology*, *55*(1), 241–247.
- Cossu, R., and M. G. Wells (2013), The evolution of submarine channels under the influence of Coriolis forces: Experimental observations of flow structures, *Terra Nova*, *25*, 65–71, doi:10.1111/ter.12006.
- Darby, S. E., and J. Peakall (2012), Modelling the equilibrium bed topography of submarine meanders that exhibit reversed secondary flows, *Geomorphology*, *164*, 99–109.
- Dorrell, R. M., S. E. Darby, J. Peakall, E. J. Sumner, D. R. Parsons, and R. B. Wynn (2013), Superelevation and overspill control secondary flow dynamics in submarine channels, *J. Geophys. Res. Oceans*, *118*, 3895–3915, doi:10.1002/jgrc.20277.
- Dorrell, R. M., S. E. Darby, J. Peakall, E. J. Sumner, D. R. Parsons, and R. B. Wynn (2014), The critical role of stratification in submarine channels: Implications for channelization and long runout of flows, *J. Geophys. Res. Oceans*, *119*, 2620–2641, doi:10.1002/2014JC009807.
- Ezz, H., and J. Imran (2014), Curvature-induced secondary flow in submarine channels, *Environ. Fluid Mech.*, *14*, 343–370, doi:10.1007/s10652-014-9345-4.
- Ezz, H., A. Cantelli, and J. Imran (2013), Experimental modeling of depositional turbidity currents in a sinuous submarine channel, *Sediment. Geol.*, *290*, 175–187.
- Friedkin, J. F. (1945), *A Laboratory Study of the Meandering of Alluvial Rivers*, 40 pp., U.S. Waterways Experimental Station, Vicksburg, Miss.
- Giorgio Serchi, F. G., J. Peakall, D. B. Ingham, and A. D. Burns (2011), A unifying computational fluid dynamics investigation on the river-like to river-reversed secondary circulation in submarine channel bends, *J. Geophys. Res.*, *116*, C06012, doi:10.1029/2010JC006361.
- Huang, H., J. Imran, and C. Pirmez (2012), The depositional characteristics of turbidity currents in submarine sinuous channels, *Mar. Geol.*, *329–331*, 93–102.

- Imran, J., G. Parker, and C. Pirmez (1999), A nonlinear model of flow in meandering submarine and subaerial channels, *J. Fluid Mech.*, *400*(1), 295–331.
- Janocko, M., M. B. J. Cartigny, W. Nemec, and E. W. M. Hansen (2013), Turbidity current hydraulics and sediment deposition in erodible sinuous channels: Laboratory experiments and numerical simulations, *Mar. Pet. Geol.*, *41*, 222–249.
- Kassem, A., and J. Imran (2004), Three-dimensional modeling of density current. II. Flow in sinuous confined and unconfined channels, *J. Hydraul. Res.*, *42*(6), 591–602.
- Keevil, G., J. Peakall, J. L. Best, and K. J. Amos (2006), Flow structure in sinuous submarine channels: Velocity and turbulence structure of an experimental submarine channel, *Mar. Geol.*, *229*(3), 241–257.
- Komar, P. D. (1969), The channelized flow of turbidity currents with application to Monterey deep-sea fan channel, *J. Geophys. Res.*, *74*, 4544–4558, doi:10.1029/JC074i018p04544.
- Lacy, J. R., and S. G. Monismith (2001), Secondary currents in a curved, stratified, estuarine channel, *J. Geophys. Res.*, *106*, 31,283–31,302, doi:10.1029/2000JC000606.
- Leopold, L. B. (1982), Water surface topography in river channels and implications for meander development, in *Gravel-Bed Rivers*, pp. 359–383, John Wiley, New York.
- Mohrig, D., and J. Buttle (2007), Deep turbidity currents in shallow channels, *Geology*, *35*, 155–158.
- Nakajima, T., J. Peakall, W. D. McCaffrey, D. A. Paton, and P. J. P. Thompson (2009), Outer-bank bars: A new intra-channel architectural element within sinuous submarine slope channels, *J. Sediment. Res.*, *79*(12), 872–886.
- Nelson, J. M., and J. D. Smith (1989), Flow in meandering channels with natural topography, in *River Meandering, Water Resour. Monogr.*, vol. 12, edited by S. Ikeda and G. Parker, pp. 69–102, AGU, Washington, D. C.
- Nidzieko, N. J., J. L. Hench, and S. G. Monismith (2009), Lateral circulation in well-mixed and stratified estuarine flows with curvature, *J. Phys. Oceanogr.*, *39*, 831–851, doi:10.1175/2008JPO4017.1.
- Parsons, D. R., J. Peakall, A. E. Aksu, R. D. Flood, R. N. Hiscott, S. Besiktepe, and D. Moulard (2010), Gravity-driven flow in a submarine channel bend: Direct field evidence of reversed helical flow reversal, *Geology*, *38*, 1063–1066, doi:10.1130/G31121.1.
- Peakall, J., B. McCaffrey, and B. Kneller (2000), A process model for the evolution, morphology, and architecture of sinuous submarine channels, *J. Sediment. Res.*, *70*, 434–448.
- Peakall, J., K. J. Amos, G. Keevil, P. W. Bradbury, and S. Gupta (2007), Flow processes and sedimentation in submarine channel bends, *Mar. Pet. Geol.*, *24*(6), 470–486.
- Pirmez, C., and J. Imran (2003), Reconstruction of turbidity currents in Amazon Channel, *Mar. Pet. Geol.*, *20*(6), 823–849.
- Pyles, D. R., M. Tomasso, and D. C. Jennette (2012), Flow processes and sedimentation associated with erosion and filling of sinuous submarine channels, *Geology*, *40*(2), 143–146.
- Seim, H. E., and M. C. Gregg (1997), The importance of aspiration and channel curvature in producing strong vertical mixing over a sill, *J. Geophys. Res.*, *102*, 3451–3472, doi:10.1029/96JC03415.
- Straub, K. M., D. Mohrig, B. McElroy, J. Buttle, and C. Pirmez (2008), Interactions between turbidity currents and topography in aggrading sinuous submarine channels: A laboratory study, *Geol. Soc. Am. Bull.*, *120*, 368–385.
- Straub, K. M., D. Mohrig, J. Buttle, B. McElroy, and C. Pirmez (2011), Quantifying the influence of channel sinuosity on the depositional mechanics of channelized turbidity currents: A laboratory study, *Mar. Pet. Geol.*, *28*, 744–760.
- Sumner, E. J., J. Peakall, R. M. Dorrell, D. R. Parsons, R. B. Wynn, S. E. Darby, S. D. McPhail, J. Perrett, A. Webb, and D. White (2014), Driven around the bend: Spatial evolution and controls on the orientation of helical bend flow in a natural submarine gravity current, *J. Geophys. Res.*, *119*, 898–913, doi:10.1002/2013JC009008.
- Wells, M., and R. Cossu (2013), The possible role of Coriolis forces in structuring large-scale sinuous patterns of submarine channel-levee systems, *Philos. Trans. R. Soc. A*, *371*, 20120366, doi:10.1098/rsta.2012.0366.