

EQUIVARIANT VECTOR BUNDLES OVER CLASSIFYING SPACES FOR PROPER ACTIONS

DIETER DEGRIJSE AND IAN J. LEARY

ABSTRACT. Let G be an infinite discrete group and let \underline{EG} be a classifying space for proper actions of G . Every G -equivariant vector bundle over \underline{EG} gives rise to a compatible collection of representations of the finite subgroups of G . We give the first examples of groups G with a cocompact classifying space for proper actions \underline{EG} admitting a compatible collection of representations of the finite subgroups of G that does not come from a G -equivariant (virtual) vector bundle over \underline{EG} . This implies that the Atiyah-Hirzebruch spectral sequence computing the G -equivariant topological K-theory of \underline{EG} has non-zero differentials. On the other hand, we show that for right angled Coxeter groups this spectral sequence always collapses at the second page and compute the K-theory of the classifying space of a right angled Coxeter group.

1. INTRODUCTION

Let G be an infinite discrete group and \mathcal{F} be the family of finite subgroups of G . Recall that the orbit category $\mathcal{O}_{\mathcal{F}}G$ is a category whose objects are the transitive G -sets G/H , for all $H \in \mathcal{F}$, and whose morphism are all G -equivariant maps between the objects. A classifying space for proper actions of G , denoted by \underline{EG} , is a proper G -CW-complex such that the fixed point set \underline{EG}^H is contractible for every $H \in \mathcal{F}$. The space \underline{EG} is said to be cocompact if the orbit space $G \backslash \underline{EG} = \underline{BG}$ is compact. Many interesting classes of groups G have cocompact models for \underline{EG} , for example cocompact lattices in Lie groups, mapping class groups of surfaces, $\text{Out}(F_n)$, CAT(0)-groups and word-hyperbolic groups. We refer the reader to [9] for more examples and details.

Now assume G is an infinite discrete group admitting a cocompact classifying space for proper actions \underline{EG} . If

$$\xi: E \rightarrow \underline{EG}$$

is a G -equivariant complex vector bundle over \underline{EG} (see Definition 2.3) and x is a point of \underline{EG} , then the fiber $\xi^{-1}(x)$ is a complex representation of the finite isotropy group G_x . The connectivity of the fixed point sets of \underline{EG} ensures that these representations are compatible (see Definition 2.1) with one another as x and hence G_x varies. Therefore, every G -equivariant complex vector bundle over \underline{EG} gives rise to a compatible collection of complex representations of the finite subgroups of G , and hence to an element of

$$\lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} R(H).$$

Date: May 19, 2016.

The first author was supported by the Danish National Research Foundation through the Centre for Symmetry and Deformation (DNRF92).

Here, $\lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} R(H)$ is the limit over the orbit category $\mathcal{O}_{\mathcal{F}}G$ of the contravariant representation ring functor

$$R(-): \mathcal{O}_{\mathcal{F}}G \rightarrow \text{Ab} \quad G/H \mapsto R(H).$$

Denoting the Grothendieck group of the abelian monoid of isomorphism classes of complex G -vector bundles over $\underline{E}G$ by $K_G^0(\underline{E}G)$, one obtains a map

$$\varepsilon_G: K_G^0(\underline{E}G) \rightarrow \lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} R(H)$$

that maps a formal difference of (isomorphism classes) vector bundles (i.e. a virtual vector bundle) to a formal difference of (isomorphism classes) of compatible collections of representations of the finite subgroups of G . We say a compatible collection of representations of the finite subgroups of G can be realized as a (virtual) G -equivariant vector bundle over $\underline{E}G$ if there exists a (virtual) G -equivariant vector bundle over $\underline{E}G$ that maps to this collection under ε_G . One can also look at the corresponding situation for real (orthogonal) vector bundles and real (orthogonal) representations and obtain the map

$$\varepsilon_G: KO_G^0(\underline{E}G) \rightarrow \lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} RO(H).$$

The maps ε_G are equal to the edge homomorphisms of certain Atiyah-Hirzebruch spectral sequences converging to $K_G^*(\underline{E}G)$ and $KO_G^*(\underline{E}G)$ (see (1) and (2)). Lück and Oliver proved that (see Proposition 2.5) the map ε_G (real or complex) is rationally surjective, meaning that a high enough multiple of every element in the target of ε_G is contained in the image of ε_G . In the last paragraph of [12, p. 596] Lück and Oliver ask for an example of a group G admitting a cocompact classifying space for proper actions $\underline{E}G$ such that ε_G is not surjective. In Section 3 of this paper we give the first example of such a group in the complex case. In Section 4 we give the first example of such a group in the real case. We also construct examples of groups G admitting a cocompact $\underline{E}G$ with the following weaker property: G admits a compatible collection of representations for its finite subgroups that cannot be realized as a G -vector bundle over $\underline{E}G$. However, for these examples we cannot exclude the possibility that there exists a virtual vector bundle that maps to this collection of representations under ε_G . On the other hand, these examples are more explicit and lower dimensional.

In the final section we show that for a right angled Coxeter group W , every compatible collection of representations of the finite subgroups of W can be realized as a W -equivariant vector bundle over $\underline{E}W$, so that the map

$$\varepsilon_W: K_W^0(\underline{E}W) \rightarrow \lim_{W/H \in \mathcal{O}_{\mathcal{F}}W} R(H).$$

is always surjective. Moreover, we show that this map is actually an isomorphism and that (see Theorem 2.4)

$$K_W^1(\underline{E}W) = 0.$$

Using a version of the Atiyah-Segal completion theorem for infinite discrete groups proven by Lück and Oliver, we use these results to compute the complex K-theory of BW , the classifying space of W (see Corollary 5.6).

ACKNOWLEDGEMENT

We thank the referee, whose comments on an earlier version of this article were extremely helpful.

2. G -VECTOR BUNDLES AND ISOTROPY REPRESENTATIONS

Let G be a discrete group and let Γ be a Lie group. Let \mathcal{S} be a family of finite subgroups of G , i.e. any collection of finite subgroups of G that is closed under conjugation and passing to subgroups. The orbit category $\mathcal{O}_{\mathcal{S}}G$ is a category whose objects are the transitive G -sets G/H , for all $H \in \mathcal{S}$, and whose morphism are all G -equivariant maps between the objects.

Definition 2.1. [12, p. 590] Let X be a G -CW-complex. A (G, Γ) -bundle over X is a Γ -principal bundle $p: E \rightarrow X$, where E is a left G -space such that p is G -equivariant and such that the left G -action and the right Γ -action on E commute. We denote the set of isomorphism classes of (G, Γ) -bundles over X by $\text{Bld}_{(G, \Gamma)}(X)$. For $H \in \mathcal{F}$, let

$$\text{Rep}_{\Gamma}(H) = \text{Hom}(H, \Gamma)/\text{Inn}(\Gamma).$$

One can consider $\text{Rep}_{\Gamma}(-)$ as a contravariant functor from $\mathcal{O}_{\mathcal{S}}G$ to Sets. An element of the limit

$$A = ([\alpha_H])_{H \in \mathcal{S}} \in \lim_{G/H \in \mathcal{O}_{\mathcal{S}}G} \text{Rep}_{\Gamma}(H)$$

is called an \mathcal{S} -compatible collection of Γ -representations. Given such an element A , let \mathcal{S}_A be the family of subgroups of $G \times \Gamma$ consisting of conjugates of the subgroups of the form

$$\{(h, \alpha_H(h)) \mid h \in H\}$$

for all $H \in \mathcal{S}$ and let $E_{\mathcal{S}}(G, A)$ be the universal $G \times \Gamma$ -CW-complex for the family \mathcal{S}_A .

Lemma 2.2. [12, Lemma 2.4] For every \mathcal{S} -compatible collection of Γ -representations $A = ([\alpha_H])_{H \in \mathcal{S}}$ there exists a G -CW-complex $B_{\mathcal{S}}(G, A)$ with isotropy in \mathcal{S} satisfying the following properties.

- The quotient map

$$\pi: E_{\mathcal{S}}(G, A) \rightarrow \Gamma \backslash E_{\mathcal{S}}(G, A) = B_{\mathcal{S}}(G, A)$$

is a (G, Γ) -bundle over the G -CW-complex $B_{\mathcal{S}}(G, A)$.

- The (G, Γ) -bundle $\pi: E_{\mathcal{S}}(G, A) \rightarrow B_{\mathcal{S}}(G, A)$ is universal in the sense that for every G -CW-complex X with isotropy in \mathcal{S} there is an isomorphism

$$[X, \mathcal{B}_{\mathcal{S}}(G, A)]_G \xrightarrow{\cong} \text{Bld}_{(G, \Gamma)}(X)$$

given by pulling back the universal bundle π along a G -map $X \rightarrow B_{\mathcal{S}}(G, A)$.

- For every $S \in \mathcal{S}$, the fixed point set $B_{\mathcal{F}}(G, A)^H$ is homotopy equivalent to $BC_{\Gamma}(\alpha_H)$, the classifying space of the centralizer of the image of α_H in Γ .

If $\Gamma = \text{U}(n)$ ($\Gamma = \text{O}(n)$) and $\mathcal{S} = \mathcal{F}$, the family of all finite subgroups of G , then $\text{Rep}_{\Gamma}(H)$ is the set of isomorphism classes of n -dimensional complex (real) representations of H . In this case, an element of the limit

$$A = ([\alpha_H])_{H \in \mathcal{F}} \in \lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} \text{Rep}_{\Gamma}(H)$$

is called is called a compatible collection of complex (real) n -dimensional representations of the finite subgroups of G . For $H \in \mathcal{F}$, let $R(H)$ ($RO(H)$) be the complex (real) representation ring of H , i.e. the Grothendieck group of the abelian cancellative monoid of isomorphism classes of finite dimensional complex (real) representations of H . Note that $\text{Rep}_{\text{U}(n)}(H)$ is

naturally a subset of $R(H)$ and $\text{Rep}_{\text{O}(n)}(H)$ is naturally a subset of $RO(H)$. One can consider $R(-)$ as a functor from $\mathcal{O}_{\mathcal{F}}G$ to Ab . An element of the inverse limit

$$\alpha = ([\alpha_H])_{H \in \mathcal{F}} \in \lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} R(H)$$

is called a *compatible collection of complex virtual representations* of the finite subgroups of G . One has a natural embedding

$$\lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} \text{Rep}_{\text{U}(n)}(H) \subset \lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} R(H).$$

The analogous statements for $\text{O}(n, \mathbb{R})$ and RO also hold.

Now let X be a proper cocompact G -CW-complex, i.e. X has finite isotropy and the orbit space $G \backslash X$ has a finite number of cells, such that for every $H \in \mathcal{F}$, the fixed point set X^H is non-empty and connected.

Definition 2.3 ([18]). A complex (real) G -vector bundle over X is a complex (real) vector bundle $\pi: E \rightarrow X$ such that π is G -equivariant and each $g \in G$ acts on E and X via a bundle isomorphism. An isomorphism of G -vector bundles over X is just an isomorphism of vector bundles that is G -equivariant. The set of isomorphism classes of complex (real) G -vector bundles over X will be denoted by $\text{Bdl}_G(X)$ ($\text{OBdl}_G(X)$). For every $x \in X$, the fiber $\pi^{-1}(x)$ is denoted by E_x . We refer the reader to [12, Section 1] and [20, Section I.9] for elementary properties of G -vector bundles over proper (cocompact) G -CW complexes.

Theorem 2.4. [12, Th. 3.2 and 3.15] *There exists a 2-periodic (8-periodic) equivariant cohomology theory $\text{K}_G^*(X, A)$ ($\text{KO}_G^*(X, A)$) on the category of proper G -CW-pairs such that when X is cocompact, $\text{K}_G^0(X)$ ($\text{KO}_G^0(X)$) is the Grothendieck group of the abelian monoid of isomorphism classes of complex (real) G -vector bundles over X . In particular, for every $H \in \mathcal{F}$, $\text{K}_G^0(G/H)$ ($\text{KO}_G^0(G/H)$) is canonically isomorphic to $R(H)$ ($RO(H)$).*

As usual (see [13, Section 6] and [5, Th. 4.7]), the skeletal filtration of X induces Atiyah-Hirzebruch spectral sequences

$$(1) \quad E_2^{p,q} = \text{H}_G^p(X, \text{K}_G^q(G/-)) \implies \text{K}_G^{p+q}(X).$$

and

$$(2) \quad E_2^{p,q} = \text{H}_G^p(X, \text{KO}_G^q(G/-)) \implies \text{KO}_G^{p+q}(X)$$

where $\text{H}_G^p(X, -)$ denotes Bredon cohomology of X (see [2]).

Proposition 2.5. [13, Prop 5.8] *If X is a cocompact G -CW complex then the spectral sequences (1) and (2) above rationally collapse, meaning that the images of all differentials in these spectral sequences consist of torsion elements.*

By our assumptions on X , the zeroth Bredon cohomology group $\text{H}_G^0(X, R(-))$ (resp. $\text{H}_G^0(X, RO(H))$), equals the limit of the functor $R(-)$ (resp. $RO(-)$), over the orbit category $\mathcal{O}_{\mathcal{F}}G$. Consider the edge homomorphisms

$$\varepsilon_G: \text{K}_G^0(X) \rightarrow \text{H}_G^0(X, R(-))$$

and

$$\varepsilon_G: \text{KO}_G^0(X) \rightarrow \text{H}_G^0(X, RO(-))$$

of the spectral sequences (1) and (2). If $[\pi]$ is the isomorphism class of an n -dimensional complex G -vector bundle $\pi: E \rightarrow X$, then $\varepsilon_G([\pi])$ equals

$$([E_{e_H}])_{H \in \mathcal{F}} \in \lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} \text{Rep}_{U(n)}(H) \subset \mathbf{H}_G^0(X, R(-))$$

where $[E_{e_H}]$ denotes the isomorphism class in $R(H)$ of the H -representation E_{e_H} . The corresponding statement for real G -vector bundles also holds. Note that it follows from Proposition 2.5 that a suitable multiple of every compatible collection of (virtual) real or complex representations of the finite subgroups of G is contained in the image of the edge homomorphism ε_G .

Recall that the classifying space for proper actions $\underline{E}G$ is a terminal object in the homotopy category of proper G -CW complexes (e.g. [9, Th. 1.9]). Hence, if X is any proper cocompact G -CW complex such that X^H is non-empty and connected for each $H \in \mathcal{F}$, then there exists a G -map $X \rightarrow \underline{E}G$ that is unique up to G -homotopy and induces commutative diagrams

$$\begin{array}{ccc} \mathbf{K}_G^0(X) & \longrightarrow & \lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} R(H) & \text{and} & \mathbf{KO}_G^0(X) & \longrightarrow & \lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} RO(H) \\ \uparrow & \nearrow & & & \uparrow & \nearrow & \\ \mathbf{K}_G^0(\underline{E}G) & & & & \mathbf{KO}_G^0(\underline{E}G) & & \end{array}$$

Hence, if a compatible collection α of virtual representations can be realized as a virtual G -vector bundle over $\underline{E}G$, it can also be realized as a virtual G -vector bundle over X .

3. COMPLEX VECTOR BUNDLES

The purpose of this section is to construct a group G with a cocompact classifying space for proper actions $\underline{E}G$ admitting a compatible collection of complex representations of the finite subgroups of G that cannot be realized as G -equivariant virtual complex vector bundle over $\underline{E}G$, i.e. so that the edge homomorphism

$$\varepsilon_G: \mathbf{K}_G^0(\underline{E}G) \rightarrow \lim_{G/H \in \mathcal{O}_{\mathcal{F}}G} R(H).$$

is not surjective.

Let $F = C_4 \rtimes C_2$ be the dihedral group of order 8 where σ is generator for C_4 and ε is a generator of C_2 . Let $H = \langle \sigma^2 \rangle$ be the center of F , which has order two and denote the n -skeleton of the universal F/H -space $X = E(F/H)$ by X^n . We let F act on X and X^n via the projection onto F/H . Consider the complex 1-dimensional representation

$$\lambda: H = \langle \sigma^2 \rangle \rightarrow \mathbf{U}(1) = S^1 : \sigma^2 \mapsto -1.$$

Lemma 3.1. *The isomorphism class $[\lambda]$ is contained in $R(H)^{F/H}$. For $k \in \mathbb{Z}$, the multiple $k[\lambda]$ is contained in the image of the restriction map $\text{res}: R(F) \rightarrow R(H)$ if and only if k is even.*

Proof. Since H is the center of F it follows that the conjugation action of F/H on $R(H)$ is trivial, hence $[\lambda] \in R(H)^{F/H} = R(H)$. One easily verifies that the representation

$$\tau: F \rightarrow \mathbf{U}(2)$$

defined by

$$\tau(\sigma) = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \quad \text{and} \quad \tau(\varepsilon) = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$$

satisfies $\text{res}([\tau]) = 2[\lambda]$. Hence, $k[\lambda]$ is contained in the image of res for every even $k \in \mathbb{Z}$. Note that, as a free abelian group, $R(H)$ is generated by $[\lambda]$ and the isomorphism class of the 1-dimensional complex trivial representation $[\text{tr}]$ (e.g. see [19]). Now suppose k is odd and there exists an element $[\mu] - [\rho] \in R(F)$ such that $\text{res}([\mu] - [\rho]) = k[\lambda]$. There are integers l, m, n and such that $\text{res}([\mu]) = l[\text{tr}] + m[\lambda]$, $\text{res}([\rho]) = l[\text{tr}] + n[\lambda]$ and $m - n = k$. By changing the representative of $[\mu]$, we may also assume that

$$\mu: F \rightarrow \text{U}(l + m)$$

where $\mu(\sigma)$ is a diagonal matrix. Since $\mu(\sigma^2)$ has an m -dimensional eigenspace with eigenvalues -1 and an l -dimensional eigenspace with eigenvalue 1 , it follows that $\mu(\sigma)$ has an s -dimensional eigenspace with eigenvalue i and a t -dimensional eigenspace with eigenvalue $-i$ such that $s + t = m$. Moreover, $\mu(\sigma^3)$ has an s -dimensional eigenspace with eigenvalue $-i$ and a t -dimensional eigenspace with eigenvalue i . Since σ and σ^3 are conjugate in F , it follows that $s = t$ proving that m is even. A similar argument shows that n is also even. But this contradicts the fact that $k = m - n$ is odd. Hence, there does not exist an element $[\mu] - [\rho] \in R(F)$ such that $\text{res}([\mu] - [\rho]) = k[\lambda]$, if k is odd. \square

The following lemma uses the notation introduced above and will be cited in the next section.

Lemma 3.2. *Every F -equivariant complex line bundle over X^3 is isomorphic to the pullback of an F -equivariant complex line bundle over $E(F/H)$ along the inclusion $i: X^3 \rightarrow E(F/H)$.*

Proof. Let \mathcal{S} be the family of subgroups of F containing only H and the trivial subgroup. Note that isomorphism classes of F -equivariant complex line bundles correspond to isomorphism classes of $(F, S^1 = \text{U}(1))$ -bundles. Let $\pi: E \rightarrow X^3$ be an F -equivariant complex line bundle over and let $[\alpha_H: H \rightarrow \text{U}(1) = S^1]$ be the isomorphism class in $\text{Rep}_{S^1}(H)$ of the H -representation induced on the fibers of π . If we set $\alpha_{\{e\}}: \{e\} \rightarrow S^1$, then $A = ([\alpha_K])_{K \in \mathcal{S}} \in \lim_{K \in \mathcal{S}} \text{Rep}_{S^1}(K)$. It follows from Lemma 2.2 for $\Gamma = S^1$, that in order to show that π is the pullback of an F -equivariant complex line bundle over $E(F/H)$ along the inclusion $i: X^3 \rightarrow E(F/H)$, it suffices to show that every F -map from X^3 to $B_{\mathcal{S}}(F, A)$ can be extended to an F -map from $E(F/H)$ to $B_{\mathcal{S}}(F, A)$. Here $B_{\mathcal{S}}(F, A)$ is homotopy equivalent to $BS^1 = \mathbb{C}P^\infty$ for all $K \in \mathcal{S}$, again by Lemma 2.2. It follows from Bredon's equivariant obstruction theory (see [2, Section II.1], [15, Th. I.5.1]) that the potential obstructions for extending such a map lie in the relative Bredon cohomology groups $H_F^{n+1}(E(F/H), X^3; \pi_n(B_{\mathcal{S}}(F, A)^-))$ for $n \geq 3$. Since $\pi_n(\mathbb{C}P^\infty)$ is zero unless $n = 2$, the lemma is proven. \square

The idea for the following lemma is contained in [12, p 596].

Lemma 3.3. *There exists an $n \geq 1$ such that $[\lambda]$ is not contained in the image of the edge homomorphism*

$$K_F^0(X^n) \rightarrow R(H)^{F/H}.$$

Proof. By [7, Theorem 5.1] for $X = \{*\}$, $\mathcal{F} = \{e, H\}$ and $E\mathcal{F} = E(F/H)$, there are maps

$$\alpha_n: R(F)/I^n \rightarrow K_F^0(X^n)$$

that induce a map of inverse systems from $\{R(F)/I^n\}_{n \geq 0}$ to $\{K_F^0(X^n)\}_{n \geq 0}$ that induces an isomorphism of pro-rings. Here I is the kernel of the restriction map $R(F) \rightarrow R(H)$. This implies that for sufficiently large $n \geq 1$ there exists a map $\beta_1: K_F^0(X^n) \rightarrow R(F)/I$ making the following diagram commute

$$\begin{array}{ccc}
 R(F)/I^n & \xrightarrow{\alpha_n} & K_F^0(X^n) \\
 \downarrow & \nearrow \beta_1 & \downarrow \\
 R(F)/I & \xrightarrow{\alpha_1} & K_F^0(X^1)
 \end{array}
 \begin{array}{c}
 \\
 \searrow \varepsilon_F \\
 \\
 \nearrow \varepsilon_F \\
 \\
 \end{array}
 R(H)^{F/H}$$

This shows that the image of the restriction map

$$R(F) \rightarrow R(H)^{F/H}$$

coincides with the image of the edge homomorphism

$$K_F^0(X^n) \rightarrow R(H)^{F/H}.$$

Since $[\lambda]$ does not lie in the image of $R(F) \rightarrow R(H)^{F/H}$ by Lemma 3.1, the lemma follows. \square

Let $n \geq 3$. By [8, Th. A & Th. 8.3] there exists a compact n -dimensional locally CAT(0)-cubical complex T_{X^n} equipped with a free cellular F/H -action and an F/H -equivariant map $t_{X^n}: T_{X^n} \rightarrow X^n$ that induces an isomorphism

$$(3) \quad \mathcal{H}_F^*(X^n) \xrightarrow{\cong} \mathcal{H}_F^*(T_{X^n})$$

for any equivariant cohomology theory $\mathcal{H}_?^*(\cdot)$ (e.g. see [11, section 1]). (We remark that [8, Th. 8.3] is stated for equivariant *homology* theories, but the analogous statement holds for equivariant *cohomology* theories by essentially the same proof.) The action of F on T_{X^n} in the above is via the projection $F \rightarrow F/H$. Now let Y^n be the universal cover of T_{X^n} and let Γ_n be the group of self-homeomorphisms of Y^n that lift the action of F/H on T_{X^n} . Since F/H acts freely on T_{X^n} , Γ_n acts freely on Y^n . We conclude that Y^n is an n -dimensional CAT(0)-cubical complex on which Γ_n acts freely, cocompactly and cellularly. Since Y_n is contractible, this implies that Γ_n is torsion-free. By construction there is a surjection $\Gamma_n \rightarrow F/H$ whose kernel N_n is the torsion-free group of deck transformation of the covering $Y^n \rightarrow T_{X^n}$. Now define the group G_n to be the pullback of $\pi_n: \Gamma_n \rightarrow F/H$ along $F \rightarrow F/H$. Then G_n acts on Y^n via the quotient map $G_n \rightarrow G_n/H = \Gamma_n$ and fits into the short exact sequence

$$1 \rightarrow N_n \rightarrow G_n \xrightarrow{p_n} F \rightarrow 1.$$

Note that the only non-trivial finite subgroup of G_n is $H \cong C_2$ and that since N_n acts freely on Y^n , the G_n -equivariant quotient map $Y^n \rightarrow N_n \backslash Y^n = T_{X^n}$ induces an isomorphism ([12, Lemma 3.5])

$$(4) \quad K_F^*(T_{X^n}) \xrightarrow{\cong} K_{G_n}^*(Y^n).$$

Applying (3) and (4) to the composition $Y^n \rightarrow T_{X^n} \rightarrow X^n$ and the equivariant cohomology theories $K_?^*(\cdot)$ and $H_?^*(\cdot, R(-))$ with $* = 0$, we obtain a commutative diagram

$$\begin{array}{ccc} K_F^0(X^n) & \xrightarrow{\cong} & K_{G_n}^0(Y^n) \\ \downarrow \varepsilon_F & & \downarrow \varepsilon_{G_n} \\ R(H)^{F/H} & \xrightarrow{\cong} & \lim_{G_n/S \in \mathcal{O}_{\mathcal{F}} G_n} R(S). \end{array}$$

The fact that this diagram commutes can be seen as follows. Using equivariant cellular approximation, we may assume that the map $X^n \rightarrow Y^n$ is cellular. By considering the inclusion of zero-skeleta in n -skeleta, naturality yields a commutative diagram

$$\begin{array}{ccc} K_F^0(X^n) & \xrightarrow{\cong} & K_{G_n}^0(Y^n) \\ \downarrow & & \downarrow \\ K_F^0(X^0) & \longrightarrow & K_{G_n}^0(Y^0). \end{array}$$

The edge homomorphism $\varepsilon_F: K_F^0(X^n) \rightarrow R(H)^{F/H} \subseteq K_F^0(X^0)$ coincides by construction with $K_F^0(X^n) \rightarrow K_F^0(X^0)$ once we restrict the codomain, and similarly for ε_{G_n} . Therefore, commutativity follows.

Since we proved in Lemma 3.3 that, for n large enough, the isomorphism class of λ does not lie in the image of the edge homomorphism

$$K_F^0(X^n) \rightarrow R(H)^{F/H}$$

it follows from the commutative diagram above that the compatible system of representations

$$(\lambda \circ p_{n|S})_{S \in \mathcal{F}} \in \lim_{G_n/S \in \mathcal{O}_{\mathcal{F}} G_n} R(S) = H_{\mathcal{F}}^0(G_n, R(-)).$$

does not lie in the image of the edge homomorphism

$$\varepsilon_{G_n}: K_{G_n}^0(Y^n) \rightarrow \lim_{G_n/S \in \mathcal{O}_{\mathcal{F}} G_n} R(S).$$

Recall from [3] that non-empty CAT(0)-cube complexes are contractible and that the fixed point set for a finite group action on a CAT(0)-cube complex is contractible. Since G_n acts cellularly properly and cocompactly on the CAT(0)-cube complex Y_n , we deduce that Y_n is a cocompact model for $\underline{E}G_n$. To summarize, we have constructed a group $G = G_n$ with a cocompact classifying space for proper actions $\underline{E}G$ admitting a compatible collection of complex representations of the finite subgroups of G that cannot be realized as G -equivariant virtual complex vector bundle over $\underline{E}G$.

We remark that Wolfgang Lück has shown us another quite different way to find a finite group F and an F -CW-complex X that satisfy Lemma 3.3; any such pair could be used to construct a group with similar properties to the group $G = G_n$.

4. REAL VECTOR BUNDLES

One could apply the techniques of the previous section in the real setting to obtain a group G with cocompact classifying space for proper actions $\underline{E}G$ so that the edge homomorphism

$$\varepsilon_G: KO_G^0(\underline{E}G) \rightarrow \lim_{G/H \in \mathcal{O}_{\mathcal{F}} G} RO(H)$$

is not surjective. Here one would need the real version of [7, Theorem 5.1], which also holds as explained in the paragraph below [7, Theorem 5.1].

Instead we give an explicit description of a group G that admits \mathbb{R}^2 as a cocompact model for $\underline{E}G$ and admits a compatible collection of real representations of its finite subgroups that cannot be realized as a real G -vector bundle over \mathbb{R}^2 .

We start by describing a related group Γ that is a 2-dimensional crystallographic group, or wallpaper group; this group is known as $p2gg$, but we will describe it explicitly. Endow \mathbb{R}^2 with the CW-structure coming from the standard tessellation by unit squares with vertices at \mathbb{Z}^2 , and let Γ be the group of automorphisms of this CW-structure that preserves the pattern shown in Figure 1. The stabilizer of a 2-cell is clearly trivial, and so the 2-cells form a single free Γ -orbit. There are two orbits of 1-cells, the vertical and horizontal edges, and again each orbit is free. There are two orbits of 0-cells, and the stabilizer of a 0-cell is cyclic of order two, generated by the rotation of order two fixing the point. Since the stabilizer of each cell acts trivially on that cell, the given CW-structure makes \mathbb{R}^2 into a Γ -CW-complex.

The translation subgroup T of Γ has index four, and consists of the elements $(x, y) \mapsto (x + 2m, y + 2n)$. The orientation-preserving subgroup N of Γ has index two, and consists of T together with the rotations through π about some point of \mathbb{Z}^2 , which are of the form $(x, y) \mapsto (2m - x, 2n - y)$. Finally the elements of $\Gamma - N$ are the glide reflections whose axes bisect the sides of the 2-cells: $(x, y) \mapsto (2m + 1 - x, 2n + 1 + y)$ and $(x, y) \mapsto (2m + 1 + x, 2n + 1 - y)$. The quotients $T \backslash \mathbb{R}^2$, $N \backslash \mathbb{R}^2$ and $\Gamma \backslash \mathbb{R}^2$ are respectively a torus consisting of four squares, an S^2 obtained by identifying the boundaries of two squares, and a copy of $\mathbb{R}P^2$ obtained by identifying the edges of a square in pairs. The fact that $\Gamma - N$ contains no torsion elements is reflected in the fact that Γ/N acts freely on the sphere $N \backslash \mathbb{R}^2$.

Now let F be a copy of C_4 and let $H \cong C_2$ be the index two subgroup of F . The group G is defined as the pullback of the two maps $\Gamma \rightarrow \Gamma/N \cong C_2$ and $F \rightarrow F/H \cong C_2$. By construction the group G admits \mathbb{R}^2 as a cocompact model for $\underline{E}G$, and fits into a short exact sequence

$$1 \rightarrow N \rightarrow G \xrightarrow{p} F \rightarrow 1$$

such that every finite subgroup of G maps onto a subgroup of H under p .

Now let

$$\lambda : H \rightarrow O(1, \mathbb{R}) = C_2$$


be the 1-dimensional real sign representation of H , i.e. λ is the identity map. The isomorphism class $[\lambda]$ is clearly contained in $RO(H)^{F/H}$, since F is abelian.

Lemma 4.1. *The isomorphism class $k[\lambda]$ is contained in the image of the restriction map*

$$RO(F) \rightarrow RO(H)^{F/H}.$$

if and only if k is even.

Proof. Recall that the irreducible real representations of C_4 are up to isomorphism the one-dimensional trivial representation, the one dimensional sign representation of $F/H = C_2$ and one 2-dimensional faithful representation in which the elements of order four act as rotations by $\pm \frac{\pi}{2}$. The restriction of the first two of the representations to H gives the trivial one dimensional representation of H , while the restriction to H of the third is $\lambda \oplus \lambda$. We therefore conclude that the image of $RO(F) \rightarrow RO(H)^{F/H}$ consists of element of the form $2n[\lambda] + m[\text{tr}]$, where tr is the trivial one dimensional representation of H and $n, m \in \mathbb{Z}$.

FIGURE 1. A wallpaper pattern for $\Gamma = p2gg$

This shows that $k[\lambda]$ is contained in the image of the restriction map $RO(F) \rightarrow RO(H)^{F/H}$ if and only if k is even. \square

Lemma 4.2. *Let F act on the infinite dimensional sphere S^∞ by first projecting onto $F/H = C_2$ and then acting via the antipodal map. View S^2 as the 2-skeleton of S^∞ . Every F -equivariant orthogonal real line bundle over S^2 is isomorphic to the pullback of an F -equivariant orthogonal real line bundle over S^∞ along the inclusion $S^2 \rightarrow S^\infty$.*

Proof. Let \mathcal{S} be the family of subgroups of F containing H and the trivial subgroup. Note that isomorphism classes of F -equivariant orthogonal real line bundles correspond to isomorphism classes of (F, C_2) -bundles. Now let ξ be an (F, C_2) -bundle over S^2 with fibers $A = (\xi_S) \in \lim_{S \in \mathcal{S}} \text{Rep}_{C_2}(S)$. By Lemma 2.2, it suffices to show that every F -map $f: S^2 \rightarrow B_{\mathcal{S}}(F, A)$ can be extended to an F -map $\tilde{f}: S^\infty \rightarrow B_{\mathcal{S}}(F, A)$. Again by Lemma 2.2, $B_{\mathcal{S}}(F, A)^S \cong BC_2 = \mathbb{R}P^\infty$ for all $S \in \mathcal{S}$. It follows from Bredon's equivariant obstruction theory (see [2, Section II.1],[15, Th. I.5.1]) that the potential obstructions for extending such a map lie in the relative Bredon cohomology groups $H_F^{n+1}(S^\infty, S^2; \pi_n(B_{\mathcal{S}}(F, A)^-))$ for $n \geq 2$. Since $\pi_n(\mathbb{R}P^\infty)$ is zero unless $n = 1$, the lemma is proven. \square

Lemma 4.3. *Let F act on S^2 by first projecting onto $F/H = C_2$ and then acting via the antipodal map. There does not exist a real F -vector bundle $\xi: E \rightarrow S^2$ such that the representation of H on the fibers of ξ is isomorphic to λ .*

Proof. Consider the infinite dimensional sphere S^∞ as a the universal C_2 -space EC_2 , where C_2 acts via the antipodal map and let F act on S^∞ via first projection onto $F/H = C_2$ and then acting via C_2 . Now assume that there exists a real F -vector bundle $\xi: E \rightarrow S^2$ such that the representation of H on the fibers of ξ is isomorphic to λ . By Lemma 4.2 there exists a real F -vector bundle $\xi: E \rightarrow S^\infty$ such that the representation of H on the fibers of ξ is isomorphic to λ . By pulling back this bundle along the inclusion $S^2 \rightarrow S^\infty$, there also exists a real F -vector bundle $\xi: E \rightarrow S^2$ such that the representation of H on the fibers of ξ is isomorphic to λ , for every $n \geq 2$.

By the real version of [7, Theorem 5.1] (see comments below [7, Theorem 5.1]), there are maps

$$\alpha_n: RO(F)/I^n \rightarrow KO_F^0(S^n)$$

that induce a map of inverse systems from $\{RO(F)/I^n\}_{n \geq 0}$ to $\{KO_F^0(S^n)\}_{n \geq 0}$ that in turn induces an isomorphism of pro-rings. Here I is the kernel of the restriction map $RO(F) \rightarrow RO(H)$. This implies that for sufficiently large $n \geq 1$ there exists a map $\beta_1: KO_F^0(S^n) \rightarrow RO(F)/I$ making the following diagram commute

$$\begin{array}{ccc} RO(F)/I^n & \xrightarrow{\alpha_n} & KO_F^0(S^n) \\ \downarrow & \searrow \beta_1 & \downarrow \\ RO(F)/I & \xrightarrow{\alpha_1} & KO_F^0(S^1) \end{array} \quad \begin{array}{c} \xrightarrow{\varepsilon_F} \\ \xrightarrow{\varepsilon_F} \end{array} \quad \begin{array}{c} \\ RO(H)^{F/H} \end{array}$$

This shows that the image of the restriction map

$$RO(F) \rightarrow RO(H)^{F/H}$$

coincides with the image of the edge homomorphism

$$KO_F^0(S^n) \rightarrow RO(H)^{F/H},$$

implying that the H -representations coming from the fibers of any real F -vector bundle over S^n can be extended to virtual F -representations. However, since λ does not lie in the image of $RO(F) \rightarrow RO(H)$ by Lemma 4.1 we arrive at a contradiction and conclude that there does not exist a real F -vector bundle $\xi: E \rightarrow S^2$ such that the representation of H on the fibers of ξ is isomorphic to λ . \square

Consider the projection $p: G \rightarrow F$ and the compatible system of real orthogonal representations

$$([\lambda \circ p|_S])_{S \in \mathcal{F}} \in \varinjlim_{G/S \in \mathcal{O}_F G} RO(S) = H_G^0(\underline{E}G, RO(-)),$$

and assume that there exists a real G -vector bundle $\xi: E \rightarrow \mathbb{R}^2$ that realizes it. Since the kernel of $p: G \rightarrow F$ is N , it follows from the lemma below and our observations above that $N \setminus \xi: N \setminus E \rightarrow N \setminus X$ is an F -vector bundle over S^2 , where F acts on S^2 via projection onto $F/H = C_2$, followed by the antipodal map. Moreover, the representation of H on the fibers of $N \setminus \xi$ is by construction exactly λ . This however contradicts Lemma 4.3, so we conclude that there does not exist a real G -vector bundle $\xi: E \rightarrow \mathbb{R}^2$ that realizes the compatible system of real orthogonal representations $(\lambda \circ p|_S)_{S \in \mathcal{F}}$.

Lemma 4.4. *Let G be any discrete group with normal subgroup N and let X be a proper G -CW-complex. If $\xi: E \rightarrow X$ is a G -vector bundle over X such that $N \cap G_x$ acts trivially on $\xi^{-1}(x)$ for every $x \in X$, then*

$$N \setminus \xi: N \setminus E \rightarrow N \setminus X$$

is a G/N -vector bundle over $N \setminus X$.

Proof. Denote the projection $G \rightarrow G/N = Q$ by π . Let us first consider the case where ξ is trivial (trivial in the sense of [10, Section 6.1.]), i.e. assume ξ is a pullback

$$\begin{array}{ccc} G \times_H V & \longrightarrow & G/H \\ r \uparrow & & \uparrow p \\ E & \xrightarrow{\xi} & X \end{array}$$

of the G -vector bundle $G \times_H V \rightarrow G/H$ along the G -map $p: X \rightarrow G/H$ where H is some finite subgroup of G and V is a finite dimensional real H -representation such that $H \cap N$ acts trivially on V . Consider the pullback diagram

$$\begin{array}{ccc} Q \times_{\pi(H)} V & \longrightarrow & Q/\pi(H) \\ w \uparrow & & \uparrow N \setminus p \\ P & \xrightarrow{q} & N \setminus X \end{array}$$

of the Q -vector bundle $Q \times_{\pi(H)} V \rightarrow Q/\pi(H)$ along the Q -map $N \setminus p: N \setminus X \rightarrow Q/\pi(H)$. We define the map

$$\psi: N \setminus E \rightarrow P: \overline{(g, v, x)} \mapsto (\pi(g), v, \bar{x}).$$

It is easy to check that ψ yields a well-defined morphism of Q -equivariant bundles over $N \setminus X$. Moreover, since ψ is a fiberwise linear map of Q -vector bundles that is a fiberwise isomorphism, it follows that ψ is a homeomorphism.

Now consider the general case. Let $\bar{x} \in N \setminus X$. Since $\xi: E \rightarrow X$ is locally trivial, $x \in X$ has an open G -neighbourhood U such that there is a G -map $p: U \rightarrow G/H$ where H is finite subgroup of G and $\xi|_U$ is (homeomorphic to) the pullback

$$\begin{array}{ccc} G \times_H V & \longrightarrow & G/H \\ \uparrow & & \uparrow p \\ \xi|_U & \longrightarrow & U \end{array}$$

of the G -vector bundle $G \times_H V \rightarrow G/H$ along the G -map $p: U \rightarrow G/H$. By the above, the quotient diagram

$$\begin{array}{ccc} Q \times_{\pi(H)} V & \longrightarrow & Q/\pi(H) \\ \uparrow & & \uparrow N \setminus p \\ N \setminus \xi|_U & \xrightarrow{N \setminus \xi} & N \setminus U \end{array}$$

is a pullback diagram. Since $N \setminus U$ is an open Q -neighbourhood of \bar{x} , it follows that $N \setminus \xi: N \setminus E \rightarrow N \setminus X$ is a Q -vector bundle. \square

We finish this section by explaining how a similar approach to the one above can be used to produce a group G admitting a three dimensional cocompact model for $\underline{E}G$ that has a compatible system of one-dimensional complex representations that cannot be realized as a complex G -vector bundle over $\underline{E}G$. As in Section 3, let $F = C_4 \rtimes C_2$ be the dihedral group of order 8 where σ is a generator for C_4 . Let $H = \langle \sigma^2 \rangle$ be the center of F , which has order two

and denote the 3-skeleton of the universal F/H -space $X = E(F/H)$ by X^3 . We let F act on X and X^3 via the projection onto F/H . Consider the complex 1-dimensional representation

$$\lambda: H = \langle \sigma^2 \rangle \rightarrow \mathrm{U}(1) = S^1 : \sigma^2 \mapsto -1.$$

By [8, Th. A & Th. 8.3] there exists a compact 3-dimensional locally CAT(0)-cubical complex T_{X^3} equipped with a free cellular F/H -action, an F/H -equivariant map $t_{X^3}: T_{X^3} \rightarrow X^3$ and an isometric cellular involution τ on T_{X^3} that commutes with the F/H -action on T_{X^3} and the map t_{X^3} such the induced F/H -equivariant map

$$\langle \tau \rangle \backslash T_{X^3} \rightarrow X^3$$

is a homotopy equivalence. Note that F/H acts freely on $\langle \tau \rangle \backslash T_{X^3}$ since it acts freely on X^3 . Hence T_{X^3} is also the 3-skeleton of a universal F/H -space Z . So we may continue assuming that $Z = X$ and $\langle \tau \rangle \backslash T_{X^3} = X^3$.

Now let Y be the universal cover of T_{X^3} and let Γ be the group of self-homeomorphism of Y that lifts the action of $F/H \oplus \langle \tau \rangle$ on T_{X^3} . Then Y is a 3-dimensional CAT(0)-cubical complex on which Γ acts properly, compactly and cellularly. By construction there is a surjection $\alpha: \Gamma \rightarrow F/H \oplus \langle \tau \rangle$ whose kernel $\mathrm{Ker}(\alpha)$ is the torsion-free group of deck transformations of $Y \rightarrow T_{X^3}$. Let π denote the composition of α with the projection of $F/H \oplus \langle \tau \rangle$ onto F/H . Since F/H acts freely on T_{X^3} and every finite subgroup of Γ must fix a point of Y since Y is CAT(0), it follows that every finite subgroup of Γ is contained in the kernel of π , which we denote by N . Now define the group G to be the pullback of $\pi: \Gamma \rightarrow F/H$ along $F \rightarrow F/H$. Then G acts on Y via the quotient map $G \rightarrow G/H = \Gamma$ that fits into the short exact sequence

$$1 \rightarrow N \rightarrow G \xrightarrow{p} F \rightarrow 1.$$

such that p maps all the finite subgroup of G onto a finite subgroup of H and $N \backslash Y = X^3$.

Let \mathcal{F} be the family of finite subgroups of G , note that Y is a three dimensional cocompact model for $\underline{E}G$ and suppose that there exists a G -vector $\xi: E \rightarrow Y$ whose fibers give rise to the compatible system of representations

$$([\lambda \circ p|_S])_{S \in \mathcal{F}} \in \lim_{G/S \in \mathcal{O}_{\mathcal{F}}G} R(S).$$

Applying Lemma 4.4, we obtain an F -equivariant complex line bundle $N \backslash \xi: N \backslash E \rightarrow X$ such that the representation of H on the fibers of $N \backslash \xi$ is isomorphic to λ . By Lemma 3.2, this bundle can be extended to an F -equivariant complex line bundle over $X = E(F/H)$. We now continue in a similar fashion as in the proof of Lemma 4.3 to conclude that $[\lambda]$ is contained in the image of the restriction map $R(F) \rightarrow R(H)^{F/H}$, which contradicts Lemma 3.1. We conclude that the bundle ξ cannot exist.

5. RIGHT ANGLED COXETER GROUPS

Let Γ be a finite graph. We denote the vertex set of Γ by $S = V(\Gamma)$ and the set edges of Γ by $E(\Gamma) \subseteq V(\Gamma) \times V(\Gamma)$. The right angled Coxeter group determined by Γ is the Coxeter group W with presentation

$$W = \langle S \mid s^2 \text{ for all } s \in V(\Gamma) \text{ and } (st)^2 \text{ if } (s, t) \in E(\Gamma) \rangle.$$

Note that W fits into the short exact sequence

$$1 \rightarrow N \rightarrow W \xrightarrow{p} F = \bigoplus_{s \in S} C_2 \rightarrow 1$$

where p takes $s \in S$ to the generator of the C_2 -factor corresponding to s . A subset $J \subseteq S$ is called spherical if the subgroup $W_J = \langle J \rangle$ is finite (and hence isomorphic to $\bigoplus_{s \in J} C_2$). The empty subset of S is by definition spherical. We denote the poset of spherical subsets of S ordered by inclusion by \mathcal{S} . If $J \in \mathcal{S}$, then W_J is called a spherical subgroup of W , while a coset wW_J is called spherical coset. We denote the poset of spherical cosets, ordered by inclusion, by $W\mathcal{S}$. Note that W acts on $W\mathcal{S}$ by left multiplication, preserving the ordering. The Davis complex Σ of W is the geometric realization of $W\mathcal{S}$. One easily sees that Σ is a proper cocompact W -CW-complex. Since Σ admits a complete CAT(0)-metric such that W acts by isometries, it follows that Σ is a cocompact model for \underline{EW} (see [4, Th. 12.1.1 & Th. 12.3.4]). A consequence of this fact is that every finite subgroup of W is subconjugate to some spherical subgroup of W . This implies that the group N defined above is torsion-free. Since the quotient space $W \backslash \Sigma$ is homeomorphic to the geometric realization of the poset \mathcal{S} , which is contractible since it has a minimal element, another consequence is that the quotient $\underline{BW} = W \backslash \underline{EW}$ is contractible. We refer the reader to [4] for more details and information about these groups and the spaces on which they act.

Let \mathcal{F} be the family of finite subgroups of W . Given an abelian group A , we denote by

$$\underline{A}: \mathcal{O}_{\mathcal{F}}W \rightarrow \text{Ab}$$

the trivial functor that takes all objects to A and all morphism to the identity map. One can verify that

$$(5) \quad H_W^*(\underline{EW}, \underline{A}) \cong H^*(\underline{BW}, A).$$

Lemma 5.1. *Let $A = ([p|_H])_{H \in \mathcal{F}} \in \lim_{W/H \in \mathcal{O}_{\mathcal{F}}W} \text{Rep}_F(H)$. For every $k \geq 0$, the contravariant functor*

$$\mathcal{O}_{\mathcal{F}}W \rightarrow \text{Ab}: W/H \mapsto \pi_k(B_{\mathcal{F}}(W, A)^H)$$

equals the trivial functor $\pi_k(\underline{BF})$.

Proof. Let EF be a contractible F -CW-complex with free F -action and consider the product space $\underline{EW} \times EF$. This space becomes a $(W \times K)$ -CW-complex by letting $(w, f) \in W \times F$ act on $(x, y) \in \underline{EW} \times EF$ as

$$(w, f) \cdot (x, y) = (w \cdot x, p(w)f \cdot y).$$

One checks that with this action $\underline{EW} \times EF$ is a model for $E_{\mathcal{F}}(W, A)$, i.e. $(\underline{EW} \times EF)^K$ is contractible when $K \in \mathcal{F}_A$ and empty otherwise. By definition, it follows that $\underline{EW} \times BF$ is a model $B_{\mathcal{F}}(W, A)$, where W acts on trivially on the second coordinate. Since \underline{EW}^H is contractible for every $H \in \mathcal{F}$, the lemma follows easily. \square

Let Γ be either the orthogonal group $O(n, \mathbb{R})$ or the unitary group $U(n)$.

Lemma 5.2. *Every element of*

$$\lim_{W/H \in \mathcal{O}_{\mathcal{F}}W} \text{Rep}_{\Gamma}(H)$$

is of the form $([\lambda \circ p|_H])_{H \in \mathcal{F}}$ for some group homomorphism $\lambda: F \rightarrow \Gamma$.

Proof. Every finite subgroup H of W is isomorphic to a finite direct sum of C_2 's. Since every element of order 2 in Γ is conjugate in Γ to a diagonal matrix with ± 1 on the diagonal and commuting matrices can be simultaneously diagonalized (e.g. see [6, Th. 1.3.12]), it follows that the image of every homomorphism $H \rightarrow \Gamma$ is conjugate to a finite subgroup of Γ consisting

of diagonal matrices. Hence, every element of $\lim_{W/H \in \mathcal{O}_{\mathcal{F}}W} \text{Rep}_{\Gamma}(H)$ is of the form $([\alpha_H])_{H \in \mathcal{F}}$ where $\alpha_H: H \rightarrow \Gamma$ is a homomorphism whose image lands in the finite abelian subgroup of Γ consisting of diagonal matrices. Since every finite subgroup of W is subconjugate to a spherical subgroup W_J , the compatibility of the representations tells us that $([\alpha_H])_{H \in \mathcal{F}}$ is completely determined by the homomorphisms $\alpha_{\langle s \rangle}: \langle s \rangle \rightarrow \Gamma$, for $s \in S$. Since the images of the $\alpha_{\langle s \rangle}$ are diagonal, they commute. Therefore, one can define the homomorphism

$$\lambda: F = \bigoplus_{s \in S} C_2 \rightarrow \Gamma: (\sigma_s)_{s \in S} \mapsto \sum_{s \in S} \alpha_{\langle s \rangle}(\sigma_s).$$

The compatibility of the representations implies that

$$([\lambda \circ p|_H])_{H \in \mathcal{F}} = ([\alpha_H])_{H \in \mathcal{F}},$$

proving the lemma. \square

The following theorem applies to both complex and real representations and vector bundles.

Theorem 5.3. *Let W be a right angled Coxeter group. Every compatible collection of representations of the finite subgroups of W can be realized as a W -equivariant vector bundle over the Davis complex $\Sigma = \underline{E}W$.*

Proof. Consider $A = ([p|_H])_{H \in \mathcal{F}} \in \lim_{W/H \in \mathcal{O}_{\mathcal{F}}W} \text{Rep}_F(H)$. It follows from Lemma 2.2 that the existence of a (W, A) -bundle over Σ follows from the existence a W -map $\Sigma \rightarrow B_{\mathcal{F}}(W, A)$. Since by Lemma 5.1, the contravariant functor

$$\pi_k(B_{\mathcal{F}}(W, A)^{-}): \mathcal{O}_{\mathcal{F}}(W) \rightarrow \text{Ab}: W/H \mapsto \pi_k(B_{\mathcal{F}}(W, A)^H)$$

equals the trivial functor $\underline{\pi}_k(B_{\mathcal{F}})$ for all $k \geq 0$, it follows from (5) and the contractibility of $\underline{B}W$ that the Bredon cohomology groups

$$\mathbb{H}_W^{k+1}(\Sigma, \pi_k(B_{\mathcal{F}}(W, A)^{-}))$$

are zero for all $k \geq 0$. Since there certainly exists a W -map from the 0-skeleton of Σ to $B_{\mathcal{F}}(W, A)$, it follows from Bredon's equivariant obstruction theory that there exists a W -map $\Sigma \rightarrow B_{\mathcal{F}}(W, A)$.

Now consider a compatible collection of representations of the finite subgroups of W . By Lemma 5.2, this collection is of the form

$$([\lambda \circ p|_H])_{H \in \mathcal{F}} \in \lim_{W/H \in \mathcal{O}_{\mathcal{F}}W} \text{Rep}_{\Gamma}(H)$$

for some group homomorphism $\lambda: F \rightarrow \Gamma$. Letting $A = ([p|_H])_{H \in \mathcal{F}}$, it follows from the above that there exists a (W, A) -bundle $\xi: E \rightarrow \Sigma$. If $\Gamma = \text{O}(n, \mathbb{R})$ then

$$\xi: E \times_F \mathbb{R}^n \rightarrow \Sigma$$

is a real W -vector bundle over Σ that realizes $([\lambda \circ p|_H])_{H \in \mathcal{F}}$, and if $\Gamma = \text{U}(n)$ then

$$\xi: E \times_F \mathbb{C}^n \rightarrow \Sigma$$

is a complex W -vector bundle over Σ that realizes $([\lambda \circ p|_H])_{H \in \mathcal{F}}$. Here F acts on \mathbb{R}^n or \mathbb{C}^n via the map λ . \square

Lemma 5.4. *If W is a right angled Coxeter group, then $\mathbb{H}_W^n(\Sigma, R(-)) = 0$ for all $n > 0$, and $\mathbb{H}_W^0(\Sigma, R(-))$ is free abelian of rank equal to the number of spherical subgroups of W .*

Proof. This is proven in much the same way as the corresponding result for homology in [17]. In more detail, one uses the cubical structure on Σ , in which there is one orbit of n -cubes with stabilizer isomorphic to $(C_2)^n$ for each n -tuple of commuting elements of S . (For each $n \geq 0$, for each spherical subgroup $W_J \cong (C_2)^n$ and for each $w \in W$, the subposet consisting of all special cosets contained in wW_J is order isomorphic to the poset of faces of an n -cube. Furthermore this isomorphism is equivariant for the stabilizer subgroup $wW_Jw^{-1} \cong (C_2)^n$, acting on the n -cube as the group generated by reflections in its coordinate planes. The realizations of these subposets are the cubes that make up the cubical structure on Σ . For more details concerning the cubical structure on Σ see [4, Ch. 1.1-1.2 or Ch. 7].) Since the stabilizer of a cube of strictly positive dimension acts non-trivially on the cube, this cubical structure is not a W -CW-structure on Σ . However, its barycentric subdivision is a simplicial complex naturally isomorphic to the realization of the poset WS as described in the introduction to this section.

Let Σ^n denote the n -skeleton of Σ with the cubical structure. Firstly, Σ^0 consists of a single free W -orbit of vertices, so $H_W^*(\Sigma^0; R(-))$ is isomorphic to the ordinary cohomology of a point; since W acts freely the calculation reduces to an equivariant cohomology calculation for the trivial group action.

Let $I = [-1, 1]$ be an interval, with C_2 acting by $x \mapsto -x$ (i.e., swapping the ends of the interval). Note that I is equivariantly isomorphic to the Davis complex for the Coxeter group C_2 . Let ∂I denote the two end points $\{-1, 1\}$. Make I into a C_2 -CW-complex, for example by taking three 0-cells in two orbits at the points $-1, 0$ and 1 , and one free orbit of 1-cells consisting of the two intervals $[-1, 0]$ and $[0, 1]$. The cellular C_2 -Bredon cochain complex for the pair $(I, \partial I)$ with coefficients in $R(-)$ is a cochain complex of free abelian groups in which the degree zero term has rank two, the degree one term has rank one, and all other terms are trivial. A direct computation with this cochain complex shows that $H_{C_2}^m(I, \partial I; R(-))$ is isomorphic to \mathbb{Z} for $m = 0$ and is zero for $m > 0$.

Next consider I^n with C_2^n acting as the direct product of n copies of the above action of C_2 on I . This is the Davis complex for the Coxeter group C_2^n . Since the representation ring of a direct product of finite groups is naturally identified with the tensor product of the representation rings [19, Ch. 3.2], the C_2^n -Bredon cochain complex for the pair $(I^n, \partial I^n)$ with coefficients in $R(-)$ is naturally isomorphic to the tensor product of n copies of the C_2 -Bredon cochain complex for $(I, \partial I)$ with coefficients in $R(-)$. (If one wants to think about this cochain complex geometrically, it arises from the $(C_2)^n$ -CW-structure on I^n in which the cells are the direct products of the cells arising in the C_2 -CW-structure on I .) Since these cochain complexes consist of finitely generated free abelian groups, there is a Künneth formula as described in for example [16, Thrm 60.3]. Since $H_{C_2}^*(I, \partial I; R(-))$ is free abelian the Künneth formula implies that

$$H_{C_2^n}^*(I^n, \partial I^n, R(-)) \cong \bigotimes_{i=1}^n H_{C_2}^*(I, \partial I; R(-)).$$

It follows that for each n , $H_{C_2^n}^m(I^n, \partial I^n; R(-))$ is isomorphic to \mathbb{Z} for $m = 0$ and is zero for $m > 0$.

From these computations, it follows easily that $H_W^m(\Sigma^n, \Sigma^{n-1}; R(-))$ is zero for $m > 0$ and is isomorphic to a direct sum of copies of \mathbb{Z} indexed by the W -orbits of n -cubes in Σ . By induction on n one sees that $H_W^m(\Sigma^n; R(-))$ is zero for $m > 0$ and isomorphic to a direct sum of copies of \mathbb{Z} indexed by the W -orbits of cubes of dimension at most n for $m = 0$. The

claimed result follows, since the W -orbits of cubes in Σ are in bijective correspondence with the spherical subgroups of W . \square

Before stating our theorem concerning $K_W^*(\underline{E}W)$, we make some remarks concerning the representation ring of a direct sum of copies of the cyclic group C_2 , indexed by a (finite) set S . For any finite group G , the collection of all isomorphism types of 1-dimensional complex representations of G is an abelian group, with product given by taking the tensor product of representations. Furthermore, this group is naturally isomorphic to the group $\text{Hom}(G, \text{U}(1))$. In the case when G is abelian, every irreducible representation of G is 1-dimensional, and so $\text{Hom}(G, \text{U}(1))$ forms a basis for the additive group of the representation ring. In this way the representation ring $R(G)$ is naturally isomorphic to the integral group algebra of the group $\text{Hom}(G, \text{U}(1))$. In the case when $G = \bigoplus_{s \in S} C_2$ is a direct sum of copies of C_2 indexed by S , we may view G as a vector space over the field of two elements, in which case $\text{Hom}(G, \text{U}(1))$ may be identified with the dual space. For $s \in S$, let s^* denote the 1-dimensional representation of G with the properties that $s^*(s) = -1$ and $s^*(t) = 1$ for $t \in S - \{s\}$. Let S^* denote the set of these representations: $S^* := \{s^* \mid s \in S\}$. In terms of vector spaces over the field of two elements, $S^* \subseteq \text{Hom}(G, \text{U}(1))$ is the dual basis to the set $S \subseteq G$. The set S^* generates the representation ring of G , giving rise to the following presentation:

$$R(G) = \mathbb{Z}[S^*]/(s^{*2} - 1 \mid s \in S),$$

in which the monomials $s_1^* s_2^* \cdots s_k^*$ for all subsets $\{s_1, \dots, s_k\} \subseteq S$ correspond to the irreducible representations.

Suppose now that J is a subset of S . The inclusion $J \subseteq S$ identifies $H = \bigoplus_{s \in J} C_2$ with a subgroup of $G = \bigoplus_{s \in S} C_2$. The induced map $R(G) \rightarrow R(H)$ of representation rings is described easily in terms of the above ring presentation: for $s \in J$, $s^* \in R(G)$ restricts to $s^* \in R(H)$, while for $s \notin J$, $s^* \in R(G)$ restricts to $1 \in R(H)$.

Now suppose that Γ is a graph with vertex set $V(\Gamma) = S$, and let W be the right angled Coxeter group associated to Γ . The abelianization of W is naturally identified with $G = \bigoplus_{s \in S} C_2$. There is a unique equivariant map $\alpha: \underline{E}W \rightarrow *$, from the W -space $\underline{E}W$ to a point $*$, viewed as a G -space with trivial action. If J is a spherical subset of S then $W_J = \bigoplus_{s \in J} C_2$ maps isomorphically to the corresponding subgroup of $G = \bigoplus_{s \in S} C_2$. If $x \in \underline{E}W$ is a 0-cell fixed by $W_J = \bigoplus_{s \in J} C_2$, then $\alpha(x) = *$, and this map is W_J -equivariant. The induced map $\alpha^*: K_G^*(*) \rightarrow K_W^*(\underline{E}W)$, and the composite map $K_G^*(*) \rightarrow K_{W_J}^*(\{x\})$ will be used in the statement and proof of our theorem. If we identify $R(G)$ with $K_G^0(*)$ and $R(W_J)$ with $K_{W_J}^0(\{x\})$, then the composite is identified with the restriction map.

Theorem 5.5. *Let W be the right angled Coxeter group determined by a finite graph Γ , with vertex set S , and let $G = \bigoplus_{s \in S} C_2$ be the abelianization of W . The map $\alpha^*: K_G^*(*) \rightarrow K_W^*(\underline{E}W)$ is surjective in each degree. In particular, $K_W^1(\underline{E}W) = 0$ and there is a ring isomorphism*

$$K_W^0(\underline{E}W) \cong \mathbb{Z}[S^*]/(s^{*2} - 1, s^* t^* - s^* - t^* + 1 \mid s \in S = V(\Gamma), (s, t) \notin E(\Gamma)).$$

It follows that $K_W^0(\underline{E}W) \cong \mathbb{Z}^d$ as an abelian group, where d is the number of spherical subgroups of W .

Proof. Consider the Atiyah-Hirzebruch spectral sequence (1)

$$E_2^{p,q} = H_W^p(\underline{E}W, K_W^q(W/-)) \implies K_W^{p+q}(\underline{E}W)$$

where $K_W^q(W/-) = R(-)$ if q is even and $K_W^q(W/-) = 0$ if q is odd (see [12, Th. 3.2]). In the lemma above, we proved that $H_W^k(\Sigma, R(-)) = 0$ for $k > 0$. It therefore follows that

$$K_W^n(\underline{EW}) = \begin{cases} H_W^0(\underline{EW}, R(-)) = \lim_{W/H \in \mathcal{O}_{\mathcal{F}W}} R(H) & \text{if } n = 0 \\ 0 & \text{if } n = 1. \end{cases}$$

Let I be the ideal

$$(s^{*2} - 1, s^*t^* - s^* - t^* + 1 \mid s \in S, (s, t) \notin E(\Gamma))$$

in the polynomial ring $\mathbb{Z}[S^*]$. Note that as an abelian group $\mathbb{Z}[S^*]/I$ is free, with basis elements the commuting products $s_1^* \dots s_k^*$, for all $J = \{s_1, \dots, s_k\} \in \mathcal{S}$ (The case $J = \emptyset$ corresponds to the unit of $\mathbb{Z}[V(\Gamma)]/I$). This shows that

$$\mathbb{Z}[S^*]/I \cong \mathbb{Z}^d$$

as an abelian group, where d is the number of spherical subgroups of W .

We claim there is an isomorphism of rings

$$\lim_{W/H \in \mathcal{O}_{\mathcal{F}W}} R(H) \cong \mathbb{Z}[S^*]/I.$$

Since every finite subgroup of W is subconjugate to a spherical subgroup of W , it follows that

$$\lim_{W/H \in \mathcal{O}_{\mathcal{F}W}} R(H) \cong \lim_{J \in \mathcal{S}} R(W_J)$$

as rings. By the remarks in the paragraph preceding the statement of the theorem, there are ring isomorphisms

$$R(W_J) = \mathbb{Z}[J^*]/(s^{*2} - 1 \mid s \in J), \quad R(G) = \mathbb{Z}[S^*]/(s^{*2} - 1 \mid s \in S),$$

which are natural for inclusions $J \subseteq J' \subseteq S$. From this it follows that the natural ring homomorphism

$$\rho: R(G) \rightarrow \lim_{W/H \in \mathcal{O}_{\mathcal{F}W}} R(H)$$

is surjective, and that $\lim_{W/H \in \mathcal{O}_{\mathcal{F}W}} R(H)$ is isomorphic to the ring described in the statement; in particular its additive group is free abelian of the same rank as $K_W^0(\underline{EW})$. Since ρ factors through $K_W^0(\underline{EW})$, the claimed isomorphism follows. \square

Before stating our corollary concerning $K^*(BW)$, we recall some facts from [1] concerning $K^*(BG)$, where as above $G = \bigoplus_{s \in S} C_2$. For any finite group H , Atiyah showed that $K^i(BH) = 0$ for i odd, and that $K^{2i}(BH)$ is naturally isomorphic to the completion of the representation ring $R(H)$ at its augmentation ideal. To discuss the case of G , it is convenient to take new generators for $R(G)$; replace the irreducible representation s^* by the degree zero virtual representation $\bar{s} = s^* - 1$. With respect to these generators one obtains the presentation

$$R(G) = \mathbb{Z}[\bar{S}]/(\bar{s}(\bar{s} + 2) \mid s \in S),$$

where $\bar{S} = \{\bar{s} \mid s \in S\}$. If $H = \bigoplus_{s \in J} C_2$, then of course there is a similar description of $R(H)$, which is natural for the inclusion $J \subseteq S$. Note that if $s \notin J$, then the image of \bar{s} under the restriction map $R(G) \rightarrow R(H)$ is zero.

Completing $R(G)$, as described above, with respect to its augmentation ideal gives rise to the following presentation for the ring $K^0(BG)$:

$$K^0(BG) = \mathbb{Z}[[\bar{S}]]/(\bar{s}(\bar{s} + 2) \mid s \in S),$$

which is natural for the inclusion $J \subseteq S$, and so also describes the induced map $K^0(BG) \rightarrow K^0(BH)$. The additive group of this ring is the direct sum of one copy of \mathbb{Z} , generated by 1, and for each non-empty subset $J \subseteq S$, one copy of the 2-adic integers, \mathbb{Z}_2 , consisting of the set of power series in the element $\prod_{s \in J} \bar{s}$ with zero constant term.

Corollary 5.6. *Let W be the right angled Coxeter group determined by a finite graph Γ with vertex set $S = V(\Gamma)$, and let $G = \bigoplus_{s \in S} C_2$ be the abelianization of W . The induced map $K^*(BG) \rightarrow K^*(BW)$ is surjective in each degree. In particular $K^1(BW) = 0$ and there is a ring isomorphism*

$$K^0(BW) \cong \mathbb{Z}[[\bar{S}]]/(\bar{s}(\bar{s} + 2), \bar{s}\bar{t} \mid s \in S, (s, t) \notin E(\Gamma)).$$

Here, $\mathbb{Z}[[\bar{S}]]$ is the formal power series ring with \mathbb{Z} coefficients in the variables $\bar{S} = \{\bar{s} \mid s \in S\}$.

Proof. The version of the Atiyah-Segal completion theorem that is proven for infinite discrete groups admitting a cocompact model for the classifying space for proper actions in [12, Theorem 4.4.(b)] implies that

$$K^n(BW) = K_W^n(\underline{EW})_J,$$

where the ideal J is the kernel of the augmentation map $K_W^n(\underline{EW}) \rightarrow \mathbb{Z}$ that maps vector bundles to their dimension. Changing variables in the above theorem to $\bar{s} = s^* - 1$, we see that $K^i(BW) = 0$ for i odd and that $K^0(BW)$ is the completion of the ring

$$\mathbb{Z}[\bar{S}]/(\bar{s}(\bar{s} + 2), \bar{s}\bar{t} \mid s \in S, (s, t) \notin E(\Gamma))$$

with respect to the ideal generated by the set $\bar{S} = \{\bar{s} \mid s \in S\}$. This completion is the ring described in the statement. \square

There is an alternative proof of Corollary 5.6 that does not use Theorem 5.5 or results from [12]. Instead one uses a description of W as a free product with amalgamation. If the graph Γ is not a complete graph, then there is an expression $\Gamma = \Gamma_1 \cup \Gamma_2$, $\Gamma_3 = \Gamma_1 \cap \Gamma_2$, in which each Γ_i is a full subgraph of Γ and has fewer vertices than Γ . This gives an expression for W as a free product with amalgamation $W = W_1 *_{W_3} W_2$. From this one obtains a Mayer-Vietoris sequence that can be used to compute $K^*(BW)$. To establish Corollary 5.6, one shows by induction on $|S|$ that $K^*(BW)$ is as described and that for each $J \subseteq S$, the map $K^*(BW) \rightarrow K^*(BW_J)$ is a split surjection.

REFERENCES

- [1] Atiyah, M. F., *Characters and cohomology of finite groups*, Inst. Hautes Études Sci. Publ. Math. 9 (1961) 23–64.
- [2] Bredon, G. E., *Equivariant cohomology theories*, Lecture Notes in Mathematics 34, Springer (1967).
- [3] Bridson, M. R. and Haefliger, A., *Metric spaces of non-positive curvature* Springer Verlag Vol. 319 (1999).
- [4] Davis, M. W., *The geometry and topology of Coxeter groups*, Princeton University Press, vol. 32 (2008).
- [5] Davis, J. F. and Lück, W., *Spaces over a category and assembly maps in isomorphism conjectures in K- and L- theory*, K-theory 15 (1998), 201–252.
- [6] Horn, R. and Johnson, C. *Matrix analysis*, Cambridge university press (2012).
- [7] Jackowski, S., *Families of subgroups and completion*, Journal of Pure and Applied Algebra 37 (1985), 167–179.
- [8] Leary, I. J., *A metric Kan-Thurston theorem*, Journal of Topology 6(1) (2013), 251–284.
- [9] Lück, W., *Survey on classifying spaces for families of subgroups*, Infinite Groups: Geometric, Combinatorial and Dynamical Aspects, Springer (2005), 269–322.
- [10] Lück, W., *The Burnside ring and equivariant stable cohomotopy for infinite groups*, Journal of Pure and Applied Mathematics Quarterly 1(3) (2005), 479–541.

- [11] Lück, W, *Equivariant cohomological Chern characters*, International Journal of Algebra and Computation 15 (2005), 1025–1052.
- [12] Lück, W, and Oliver, B., *The completion theorem in K-theory for proper actions of a discrete group*, Topology 40 (2001), 585–616.
- [13] Lück, W, and Oliver, B., *Chern characters for the equivariant K-theory of proper G-CW-complexes*, Cohomological methods in homotopy theory (2001), 217–247.
- [14] Lück, W, and Uribe, B., *Equivariant principal bundles and their classifying spaces*, Algebraic and Geometric Topology 14 (2014), 1925-1995.
- [15] May, J.P., *Equivariant homotopy and cohomology theory*, CBMS regional conferences series in mathematics 91.
- [16] Munkres, J. R., *Elements of Algebraic Topology*, Addison-Wesley (1984).
- [17] Sánchez-García, R., *Equivariant K-homology for some Coxeter groups*, Journal of the London Mathematical Society, 75(3), (2007), 773–790.
- [18] Segal, G., *Equivariant K-theory*, Publ. IHES 34 (1968) 129–151.
- [19] Serre, J.-P., *Linear representations of finite groups*, Graduate Text in Mathematics, Springer, Vol 42 (2012).
- [20] Dieck, T., *Transformation groups*, Studies in Math. 8, de Gruyter (1987).