

This Provisional PDF corresponds to the article as it appeared upon acceptance. Fully formatted PDF and full text (HTML) versions will be made available soon.

The geography of maternal and newborn health: the state of the art

International Journal of Health Geographics (2015) 14:19

doi:10.1186/s12942-015-0012-x

Steeve Ebener (steeve.ebener@gaiatech.org)
Maria Guerra-Arias (maria.guerra.arias@integrate.es)
James Campbell (campbellj@who.int)
Andrew J. Tatem (andy.tatem@gmail.com)
Allisyn Moran (amoran@usaid.gov)
Fiifi Amoako Johnson (faj100@soton.ac.uk)
Helga Fogstad (helga.fogstad@norad.no)
Karin Stenberg (stenbergk@who.int)
Sarah Neal (s.neal@soton.ac.uk)
Patricia Bailey (pbailey@fhi360.org)
Reid Porter (reidporter@gmail.com)
Zoe Matthews (z.matthews@soton.ac.uk)

Published online: 27 May 2015

ISSN 1476-072X

Article type Review

Submission date 17 March 2015

Acceptance date 13 May 2015

Article URL <http://dx.doi.org/10.1186/s12942-015-0012-x>

For information about publishing your research in BioMed Central journals, go to
<http://www.biomedcentral.com/info/authors/>

© 2015 Ebener *et al.*

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

The geography of maternal and newborn health: the state of the art

Steeve Ebener^{1*}

* Corresponding author

Email: steeve.ebener@gaia-geosystems.org

Maria Guerra-Arias²

Email: maria.guerra.arias@integrare.es

James Campbell³

Email: campbellj@who.int

Andrew J. Tatem⁴

Email: andy.tatem@gmail.com

Allisyn Moran⁵

Email: amoran@usaid.gov

Fiifi Amoako Johnson⁶

Email: faj100@soton.ac.uk

Helga Fogstad⁷

Email: helga.fogstad@norad.no

Karin Stenberg⁸

Email: stenbergk@who.int

Sarah Neal⁶

Email: s.neal@soton.ac.uk

Patricia Bailey⁸

Email: pbailey@fhi360.org

Reid Porter⁹

Email: reidmporter@gmail.com

Zoe Matthews⁶

Email: z.matthews@soton.ac.uk

¹ Gaia GeoSystems, Muscat, Oman

² ICS Integrare, Barcelona, Spain

³ WHO, Geneva, Switzerland

⁴ WorldPop, University of Southampton, Southampton, UK

⁵ USAID, Washington, DC, USA

⁶ University of Southampton, Southampton, UK

⁷ NORAD, Oslo, Norway

⁸ AMDD and FHI 360, Durham, USA

⁹ The University of Texas at Austin, Austin, USA

Abstract

As the deadline for the millennium development goals approaches, it has become clear that the goals linked to maternal and newborn health are the least likely to be achieved by 2015. It is therefore critical to ensure that all possible data, tools and methods are fully exploited to help address this gap. Among the methods that are under-used, mapping has always represented a powerful way to 'tell the story' of a health problem in an easily understood way. In addition to this, the advanced analytical methods and models now being embedded into Geographic Information Systems allow a more in-depth analysis of the causes behind adverse maternal and newborn health (MNH) outcomes. This paper examines the current state of the art in mapping the geography of MNH as a starting point to unleashing the potential of these under-used approaches. Using a rapid literature review and the description of the work currently in progress, this paper allows the identification of methods in use and describes a framework for methodological approaches to inform improved decision-making. The paper is aimed at health metrics and geography of health specialists, the MNH community, as well as policy-makers in developing countries and international donor agencies.

Keywords

Maternal health, Newborn health, Geography, GIS, Inequalities, Millennium development goals

Introduction

Maternal and newborn health (MNH) care refers to 'those activities whose primary purpose is to restore, improve and maintain the health of women and their newborn during pregnancy, childbirth and the 7-day postnatal period' ¹[1]. World Health Organization (WHO) guidance on the activities that should be included in MNH care includes the essential packages of interventions that address pregnancy, childbirth, postpartum care to mothers and care to newborns [2]. MNH outcomes are subject to wide geographic inequities, both at the regional, national and sub-national levels. The location of services is a key determinant of women's access to MNH care [3–6], thus the application of geographic approaches and geographic information systems (GIS) to MNH is critical to assess the situation and needs and guide policy decisions, leading to improved services and care and thus to improved MNH outcomes [7, 8].

The health and survival of women and their newborn babies has been a key priority in public health for decades, as highlighted by two of the eight ‘Millennium Development Goals’ (MDGs), from the Millennium Declaration endorsed by 189 member states in the year 2000 [9]. Goal number five of these international priorities aims to cut maternal mortality by three-quarters from 1990 to 2015 and goal four focuses on reducing newborn and child mortality by two-thirds. However, as the deadline for these goals approaches, it has become clear that of the eight original goals (which also target poverty, malnutrition, educational levels, environmental conditions and sanitation) goals four and five are the least likely to be achieved in the time span set. Women still die in childbirth—289 000 per year in 2013 [10]—especially in low income countries and although child mortality is dropping, neonatal deaths have not decreased at the same rate, now representing 44 % of all deaths under 5 years [11].

In recognition of poor progress and a lack of information, the UN Secretary-General commissioned a new Global Strategy for Women’s and Children’s Health in 2010 [12] to accelerate progress in reducing maternal and newborn mortality ahead of the rapidly approaching MDG deadline. As a result of this, a Commission on Information and Accountability for Women’s and Children’s Health was established [13] to systematically track progress as well as collect health system data that support the survival and wellbeing of pregnant women and their newborns. These data include indicators that measure the coverage of MNH services, including antenatal care and skilled birth attendance.² Availability of these services is an important step to tackling the problem, but services also need to be accessible to women, acceptable to them and their families and crucially, of a sufficient quality. Finally, women must be able and enabled to make the decision to seek care, otherwise the availability, accessibility, acceptability and quality of services becomes irrelevant [14, 15].

Thus far, the measurement of progress has focused on aggregate indicators at the country level, but increasingly there are concerns that even where progress has been made, it is subject to wide inequalities [16]. Even in countries where national maternal and newborn mortality rates have declined, there may be subgroups where survival rates and access to services have not changed or have worsened. Investigation of sub-national situations through a geographical analysis is now increasingly required.

This paper arises from a workshop held in March 2013 entitled ‘Geography 4 MNH’³, to explore and describe the current “State of the Art” of MNH mapping. The workshop, hosted by Instituto de Cooperación Social Integrare and the University of Southampton, enabled collaboration across research and development agencies to identify existing published and grey literature, describe current work in this field and develop a framework to accelerate the use of this capacity in priority countries with high maternal and neonatal mortality, in order to improve MNH outcomes.

The urgency with which the MNH community must embrace the geographical interpretation of existing databases is particularly critical, both in advance of the endpoint for the MDGs and to inform the post-2015 development agenda for health and the new generation of goals and information requirements over the period 2015–2035 [17]. Successful examples of the application of geographic approaches and use of geospatial data to improve MNH services show the potential of these technologies when integrated within national health strategies. Examples include the use in Bangladesh of an interactive EmONC monitoring GIS system that tracks quality and gaps in services [7]; the use of GIS to evaluate the impact of a community-based health initiative in rural Ghana [8]; and the UK’s teenage pregnancy

reduction strategy, which used ward level data of conceptions in young women under 18 years to assist in targeting interventions where most needed [18].

Review

What has been done so far – a rapid review of previous studies

A rapid review was conducted of the published and grey literature on geographical approaches related to the health and survival of women and children. To identify key sources, an exercise was conducted with technical and academic experts, who were invited to contribute relevant studies to be reviewed. Subsequently, a “snowballing technique” was used to identify further valuable references from these initial sources and so on. The selection was based on a review of title, abstract and key words related to geography and MNH, including: distance, emergency obstetric care, geographic information systems, health facilities, health services accessibility, maternal health, maternal health services, newborn, pregnancy, spatial analysis, *etc.* The start date for inclusion was the year 2000. As well as the articles identified through the “snowballing technique”, additional studies were identified independently by the authors and added to the review.

In total, 33 studies were identified with a focus on MNH. Studies that focus on particular health problems appear to be less common than those that focus on overall disease surveillance, access or risk analysis. A systematic review in 2011 found 621 articles relating to the use of GIS in health, out of which 227 related to disease surveillance, 189 to risk analysis (usually linked to environmental health), 138 to health access and planning, 115 on community profiling and 17 on general or methodological issues [19].

An extraction table was used to compile key information from each of the 33 retrieved papers, namely: year of publication, region and country of implementation, geographical scope, objectives and themes covered, major findings, geographical data layers included, other data collected and GIS approaches used. Table 1 presents selected dimensions of the literature retrieved.

Table 1 Selected information from a rapid literature review on mapping for MNH

Publication	Region	Scope	Theme	GIS approaches used
Sudhof <i>et al.</i> , 2013 [39]	Africa	Sub-district	Geographic access to MNH care	Spatial modelling (travel time to facilities along road networks)
Bowie, 2013 [40]	Africa	National	Geographic access to MNH care	Spatial modelling (travel time to facilities)
Ismaeel & Jabar, 2013 [41]	Middle East	City	Support system for MNH care	Spatial analysis (straight-line distance to facilities)
Echoka <i>et al.</i> , 2013 [42]	Africa	Sub-national	Geographic access to MNH care	Spatial analysis (straight-line distance to facilities)
Lohela <i>et al.</i> , 2012 [43]	Africa	National (rural)	Impact of geographical access on newborn mortality	Spatial analysis (straight-line distance to facilities)
Gething <i>et al.</i> , 2012 [44]	Africa	National	Geographic access to MNH care	Spatial modelling (travel time to facilities along road networks)
Kyei <i>et al.</i> , 2012 [45]	Africa	National (rural)	Impact of geographic access on ANC utilization	Spatial analysis (straight-line distance to facilities)
Pilkington <i>et al.</i> , 2012 [46]	Europe	National	Impact of geographical access on delivery	Spatial analysis (distance to facilities along road networks)
Bailey <i>et al.</i> , 2011 [47]	Africa	Sub-national	Geographic access to MNH care	Spatial modelling (travel time to facilities along road networks)
Gjesfjeld & Jung, 2011 [48]	North America	Sub-national	Geographic access to MNH care	Spatial analysis (straight-line distance to facilities)
Fisher & Myers, 2011 [49]	Asia	Sub-national	Geographic access to MNH care	Spatial modelling (travel time to facilities along road networks)
Gabrysch <i>et al.</i> , 2011 [50]	Africa	National (rural)	Impact of geographical access on use of services.	Spatial analysis (straight-line distance to facilities)
Cordivano, 2011 [51]	North America	City	Impact on maternity ward closure	Spatial analysis (straight-line distance to facilities, analysis of facility based buffers to determine catchment areas)
Redshaw <i>et al.</i> , 2011 [52]	Europe	National	Maternity care organization	Thematic mapping (density of facilities)
Massey, 2011 [53]	Africa	National	Geographic patterns of MNH	Thematic mapping (analysis of hot-spots of adverse outcomes)
Simoes & Almeida, 2011 [54]	Latin America	City	Impact of geographic access on MMR.	Spatial analysis (distance to facilities along road networks)
Zahinos Ruiz, 2010 [55]	Africa	Sub-national	Assess effectiveness (MWHs)	Spatial analysis (investigation of facility based buffers to determine catchment areas)

Saugene, 2010 [56]	Africa	Sub-national	Planning of ambulance routes	Spatial modelling (travel time to facilities along road networks)
Malqvist <i>et al.</i> , 2010 [57]	Asia	Sub-national	Impact of geographic access on mortality and care utilization.	Spatial analysis (straight-line distance to facilities)
Pilkington <i>et al.</i> , 2008 [58]	Europe	National	Impact on maternity unit closure	Spatial analysis (straight-line distance to facilities and, analysis of facility based buffers to determine catchment areas)
Dummer & Parker, 2004 [59]	Europe	Sub-national	Accessibility and infant death risk	Spatial modelling (travel time to facilities along road networks)
Balk <i>et al.</i> , 2003 [20]	Africa	Region	Impact of geography on child/infant mortality	Spatial analysis (straight-line distance to facilities and analysis of facility based buffers to determine catchment areas)
Ayeni, 2002 [60]	Africa	City	Birth weight analysis	Thematic mapping (overlaid different map layers and small area analysis)
Leewannapasai <i>et al.</i> , 2001 [61]	Asia	Sub-district	Geographic access	Thematic mapping (overlaid different map layers)
Jain <i>et al.</i> , 2015 [62]	Asia	Sub-national	Impact of geographical access on institutional deliveries	Spatial analysis (distance to facilities along road networks)
Tatem <i>et al.</i> , 2014 [63]	Asia and Africa	National	Geographic distribution of women of child-bearing age, pregnancies, births and facilities	Thematic mapping (overlaid different map layers)
Heard <i>et al.</i> , 2004 [64]	Africa	National	Impact of geographic access on service utilization	Spatial analysis (straight-line distances to facilities)
Blanford <i>et al.</i> , 2012 [65]	Africa	National	Geographic access	Spatial modelling (travel time to facilities along road networks)
Nesbitt <i>et al.</i> , 2014 [66]	Africa	Sub-national	Accuracy of different measures of geographic access as proxies for access to services.	Spatial analysis (straight-line distance to facilities, distance to facilities along road network) and spatial modelling (travel time to facilities along road networks).
Chong <i>et al.</i> , 2013 [67]	Oceania	City (sub-region)	Use of geographic methods to identify high-risk communities	Thematic mapping (spatial clusters of at risk patients)
Amoako Johnson <i>et al.</i> , 2015 [8]	Africa	National (rural)	Impact of geographical access on use of skilled birth attendance.	Spatial analysis (distance to facilities along road networks)
Ravelli <i>et al.</i> , 2010 [4]	Europe	National	Impact of geographical access on neonatal outcomes.	Spatial modelling (travel time to facilities along road networks)
Okwaraji <i>et al.</i> , 2012 [3]	Africa	Sub-national	Impact of geographical access on under 5 mortality	Spatial modelling (travel time to facilities)

This review shows a clear increase in the number of GIS for MNH publications since 2010. The studies cover most regions of the world. As expected, in view of its high maternal and infant mortality levels [10] Africa has received the greatest research focus with 19 studies. Asia (including South, South-East and East Asia) and Europe follow with 5 studies each.

The majority of the studies conduct their analysis at the national or sub-national level. One of the studies [20] covers the broader geographical region of West Africa, but individual analyses are conducted at the level of each country in the region.

The most common study objective is to examine geographic access to health services.⁴ A variety of different GIS techniques are used to map, extract or model MNH related indicators, from simple thematic mapping, spatial analysis (straight line distance, buffers) or spatial modelling involving travel time cost surfaces, which take transport networks and other geographical features into account.

Work in progress

One of the key purposes of convening parties actively mapping the geography of MNH was to identify the work in progress that may influence the ‘state of the art’. Table 2 provides examples of on-going work, looking at the who?, what? and where? for each project (full descriptions available on request).

Table 2 Examples of work in progress on mapping for MNH

Who?	What	Where?	
Saving Mothers Giving Life (SMGL)/ Centers for Disease Control (CDC)	Thematic Mapping	Maps on maternal mortality and access to facilities	Uganda
Evidence for Action (E4A) [68]/ White Ribbon Alliance (WRA) [69]	Thematic Mapping	Thematic mapping of selected indicator at the sub-national level (2013)	Ethiopia
Countdown to 2015 [70]	Thematic Mapping	Atlas of Birth, extensive mapping. Country case study – focus on maternal mortality reduction	Ghana Bangladesh
United Nations Children’s Fund (UNICEF)	Thematic Mapping	Sharpening Maternal, Newborn and Child Health (MNCH) plans. Using Dev Info maps for program planning	Senegal
USAID	Thematic Mapping, Spatial analysis and modeling	Emergency Obstetric and Newborn Care (EmONC) availability and accessibility coverage (In collaboration with FHI 360, MEASURE Evaluation and AMDD)	Mozambique and Ethiopia
USAID	Spatial analysis	MNH Mapping Modelling sub-national estimates from 2010 Bangladesh Maternal Mortality Survey; Other work under development	Bangladesh
Health 4+ (H4+)/ Technical Working Group and Secretariat + Muskoka [71]	Spatial analysis and modelling	High Burden Countries Initiative (HBCI) Midwifery Workforce Assessment (2012–13). Determining sub-national population needs (through a proxy of the estimated number of pregnancies) in relation to the supply of EmOC facilities and the MNH workforce.	Afghanistan, Bangladesh, Benin, Democratic Republic of Congo, Ethiopia, Ghana, Guinea, India (2 States), Mozambique (in collaboration with USAID), Nigeria, Tanzania, Togo
USAID	Spatial analysis and modelling	Analysis of EmOC availability and accessibility coverage; feasibility of real time mapping.	Malawi
Southampton University	Spatial analysis and modelling	EmOC availability and accessibility coverage.	Ghana
World Health Organization (WHO)	Spatial analysis and modelling	EmOC availability and accessibility coverage as well as scaling up costing analysis	Burkina Faso, Cambodia, Laos, Malawi, Philippines

The work in progress is consistent with the findings of the literature review, both in terms of countries studied and methods employed. Several partners are involved in these activities, creating some overlap in the countries being covered and calling for coordination.

These different projects span a wide range of methods: from the use of thematic maps to visualize indicators at different levels of disaggregation, to the use of complex spatial analysis and modeling techniques in order to assess the accessibility of MNH services. In each case, the power of GIS is used to represent, extract or produce new evidence of interest to the decision-making process (see the next section for a description of the benefits of each method).

Of particular interest is that several of the multi-country initiatives (H4+, USAID and WHO) have independently adopted the Tanahashi framework of effective coverage [14] to guide their technical analyses. This framework considers the dimensions of availability, accessibility, acceptability and quality of services when measuring population and facilitates the identification of bottlenecks and solutions, as appropriate to the geography. This convergence between geography and therefore indirectly GIS and the Tanahashi framework is encouraging and provides scope for ‘good practise’ to be documented and disseminated. Another encouraging finding is that professional learning and capacity-building are incorporated within some of the activities with a longer duration, *e.g.* the multi-country work supported by WHO and Evidence for Action (E4A). Knowledge transfer on the methods, analysis and policy implications will be critical to integrate the geography of MNH into national policy and planning cycles in the future.

The field of Small Area Estimation (SAE) is also emerging in the geographical analysis of MNH and allows the estimation of measures for very small geographies by linking the variable of interest with covariate or contextual information from census, administrative or geographic information systems data to derive model-based estimates for small areas. If data on the variable of interest are available only for a limited number of individuals in the small area, the covariate information is available for all individuals in the area and is assumed to explain part of the between area variability [21, 22]. Two broad types of small-area models exist. Area-level models are used when the auxiliary information is available only at the area level [22] and nested error unit-level models are used the auxiliary information is available at the individual level and is related to the variables of interest at the unit-level [23]. Such methods have been employed widely in high income countries and for economic variables such as poverty indices [24, 25] – but less so for health variables [26, 27]. Amoako-Johnson *et al.* [26] apply SAE for skilled attendance and unmet need for contraception and Ahmed *et al.* [28] have reviewed maternal mortality ratios in Bangladesh.

In demographic and health research, particularly in the estimation of maternal and newborn healthcare indicators there are methodological as well as data challenges that limit the application of small area estimation techniques [29–32]. Despite these limitations, more and more low and middle income countries are using this type of analysis to reduce the extent of inequality in access to health care as, in these countries, local level statistics on healthcare provision and outcomes are rare if not non-existent for aid planning, monitoring and evaluation of programs.

Moving towards comprehensive GIS for MNH

The review of the literature and work in progress presented here has identified three types of GIS methods, namely:

1. Thematic mapping (creation of maps to convey information about a topic or theme);
2. Spatial analyses (extraction or creation of new information from spatial data);
3. Spatial modelling (spatial analysis that includes the use of a mathematical model to simulate natural or anthropogenic phenomena).

Each of these methods adds specific value to the policy discussion. Thematic maps are powerful instruments that allow visualizing sub national information from a geographic perspective. Spatial analysis techniques, including spatial modelling, are advanced methods that are used to provide a more in-depth analysis and understanding of the health systems factors and behaviours behind MNH related health outcomes.

These methods are listed not only in order of increasing complexity but also by their reliance on data in terms of volume, completeness (having all the data on the map), timeliness (chronologic consistency between the different sources of data) and accuracy (precise location of each geographic object), and the advanced level of GIS skills required to apply them.

The possibility for countries to benefit from any of these applications will be influenced by the availability and quality of data and technical resources. Thus, an appropriate and adequately financed institutional framework is needed, to sustain and improve the availability and quality of the data and to further enhance the required technical skills and equipment. Table 3, while not exhaustive, provides an overview of the components that would support such an institutional framework to advance the application and implementation of GIS in MNH.

Table 3 Required data, data quality issues and required GIS skills, by GIS approach to mapping for MNH

	Required data	Important GIS data quality issues to be addressed	Required GIS skills
Thematic mapping	Statistical data (indicators) and GIS data containing the extension of the geographic objects to which the statistical data are attached	Completeness and timeliness	Basic GIS training (Knowledge for producing relevant thematic maps)
Spatial analysis	Location of the different geographic objects necessary to create or extract the new information (location of the households, health facilities,...)	Completeness, timeliness and accuracy	Intermediate GIS training (GIS data management, spatial analysis functions)
Spatial modelling (geographic access)	Location of the health facilities providing the concerned MNH care; Spatial distribution of the population in demand (pregnant women, births,...); Road network; hydrographic network; Digital Elevation Model (DEM); Land cover and coverage capacity of each facility if human resources taken into account in the analysis		Advanced GIS training (GIS data management and manipulation, use of advanced GIS functions)

Additional considerations from the authors' experience to complement this table include:

- 1) The lowest available level of disaggregation should be used to avoid masking potential pockets of heterogeneity in aggregate data;

- 2) Finding accurate GIS data is often a challenge as indicators might have been collected according to different types of divisions (administrative, statistical or even health) and/or at different periods in time;
- 3) A particular challenge resides in the use of MNH indicators that would be available at the health facility level (presence of EmONC signal functions, for example). A master list containing all national health facilities (rather than a sample) as well as their geographic location (latitude/longitude) is an absolute necessity. Ensuring the use of such registry as well as the integration of geography in any health facility based survey would also contribute to facilitating the use of GIS as well as the comparison between maternal/neonatal health services provision and outcomes [33];
- 4) At all levels it is important that the GIS technician works as part of a team of experts to produce timely, reliable and relevant results for decision-making.

Table 3 emphasises the importance of having a strong health information system in place but also for this system to integrate and contextualise both geography and time effectively and accurately. The former requires complete and up-to-date registries, to be established and maintained for each of the geographic objects to which MNH related statistics can be attached (administrative, statistical, health divisions, health facilities). In addition, advanced techniques such as spatial analysis or spatial modelling require the use of GIS data under the mandate of institutions outside of the health sector (for example, National Statistical Office, National Mapping Agency and/or National Road Authority). All these institutions therefore need to use compatible data based on common standards.

The use of GIS also requires that the health sector has the necessary capacity (human resources, software and hardware) in place. While this capacity is increasing, thanks partly to the availability of open source software, in a large number of Ministries of Health it remains at a basic level and limits analysis to thematic mapping. Sustained implementation and continued expertise in countries are needed. Good news on this comes from non-health sectors such as agriculture – where GIS expertise is being utilized more and more frequently and from Health Management Information Systems development – where growing IT teams are increasingly embedding GIS within the standard monitoring tools. The need to support and harness GIS systems and expertise is also explicit in post –2015 plans for statistical development in countries – for example in the forthcoming ‘road map for health measurement’ to be discussed at a summit in Washington June 2015 [34], and among the new leaders of the Inter Agency Group for the SDGs [35], which is being led by national statistical agencies and whose needs for capacity development are increasingly being heard.

Despite these challenges the emphasis on reducing maternal and newborn mortality is an opportunity for new synergies. The health sector can use MNH as the driver to improve GIS capacity, thus benefiting all health initiatives. This win-win situation would be possible through an institutional framework in which:

- 1) The health sector is actively involved in discussions aimed at standardising their own geospatial information. The forum for this is generally referred to as the National Spatial Data Infrastructure (NSDI);
- 2) International health partners supporting MNH program activities strengthen the existing GIS capacity through a coordinated approach; independently from specific health program support;
- 3) Institutions involved in the strengthening of health information systems include the

integration of geographic and time dimensions in their approach;

4) MNH oriented programs systematically consider geographic analysis and GIS as a tool to improve the implementation and monitoring of MNH programs.

The institutional framework would ideally be developed and taken forward as an integral component of the UN Secretary-General's Every Woman, Every Child initiative, with linkages to the Commission on Information and Accountability for Women's and Children's Health (CoIA) and the independent Expert Review Group (iERG).

Conclusions

Despite an encouraging trend of increased interest in applying GIS technology for improving MNH outcomes in recent years, much remains to be done to develop national capacity in low- and middle-income settings, to perform geographical analysis.

This state of the art serves as a baseline to raise awareness of the potential that GIS methods have to offer to MNH and provides the components of an institutional framework that we recommend be included in the further work of the CoIA and the iERG. It calls for better communication and coordination among institutions working at improving the GIS capacity of the health sector, as well as between the health sector and other sectors. Improvements should ideally be independent of particular health priorities, as geography has a shared value to all efforts targeting improved health and population outcomes.

The authors of this paper encourage institutions and individuals working or interested in this area, to make themselves known to the Geography of MNH platform in order to strengthen coordination mechanisms and disseminate global public goods.

Abbreviations

CoIA, Commission on Information and Accountability for Women's and Children's Health; GIS, Geographic information systems; iERG, Independent Expert Review Group; MDG, Millennium development goals; MNH, Maternal and newborn health; NSDI, National spatial data infrastructure; SAE, Small area estimation; WHO, World Health Organization

Endnotes

¹ The neonatal period for the newborn baby is normally defined as the first 28 days of life, and the postpartum health of a woman is similarly considered for a month after the birth. However – most of the adverse outcomes related to the birth occur during the first 7 days.

² The 'Countdown to 2015' initiative collects these service data – and Countdown is the official agency for tracking indicators for the Commission on Information and Accountability. Skilled birth attendance – a key MDG indicator – is a crucial human resources indicator [3, 36]

³ For more information: http://integrare.es/mapping-for-maternal-and-newborn-health-innovative-application-of-gis-2/?goback=.gde_4885650_member_229843409

⁴ The purpose of this paper is not to compare methods that investigate geographical access to services. Please refer to Higgs [37] or Guagliardo [38] for a comparison and description of several of these methods.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SE conceptualized the paper, designed the structure of the paper and was the lead for research and writing of all sections. MGA conducted the literature review and was involved in researching and writing the paper. JC conceptualized the paper and conducted research and writing of all sections. AT was involved in conceptualizing the paper, providing research inputs regarding GIS methodology and commenting on revisions. AM also collaborated in the conceptualization of the paper, collaborated in writing the “work in progress” section and commented on revisions. FAJ provided research and writing inputs for the “work in progress” section. HF and RP participated in designing the concept for the paper and commented on revisions. KS reviewed the drafts of the paper and provided inputs and revisions on the “work in progress” section and tables. SN was involved in revising the manuscript. PB provided research, writing inputs and revisions of the manuscript. ZM was an architect of the main concepts in the paper and was involved with the literature review and writing all sections of the paper. All authors read and approved the final manuscript.

Acknowledgements

This paper is an output of the ‘Mapping for MNH’ research project co-managed by ICS Integrare (Spain/UK) and the University of Southampton (UK) and WorldPop project (www.worldpop.org), with funding from the Norwegian Agency for Development (Norad). Our appreciation is extended to Norad for their financial support.

Individuals participating in the Platform activities acknowledge support from the Science and Technology Directorate, Department of Homeland Security; the Fogarty International Center, National Institutes of Health; NIH/NIAID; the Bill and Melinda Gates Foundation; the UK Department for International Development on the programme ‘Evidence for Action’; the Economic & Social Research Council (ESRC) (Res-167-25-0343); and the United Nations Population Fund.

References

1. Powell-Jackson T, Borghi J, Mueller DH, Patouillard E, Mills A. Countdown to 2015: tracking donor assistance to maternal, newborn and child health. *Lancet*. 2006;368(9541):1077–87.
2. WHO. WHO Recommended Interventions for Improving Maternal and Newborn Health. Geneva: World Health Organization; 2009.

3. Okwaraji YB, Cousens S, Berhane Y, Mulholland K, Edmond K. Effect of geographical access to health facilities on child mortality in rural Ethiopia: a community based cross sectional study. *PLoS One*. 2012;7(3):e33564.
4. Ravelli ACJ, Jager KJ, de Groot MH, Erwich JJHM, Rijninks-van Driel GC, Tromp M, et al. Travel time from home to hospital and adverse perinatal outcomes in women at term in the Netherlands. *BJOG*. 2011;118(4):457–65.
5. Gabrysch S, Campbell OMR. Still too far to walk: literature review of the determinants of delivery service use. *BMC Pregnancy Childbirth*. 2009;9:34.
6. Thaddeus S, Maine D. Too far to walk: maternal mortality in context. *Soc Sci Med*. 1994;38(8):1091–110.
7. Ahmed N. Study reveals inadequate access to EmONC facilities in 24 districts of Banglades. icddr,b. 2014. <http://www.icddrb.org/media-centre/news?id=634&task=view>. Accessed 1 May 2015.
8. Amoako-Johnson F, Frempong-Ainguah F, Matthews Z, Harfoot AJP, Nyarko P, Baschieri A, et al. Evaluating the impact of the community-based health planning and services initiative on uptake of skilled birth care in Ghana. *PLoS One*. 2015;10(3):e0120556.
9. United Nations. The Millennium Development Goals Report 2014. New York: United Nations; 2014. http://www.un.org/millenniumgoals/2014_MDG_report/MDG_2014_English_web.pdf. Accessed 1 Jan 2015.
10. WHO, UNICEF, UNFPA, World Bank, UN Population Division. Trends in Maternal Mortality: 1990–2013. Geneva: World Health Organization; 2014. <http://www.who.int/reproductivehealth/publications/monitoring/maternal-mortality-2013/en/>. Accessed 1 Jan 2015.
11. UNICEF, WHO, World Bank, UN Population Division. Levels and Trends in Child Mortality. New York: United Nations Children's Fund; 2014. http://www.childmortality.org/files_v17/download/UNICEF%202014%20IGME%20child%20mortality%20Report_Final.pdf. Accessed 1 Jan 2015.
12. UN Secretary-General. Global Strategy for Women's and Children's Health. New York: The Partnership for Maternal, Newborn and Child Health; 2010. http://www.who.int/pmnch/activities/advocacy/fulldocument_globalstrategy/en/. Accessed 1 Jan 2015.
13. WHO. Commission on Information and Accountability for Women's and Children's Health. About. 2013. http://www.who.int/woman_child_accountability/about/coia/en/. Accessed 1 Jan 2015.
14. Tanahashi T. Health service coverage and its evaluation. *Bull World Health Organ*. 1978;56(2):295–303.

15. Van Lerberghe W, Matthews Z, Achadi E, Ancona C, Campbell J, Channon A, et al. Country experience with strengthening of health systems and deployment of midwives in countries with high maternal mortality. *Lancet*. 2014;384(9949):1215–25.
16. Bhutta ZA, Reddy S. Achieving equity in global health: so near and yet so far. *J Am Med Assoc*. 2012;307(19):2035–6.
17. Nations U. A new Global Partnership: Eradicate Poverty and Transform Economies through Sustainable Development. New York: United Nations; 2013.
18. Department for Education and Skills. Teenage Pregnancy: Accelerating the Strategy to 2010. 2006. <http://webarchive.nationalarchives.gov.uk/20130401151715/http://www.education.gov.uk/publications/eOrderingDownload/DFES-03905-2006.pdf>. Accessed 1 May 2015.
19. Nykiforuk CIJ, Flaman LM. Geographic information systems (GIS) for health promotion and public health: a review. *Health Promot Pract*. 2011;12(1):63–73.
20. Balk D, Pullum T, Storeygard A, Greenwell F, Neuman M. Spatial Analysis of Childhood Mortality in West Africa. Calverton: ORC Macro and Center for International Earth Science Information Network (CIESIN), Columbia University; 2003.
21. Pfeffermann D. Small area estimation: new developments and directions. *Int Stat Rev*. 2002;70(1):125–43.
22. Fay RE, Herriot RA. Estimates of income for small places: an application of James-Stein procedures to census data. *J Am Stat Assoc*. 1979;74(366):269–77.
23. Battese GE, Harter RM, Fuller WA. An error-components model for prediction of county crop areas using survey and satellite data. *J Am Stat Assoc*. 1988;83(401):28–36.
24. Elbers C, Lanjouw JO, Lanjouw P. Micro-level estimation of poverty and inequality. *Econometrica*. 2003;71(1):355–64.
25. Demombynes G, Elbers C, Lanjouw JO, Lanjouw P. How good a map? Putting small area estimation to the test. The World Bank. 2007. <http://elibrary.worldbank.org/doi/book/10.1596/1813-9450-4155>. Accessed 1 Jan 2015.
26. Amoako Johnson F, Padmadas SS, Chandra H, Matthews Z, Madise NJ. Estimating unmet need for contraception by district within Ghana: an application of small-area estimation techniques. *Popul Stud (NY)*. 2012;66(2):105–22.
27. Fujii T. Micro-Level Estimation of Child Malnutrition Indicators and its Application in Cambodia. The World Bank. 2005. <http://elibrary.worldbank.org/doi/book/10.1596/1813-9450-3662>. Accessed 1 Jan 2015.
28. Ahmed S, Hill K. Maternal mortality estimation at the subnational level: a model-based method with an application to Bangladesh. *Bull World Health Organ*. 2011;89(1):12–21.

29. McCullagh P, Nelder JA, McCullagh P. *Generalized Linear Models*. Vol. 2. London: Chapman and Hall; 1989.

30. Chandra H, Sud UC. Small area estimation for zero-inflated data. *Commun Stat - Simul Comput*. 2012;41(5):632–43.

31. Chandra H, Chambers R. Small area estimation for skewed data in presence of zeros. *Bull Calcutta Stat Assoc*. 2011;63:249–52.

32. Pfeffermann D, Terryn B, Moura FAS. Small area estimation under a two-part random effects model with application to estimation of literacy in developing countries. *Surv Methodol*. 2008;34(2):235–49.

33. DeJong J, Akik C, El Kak F, Osman H, El-Jardali F. The safety and quality of childbirth in the context of health systems: mapping maternal health provision in Lebanon. *Midwifery*. 2010;26(5):549–57.

34. MA4Health. MA4 health roadmap. 2015 <http://ma4health.haccess.org/roadmap>. Accessed 1 May 2015.

35. United Nations Economic and Social Council. Report of the Friends of the Chair Group on Broader Measures of Progress. Forty-Sixth Session of the Statistical Commission. New York: United Nations; 2015.

36. WHO, UNICEF. Countdown to 2015: Building a Future for Women and Children. The 2012 Report. Geneva: WHO; UNICEF; 2012.

37. Higgs G. A Literature Review of the use of GIS-based Measures of Access to Health Care Services. *Heal Serv Outcomes Res Methodol*. 2004;5(2):119–39.

38. Guagliardo MF. Spatial accessibility of primary care: concepts, methods and challenges. *Int J Health Geogr*. 2004;3(1):3.

39. Sudhof L, Amoroso C, Barebwanuwe P, Munyaneza F, Karamaga A, Zambotti G, et al. Local use of geographic information systems to improve data utilisation and health services: mapping caesarean section coverage in rural Rwanda. *Trop Med Int Heal*. 2013;18(1):18–26.

40. Bowie C. Recent trends in geographical access to health care facilities in Malawi. Blantyre: Division of Community Health, College of Medicine, University of Malawi; 2013. <http://www.medcol.mw/commhealth/publications/national%20research/Recent%20Trends%20in%20Geographical%20Access%20to%20Emergency%20Obstetric%20Care%20in%20Malawi6.pdf>. Accessed 1 May 2015.

41. Ismaeel AG, Jabar EK. Effective system for pregnant women using mobile GIS. *Int J Comput Appl*. 2013;64(11):1–7.

42. Echoka E, Kombe Y, Dubourg D, Makokha A, Evjen-Olsen B, Mwangi M, et al. Existence and functionality of emergency obstetric care services at district level in Kenya: theoretical coverage versus reality. *BMC Health Serv Re*. 2013;13(1):113.

43. Lohela TJ, Campbell OMR, Gabrysch S. Distance to care, facility delivery and early neonatal mortality in Malawi and Zambia. *PLoS One*. 2012;7(12):e52110.

44. Gething PW, Johnson FA, Frempong-Ainguah F, Nyarko P, Baschieri A, Aboagye P, et al. Geographical access to care at birth in Ghana: a barrier to safe motherhood. *BMC Public Health*. 2012;12:991.

45. Kyei NN, Campbell OMR, Gabrysch S. The influence of distance and level of service provision on antenatal care use in rural Zambia. *PLoS One*. 2012;7(10):1–6.

46. Pilkington H, Blondel B, Drewniak N, Zeitlin J. Choice in maternity care: associations with unit supply, geographic accessibility and user characteristics. *Int J Health Geogr*. 2012;11:35.

47. Bailey PE, Keyes EB, Parker C, Abdullah M, Kebede H, Freedman L. Using a GIS to model interventions to strengthen the emergency referral system for maternal and newborn health in Ethiopia. *Int J Gynecol Obstet [Internet] Int Fed Gynecol Obstet*. 2011;115(3):300–9.

48. Gjesfjeld CD, Jung J-K. How far?: Using geographical information systems (GIS) to examine maternity care access for expectant mothers in a rural state. *Soc Work Health Care*. 2011;50(9):682–93.

49. Fisher RP, Myers B. Free and simple GIS as appropriate for health mapping in a low resource setting: a case study in eastern Indonesia. *Int J Health Geogr*. 2011;10(1):15.

50. Gabrysch S, Cousens S, Cox J, Campbell OMR. The influence of distance and level of care on delivery place in rural Zambia: a study of linked national data in a geographic information system. *PLoS Med*. 2011;8(1):e1000394.

51. Cordivano S. Maternity ward closures in Philadelphia: using GIS to measure disruptions in essential health services. *J Map Geogr Libr*. 2011;7(3):282–303.

52. Redshaw M, Rowe R, Schroeder L, Macfarlane A, Newburn M, Sandall J, et al. Mapping maternity care: the configuration of maternity care in England. Birthplace in England research programme. Final report part 3. NIHR service delivery and organisation programme. 2011.

53. Massey P. Reducing maternal mortality in Senegal: using GIS to identify priority regions for the expansion of human resources for health. *World Health Popul*. 2011;13(2):13–22.

54. Simões PP, Almeida RMVR. Geographic accessibility to obstetric care and maternal mortality in a large metropolitan area of Brazil. *Int J Gynaecol Obstet*. 2011;112(1):25–9.

55. Zahinos Ruiz I. Effectiveness Assessment of Maternity Waiting Homes in Increasing Coverage of Institutional Deliveries using Geographical Information Systems in Six Districts of Cabo Delgado Province (Mozambique). Pretoria: University of South Africa; 2010.

56. Saugene ZB. Achieving the Health Millennium Developing Goals using GIS Knowledge Networks: The Case of Maternal Health in Mozambique. GSDI 12 World conference, Singapore, October 19, 2010. <http://www.gsdi.org/gsdiconf/gsdi12/papers/107.pdf>. Accessed 1 Jan 2015.

57. Malqvist M, Sohel N, Do TT, Eriksson L, Persson LA. Distance decay in delivery care utilisation associated with neonatal mortality. A case referent study in northern Vietnam. *BMC Public Health*. 2010;10:762.

58. Pilkington H, Blondel B, Carayol M, Breart G, Zeitlin J. Impact of maternity unit closures on access to obstetrical care: the French experience between 1998 and 2003. *Soc Sci Med*. 2008;67(10):1521–9.

59. Dummer TJB, Parker L. Hospital accessibility and infant death risk. *Arch Dis Child*. 2004;89(3):232–4.

60. Ayeni B. An application of GIS to the analysis of maternal and child healthcare delivery in Ibadan, Nigeria. Nigeria: University of Ibadan; 2002. <http://www.uiowa.edu/~gishlh/PDFs/ibadanreport.pdf>. Accessed 1 Jan 2015.

61. Leewannapaisai S, Durongdej S, Sowana A. Application of Geographic Information Systems (GIS) for Analysing and Planning of Reproductive Health Services in Pattani Province. 2001. <http://geospatialworld.net/Paper/Application/ArticleView.aspx?aid=850>. Accessed 1 Jan 2015.

62. Jain AK, Sathar ZA, ul Haque M. The constraints of distance and poverty on institutional deliveries in Pakistan: evidence from georeference-linked data. *Stud Fam Plan*. 2015;46(1):21–39.

63. Tatem AJ, Campbell J, Guerra Arias M, de Bernis L, Moran A, Matthews Z. Mapping for maternal and newborn health: the distributions of women of childbearing age, pregnancies and births. *Int J Health Geogr*. 2014;13(1):2.

64. Heard NJ, Larsen U, Hozumi D. Investigating access to reproductive health services using GIS: proximity to services and the use of modern contraceptives in Malawi. *Afr J Reprod Health*. 2004;8(2):164–79.

65. Blanford JI, Kumar S, Luo W, MacEachren AM. It's a long, long walk: accessibility to hospitals, maternity and integrated health centers in Niger. *Int J Health Geogr*. 2012;11:24.

66. Nesbitt RC, Gabrysich S, Laub A, Soremekun S, Manu A, Kirkwood BR, et al. Methods to measure potential spatial access to delivery care in low- and middle-income countries: a case study in rural Ghana. *Int J Health Geogr*. 2014;13:25.

67. Chong S, Nelson M, Byun R, Harris L, Eastwood J, Jalaludin B. Geospatial analyses to identify clusters of adverse antenatal factors for targeted interventions. *Int J Health Geogr*. 2013;12(1):46.

68. Evidence for Action. <http://www.evidence4action.net/> (2015). Accessed 1 Jan 2015.

69. White Ribbon Alliance. <http://www.whiteribbonalliance.org/> (2013). Accessed 1 Jan 2015.

70. Countdown to 2015. <http://www.countdown2015mnch.org/> (2015). Accessed 1 Jan 2015.

71. Foreign Affairs, Trade and Development Canada: The Muskoka Initiative: Background. <http://www.acdi-cida.gc.ca/acdi-cida/ACDI-CIDA.nsf/En/FRA-119133138-PQT> (2012). Accessed 1 Jan 2015.