



Cite this: *Chem. Commun.*, 2015, 51, 9555

Received 16th April 2015,
Accepted 8th May 2015

DOI: 10.1039/c5cc03184b

www.rsc.org/chemcomm

Unique Group 1 cations stabilised by homoleptic neutral phosphine coordination†

Marina Carravetta, Maria Concistre, William Levason, Gillian Reid* and Wenjian Zhang

Homoleptic coordination of the neutral diphosphines $\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2$ and $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$ to the hard Li^+ and Na^+ cations is achieved using $\text{Li}[\text{Al}(\text{OC}(\text{CF}_3)_3)_4]$ and $\text{Na}[\text{B}(\text{3,5-}(\text{CF}_3)_2\text{-C}_6\text{H}_3)_4]$ as 'naked' cation sources. Crystallographic, solid state and solution multinuclear NMR studies confirm distorted octahedral coordination solely *via* three chelating diphosphines in these unique species.

Neutral phosphine ligands, PR_3 ($\text{R} = \text{alkyl, aryl}$), are ubiquitous in transition metal chemistry, owing to their capacity to tune the electronic and steric properties, and hence the reactivity, of the complexes, and to the strong σ -donor properties of the soft phosphine donor functions. This has led to wide utilisation of phosphine co-ligands in many transition metal reagents and catalysts.¹ Phosphine complexes of many p-block acceptors have also developed substantially in recent years.² However, complexes involving coordination of neutral phosphine ligands towards the strongly electropositive s-block elements, particularly the Group 1 cations, have remained extremely elusive, and there are no reported examples with exclusively PR_3 coordination. This is no doubt in part due to the high affinity of the alkali metal and alkaline earth cations for hard, electronegative Lewis bases such as water, alkoxide, amide *etc.*, their high lability, as well as the high lattice energies often associated with many Group 1 and 2 salts, which severely limit their solubilities in non-competitive organic media. Thus, to-date there has been only one reported example of a neutral phosphine co-ligand coordinated to an alkali metal cation, the organometallic silylamide dimer $[\text{Li}\{\text{N}(\text{Ar})\text{CC}(\text{R})\text{-Si}(\text{R})_2\text{NAr}\}\{\mu\text{-Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2\}]_2$ (mean $d(\text{Li-P}) = 2.650(3) \text{ \AA}$),³ and two structurally authenticated species with PR_3 coordination to alkaline earth ions; $[\text{BeCl}_2(\kappa^1\text{-Ph}_2\text{PCH}_2\text{PPh}_2)_2]$ ⁴ and the dinuclear $[\text{Be}_2\text{Cl}_2(\mu\text{-Cl})_2(\text{PCy}_3)_2]$,⁵ both containing distorted tetrahedral $\text{Be}(\text{II})$.

A small number of anionic ligands bearing phosphine functions have been coordinated to s-block cations, including $[\text{Mg}\{\text{C}_6\text{H}_3\text{-2,6-}(\text{CH}_2\text{PMe}_2)_2\}]_2$, in which the Mg-P bonds are also stabilised by the anionic pincer ligand framework,⁶ $[\{\text{Li}(2\text{-PPh}_2\text{-C}_6\text{H}_4)\}_2(\text{OEt}_2)_2]$ ($d(\text{Li-P}) = 2.69\text{--}2.75 \text{ \AA}$),⁷ hindered alkoxy-phosphine complexes, including $[\text{Li}(\mu\text{-OC}^t\text{Bu}_2\text{CH}_2\text{PR}_2)]_2$ ($\text{R} = \text{Me}$ or Ph) and $[\text{Li}(\mu\text{-OC}^t\text{Bu}_2\text{CH}_2\text{PPh}_2)_2\text{Li}(\text{OC}^t\text{Bu}_2)]$ ($d(\text{Li-P}) = 2.50\text{--}2.65 \text{ \AA}$),⁸ $[\text{Na}(\text{H}_2\text{Al}\{\text{P}(\text{SiMe}_3)_2\}_2)(\text{dme})_2]$,⁹ $[\text{Li}(\text{solvent})_x\{\text{Ph}_2\text{B}(\text{CH}_2\text{P}^i\text{Pr}_2)_2\}]$ (solvent = thf, $x = 2$; Et_2O , $x = 1$) ($d(\text{Li-P}) = 2.596(3)$, $2.608(3) \text{ \AA}$),¹⁰ as well as a small number of Li^+ complexes with (phosphinomethyl)aluminate ligands.¹¹ The negative charge on the anionic ligands in these species brings a significant electrostatic component to the bonding, and contrasts the covalent metal-phosphine bonding present in the d- and p-block acceptor ions.

In recent work we reported¹² that complexes of Na^+ with polyamines and aza macrocycles, including the $[\text{Na}(\text{Me}_3\text{-tacn})_2]^+$ sandwich cation and the distorted five-coordinate $[\text{Na}(\text{thf})(\text{Me}_4\text{-cyclam})]^+$ cation ($\text{Me}_3\text{-tacn} = 1,4,7\text{-trimethyl-1,4,7-triazacyclononane}$, $\text{Me}_4\text{-cyclam} = 1,4,8,11\text{-tetramethyl-1,4,8,11-tetraazacyclotetradecane}$), could be prepared readily by using $\text{Na}[\text{BAR}^F] \cdot 2\text{thf}$ ($[\text{BAR}^F]^- = [\text{B}\{3,5\text{-}(\text{CF}_3)_2\text{-C}_6\text{H}_3\}_4]^-$). This $[\text{BAR}^F]^-$ salt¹³ has good solubility in weak-donor solvents such as CH_2Cl_2 and toluene. Very recently we extended this chemistry by reporting the homoleptic octathia coordination to Na^+ in the macrocyclic complex, $[\text{Na}(\text{[24]aneS}_8)][\text{BAR}^F]$ ($[\text{24]aneS}_8 = 1,4,7,10,13,16,19,22\text{-octathia-cyclotetrasocane}$), containing distorted dodecahedral coordination.¹⁴ To develop this chemistry further we sought to establish whether it would be possible to induce coordination of softer, neutral phosphine ligands towards Group 1 cations without the additional stability offered by the macrocyclic frameworks employed in the aza and thioether chemistry. To achieve this we have used both the $[\text{BAR}^F]^-$ ¹⁵ and $[\text{Al}\{\text{OC}(\text{CF}_3)_3\}_4]^-$ weakly coordinating anions.^{16–19}

We describe here the first series of Group 1 cations coordinated only to neutral phosphine ligands, in the form of distorted octahedral Li^+ and Na^+ cations containing tris-diphosphine coordination.

Reaction of $\text{Li}[\text{Al}\{\text{OC}(\text{CF}_3)_3\}_4]^{19}$ with three mol. equiv. of $\text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2$ (dmpe) or $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$ (diphos) in anhydrous

Chemistry, University of Southampton, Highfield, Southampton SO17 1BJ, UK.

E-mail: G.Reid@soton.ac.uk

† Electronic supplementary information (ESI) available: A packing diagram for $[\text{Na}(\text{diphos})_3][\text{BAR}^F]$ and MAS NMR data (^7Li , ^{23}Na and ^{31}P) for the phosphine complexes reported. CCDC 1044099–1044101. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5cc03184b



toluene gives $[\text{Li}(\text{dmpe})_3][\text{Al}\{\text{OC}(\text{CF}_3)_3\}_4]$ and $[\text{Li}(\text{diphos})_3][\text{Al}\{\text{OC}(\text{CF}_3)_3\}_4]$, respectively as white solids in very good yield.‡ The corresponding $[\text{Na}(\text{dmpe})_3][\text{BAR}^{\text{F}}]$ and $[\text{Na}(\text{diphos})_3][\text{BAR}^{\text{F}}]$ salts were obtained similarly from a 3:1 diphosphine: $\text{Na}[\text{BAR}^{\text{F}}]$ ratio. Attempts to prepare the analogous K^+ complexes by reaction of the diphosphine with $\text{K}[\text{BAR}^{\text{F}}]$ in a 3:1 molar ratio failed, while the weaker donor and sterically bulkier $o\text{-C}_6\text{H}_4(\text{PPh}_2)_2$, and the diarsine, $o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2$ (the direct analogue of diphos), did not coordinate to Li^+ or Na^+ under similar conditions.

The coordination environments present in the new complexes were established unambiguously from X-ray crystallographic studies on three examples. The structure§ of $[\text{Na}(\text{dmpe})_3][\text{BAR}^{\text{F}}]$ contains discrete Na^+ cations coordinated to three chelating dmpe ligands, in a distorted octahedral environment (Fig. 1), with discrete $[\text{BAR}^{\text{F}}]^-$ anions providing charge balance. The Na–P bond distances lie in the range 2.95–3.03 Å, suggesting relatively weak coordination, and the P–Na–P angles within the five-membered chelate rings are very acute (69.8–73.4°). A similar structure is present in $[\text{Na}(\text{diphos})_3][\text{BAR}^{\text{F}}]$,§ with coordination at Na^+ through six P-donor atoms from three chelating diphos ligands, with $d(\text{Na}–\text{P}) = 2.92\text{--}3.07$ Å (Fig. 2). As in the dmpe complex, these are considerably longer than the sum of the ionic radius for Na (1.02 Å) and the covalent radius for P (1.06 Å). They compare with $[\text{Na}(\text{H}_2\text{Al}\{\text{P}(\text{SiMe}_3)_2\}_2)(\text{dme})_2]$ ($d(\text{Na}–\text{P}) = 3.052(1), 3.092(1)$ Å).⁹ The P–Na–P angles within the chelate rings are even more acute, *ca.* 65°, reflecting the smaller bite angle associated with the rigid *o*-phenylene diphosphine *cf.* the dimethylene-linked dmpe. The large $[\text{BAR}^{\text{F}}]^-$ anions remain discrete, but interleave between the cations (Fig. S1, ESI†).

The structure§ of the lithium–diphosphine complex, $[\text{Li}(\text{diphos})_3][\text{Al}\{\text{OC}(\text{CF}_3)_3\}_4]$ was also determined from a small, weakly diffracting crystal. While the weak diffraction data mean that detailed geometric comparisons require caution, the presence of three chelating neutral diphos ligands at Li^+ , giving




Fig. 1 View of the structure of the $[\text{Na}(\text{dmpe})_3]^+$ cation with numbering scheme adopted. Ellipsoids are drawn at the 50% probability level and H atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): Na1–P1 3.0287(8), Na1–P2 3.0035(8), Na1–P3 2.9621(8), Na1–P4 2.9459(8), Na1–P5 2.9718(8), Na1–P6 2.9960(8), P1–Na1–P2 69.801(18), P3–Na1–P4 73.40(2), P5–Na1–P6 71.48(2).




Fig. 2 View of the structure of the $[\text{Na}(\text{diphos})_3]^+$ cation with numbering scheme adopted. Ellipsoids are drawn at the 50% probability level and H atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): Na1–P1 2.9216(12), Na1–P2 3.0677(12), Na1–P3 3.0087(12), Na1–P4 2.9937(12), Na1–P5 2.9992(12), Na1–P6 3.0400(12), P1–Na1–P2 65.36(3), P3–Na1–P4 65.49(3), P5–Na1–P6 65.57(3).

homoleptic P_6 -coordination, is unequivocal (Fig. 3). The aluminate anion provides charge balance, but does not interact with the cation. The Li–P bond distances are considerably shorter (by *ca.* 0.4 Å) than $d(\text{Na}–\text{P})$ in these systems, while the P–Li–P angles within the chelate rings are correspondingly larger (*ca.* 75°), as expected due to the smaller ionic radius.

To investigate the properties of these unusual complexes further, we obtained the MAS NMR spectroscopic data (^{31}P , ^{23}Na and ^7Li) from the powdered solids. The NMR data are summarised in Table 1. The spectra for $[\text{Li}(\text{dmpe})_3]^+$ are shown in Fig. 4 (the other spectra are provided as ESI,† Fig. S2–S4).¶ The ^{31}P NMR data from direct excitation (Fig. 4(a)) exhibits two peaks, the main one at –54.5 ppm is attributed to the six equivalent P-donor atoms in the complex cation; the minor peak at –48.5 ppm is consistent with the chemical shift for ‘free’ dmpe




Fig. 3 View of the structure of the $[\text{Li}(\text{diphos})_3]^+$ cation with numbering scheme adopted. Ellipsoids are drawn at the 50% probability level and H atoms are omitted for clarity.



Table 1 Selected solid state and solution NMR spectroscopic data^a

Complex	$\delta^{31}\text{P}/\text{ppm}$		$\delta^7\text{Li}/\text{ppm}$		$\delta^{23}\text{Na}/\text{ppm}^c$	
	Solid	Soln ^b	Solid	Soln ^b	Solid	Soln ^b
[Li(diphos) ₃][Al{OC(CF ₃) ₃ } ₄]	-59.2	-53.0	+0.4	-0.1	—	—
[Li(dmpe) ₃][Al{OC(CF ₃) ₃ } ₄]	-54.5	-45.9	-0.7	-1.3	—	—
[Na(diphos) ₃][BAR ^F]	-61.5, -59.0 (1:1)	-50.3	—	—	+3.8	+5.5
[Na(dmpe) ₃][BAR ^F]	-57.4	-49.7	—	—	+8.9	+2.5

^a Small amounts of uncomplexed ligand (literature $\delta^{31}\text{P} = -48$ (dmpe)²⁰ and -55 (diphos)²⁰) were also observed in the solid state spectra in some of the samples, arising from some sample degradation during spectral acquisition. ^b Li complexes recorded in d⁸-toluene solution (298 K); Na complexes recorded in CD₂Cl₂ solution (298 K). ^c $\delta^{23}\text{Na}$ measured for Na[BAR^F] = -35.5 (s) ppm.




Fig. 4 Solid state MAS NMR measurements at 9.4 T and 7.1 kHz spinning frequency on [Li(dmpe)₃][Al{OC(CF₃)₃}₄]. Spinning sidebands are labelled with asterisks. (a) Direct acquisition ³¹P NMR spectrum. The spectrum is an average of 8 scans with inter-pulse delay of 60 s. The main peak at -54.5 ppm is due to the complex, while the small peak at -48.5 ppm corresponds to 'free' dmpe (due to a small amount of decomposition of the complex during data acquisition). (b) ³¹P CP MAS NMR spectrum, where the peak from the very mobile reagent ('free' dmpe) is absent. The spectrum is an average of 16 scans with inter-pulse delay of 10 s. (c) Direct excitation ⁷Li NMR spectrum. The spectrum is an average of 32 scans with inter-pulse delay of 15 s. The signal is at -0.7 ppm.

in solution (-48 ppm).²⁰ This is further confirmed by ³¹P cross-polarization (CP) MAS²¹ data (Fig. 4(b)), where the second peak is absent, in accord with the highly mobile nature of 'free' dmpe.

The chemical shift differences between 'free' and coordinated diphosphine (Δ) are small and negative for all four complexes; (L = dmpe: $\Delta = -6$ ppm for Li⁺; -9.4 ppm for Na⁺; L = diphos: $\Delta = -5$ ppm for Li⁺; ca. -6 ppm for Na⁺). These contrast with the large, positive Δ values typically observed in transition metal phosphine complexes which contain five-membered chelate rings.²² No ⁷Li-³¹P/²³Na-³¹P couplings are evident in the spectra, presumably due to the small magnitude of the J values, which fall within the line width.

Fig. 4(c) shows the ⁷Li NMR data for [Li(dmpe)₃][Al{OC(CF₃)₃}₄], showing a single peak at -0.7 ppm.

Solution ¹H and ³¹P{¹H} NMR spectra (d⁸-toluene or CD₂Cl₂) on the four compounds also show very small coordination

shifts, although in these spectra the resonances are closer to the respective 'free' ligand. These, as well as the ⁷Li and ²³Na solution spectra, are essentially unchanged upon cooling to 183 K (it seems likely that the low temperature-limiting spectrum is not reached at the freezing point of the solvent). These observations may indicate that in solution the complexes are partially dissociated, leading to chemical shifts closer to the free ligand values. Sharp singlets are observed by ⁷Li and ²³Na NMR spectroscopy, with chemical shifts similar to those from the solid state spectra (Table 1).

These results demonstrate that unusual homoleptic neutral phosphine complexes of the Group 1 cations can be readily accessed in (non-polar) organic media through the use of the strong σ -donating bidentate ligands with 'naked' metal cation sources.

We thank the EPSRC for support through a Programme Grant (EP/I033394/1). The SCFED Project (www.scfed.net) is a multidisciplinary collaboration of British universities investigating the fundamental and applied aspects of supercritical fluids. MCA also thanks the Royal Society for a University Research Fellowship.

Notes and references

‡ Synthetic procedure. Schlenk techniques and a glove-box were used for all manipulations, which were conducted under anhydrous and anaerobic conditions.

[Li(diphos)₃][Al{OC(CF₃)₃}₄]: Li[Al{OC(CF₃)₃}₄] (121 mg, 0.125 mmol) in 3 mL toluene was added *o*-C₆H₄(PMe₂)₂ (74 mg, 0.375 mmol) with stirring. After 5 min, the volatiles were removed *in vacuo*. The white solid residue was washed with pentane and dried *in vacuo*. Yield: 181 mg 92%. Colourless single crystals were grown by slow evaporation from a toluene solution under N₂. Anal, required for C₄₆H₄₈AlF₃₆LiO₄P₆ (1568.5): C, 35.2%; H, 3.0%. Found: C, 35.2%; H, 3.0%. ¹H NMR (CD₂Cl₂, 298 K): 1.35 (s, [36H], Me), 7.37–7.43 (m, [6H], *o*-C₆H₄), 7.47–7.53 (m, [6H], *o*-C₆H₄) ppm.

[Na(diphos)₃][B{3,5-(CF₃)₂-C₆H₃}₄]: method as above, using Na[B{3,5-(CF₃)₂-C₆H₃}₄] (110 mg, 0.125 mmol) *o*-C₆H₄(PMe₂)₂ (74 mg, 0.375 mmol). White solid. Yield: 160 mg, 86%. Colourless single crystals were grown by slow evaporation from a toluene solution under N₂. Anal, required for C₆₂H₆₀BF₂₄NaP₆ (1480.7): C, 50.3%; H, 4.0%. Found: C, 50.4%; H, 4.0%. ¹H NMR (CD₂Cl₂, 298 K): 1.20 (s, [36H], Me), 7.34–7.37 (m, [6H], *o*-C₆H₄), 7.42–7.48 (m, [10H], overlapping *o*-C₆H₄ and [BAR^F]⁻ H4), 7.62 (s, [8H], [BAR^F]⁻ H2/6) ppm.

[Li(dmpe)₃][Al{OC(CF₃)₃}₄]: Li[Al{OC(CF₃)₃}₄] (121 mg, 0.125 mmol) in 5 mL toluene was added dmpe (58 mg, 0.375 mmol) with stirring. After 5 min, the volatiles were removed *in vacuo*. The white solid residue was washed with anhydrous pentane and dried *in vacuo*. Yield: 164 mg, 91%. Anal, required for C₃₄H₄₈AlF₃₆LiO₄P₆ (1424.4): C, 28.6%; H, 3.3%. Found: C, 28.7%; H, 3.0%. ¹H NMR (CD₂Cl₂, 298 K): 1.30 (br s, [36H], Me), 1.85 (br s, [12H], CH₂) ppm.

[Na(dmpe)₃][B{3,5-(CF₃)₂-C₆H₃}₄]: As above, using Na[B{3,5-(CF₃)₂-C₆H₃}₄] (110 mg, 0.125 mmol) and dmpe (58 mg, 0.375 mmol).



White solid. Yield: 143 mg, 85%. Colourless single crystals were grown by slow evaporation from a toluene solution under N₂. Anal, required for C₅₀H₆₀BF₂₄NaP₆ (1336.6): C, 44.9; H, 4.5%. Found: C, 44.8; H, 4.5%. ¹H NMR (CD₂Cl₂, 298 K): 1.01 (br s, [36H], Me), 1.45 (br s, [12H], CH₂), 7.46 (s, [4H], [BAR^F]⁻ H4), 7.62 (s, [8H], [BAR^F]⁻ H2/6) ppm.

§ Crystal data for [Na(dmpe)₃][BAR^F]: C₅₀H₆₀BF₂₄NaP₆, M_w = 1336.60, monoclinic, P2₁/c (no. 14), a = 18.4514(10), b = 13.2631(10), c = 26.133(2) Å, β = 90.826(3)°, U = 6394.6(8) Å³, Z = 4, μ = 0.277 mm⁻¹, T = 100 K, 35 720 total reflections, 12 513 unique reflections, R_{int} = 0.059, R₁ (I > 2σ(I)) = 0.068, R₁ (all data) = 0.105, wR₂ (I > 2σ(I)) = 0.169, wR₂ (all data) = 0.197.

Crystal data for [Na(diphos)₃][BAR^F]: C₆₂H₆₀BF₂₄NaP₆, M_w = 1480.72, triclinic, P1̄ (no. 2), a = 12.5084(10), b = 16.4798(10), c = 17.5637(10) Å, α = 79.047(4), β = 78.403(4), γ = 81.625(4)°, U = 3460.2(4) Å³, Z = 2, μ = 0.264 mm⁻¹, 31 294 total reflections, 13 561 unique reflections, R_{int} = 0.033, R₁ (I > 2σ(I)) = 0.053, R₁ (all data) = 0.063, wR₂ (I > 2σ(I)) = 0.143, wR₂ (all data) = 0.149.

Crystal data for [Li(diphos)₃][Al{OC(CF₃)₃}₄]: C₄₆H₄₈AlF₃₆LiO₄P₆, M_w = 1568.58, orthorhombic, Pna2₁ (no. 33), a = 15.092(2), b = 27.737(5), c = 15.405(3) Å, U = 6449.1(19) Å³, Z = 4, μ = 0.325 mm⁻¹, 38 663 total reflections, 12 559 unique reflections, R_{int} = 0.093, R₁ (I > 2σ(I)) = 0.101, R₁ (all data) = 0.150, wR₂ (I > 2σ(I)) = 0.210, wR₂ (all data) = 0.236. Data collections used a Rigaku AFC12 goniometer equipped with an enhanced sensitivity (HG) Saturn724+ detector mounted at the window of an FR-E+ SuperBright molybdenum rotating anode generator (λ₁ = 0.71073 Å) with VHF Varimax optics (70 μm focus). The crystals were held at 100 K in a nitrogen gas stream. Structure solution and refinement on the Na complexes were mostly straightforward,^{23,24} except for some disorder in the CF₃ groups of the [BAR^F]⁻ anions which was modelled satisfactorily. For [Li(diphos)₃][Al{OC(CF₃)₃}₄], despite several attempts, only very small crystals could be obtained. This led to weak diffraction, particularly at high angle, and hence higher R-factors and a less well-defined structure. The H atoms were placed in calculated positions and refined using a riding model. The H atoms on the disordered CH₂ and CH₃ group were not located. CCDC 1044099–1044101.

¶ Solid state NMR experiments. All measurements were performed on a Bruker 9.4 T magnet with a Chemagnetics Infinity console using a double-resonance 4 mm APEX probe. Solid powdered samples were transferred into 4 mm zirconium oxide thin wall rotors within the glovebox, using special end caps with o-rings to exclude air. Magic angle spinning (MAS) conditions have been applied with a spinning speed of 7.1 kHz at room temperature, using N₂ gas flow for bearing and drive. The chemical shift scales were referenced by setting at 0 ppm the signals of LiCl 1 M, NaCl 1 M and 85% H₃PO₄, respectively for ⁷Li, ²³Na and ³¹P. Spectra were recorded with direct excitation using a 90° pulse followed by acquisition, without proton decoupling. For ³¹P NMR, additional measurements were also performed with ramped CP methods with 3 ms contact time.

- 1 See for example *Comprehensive Coordination Chemistry*, ed. G. Wilkinson, R. D. Gillard and J. A. McCleverty, Pergamon Press, Oxford, 1987; *Comprehensive Coordination Chemistry II*, ed. J. A. McCleverty and T. J. Meyer, Elsevier, Oxford, 2004; P. W. N. M. van Leeuwen, *Homogeneous Catalysis: Understanding the Art*, Kluwer Academic Publishers, Dordrecht, 2004; R. H. Crabtree, *The Organometallic Chemistry of the Transition Metals*, John Wiley & Sons, 2005.
- 2 J. Burt, W. Levason and G. Reid, *Coord. Chem. Rev.*, 2014, **260**, 65.
- 3 R. J. Bowen, M. A. Fernandes, P. B. Hitchcock, M. F. Lappert and M. Lay, *J. Chem. Soc., Dalton Trans.*, 2002, 3253.
- 4 G. Frenking, N. Holzmann, B. Neumüller and K. Dehnicke, *Z. Anorg. Allg. Chem.*, 2010, **636**, 1772.
- 5 H. Braunschweig and K. Gruss, *Z. Naturforsch.*, 2011, **B66**, 55.
- 6 A. Pape, M. Lutz and G. Müller, *Angew. Chem., Int. Ed.*, 1994, **33**, 2281.
- 7 S. Harder, L. Brandsma, J. A. Kanters, A. Duisenberg and J. H. van Lenthe, *J. Organomet. Chem.*, 1991, **420**, 143.
- 8 L. M. Engelhardt, J. M. Harrowfield, M. F. Lappert, I. A. MacKinnon, B. H. Newton, C. L. Raston, B. W. Skelton and A. H. White, *J. Chem. Soc., Chem. Commun.*, 1986, 846.
- 9 C. von Hänisch and B. Rolli, *Z. Anorg. Allg. Chem.*, 2004, **630**, 1987.
- 10 J. C. Thomas and J. C. Peters, *Inorg. Chem.*, 2003, **42**, 5055.
- 11 H. H. Karsch, A. Appelt and G. Müller, *Organometallics*, 1985, **4**, 1624; H. H. Karsch, A. Appelt and G. Müller, *J. Chem. Soc., Chem. Commun.*, 1984, 1415.
- 12 M. Everett, A. Jolleys, W. Levason, D. Pugh and G. Reid, *Chem. Commun.*, 2014, **50**, 5843.
- 13 M. Brookhart, B. Grant and A. F. Volpe Jr., *Organometallics*, 1992, **11**, 3920.
- 14 M. J. D. Champion, J. M. Dyke, W. Levason, M. E. Light, D. Pugh, H. Bhakhoa, L. Rhyman, P. Ramasami and G. Reid, *Inorg. Chem.*, 2015, **54**, 2497.
- 15 S. Strauss, *Chem. Rev.*, 1993, **93**, 927.
- 16 I. Krossing and I. Raabe, *Angew. Chem., Int. Ed.*, 2004, **43**, 2066.
- 17 T. S. Cameron, A. Decken, I. Dionne, M. Fang, I. Krossing and J. Passmore, *Chem. – Eur. J.*, 2002, **8**, 3386.
- 18 I. Krossing and L. van Wullen, *Chem. – Eur. J.*, 2002, **8**, 700.
- 19 I. Raabe, K. Wagner, K. Guttsche, M. Wang, M. Grätzel, G. Santiso-Quiñones and I. Krossing, *Chem. – Eur. J.*, 2009, **15**, 1966.
- 20 R. J. Burt, J. Chatt, W. Hussain and G. J. Leigh, *J. Organomet. Chem.*, 1979, **182**, 203; E. P. Kyba, S. T. Liu and R. L. Harris, *Organometallics*, 1983, **2**, 1877.
- 21 G. Metz, X. Wu and S. O. Smith, *J. Magn. Reson., Ser. A*, 1994, **110**, 219.
- 22 P. Garrou, *Chem. Rev.*, 1981, **81**, 229.
- 23 G. M. Sheldrick, *SHELXS-97, Program for crystal structure solution*, University of Göttingen, Germany, 1997.
- 24 G. M. Sheldrick, *SHELXL-97, Program for crystal structure refinement*, University of Göttingen, Germany, 1997.

