



BRCA1 and BRCA2 genetic testing – pitfalls and recommendations for managing variants of uncertain clinical significance

Journal:	<i>Annals of Oncology</i>
Manuscript ID:	ANNONC-2014-2145.R1
Manuscript Type:	Review
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Eccles, Diana; University of Southampton and University Hospital Southampton Foundation Trust, University of Southampton Clinical Trials Unit</p> <p>Mitchell, Gillian; Peter MacCallum Cancer Centre, Familial Cancer Centre</p> <p>Monteiro, Alvaro; H. Lee Moffitt Cancer Center, Cancer Epidemiology Program</p> <p>Schmutzler, Rita; Centre for familial breast and ovarian cancer, University hospital, Genetics</p> <p>Couch, Fergus; Mayo Clinic, Department of Laboratory Medicine and Pathology</p> <p>Spurdle, Amanda; QIMR Berghofer Medical Research Institute, Molecular Cancer Epidemiology Laboratory</p> <p>Gómez García, Encarna; Maastricht University Medical Center, Clinical Genetics</p>
Keywords:	BRCA, clinical utility, Variants of Uncertain clinical Significance, ENIGMA consortium, classification, Hereditary breast and ovarian cancer
Abstract:	<p>Background</p> <p>Increasing use of BRCA1/2 testing for tailoring cancer treatment and extension of testing to tumour tissue for somatic mutation is moving BRCA1/2 mutation screening from a primarily prevention arena delivered by specialist genetic services into mainstream oncology practice. A considerable number of gene tests will identify rare variants where clinical significance cannot be inferred from sequence information alone. The proportion of Variants of Uncertain clinical Significance (VUS) is likely to grow with lower thresholds for testing and laboratory providers with less experience of BRCA. Most VUS will not be associated with a high risk of cancer but a misinterpreted VUS has the potential to lead to mismanagement of both the patient and their relatives.</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>Design</p> <p>Members of the Clinical Working Group of ENIGMA (Evidence-based Network for the Interpretation of Germline Mutant Alleles) global consortium (www.enigmaconsortium.org) observed wide variation in practices in reporting, disclosure and clinical management of patients with a VUS. Examples from current clinical practice are presented and discussed to illustrate potential pitfalls, explore factors contributing to misinterpretation, and propose approaches to improving clarity.</p> <p>Results and conclusion</p> <p>Clinicians, patients and their relatives would all benefit from an improved level of genetic literacy. Genetic laboratories working with clinical geneticists need to agree on a clinically clear and uniform format for reporting BRCA test results to non-geneticists. An international consortium of experts, collecting and integrating all available lines of evidence and classifying variants according to an internationally recognized system will facilitate reclassification of variants for clinical use.</p>

Peer Review

SCHOLARONE™
Manuscripts

1
2
3 ***BRCA1* and *BRCA2* genetic testing – pitfalls and recommendations for managing**
4 **variants of uncertain clinical significance**
5

6 D. Eccles,^{1*} G. Mitchell,^{2,3*} A.N.A. Monteiro,⁴ R. Schmutzler,⁵ F. J. Couch,⁶ A. S. Spurdle,⁷
7 E.B. Gómez-García,^{8#} on behalf of the ENIGMA Clinical Working Group.⁹
8
9

10 **Authors' affiliations:**
11

- 12
13 1) Faculty of Medicine Academic Unit of Cancer Sciences, Southampton General Hospital,
14 Southampton, UK.
15 2) Familial Cancer Centre, Peter MacCallum Cancer Centre, The University of Melbourne,
16 Melbourne, Australia.
17 3) Hereditary Cancer Program, Dept. of Medical Oncology, University of British Columbia,
18 Vancouver, Canada.
19 4) Cancer Epidemiology Program, H. Lee Moffitt Cancer Center, Tampa, USA.
20 5) Center for Hereditary Breast and Ovarian Cancer, Center for Integrated Oncology (CIO)
21 and Center for Molecular Medicine Cologne (CMMC), University Hospital Cologne,
22 Germany.
23 6) Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, Minnesota,
24 USA.
25 7) Molecular Cancer Epidemiology Laboratory, Genetics and Computational Biology
26 Division, QIMR Berghofer Medical Research Institute, BNE, Herston, QLD, Australia.
27 8) Department of Clinical Genetics, Maastricht University Medical Centre, Maastricht, the
28 Netherlands.
29 9) ENIGMA CWG contributors: see appendix.
30
31
32
33
34

35 * D. Eccles and G. Mitchell are joint first authors.
36
37

38 # Corresponding Author: Dr. Encarna B. Gómez-García
39

40 Dept. Clinical Genetics
41 Maastricht University Medical Centre
42 P. Debyelaan 25
43 6202 AZ Maastricht
44 The Netherlands
45
46 Tel: 31-43-3875855; Fax: 31-43-3875800
47
48 E-mail: Encarna.Gomezgarcia@mumc.nl
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Background

Increasing use of *BRCA1/2* testing for tailoring cancer treatment and extension of testing to tumour tissue for somatic mutation is moving *BRCA1/2* mutation screening from a primarily prevention arena delivered by specialist genetic services into mainstream oncology practice. A considerable number of gene tests will identify rare variants where clinical significance cannot be inferred from sequence information alone. The proportion of Variants of Uncertain clinical Significance (VUS) is likely to grow with lower thresholds for testing and laboratory providers with less experience of *BRCA*. Most VUS will not be associated with a high risk of cancer but a misinterpreted VUS has the potential to lead to mismanagement of both the patient and their relatives.

Design

Members of the Clinical Working Group of ENIGMA (Evidence-based Network for the Interpretation of Germline Mutant Alleles) global consortium (www.enigmaconsortium.org) observed wide variation in practices in reporting, disclosure and clinical management of patients with a VUS. Examples from current clinical practice are presented and discussed to illustrate potential pitfalls, explore factors contributing to misinterpretation, and propose approaches to improving clarity.

Results and conclusion

Clinicians, patients and their relatives would all benefit from an improved level of genetic literacy. Genetic laboratories working with clinical geneticists need to agree on a clinically clear and uniform format for reporting *BRCA* test results to non-geneticists. An international consortium of experts, collecting and integrating all available lines of evidence and classifying variants according to an internationally recognized system will facilitate reclassification of variants for clinical use.

Key words: variants of uncertain significance, VUS, *BRCA*, clinical utility, classification

Introduction

Germline inactivating variants in the tumor suppressor genes *BRCA1* and *BRCA2* confer high life time risks of breast cancer, ovarian cancer and less frequently also other cancers[1]. These pathogenic variants are conventionally called ‘mutations’ or ‘deleterious variants’ in *BRCA* genetic testing parlance and the term pathogenic variant is used here for the sake of precision. Pathogenic variants in either gene confer a high lifetime risk of developing ovarian or (another) primary breast cancer in female carriers but they explain only about 20% of familial breast cancer[2]. The more cancers, the younger the onset and the admixture of ovarian with breast cancer amongst relatives all increase the chance that a familial cluster is due to a *BRCA* gene mutation but nonetheless most familial clusters of breast cancer are not due to an inherited mutation in *BRCA1* or *BRCA2*. In addition to familial clusters *BRCA* mutations account for over 10% of patients with early onset triple negative breast cancer and over 10% of women with non-mucinous ovarian cancer with no family history[3,4]. Female carriers with pathogenic variants can make informed decisions about prophylactic surgery or intensified screening programs. High profile media coverage increases patient expectations from genetic testing[5,6]. The indications for germline genetic testing for *BRCA1* and *BRCA2* are further increasing to include directing cancer chemotherapy, novel targeted treatments and informing choices about the extent of therapeutic surgery[7-9]. Germline genetic testing for *BRCA1* and *BRCA2* for cancer risk prediction and management is routinely delivered by clinical genetics professionals but increasing demand is overwhelming the current delivery model with insufficient capacity amongst trained geneticists and genetic counsellor. Safe integration of genetic counselling and testing for HBOC into mainstream oncology is an ongoing challenge. Challenges exist at many levels but key are genetic literacy and genetic test outcome interpretation and reporting.

Variability between individuals’ genetic code is common within the general population and between individuals of different ethnic background and this intrinsic variability can lead to difficulties in interpreting some types of sequence change (see supplementary figure F1 for more information). Variants of uncertain clinical significance (VUS) represent a particular challenge since the clinical significance cannot be inferred from sequence information alone. Mis-interpretation of VUS can lead to real clinical harms for both patients and families [10-12]. Furthermore terms used in genetic test clinical reports vary (see supplementary table T1). Up to 20% of *BRCA1/2* tests will report genetic variants of uncertain clinical significance (VUS) but in a well characterised ethnic population the proportion may drop to 5% or less[13-15]. The ENIGMA consortium has received over 6,000 submissions of unique VUS, identified in over 13,000 families from over 17 countries[16]. Laboratories following generic reporting rules and with limited *BRCA* specific experience, may report more significance to variants in *BRCA* genes than is supported when multiple lines of evidence are taken into account[17]. The percentage of gene-tests resulting in a VUS is expected to increase when the extent of sequencing increases to include untranslated and deeper intronic regions and when tumour testing is offered. We restrict examples in this paper to the *BRCA* genes tested in blood samples.

Members of the Evidence-based Network for the Interpretation of Germline Mutant Alleles (ENIGMA) consortium’s Clinical Working Group have collated real case scenarios from clinical practice to illustrate the pitfalls that can arise after a VUS test result and suggest some strategies to mitigate this risk. Clinicians who are requesting *BRCA* tests need to consider these issues when using *BRCA* test results in the management of patients and their families.

I. The VUS report: why are VUS difficult to classify?

I-1- *Pitfalls associated with variability in the information content of genetic test reports*

Based on the experience of representatives from 17 countries, it is clear that there is currently no internationally accepted standard for *BRCA* testing reporting (supplementary table T1), and no agreed consistent classification system: some laboratories report variants without interpretation, some use a narrative approach and some use locally developed guidelines or published schemes[18,19]. For some types of *BRCA* gene variants, additional evidence may be essential before a variant can be clearly classified. Different lines of evidence may appear to conflict, so an integrated estimation of probability taking all available evidence into account is essential to reach a final classification for many variants[20,21].

I-1a-Multiple lines of evidence may be required to establish pathogenicity

Software packages are available to help interpret genetic variants, each with strengths and weaknesses (supplementary table T2). Understanding of how the multiple functions of the *BRCA1* and *BRCA2* proteins relate to cancer predisposition is limited and few validated functional assays are available[22]. The interpretation of functional studies is often technically complex and results may not be calibrated against clinical parameters in order to give measures of sensitivity and specificity. A major breakthrough in the field of *BRCA* variant classification has been the development of a multifactorial likelihood classification model[20,23]. It is anticipated that functional data from *BRCA1* and *BRCA2* will soon be incorporated in this model[22,24,25].

The multifactorial likelihood method uses a number of different independent features in order to establish a combined likelihood estimate that a *BRCA* variant has the characteristics of known pathogenic variants[15]. The model currently combines the prior probability of variant pathogenicity (based on evolutionary conservation and amino acid physicochemical properties) with additional estimates of pathogenicity from likelihoods based on clinical information, including variant co-occurrence with a known pathogenic mutation in the same gene, tracking of the variant with cancer affected family members (segregation) and *BRCA* tumour features. The concept underlying multiparametric methods is that empiric probabilities that a person *does* have or *does not* have a pathogenic *BRCA* variant, can be established for each available line of evidence[20]. Multiple lines of evidence can be factored together as they become available in an iterative manner to produce an increasingly accurate probability of pathogenicity estimates for an individual variant. These estimates can be used to classify variants into clinically relevant strata (supplementary table T3)[18].

I-1b- Several classification systems have been proposed and there is no international consensus on which to adopt

A universal system for classification of variants common across all countries would facilitate education amongst new users of genetic tests and minimise the risk of misinterpretation. Unfortunately multiple systems have been proposed and are currently in use with many reporting laboratories not offering a clinically relevant classification. A World Health Organization-funded expert workshop at the International Agency for Research on Cancer (IARC) in 2008 recommended that the score from a multifactorial likelihood model is used to categorize high risk cancer gene variants[26,26] based on the multifactorial likelihood estimates of variant pathogenicity [18]. This is the only published classification system that links clinical recommendations to each class (supplementary table T3). The American College of Medical Genetics (ACMG) are recommending ClinVar as a common repository for genetic variants to help standardise reporting in the USA (supplementary table T4).

I-1c- Databases reporting genetic variants

A number of web-based resources catalogue variants reported in the *BRCA1* and 2 genes (supplementary table T4). For all variant databases, ongoing curation of deposited information, both at the time of deposition and reclassification, is a significant challenge. Some databases are actively curated while others rely heavily on the classification by the submitting laboratory with varying levels of supporting data provided. Also important is the use of standardized HGVS nomenclature (<http://www.hgvs.org/>) allowing unambiguous comparison of all data on the same variant in the database and across the literature. Apart from occasional national consortia, reporting of variants is not mandated so databases cannot usually be used to derive population frequencies of variants to aid classification (e.g. laboratories tend to underreport variants that have been found several times) and many do not catalogue the supporting evidence used to determine pathogenicity. A new attempt to collate BRCA VUS for the purpose of classification is the PROMPT registry (www.promptstudy.org). Patients receiving a BRCA VUS result are informed about the registry directly by the collaborating testing laboratory (including Myriad and Ambry Genetics) and/or their clinician. The patient can submit a range of medical information, genetic test report and tumour pathology reports and opt to participate in a variety of variant-specific research studies.

Case 1. Two families were identified in separate countries with the same missense variant in *BRCA1* designated c.5212G>A, NM_007294.3 (p.Gly1738Arg). A number of women in both families were seeking risk-reducing surgery because of a strong family history of early age of onset of breast cancer in their families. The variant was a Class 3 (uncertain) variant (supplementary table T3) but it occurred at a highly conserved residue and it was recognised that more evidence may successfully confirm it was pathogenic. Predictive testing could then be offered to at risk family members to facilitate their preventive choices. Each family provided insufficient power for an informative segregation analysis but collaboration across centres in two countries with the same variant, allowed a segregation analysis to be completed and the VUS could then be re-classified as clearly pathogenic [27,27]. It took a fortuitous academic collaboration and over 3 years to re-classify this VUS. The ENIGMA consortium[16] provides a mechanism to rapidly link clinical teams looking after families with the same VUS to pool evidence. Segregation analyses will iteratively improve the estimations of the likelihood that each variant is pathogenic or not in a much more timely fashion.

I-2- Pitfalls derived from frequency and population of origin of the VUS:

Rare variants (allele frequency <0.01) are usually not classifiable by an individual laboratory due to paucity of information and lack of statistical power. When the ethnicity of the patient being tested differs from the patient groups where most testing has been done, a sequence variant may have little publically available data and be rare in that laboratory so is more likely to be considered a VUS.

Case 2: A 35 year old woman of African ancestry developed two primary breast tumours presenting at age 25 and 33. Both tumours were ER-, PR-, and Her2-. She had no family history of cancer, but testing was initiated on the basis of her personal cancer history. She was found to have two VUS in *BRCA1*; variant 1, c.5154G>T, NM_007294.3 (p.Trp1718Cys), and variant 2 in the 3' UTR region, c.*36C>G, NM_007294.3. DNA samples from relatives indicated that both variants are likely located on the same allele. Variant 2 is frequent (6-11%) in the African-American population[28,28] and could be

1
2
3 assigned to Class 1 (non-pathogenic). Variant 1 involves amino acid substitution at a highly
4 conserved residue predicted *in silico* to have a functional impact but could not then be
5 classified unambiguously due to insufficient data. The clinical dilemma for this patient thus
6 becomes whether the available information about variant 1 is sufficient to predict a high risk
7 of ovarian cancer and recommend risk reducing bilateral salpingo-oophorectomy.
8

9 A literature search identified two reports of variant 1. The first report was of 2 families from
10 Asturias, Northern Spain[29]. Contact with the authors clarified the following details, some
11 directly applicable for variant interpretation: Asturian proband 1 was diagnosed with bilateral
12 invasive ductal breast cancer at age 28 (grade III, ER-) and 30 (grade II, ER -, PR -, HER2 -);
13 Asturian proband 2 developed breast cancer at age 40 (ER and grade unknown), and probably
14 also ovarian cancer at age 63. The second report of the variant was in an African American
15 patient and family members where the variant was present in 4/4 women with cancer and
16 absent in 3/3 at-risk women without cancer. Pathogenicity was supported by the functional
17 assay showing loss of transcriptional activity[30].
18

19
20 Using multifactorial likelihood analysis[23] combining all currently available evidence: a
21 prior probability of 0.81 based on Align-GVGD prediction[31], and likelihood ratios based
22 on additional information gained about breast tumour pathology (Spurdle et al, in press), the
23 posterior probability of pathogenicity for the variant is 0.98 which places it in class 4. Thus
24 sufficient evidence is now available to estimate a substantially raised ovarian cancer risk and
25 permit a clinically sound recommendation of BSO to reduce ovarian cancer risk.
26
27

28 **II Why is genetic counselling around VUS complex?**

29
30 Although it is expected that most Class 3 variants will represent non-pathogenic variants, it is
31 critical that we improve and accelerate clinical annotation and classification.
32

33 *II-1- Limited genetic literacy amongst clinical professionals*

34 Disclosure of a VUS result can test the skills of even highly experienced genetic counselling
35 teams. There can be considerable variation in experience with molecular genetics and with
36 clinical *BRCA* gene testing leading to diverse clinical management recommendations. This
37 situation is compounded by complex and variable presentation of diagnostic reports from
38 laboratories and limited genetic literacy amongst treating clinicians, and exacerbated further
39 by the patient's perception, prior experiences and beliefs.
40
41

42 Case 3: A *BRCA2* missense variant p.Thr3033Ile (c.9098C>T, NM_000059.3) was identified
43 in a 53 year old woman with ovarian cancer and no family history of breast or ovarian cancer.
44 A Class 3 report was issued as little supporting evidence was available. The managing
45 clinician treated the variant as clinically significant and offered genetic testing to the patient's
46 daughter and sister. The patient's sister did not carry the variant, but the daughter did. The
47 daughter was given the same cancer risk management advice as a pathogenic variant carrier
48 including risk-reducing surgery. Additional information from a *BRCA2* functional assay was
49 subsequently used to reclassify this variant to Class 2 (likely of no clinical significance)[32].
50 Cancer risk estimates for the daughter and sister should have been based on family history
51 alone. Since there was no family history, testing in relatives could not even provide useful
52 additional information about segregation.
53
54

55 This case illustrates the need to ensure improved genetic literacy amongst non-genetics
56 professionals ordering DNA tests, particularly the uncertainty around VUS results, the
57
58
59
60

1
2
3 implications of testing in other family members and the need for a clear pathway for referral
4 to colleagues experienced in the interpretation and follow up of variant results[33].
5
6

7 *II-2- Perceptions of cancer risks associated with VUS*

8
9 A genetic test for a disease like breast cancer that uncovers a rare variant may lead to
10 both patients and clinicians to think, “what is the chance this rare variant has nothing to do
11 with this clinical presentation?” and convince themselves it cannot be coincidental. The
12 decision to undertake preventive surgery is complex and patients are likely to have had
13 personal and family experiences which may have strongly contributed to their risk
14 management decision. Class 3 VUS reports discussed with counselees are too frequently
15 inaccurately perceived typically leading to overestimation of cancer risks, adverse
16 psychological outcomes and more radical medical decisions[34,35]. Furthermore, “almost the
17 same large number of counselees with an unclassified variant decided to have preventive
18 surgery as pathogenic mutation carriers”[36].
19

20 Case 4: A *BRCA1* VUS was identified some years previously in a woman who developed
21 breast cancer at the age of 45 in the setting of a strong family history of early onset breast and
22 ovarian cancer. The family believed the variant must be the answer to the strong family
23 history despite the inconclusive evidence. Several family members came forward for
24 predictive genetic testing which they incorrectly believed was available. Genetic testing was
25 repeated some years later and a clearly pathogenic large exonic deletion in *BRCA1* was then
26 identified on the opposite allele confirming irrefutably that the original VUS could not be
27 pathogenic and demonstrating that new or optimised mutation screening approaches will
28 uncover missed mutations in some families.
29

30 This case illustrates the importance of understanding a “non-informative” or inconclusive test
31 aimed at detecting a genetic predisposition to cancer. It is important to be aware that mutation
32 testing is less than 100% sensitive and a negative *BRCA* test result does not exclude an
33 underlying hereditary cause. *BRCA1* or *BRCA2* gene mutations only account for an estimated
34 20-30% of familial clustering of breast cancer so cancer patients negative for a *BRCA* test but
35 with a strong family history should be referred for specialist assessment and advice about risk
36 management in the cancer genetics clinic.
37

38 The testing clinician has a responsibility to ensure that the person being tested accurately
39 understands the test result. A helpful starting point is to estimate the probability of a
40 deleterious *BRCA* mutation given the family history and tumour characteristic[37]. It is worth
41 bearing in mind that with a low *a priori* probability of finding a pathogenic variant, the most
42 likely outcome from testing would be no pathogenic variant found, the second most likely
43 outcome would be a VUS, and the least likely outcome a pathogenic variant. It is good
44 practice before embarking on a *BRCA* gene test to discuss with the patient the likelihood of
45 these possible outcomes. The test result must be interpreted in the context of the family
46 history and clear guidelines should be agreed for referral of cases to a specialist genetic clinic
47 (supplementary figure F2). Finally, the experience gathered with VUS in *BRCA1/2* is
48 relevant to developing ethical norms and policy issues, including duty to recontact patients,
49 as cancer gene panel, exome and genome sequencing become increasingly commonplace[38].
50

51 **Recommendations**

52
53 We believe some essential elements are necessary for the optimal clinical use and
54 interpretation of VUS in mainstream medical practice.
55

56 a) Variant reporting and classification
57
58
59
60

- An internationally accepted terminology and a clinically relevant classification to report and discuss *BRCA* test results.
- A framework for clinicians, clinical scientists and research groups to work together towards classification of VUS.
- Reporting of sequence variants to a single, transparently and expertly curated database providing clinically relevant classification for each reported *BRCA* variant based on clearly stated lines of evidence and freely available to all providers of genetic testing.

b) Risk communication

- Improve the genetics literacy of medical providers by structured training and integrating genomics into undergraduate, basic and specialist medical training curriculae.
- Access and clear referral guidelines to specialist clinical genetics services for patients with a pathogenic variant, a VUS or a strong family history and no detectable genetic cause.
- A close working relationship between genetic diagnostic laboratories, genetic specialists and cancer clinicians delivering an integrated care pathway for patients and their families.

c) Data review

- A Standard Operating Procedure within reporting laboratories to review variant classification each time new evidence emerges.
- A clearly agreed process for reporting updated classifications (laboratories) and recontacting patients (clinicians) if a VUS becomes classified into clinically actionable or definitively non-pathogenic.

Concluding remarks

VUS identified in *BRCA* genes represent a major clinical challenge. Individuals with significant family history, a pathogenic variant or a VUS should be referred to a genetic specialist service. Patients with no pathogenic variant or a VUS should be managed based on the family history only. Standardised reporting and better genetic literacy must be implemented to safely introduce genetics into mainstream oncology. In conclusion, concerted action between the clinical and research communities is the best approach to optimally managing *BRCA* variants for maximum patient benefit. The international ENIGMA consortium is focussed on improving the interpretation of *BRCA* VUS and incorporates both clinical and research expertise. Through a collaborative approach and a global and unified source of data and variant classification, important advances in this complex field will continue to benefit patients.

Conflict of Interests: the authors have declared no conflict of interests.

References

1. Mavaddat N, Peock S, Frost D et al. Cancer Risks for BRCA1 and BRCA2 Mutation Carriers: Results From Prospective Analysis of EMBRACE. *J Natl Cancer Inst* 2013;105:812-22.
2. Mavaddat N, Antoniou AC, Easton DF et al. Genetic susceptibility to breast cancer. *Mol Oncol* 2010;4:174-91.
3. Risch HA, McLaughlin JR, Cole DE et al. Prevalence and penetrance of germline BRCA1 and BRCA2 mutations in a population series of 649 women with ovarian cancer. *Am J Hum Genet* 2001;68:700-10.
4. Couch FJ, Hart SN, Sharma P et al. Inherited Mutations in 17 Breast Cancer Susceptibility Genes Among a Large Triple-Negative Breast Cancer Cohort Unselected for Family History of Breast Cancer. *J Clin Oncol* 2015;33(4):304-11
5. Evans DGR, Barwell J, Eccles DM et al. The Angelina Jolie effect: how high celebrity profile can have a major impact on provision of cancer related services. *Breast Cancer Research* 2014;16:442-7.
6. James PA, Mitchell G, Bogwitz M et al. The Angelina Jolie effect. *Med J Aust* 2013;199:646.
7. Alsop K, Fereday S, Meldrum C et al. BRCA mutation frequency and patterns of treatment response in BRCA mutation-positive women with ovarian cancer: a report from the Australian Ovarian Cancer Study Group. *J Clin Oncol* 2012;30:2654-63.
8. Tutt A, Robson M, Garber JE et al. Oral poly(ADP-ribose) polymerase inhibitor olaparib in patients with BRCA1 or BRCA2 mutations and advanced breast cancer: a proof-of-concept trial. *Lancet* 2010;376:235-44.
9. Schwartz MD, Lerman C, Brogan B et al. Impact of BRCA1/BRCA2 counseling and testing on newly diagnosed breast cancer patients. *J Clin Oncol* 2004;22:1823-9.
10. Vadaparampil ST, Scherr CL, Cragun D et al. Pre-test genetic counseling services for hereditary breast and ovarian cancer delivered by non-genetics professionals in the state of Florida. *Clin Genet* 2014. DOI:10.1111/cge.12405.
11. Pal T, Cragun D, Lewis C et al. A statewide survey of practitioners to assess knowledge and clinical practices regarding hereditary breast and ovarian cancer. *Genet Test Mol Biomarkers* 2013;17:367-75.
12. Wideroff L, Vadaparampil ST, Greene MH et al. Hereditary breast/ovarian and colorectal cancer genetics knowledge in a national sample of US physicians. *J Med Genet* 2005;42:749-55.
13. Chenevix-Trench G, Healey S, Lakhani S et al. Genetic and histopathologic evaluation of BRCA1 and BRCA2 DNA sequence variants of unknown clinical significance. *Cancer Res* 2006;66:2019-27.
14. Eggington JM, Bowles KR, Moyes K et al. A comprehensive laboratory-based program for classification of variants of uncertain significance in hereditary cancer genes. *Clin Genet* 2014; 86: 229-37.
15. Lindor NM, Guidugli L, Wang X et al. A review of a multifactorial probability-based model for classification of BRCA1 and BRCA2 variants of uncertain significance (VUS). *Hum Mutat* 2012;33:8-21.
16. Spurdle AB, Healey S, Devereau A et al. ENIGMA--evidence-based network for the interpretation of germline mutant alleles: an international initiative to evaluate risk

1
2
3 and clinical significance associated with sequence variation in BRCA1 and BRCA2 genes.
4 Hum Mutat 2012;33:2-7.

5 17. Spurdle AB, Whiley PJ, Thompson B et al. BRCA1 R1699Q variant
6 displaying ambiguous functional abrogation confers intermediate breast and ovarian cancer
7 risk. J Med Genet 2012;49:525-32.

8 18. Plon SE, Eccles DM, Easton D et al. Sequence variant classification and
9 reporting: recommendations for improving the interpretation of cancer susceptibility genetic
10 test results. Hum Mutat 2008;29:1282-91.

11 19. Moghadasi S, Hofland N, Wouts JN et al. Variants of uncertain significance in
12 BRCA1 and BRCA2 assessment of in silico analysis and a proposal for communication in
13 genetic counselling. J Med Genet 2013;50:74-9.

14 20. Goldgar DE, Easton DF, Deffenbaugh AM et al. Integrated evaluation of DNA
15 sequence variants of unknown clinical significance: application to BRCA1 and BRCA2. Am
16 J Hum Genet 2004;75:535-44.

17 21. Walker LC, Whiley PJ, Couch FJ et al. Detection of splicing aberrations
18 caused by BRCA1 and BRCA2 sequence variants encoding missense substitutions:
19 implications for prediction of pathogenicity. Hum Mutat 2010;31:E1484-E1505.

20 22. Millot GA, Carvalho MA, Caputo SM et al. A guide for functional analysis of
21 BRCA1 variants of uncertain significance. Hum Mutat 2012;33:1526-37.

22 23. Goldgar DE, Easton DF, Byrnes GB et al. Genetic evidence and integration of
23 various data sources for classifying uncertain variants into a single model. Hum Mutat
24 2008;29:1265-72.

25 24. Guidugli L, Carreira A, Caputo SM et al. Functional assays for analysis of
26 variants of uncertain significance in BRCA2. Hum Mutat 2014;35:151-64.

27 25. Iversen ES, Jr., Couch FJ, Goldgar DE et al. A computational method to
28 classify variants of uncertain significance using functional assay data with application to
29 BRCA1. Cancer Epidemiol Biomarkers Prev 2011;20:1078-88.

30 26. Tavtigian SV, Greenblatt MS, Goldgar DE et al. Assessing pathogenicity:
31 overview of results from the IARC Unclassified Genetic Variants Working Group. Hum
32 Mutat 2008;29:1261-4.

33 27. Anagnostopoulos T, Pertesi M, Konstantopoulou I et al. G1738R is a BRCA1
34 founder mutation in Greek breast/ovarian cancer patients: evaluation of its pathogenicity and
35 inferences on its genealogical history. Breast Cancer Res Treat 2008;110:377-85.

36 28. Newman B, Mu H, Butler LM et al. Frequency of breast cancer attributable to
37 BRCA1 in a population-based series of American women. JAMA 1998;279:915-21.

38 29. Blay P, Santamaria I, Pitiot AS et al. Mutational analysis of BRCA1 and
39 BRCA2 in hereditary breast and ovarian cancer families from Asturias (Northern Spain).
40 BMC Cancer 2013;13:243.

41 30. Mirkovic N, Marti-Renom MA, Weber BL et al. Structure-based assessment
42 of missense mutations in human BRCA1: implications for breast and ovarian cancer
43 predisposition. Cancer Res 2004;64:3790-7.

44 31. Tavtigian SV, Byrnes GB, Goldgar DE et al. Classification of rare missense
45 substitutions, using risk surfaces, with genetic- and molecular-epidemiology applications.
46 Hum Mutat 2008;29:1342-54.

47 32. Guidugli L, Pankratz VS, Singh N et al. A classification model for BRCA2
48 DNA binding domain missense variants based on homology-directed repair activity. Cancer
49 Res 2013;73:265-75.

50 33. Lindor NM, Goldgar DE, Tavtigian SV et al. BRCA1/2 sequence variants of
51 uncertain significance: a primer for providers to assist in discussions and in medical
52 management. Oncologist 2013;18:518-24.
53
54
55
56
57
58
59
60

1
2
3 34. Petrucelli N, Lazebnik N, Huelsman KM et al. Clinical interpretation and
4 recommendations for patients with a variant of uncertain significance in BRCA1 or BRCA2:
5 a survey of genetic counseling practice. *Genet Test* 2002;6:107-13.

6 35. Vos J, Gomez-Garcia E, Oosterwijk JC et al. Opening the psychological black
7 box in genetic counseling. The psychological impact of DNA testing is predicted by the
8 counselees' perception, the medical impact by the pathogenic or uninformative BRCA1/2-
9 result. *Psychooncology* 2012;21:29-42.

10 36. Vos J, Otten W, van AC et al. The counselees' view of an unclassified variant
11 in BRCA1/2: recall, interpretation, and impact on life. *Psychooncology* 2008; 17: 822-30.

12 37. Gomez-Garcia EB, Ambergen T, Blok MJ et al. Patients with an unclassified
13 genetic variant in the BRCA1 or BRCA2 genes show different clinical features from those
14 with a mutation. *J Clin Oncol* 2005;23:2185-90.

15 38. Cheon JY, Mozersky J, Cook-Deegan R. Variants of uncertain significance in
16 BRCA: a harbinger of ethical and policy issues to come? *Genome Med* 2014; 6: 121.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Supplementary material:

4
5 Supplementary figure (F1): Genetic variability information

6
7 Supplementary figure (F2): Suggested clinical pathway for patients when genetic testing is
8 initiated by the oncology clinic.
9

10
11 Supplementary table (T1). Types of sequence variants reported in *BRCA* mutation detection
12 tests according to risk relevant for clinical management

13
14 Supplementary table (T2). Examples of software tools available for clinical variant evaluation
15 online

16
17 Supplementary table (T3). 5-tier classification *BRCA1/2* VUS

18
19 Supplementary table (T4). Commonly used online database resources that provide some
20 interpretation of *BRCA* sequence variants
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Appendix:**Name and affiliations of the ENIGMA CWG contributors:** (excluding those named in the authors' list)

- R. Driessen: Familial Cancer Centre, Peter MacCallum Cancer Centre, The University of Melbourne, Melbourne, Australia.
- N.M. Lindor: Department of Health Sciences Research, Mayo Clinic, Scottsdale, AZ, USA.
- M.J. Blok: Department of Clinical Genetics, Maastricht University Medical Centre, Maastricht, the Netherlands.
- P. Moller: Research Group Inherited Cancer, Department of Medical Genetics, University Hospital, Oslo, Norway.
- M. de la Hoya: Molecular Oncology Laboratory, Hospital Clínico San Carlos (IdISSC), Madrid, Spain.
- T. Pal: Cancer Epidemiology Program, H. Lee Moffitt Cancer Center, Tampa, USA.
- S. Domchek and K. Nathanson: Basser Research Center and Abramson Cancer Center University of Pennsylvania Philadelphia, PA, USA.
- C. Van Asperen: Department of Clinical Genetics, Leiden University Medical Center, Leiden, the Netherlands
- O. Diez: Oncogenetics Group, University Hospital Vall d'Hebron; Vall d'Hebron Institute of Oncology (VHIO), Barcelona, Spain.
- K. Rheim: Center for Hereditary Breast and Ovarian Cancer, Center for Integrated Oncology (CIO) and Center for Molecular Medicine Cologne (CMMC), University Hospital Cologne, Germany.
- D. Stoppa-Lyonnet: Genetis Service, Curie Institute, University of Paris Descartes, Paris, France
- M. Parsons: Molecular Cancer Epidemiology Laboratory, Genetics and Computational Biology Division, QIMR Berghofer Medical Research Institute, BNE, Herston, QLD, Australia.
- D. Goldgar: Department of Dermatology, Huntsman Cancer Institute, University of Utah School of Medicine; Salt Lake City, Utah, USA.

1
2
3
4
5
6
7 ***BRCA1* and *BRCA2* genetic testing – pitfalls and recommendations for managing**
8 **variants of uncertain clinical significance**

9
10 D. Eccles,^{1*} G. Mitchell,^{2,3*} A.N.A. Monteiro,⁴ R. Schmutzler,⁵ F. J. Couch,⁶ A. S. Spurdle,⁷
11 E.B. Gómez-García,^{8#} on behalf of the ENIGMA Clinical Working Group.⁹

12
13 **Authors' affiliations:**

- 14
15 1) Faculty of Medicine Academic Unit of Cancer Sciences, Southampton General Hospital,
16 Southampton, UK.
17 2) Familial Cancer Centre, Peter MacCallum Cancer Centre, The University of Melbourne,
18 Melbourne, Australia.
19 3) Hereditary Cancer Program, Dept. of Medical Oncology, University of British Columbia,
20 Vancouver, Canada.
21 4) Cancer Epidemiology Program, H. Lee Moffitt Cancer Center, Tampa, USA.
22 5) Center for Hereditary Breast and Ovarian Cancer, Center for Integrated Oncology (CIO)
23 and Center for Molecular Medicine Cologne (CMMC), University Hospital Cologne,
24 Germany.
25 6) Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, Minnesota,
26 USA.
27 7) Molecular Cancer Epidemiology Laboratory, Genetics and Computational Biology
28 Division, QIMR Berghofer Medical Research Institute, BNE, Herston, QLD, Australia.
29 8) Department of Clinical Genetics, Maastricht University Medical Centre, Maastricht, the
30 Netherlands.
31 9) ENIGMA CWG contributors: see appendix.
32
33

34
35 * D. Eccles and G. Mitchell are joint first authors.

36
37 # Corresponding Author: Dr. Encarna B. Gómez-García
38 Dept. Clinical Genetics
39 Maastricht University Medical Centre
40 P. Debyelaan 25
41 6202 AZ Maastricht
42 The Netherlands
43
44 Tel: 31-43-3875855; Fax: 31-43-3875800
45
46 E-mail: Encarna.Gomezgarcia@mumc.nl
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Background

Increasing use of *BRCA1/2* testing for tailoring cancer treatment and extension of testing to tumour tissue for somatic mutation is moving *BRCA1/2* mutation screening from a primarily prevention arena delivered by specialist genetic services into mainstream oncology practice. A considerable number of gene tests will identify rare variants where clinical significance cannot be inferred from sequence information alone. The proportion of Variants of Uncertain clinical Significance (VUS) is likely to grow with lower thresholds for testing and laboratory providers with less experience of BRCA. Most VUS will not be associated with a high risk of cancer but a misinterpreted VUS has the potential to lead to mismanagement of both the patient and their relatives.

Design

Members of the Clinical Working Group of ENIGMA (Evidence-based Network for the Interpretation of Germline Mutant Alleles) global consortium (www.enigmaconsortium.org) observed wide variation in practices in reporting, disclosure and clinical management of patients with a VUS. Examples from current clinical practice are presented and discussed to illustrate potential pitfalls, explore factors contributing to misinterpretation, and propose approaches to improving clarity.

Results and conclusion

Clinicians, patients and their relatives would all benefit from an improved level of genetic literacy. Genetic laboratories working with clinical geneticists need to agree on a clinically clear and uniform format for reporting *BRCA* test results to non-geneticists. An international consortium of experts, collecting and integrating all available lines of evidence and classifying variants according to an internationally recognized system will facilitate reclassification of variants for clinical use.

Key words: variants of uncertain significance, VUS, *BRCA*, clinical utility, classification

Introduction

Germline inactivating variants in the tumor suppressor genes *BRCA1* and *BRCA2* confer high life time risks of breast cancer, ovarian cancer and less frequently also other cancers[1][14]. These pathogenic variants are conventionally called ‘mutations’ or ‘deleterious variants’ in *BRCA* genetic testing parlance and the term pathogenic variant is used here for the sake of precision. Pathogenic variants in either gene confer a high lifetime risk of developing ovarian or (another) primary breast cancer in female carriers but they explain only about 20% of familial breast cancer[2]. The more cancers, the younger the onset and the admixture of ovarian with breast cancer amongst relatives all increase the chance that a familial cluster is due to a *BRCA* gene mutation but nonetheless most familial clusters of breast cancer are not due to an inherited mutation in *BRCA1* or *BRCA2*, can be found in about 30% of multiple case breast cancer families, but are also In addition to familial clusters the underlying cause in *BRCA* mutations account for over 10% of patients with early onset triple negative breast cancer and over 10% of women with non-mucinous ovarian cancer with no family history[3,4][2,3]. Germline genetic testing for *BRCA1* and *BRCA2* for cancer risk prediction and management is delivered by clinical genetics professionals. Female carriers with pathogenic variants who are still free of disease can make informed decisions about prophylactic surgery or intensified screening programs. High profile media coverage increases patient expectations from genetic testing[5,6][4,5]. The indications for germline genetic testing for *BRCA1* and *BRCA2* are further increasing to include directing cancer chemotherapy, novel targeted treatments and informing choices about the extent of therapeutic surgery[7-9][6-8]. Germline genetic testing for *BRCA1* and *BRCA2* for cancer risk prediction and management is routinely delivered by clinical genetics professionals but increasing demand is overwhelming the current delivery model with insufficient capacity amongst trained geneticists and genetic counsellor. Safe i-There are insufficient trained geneticists and genetic counsellors to meet the, therefore integration of genetic counselling with testin and testingg for HBOC into mainstream oncology is an ongoing challenge. Challenges exist at many levels but key are genetic literacy and genetic test outcome interpretation and reporting.

Variability between individuals’ genetic code is common within the general population and between individuals of different ethnic background and this intrinsic variability can lead to difficulties in interpreting certain types of gsequence change (see supplementary figure 1 for more information). Currently, terms used in genetic test clinical reports vary (Supplementary table 1). Variants of uncertain clinical significance (VUS) including predicted missense alterations, in-frame deletions or insertions, and intronic variants outside of conserved splice site motifs, represent a particular challenge since the clinical significance cannot be inferred from sequence information alone. Mis-interpretation of VUS can lead to real clinical harms for both patients and families [10-12][9-11]. Furthermore terms used in genetic test clinical reports vary (see supplementary table 1).

Approximately 5-15Up to 20.% of *BRCA1/2* tests will report genetic variants of uncertain clinical significance (VUS) but in a well characterised ethnic population the proportion may drop to 5% or less[13,14][15][12]. The ENIGMA consortium has received over 6,000 submissions of unique VUS, identified in over 13,000 families from over 17 countries[16][13]. Laboratories following generic reporting rules and with limited *BRCA* specific experience, may report more significance to variants in *BRCA* genes than is supported when multiple lines of evidence are taken into account[17][14]. The percentage of gene-tests resulting in a VUS is expected to increase when the extent of sequencing increases to include untranslated and deeper intronic regions and when tumour testing is offered. We

Field Code Changed

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Field Code Changed

Field Code Changed

Field Code Changed

Formatted: Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Field Code Changed

Field Code Changed

Field Code Changed

~~restrict examples in this paper to the *BRCA* genes tested in blood samples but an additional challenge arises when genetic tests are sequencing lower penetrance genes on targeted gene panels. In lower penetrance genes, even unequivocally protein damaging variants have very different clinical implications compared to highly penetrant deleterious *BRCA* gene mutations.~~

Members of the Evidence-based Network for the Interpretation of Germline Mutant Alleles (ENIGMA) consortium's Clinical Working Group have collated real case scenarios from clinical practice to illustrate the pitfalls that can arise after a VUS test result and suggest some strategies to mitigate this risk. Clinicians who are requesting *BRCA* tests need to consider these issues when using *BRCA* test results in the management of patients and their families.

I. The VUS report: why are VUS difficult to classify?

I-1- Pitfalls associated with variability in the information content of genetic test reports

Based on the experience of representatives from 17 countries, it is clear that there is currently no internationally accepted standard for *BRCA* testing reporting (supplementary table 1), and no agreed consistent classification system: some laboratories report variants without interpretation, some use a narrative approach and some use locally developed guidelines or published schemes^{[18,19][15,16]}. For some types of *BRCA* gene variants, additional evidence may be essential before a variant can be clearly classified. Different lines of evidence may appear to conflict, so an integrated estimation of probability taking all available evidence into account is essential to reach a final classification for many variants^{[20,21][17,18]}.

Field Code Changed

Field Code Changed

I-1a-Multiple lines of evidence may be required to establish pathogenicity

Software packages are available to help interpret genetic variants, each with strengths and weaknesses (supplementary table 2). Understanding of how the multiple functions of the BRCA1 and BRCA2 proteins relate to cancer predisposition is limited and few validated functional assays are available^{[22][19]}. The interpretation of functional studies is often technically complex and results may not be calibrated against clinical parameters in order to give measures of sensitivity and specificity. A major breakthrough in the field of *BRCA* variant classification has been the development of a multifactorial likelihood classification model^{[20,23][17,20]}. It is anticipated that functional data from BRCA1 and BRCA2 will soon be incorporated in this model^{[22,24,25][19,21,22]}.

Field Code Changed

Field Code Changed

Field Code Changed

The multifactorial likelihood method uses a number of different independent features in order to establish a combined likelihood estimate that a *BRCA* variant has the characteristics of known pathogenic variants^{[15][12]}. The model currently combines the prior probability of variant pathogenicity (based on evolutionary conservation and amino acid physicochemical properties) with additional estimates of pathogenicity from likelihoods based on clinical information, including variant co-occurrence with a known pathogenic mutation in the same gene, tracking of the variant with cancer affected family members (segregation) and *BRCA* tumour features. The concept underlying multiparametric methods is that empiric probabilities that a person *does* have or *does not* have a pathogenic *BRCA* variant, can be established for each available line of evidence^{[20][17]}. Multiple lines of evidence can be factored together as they become available in an iterative manner to produce an increasingly accurate probability of pathogenicity estimates for an individual variant.

Field Code Changed

Field Code Changed

These estimates can be used to classify variants into clinically relevant strata (supplementary table 3)^{[18][H5]}.

I-1b- Several classification systems have been proposed and there is no international consensus on which to adopt

A universal system for classification of variants common across all countries would facilitate education amongst new users of genetic tests and minimise the risk of misinterpretation. Unfortunately multiple systems have been proposed and are currently in use with many reporting laboratories not offering a clinically relevant classification. A World Health Organization-funded expert workshop at the International Agency for Research on Cancer (IARC) in 2008 recommended that the score from a multifactorial likelihood model is used to categorize high risk cancer gene variants^{[26,26][23,23]} based on the multifactorial likelihood estimates of variant pathogenicity ^{[18][H5]}. This is the only published classification system that links clinical recommendations to each class (supplementary table 3). The American College of Medical Genetics (ACMG) are recommending ClinVar as a common repository for genetic variants to help standardise reporting in the USA (Supplementary table 4).

I-1c- Databases reporting genetic variants

A number of web-based resources catalogue variants reported in the *BRCA1* and 2 genes (supplementary table 4). For all variant databases, ongoing curation of deposited information, both at the time of deposition and reclassification, is a significant challenge. Some databases are actively curated while others rely heavily on the classification by the submitting laboratory with varying levels of supporting data provided. Also important is the use of standardized HGVS nomenclature (<http://www.hgvs.org/>) allowing unambiguous comparison of all data on the same variant in the database and across the literature. Apart from occasional national consortia, reporting of variants is not mandated so databases cannot usually be used to derive population frequencies of variants to aid classification (e.g. laboratories tend to underreport variants that have been found several times) and many do not catalogue the supporting evidence used to determine pathogenicity. [A new attempt to collate BRCA VUS for the purpose of classification is the PROMPT registry \(www.promptstudy.org\). Patients receiving a BRCA VUS result are informed about the registry directly by the collaborating testing laboratory \(including Myriad and Ambry Genetics\) and/or their clinician. The patient can submit a range of medical information, genetic test report and tumour pathology reports and opt to participate in a variety of variant-specific research studies.](http://www.promptstudy.org)

[Case 1.](#) Two families were identified in separate countries with the same missense variant in *BRCA1* designated c.5212G>A, NM_007294.3 (p.Gly1738Arg). A number of women in both families were seeking risk-reducing surgery because of a strong family history of early age of onset of breast cancer in their families. The variant was a Class 3 (uncertain) variant (supplementary table 3) but it occurred at a highly conserved residue and it was recognised that more evidence may successfully confirm it was pathogenic. Predictive testing could then be offered to at risk family members to facilitate their preventive choices. Each family provided insufficient power for an informative segregation analysis but collaboration across centres in two countries with the same variant, allowed a segregation analysis to be completed and the VUS could then be re-classified as clearly pathogenic ^{[27,27][24,24]}. It took a fortuitous academic collaboration and over 3 years to re-classify this VUS. The ENIGMA consortium^{[16][H3]} provides a mechanism to rapidly link clinical teams looking after families with the same VUS to pool evidence. Segregation analyses will iteratively

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

improve the estimations of the likelihood that each variant is pathogenic or not in a much more timely fashion.

I-2- Pitfalls derived from frequency and population of origin of the VUS:

Rare variants (allele frequency <0.01) are usually not classifiable by an individual laboratory due to paucity of information and lack of statistical power. When the ethnicity of the patient being tested differs from the patient groups where most testing has been done, a sequence variant may have little publically available data and be rare in that laboratory so is [more](#) likely to be considered a VUS.

Case 2: A 35 year old woman of African ancestry developed two primary breast tumours presenting at age 25 and 33. Both tumours were ER-, PR-, and Her2-. She had no family history of cancer, but testing was initiated on the basis of her personal cancer history. She was found to have two VUS in *BRCA1*; variant 1, c.5154G>T, NM_007294.3 (p.Trp1718Cys), and variant 2 in the 3' UTR region, c.*36C>G, NM_007294.3. DNA samples from relatives indicated that both variants are likely located on the same allele. Variant 2 is frequent (6-11%) in the African-American population^{[28,28][25,25]} and could be assigned to Class 1 (non-pathogenic). Variant 1 involves amino acid substitution at a highly conserved residue predicted *in silico* to have a functional impact but could not then be classified unambiguously due to insufficient data. The clinical dilemma for this patient thus becomes whether the available information about variant 1 is sufficient to predict a high risk of ovarian cancer and recommend risk reducing bilateral salpingo-oophorectomy.

A literature search identified two reports of variant 1. The first report was of 2 families from Asturias, Northern Spain^{[29][26]}. Contact with the authors clarified the following details, some directly applicable for variant interpretation: Asturian proband 1 was diagnosed with bilateral invasive ductal breast cancer at age 28 (grade III, ER-) and 30 (grade II, ER -, PR -, HER2 -); Asturian proband 2 developed breast cancer at age 40 (ER and grade unknown), and probably also ovarian cancer at age 63. The second report of the variant was in an African American patient and family members where the variant was present in 4/4 women with cancer and absent in 3/3 at-risk women without cancer. Pathogenicity was supported by the functional assay showing loss of transcriptional activity^{[30][27]}.

Using multifactorial likelihood analysis^{[23][20]} combining all currently available evidence: a prior probability of 0.81 based on Align-GVGD prediction^{[31][28]}, and likelihood ratios based on additional information gained about breast tumour pathology (Spurdle et al, in press), the posterior probability of pathogenicity for the variant is 0.98 which places it in class 4. Thus sufficient evidence is now available to estimate a substantially raised ovarian cancer risk and permit a clinically sound recommendation of BSO to reduce ovarian cancer risk.

II Why is genetic counselling around VUS complex?

Although it is expected that most Class 3 variants will represent non-pathogenic variants, it is critical that we improve and accelerate clinical annotation and classification.

II-1- Limited genetic literacy amongst clinical professionals

Disclosure of a VUS result can test the skills of even highly experienced genetic counselling teams. There can be considerable variation in experience with molecular genetics and with [clinical](#) *BRCA* gene testing leading to diverse clinical management recommendations. This

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

situation is compounded by complex and variable presentation of diagnostic reports from laboratories and limited genetic literacy amongst treating clinicians, and exacerbated further by the patient's perception, prior experiences and beliefs.

Case 3: A *BRCA2* missense variant p.Thr3033Ile (c.9098C>T, NM_000059.3) was identified in a 53 year old woman with ovarian cancer and no family history of breast or ovarian cancer. A Class 3 report was issued as little supporting evidence was available. The managing clinician treated the variant as clinically significant and offered genetic testing to the patient's daughter and sister. The patient's sister did not carry the variant, but the daughter did. The daughter was given the same cancer risk management advice as a pathogenic variant carrier including risk-reducing surgery. Additional information from a *BRCA2* functional assay was subsequently used to reclassify this variant to Class 2 (likely of no clinical significance)^{[32][29]}. Cancer risk estimates for the daughter and sister should have been based on family history alone. Since there was no family history, testing in relatives could not even provide useful additional information about segregation.

This case illustrates the need to ensure improved genetic literacy amongst non-genetics professionals ordering DNA tests, particularly the uncertainty around VUS results, the implications of testing in other family members and the need for a clear pathway for referral to colleagues experienced in the interpretation and follow up of variant results^{[33][30]}.

II-2- Perceptions of cancer risks associated with VUS

A genetic test for a disease like breast cancer that uncovers a rare variant may lead to both patients and clinicians to think, "what is the chance this rare variant has nothing to do with this clinical presentation?" and convince themselves it cannot be coincidental. The decision to undertake preventive surgery is complex and patients are likely to have had personal and family experiences which may have strongly contributed to their risk management decision. Class 3 VUS reports discussed with counselees are too frequently inaccurately perceived typically leading to overestimation of cancer risks, adverse psychological outcomes and more radical medical decisions^{[34,35][31,32]}. Furthermore, "almost the same large number of counselees with an unclassified variant decided to have preventive surgery as pathogenic mutation carriers"^{[36][33]}.

Case 4: A *BRCA1* VUS was identified some years previously in a woman who developed breast cancer at the age of 45 in the setting of a strong family history of early onset breast and ovarian cancer. The family believed the variant must be the answer to the strong family history despite the inconclusive evidence. Several family members came forward for predictive genetic testing which they incorrectly believed was available. Genetic testing was repeated some years later and a clearly pathogenic large exonic deletion in *BRCA1* was then identified on the opposite allele confirming irrefutably that the original VUS could not be pathogenic and demonstrating that new or optimised mutation screening approaches will uncover missed mutations in some families.

This case illustrates the importance of understanding a "non-informative" or inconclusive test aimed at detecting a genetic predisposition to cancer. It is important to be aware that mutation testing is less than 100% sensitive and a negative *BRCA* test result does not exclude an underlying hereditary cause. *BRCA1* or *BRCA2* gene mutations only account for an estimated 20-30% of familial clustering of breast cancer so cancer patients negative for a *BRCA* test but with a strong family history should be referred for specialist assessment and advice about risk management in the cancer genetics clinic.

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

1
2
3
4
5
6
7 The testing clinician has a responsibility to ensure that the person being tested accurately
8 understands the test result. A helpful starting point is to estimate the probability of a
9 deleterious *BRCA* mutation given the family history and tumour characteristic[37][34]. It is
10 worth bearing in mind that with a low *a priori* probability of finding a pathogenic variant, the
11 most likely outcome from testing would be no pathogenic variant found, the second most
12 likely outcome would be a VUS, and the least likely outcome a pathogenic variant. It is good
13 practice before embarking on a *BRCA* gene test to discuss with the patient the likelihood of
14 these possible outcomes. The test result must be interpreted in the context of the family
15 history and clear guidelines should be agreed for referral of cases to a specialist genetic clinic
16 (supplementary figure +F2). Finally, the experience gathered with VUS in *BRCA1/2* is
17 relevant to developing ethical norms and policy issues, including duty to recontact patients,
18 as cancer gene panel, exome and genome sequencing become increasingly commonplace[38].

Field Code Changed

19 **Recommendations**

20 We believe some essential elements are necessary for the optimal clinical use and
21 interpretation of VUS in mainstream medical practice.

22 a) Variant reporting and classification

- 23 • An internationally accepted terminology and a clinically relevant classification to
24 report and discuss *BRCA* test results.
- 25 • A framework for clinicians, clinical scientists and research groups to work together
26 towards classification of VUS.
- 27 • Reporting of sequence variants to a single, transparently and expertly curated database
28 providing clinically relevant classification for each reported *BRCA* variant based on
29 clearly stated lines of evidence and freely available to all providers of genetic testing.

31 b) Risk communication

- 32 • Improve the genetics literacy of medical providers by structured training and
33 integrating genomics into undergraduate, basic and specialist medical training
34 curriculae.
- 35 • Access and clear referral guidelines to specialist clinical genetics services for patients
36 with a pathogenic variant, a VUS or a strong family history and no detectable genetic
37 cause.
- 38 • A close working relationship between genetic diagnostic laboratories, genetic
39 specialists and cancer clinicians delivering an integrated care pathway for patients and
40 their families.

41 c) Data review

- 42 • A Standard Operating Procedure within reporting laboratories to review variant
43 classification each time new evidence emerges.
- 44 • A clearly agreed process for reporting updated classifications (laboratories) and
45 recontacting patients (clinicians) if a VUS becomes classified into clinically
46 actionable or definitively non-pathogenic.

47 **Concluding remarks**

48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 VUS identified in *BRCA* genes represent a major clinical challenge. Individuals with
8 significant family history, a pathogenic variant or a VUS should be referred to a genetic
9 specialist service. Patients with no pathogenic variant or a VUS should be managed based on
10 the family history only. Standardised reporting and better genetic literacy must be
11 implemented to safely introduce genetics into mainstream oncology. In conclusion, concerted
12 action between the clinical and research communities is the best approach to optimally
13 managing *BRCA* variants for maximum patient benefit. The international ENIGMA
14 consortium is focussed on improving the interpretation of *BRCA* VUS and incorporates both
15 clinical and research expertise. Through a collaborative approach and a global and unified
16 source of data and variant classification, important advances in this complex field will
17 continue to benefit patients.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Conflict of Interests: the authors have declared no conflict of interests.

References

1. Mavaddat N, Peock S, Frost D et al. Cancer Risks for BRCA1 and BRCA2 Mutation Carriers: Results From Prospective Analysis of EMBRACE. *J Natl Cancer Inst* 2013; 105: 812-22.
2. Evans DG, Howell A, Ward D et al. Prevalence of BRCA1 and BRCA2 mutations in triple negative breast cancer. *J Med Genet* 2011; 48: 520-2.
3. Risch HA, McLaughlin JR, Cole DE et al. Prevalence and penetrance of germline BRCA1 and BRCA2 mutations in a population series of 649 women with ovarian cancer. *Am J Hum Genet* 2001; 68: 700-10.
4. Alsop K, Fereday S, Meldrum C et al. BRCA mutation frequency and patterns of treatment response in BRCA mutation positive women with ovarian cancer: a report from the Australian Ovarian Cancer Study Group. *J Clin Oncol* 2012; 30: 2654-63.
5. Tutt A, Robson M, Garber JE et al. Oral poly(ADP-ribose) polymerase inhibitor olaparib in patients with BRCA1 or BRCA2 mutations and advanced breast cancer: a proof-of-concept trial. *Lancet* 2010; 376: 235-44.
6. Schwartz MD, Lerman C, Brogan B et al. Impact of BRCA1/BRCA2 counseling and testing on newly diagnosed breast cancer patients. *J Clin Oncol* 2004; 22: 1823-9.
7. Evans DGR, Barwell J, Eccles DM et al. The Angelina Jolie effect: how high celebrity profile can have a major impact on provision of cancer related services. *Breast Cancer Research* 2014; 16: 442-7.
8. James PA, Mitchell G, Bogwitz M et al. The Angelina Jolie effect. *Med J Aust* 2013; 199: 646.
9. Vadaparampil ST, Scherr CL, Cragun D et al. Pre-test genetic counseling services for hereditary breast and ovarian cancer delivered by non-genetics professionals in the state of Florida. *Clin Genet* 2014.
10. Pal T, Cragun D, Lewis C et al. A statewide survey of practitioners to assess knowledge and clinical practices regarding hereditary breast and ovarian cancer. *Genet Test Mol Biomarkers* 2013; 17: 367-75.
11. Wideroff L, Vadaparampil ST, Greene MH et al. Hereditary breast/ovarian and colorectal cancer genetics knowledge in a national sample of US physicians. *J Med Genet* 2005; 42: 749-55.
12. Lindor NM, Guidugli L, Wang X et al. A review of a multifactorial probability-based model for classification of BRCA1 and BRCA2 variants of uncertain significance (VUS). *Hum Mutat* 2012; 33: 8-21.
13. Spurdle AB, Healey S, Devereau A et al. ENIGMA—evidence-based network for the interpretation of germline mutant alleles: an international initiative to evaluate risk and clinical significance associated with sequence variation in BRCA1 and BRCA2 genes. *Hum Mutat* 2012; 33: 2-7.
14. Spurdle AB, Whiley PJ, Thompson B et al. BRCA1 R1699Q variant displaying ambiguous functional abrogation confers intermediate breast and ovarian cancer risk. *J Med Genet* 2012; 49: 525-32.
15. Plon SE, Eccles DM, Easton D et al. Sequence variant classification and reporting: recommendations for improving the interpretation of cancer susceptibility genetic test results. *Hum Mutat* 2008; 29: 1282-91.
16. Moghadasi S, Hofland N, Wouts JN et al. Variants of uncertain significance in BRCA1 and BRCA2 assessment of in silico analysis and a proposal for communication in genetic counselling. *J Med Genet* 2013; 50: 74-9.

Formatted: German (Germany)

Formatted: French (France)

Formatted: French (France)

Formatted: French (France)

- 1
2
3
4
5
6
7 17. — Goldgar DE, Easton DF, Deffenbaugh AM et al. Integrated evaluation of DNA
8 sequence variants of unknown clinical significance: application to BRCA1 and BRCA2. *Am*
9 *J Hum Genet* 2004; 75: 535-44.
- 10 18. — Walker LC, Whaley PJ, Couch FJ et al. Detection of splicing aberrations caused by
11 BRCA1 and BRCA2 sequence variants encoding missense substitutions: implications for
12 prediction of pathogenicity. *Hum Mutat* 2010; 31: E1484-E1505.
- 13 19. — Millot GA, Carvalho MA, Caputo SM et al. A guide for functional analysis of
14 BRCA1 variants of uncertain significance. *Hum Mutat* 2012; 33: 1526-37.
- 15 20. — Goldgar DE, Easton DF, Byrnes GB et al. Genetic evidence and integration of various
16 data sources for classifying uncertain variants into a single model. *Hum Mutat* 2008; 29:
17 1265-72.
- 18 21. — Guidugli L, Carreira A, Caputo SM et al. Functional assays for analysis of variants of
19 uncertain significance in BRCA2. *Hum Mutat* 2014; 35: 151-64.
- 20 22. — Iversen ES, Jr., Couch FJ, Goldgar DE et al. A computational method to classify
21 variants of uncertain significance using functional assay data with application to BRCA1.
22 *Cancer Epidemiol Biomarkers Prev* 2011; 20: 1078-88.
- 23 23. — Tavtigian SV, Greenblatt MS, Goldgar DE et al. Assessing pathogenicity: overview of
24 results from the IARC Unclassified Genetic Variants Working Group. *Hum Mutat* 2008; 29:
25 1261-4.
- 26 24. — Anagnostopoulos T, Pertesi M, Konstantopoulou I et al. G1738R is a BRCA1 founder
27 mutation in Greek breast/ovarian cancer patients: evaluation of its pathogenicity and
28 inferences on its genealogical history. *Breast Cancer Res Treat* 2008; 110: 377-85.
- 29 25. — Newman B, Mu H, Butler LM et al. Frequency of breast cancer attributable to
30 BRCA1 in a population based series of American women. *JAMA* 1998; 279: 915-21.
- 31 26. — Blay P, Santamaria I, Pitiot AS et al. Mutational analysis of BRCA1 and BRCA2 in
32 hereditary breast and ovarian cancer families from Asturias (Northern Spain). *BMC Cancer*
33 2013; 13: 243.
- 34 27. — Mirkovic N, Marti-Renom MA, Weber BL et al. Structure based assessment of
35 missense mutations in human BRCA1: implications for breast and ovarian cancer
36 predisposition. *Cancer Res* 2004; 64: 3790-7.
- 37 28. — Tavtigian SV, Byrnes GB, Goldgar DE et al. Classification of rare missense
38 substitutions, using risk surfaces, with genetic and molecular epidemiology applications.
39 *Hum Mutat* 2008; 29: 1342-54.
- 40 29. — Guidugli L, Pankratz VS, Singh N et al. A classification model for BRCA2 DNA
41 binding domain missense variants based on homology directed repair activity. *Cancer Res*
42 2013; 73: 265-75.
- 43 30. — Petrucelli N, Lazebnik N, Huelsman KM et al. Clinical interpretation and
44 recommendations for patients with a variant of uncertain significance in BRCA1 or BRCA2:
45 a survey of genetic counseling practice. *Genet Test* 2002; 6: 107-13.
- 46 31. — Vos J, Gomez Garcia E, Oosterwijk JC et al. Opening the psychological black box in
47 genetic counseling. The psychological impact of DNA testing is predicted by the counselees'
48 perception, the medical impact by the pathogenic or uninformative BRCA1/2 result.
49 *Psychooncology* 2012; 21: 29-42.
- 50 32. — Vos J, Otten W, van AC et al. The counselees' view of an unclassified variant in
51 BRCA1/2: recall, interpretation, and impact on life. *Psychooncology* 2008; 17: 822-30.
- 52 33. — Gomez Garcia EB, Ambergen T, Blok MJ et al. Patients with an unclassified genetic
53 variant in the BRCA1 or BRCA2 genes show different clinical features from those with a
54 mutation. *J Clin Oncol* 2005; 23: 2185-90.
- 55
56
57
58
59
60

Formatted: English (U.K.)

Formatted: English (U.K.)

Formatted: Italian (Italy)

Formatted: French (France)

Formatted: English (U.K.)

Formatted: English (U.K.)

Reference List

1. Mavaddat N, Peock S, Frost D et al. Cancer Risks for BRCA1 and BRCA2 Mutation Carriers: Results From Prospective Analysis of EMBRACE. J Natl Cancer Inst 2013; 105: 812-22.
2. Mavaddat N, Antoniou AC, Easton DF et al. Genetic susceptibility to breast cancer. Mol Oncol 2010; 4: 174-91.
3. Risch HA, McLaughlin JR, Cole DE et al. Prevalence and penetrance of germline BRCA1 and BRCA2 mutations in a population series of 649 women with ovarian cancer. Am J Hum Genet 2001; 68: 700-10.
4. Couch FJ, Hart SN, Sharma P et al. Inherited Mutations in 17 Breast Cancer Susceptibility Genes Among a Large Triple-Negative Breast Cancer Cohort Unselected for Family History of Breast Cancer. J Clin Oncol 2014.
5. Evans DGR, Barwell J, Eccles DM et al. The Angelina Jolie effect: how high celebrity profile can have a major impact on provision of cancer related services. Breast Cancer Research 2014; 16: 442-7.
6. James PA, Mitchell G, Bogwitz M et al. The Angelina Jolie effect. Med J Aust 2013; 199: 646.

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Space After: 24 pt

- 1
2
3
4
5
6
7 7. Alsop K, Fereday S, Meldrum C et al. BRCA mutation frequency and patterns of
8 treatment response in BRCA mutation-positive women with ovarian cancer: a report
9 from the Australian Ovarian Cancer Study Group. J Clin Oncol 2012; 30: 2654-63.
- 10
11
12
13
14 8. Tutt A, Robson M, Garber JE et al. Oral poly(ADP-ribose) polymerase inhibitor
15 olaparib in patients with BRCA1 or BRCA2 mutations and advanced breast cancer: a
16 proof-of-concept trial. Lancet 2010; 376: 235-44.
- 17
18
19
20
21
22 9. Schwartz MD, Lerman C, Brogan B et al. Impact of BRCA1/BRCA2 counseling and
23 testing on newly diagnosed breast cancer patients. J Clin Oncol 2004; 22: 1823-9.
- 24
25
26
27 10. Vadaparampil ST, Scherr CL, Cragun D et al. Pre-test genetic counseling services for
28 hereditary breast and ovarian cancer delivered by non-genetics professionals in the
29 state of Florida. Clin Genet 2014.
- 30
31
32
33
34 11. Pal T, Cragun D, Lewis C et al. A statewide survey of practitioners to assess
35 knowledge and clinical practices regarding hereditary breast and ovarian cancer.
36 Genet Test Mol Biomarkers 2013; 17: 367-75.
- 37
38
39
40
41
42 12. Wideroff L, Vadaparampil ST, Greene MH et al. Hereditary breast/ovarian and
43 colorectal cancer genetics knowledge in a national sample of US physicians. J Med
44 Genet 2005; 42: 749-55.
- 45
46
47
48
49 13. Chenevix-Trench G, Healey S, Lakhani S et al. Genetic and histopathologic
50 evaluation of BRCA1 and BRCA2 DNA sequence variants of unknown clinical
51 significance. Cancer Res 2006; 66: 2019-27.
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7 14. Eggington JM, Bowles KR, Moyes K et al. A comprehensive laboratory-based
8 program for classification of variants of uncertain significance in hereditary cancer
9 genes. Clin Genet 2014; 86: 229-37.
10
11
12
13
14 15. Lindor NM, Guidugli L, Wang X et al. A review of a multifactorial probability-based
15 model for classification of BRCA1 and BRCA2 variants of uncertain significance
16 (VUS). Hum Mutat 2012; 33: 8-21.
17
18
19
20
21
22 16. Spurdle AB, Healey S, Devereau A et al. ENIGMA--evidence-based network for the
23 interpretation of germline mutant alleles: an international initiative to evaluate risk
24 and clinical significance associated with sequence variation in BRCA1 and BRCA2
25 genes. Hum Mutat 2012; 33: 2-7.
26
27
28
29
30
31 17. Spurdle AB, Whiley PJ, Thompson B et al. BRCA1 R1699Q variant displaying
32 ambiguous functional abrogation confers intermediate breast and ovarian cancer risk.
33 J Med Genet 2012; 49: 525-32.
34
35
36
37
38
39 18. Plon SE, Eccles DM, Easton D et al. Sequence variant classification and reporting:
40 recommendations for improving the interpretation of cancer susceptibility genetic test
41 results. Hum Mutat 2008; 29: 1282-91.
42
43
44
45
46 19. Moghadasi S, Hofland N, Wouts JN et al. Variants of uncertain significance in
47 BRCA1 and BRCA2 assessment of in silico analysis and a proposal for
48 communication in genetic counselling. J Med Genet 2013; 50: 74-9.
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7 20. Goldgar DE, Easton DF, Deffenbaugh AM et al. Integrated evaluation of DNA
8 sequence variants of unknown clinical significance: application to BRCA1 and
9 BRCA2. Am J Hum Genet 2004; 75: 535-44.
- 10
11
12
13
14 21. Walker LC, Whiley PJ, Couch FJ et al. Detection of splicing aberrations caused by
15 BRCA1 and BRCA2 sequence variants encoding missense substitutions: implications
16 for prediction of pathogenicity. Hum Mutat 2010; 31: E1484-E1505.
- 17
18
19
20
21 22. Millot GA, Carvalho MA, Caputo SM et al. A guide for functional analysis of
22 BRCA1 variants of uncertain significance. Hum Mutat 2012; 33: 1526-37.
- 23
24
25
26
27 23. Goldgar DE, Easton DF, Byrnes GB et al. Genetic evidence and integration of various
28 data sources for classifying uncertain variants into a single model. Hum Mutat 2008;
29 29: 1265-72.
- 30
31
32
33
34 24. Guidugli L, Carreira A, Caputo SM et al. Functional assays for analysis of variants of
35 uncertain significance in BRCA2. Hum Mutat 2014; 35: 151-64.
- 36
37
38
39
40 25. Iversen ES, Jr., Couch FJ, Goldgar DE et al. A computational method to classify
41 variants of uncertain significance using functional assay data with application to
42 BRCA1. Cancer Epidemiol Biomarkers Prev 2011; 20: 1078-88.
- 43
44
45
46
47 26. Tavtigian SV, Greenblatt MS, Goldgar DE et al. Assessing pathogenicity: overview of
48 results from the IARC Unclassified Genetic Variants Working Group. Hum Mutat
49 2008; 29: 1261-4.
- 50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
27. Anagnostopoulos T, Pertesi M, Konstantopoulou I et al. G1738R is a BRCA1 founder mutation in Greek breast/ovarian cancer patients: evaluation of its pathogenicity and inferences on its genealogical history. Breast Cancer Res Treat 2008; 110: 377-85.
28. Newman B, Mu H, Butler LM et al. Frequency of breast cancer attributable to BRCA1 in a population-based series of American women. JAMA 1998; 279: 915-21.
29. Blay P, Santamaria I, Pitiot AS et al. Mutational analysis of BRCA1 and BRCA2 in hereditary breast and ovarian cancer families from Asturias (Northern Spain). BMC Cancer 2013; 13: 243.
30. Mirkovic N, Marti-Renom MA, Weber BL et al. Structure-based assessment of missense mutations in human BRCA1: implications for breast and ovarian cancer predisposition. Cancer Res 2004; 64: 3790-7.
31. Tavtigian SV, Byrnes GB, Goldgar DE et al. Classification of rare missense substitutions, using risk surfaces, with genetic- and molecular-epidemiology applications. Hum Mutat 2008; 29: 1342-54.
32. Guidugli L, Pankratz VS, Singh N et al. A classification model for BRCA2 DNA binding domain missense variants based on homology-directed repair activity. Cancer Res 2013; 73: 265-75.
33. Lindor NM, Goldgar DE, Tavtigian SV et al. BRCA1/2 sequence variants of uncertain significance: a primer for providers to assist in discussions and in medical management. Oncologist 2013; 18: 518-24.

- 1
2
3
4
5
6
7 34. Petrucelli N, Lazebnik N, Huelsman KM et al. Clinical interpretation and
8 recommendations for patients with a variant of uncertain significance in BRCA1 or
9 BRCA2: a survey of genetic counseling practice. Genet Test 2002; 6: 107-13.
10
11
12
13
14 35. Vos J, Gomez-Garcia E, Oosterwijk JC et al. Opening the psychological black box in
15 genetic counseling. The psychological impact of DNA testing is predicted by the
16 counselees' perception, the medical impact by the pathogenic or uninformative
17 BRCA1/2-result. Psychooncology 2012; 21: 29-42.
18
19
20
21
22
23
24 36. Vos J, Otten W, van AC et al. The counselees' view of an unclassified variant in
25 BRCA1/2: recall, interpretation, and impact on life. Psychooncology 2008; 17: 822-
26 30.
27
28
29
30
31 37. Gomez-Garcia EB, Ambergen T, Blok MJ et al. Patients with an unclassified genetic
32 variant in the BRCA1 or BRCA2 genes show different clinical features from those
33 with a mutation. J Clin Oncol 2005; 23: 2185-90.
34
35
36
37
38 38. Cheon JY, Mozersky J, Cook-Deegan R. Variants of uncertain significance in BRCA:
39 a harbinger of ethical and policy issues to come? Genome Med 2014; 6: 121.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Formatted: Indent: Left: 0", Hanging: 0.5",
Space After: 0 pt, Line spacing: Double, Tab
stops: 0.38", Right + 0.5", Left

1
2
3
4
5
6
7 Supplementary material:

Formatted: Not Highlight

8 Supplementary figure 1: Genetic variability information

9
10 Supplementary figure 2: Suggested clinical pathway for patients when genetic testing is
11 initiated by the oncology clinic.

12
13 Supplementary table (1). Types of sequence variants reported in *BRCA* mutation detection
14 tests according to risk relevant for clinical management

Formatted: Left, Space After: 0 pt

15
16 Supplementary table (2). Examples of software tools available for clinical variant evaluation
17 online

18
19 Supplementary table 3. 5-tier classification *BRCA1/2* VUS

Formatted: Space After: 0 pt, Line spacing:
single

20
21 Supplementary table 4. Commonly used online database resources that provide some
22 interpretation of *BRCA* sequence variants

Formatted: Left, Space After: 0 pt

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Appendix:**Name and affiliations of the ENIGMA CWG contributors: (excluding those named in the authors' list)**

R. Driessen: Familial Cancer Centre, Peter MacCallum Cancer Centre, The University of Melbourne, Melbourne, Australia.

N.M. Lindor: Department of Health Sciences Research, Mayo Clinic, Scottsdale, AZ, USA.

M.J. Blok: Department of Clinical Genetics, Maastricht University Medical Centre, Maastricht, the Netherlands.

P. Moller: Research Group Inherited Cancer, Department of Medical Genetics, University Hospital, Oslo, Norway.

M. de la Hoya: Molecular Oncology Laboratory, Hospital Clínico San Carlos (IdISSC), Madrid, Spain.

T. Pal: Cancer Epidemiology Program, H. Lee Moffitt Cancer Center, Tampa, USA.

S. Domchek and K. Nathanson: Bassett Research Center and Abramson Cancer Center University of Pennsylvania Philadelphia, PA, USA.

C. Van Asperen: Department of Clinical Genetics, Leiden University Medical Center, Leiden, the Netherlands

O. Diez: Oncogenetics Group, University Hospital Vall d'Hebron; Vall d'Hebron Institute of Oncology (VHIO), Barcelona, Spain.

K. Rheim: Center for Hereditary Breast and Ovarian Cancer, Center for Integrated Oncology (CIO) and Center for Molecular Medicine Cologne (CMMC), University Hospital Cologne, Germany.

D. Stoppa-Lyonnet: Genetis Service, Curie Institute, University of Paris Descartes, Paris, France

M. Parsons: Molecular Cancer Epidemiology Laboratory, Genetics and Computational Biology Division, QIMR Berghofer Medical Research Institute, BNE, Herston, QLD, Australia.

D. Goldgar: Department of Dermatology, Huntsman Cancer Institute, University of Utah School of Medicine; Salt Lake City, Utah, USA.

Supplementary Table (1). Types of sequence variants reported in *BRCA* mutation detection tests according to risk relevant for clinical management

Clinical risk	Descriptors observed in clinical reports	Interpretation
High	<ul style="list-style-type: none"> Functionally deleterious mutation Pathogenic mutation High risk mutation Deleterious variant 	<p>Variants that result in a high lifetime risk of breast and ovarian cancer. Deleterious variants disrupt normal protein function. They include nonsense changes, out-of-frame insertions or deletions, large gene rearrangements or splicing variants altering the canonical splicing acceptor and donor sites disrupting regulatory regions, as well as some missense changes. Supportive evidence from multiple sources may be required to call a variant deleterious where predictions of functional consequences are unclear^a.</p>
Uncertain	<ul style="list-style-type: none"> Missense mutation Rare variant Variant of uncertain clinical significance (VUS, VOUS) Uncertain Variant Variant of Uncertain Variant of Unknown Clinical Significance 	<p>Variants that differ from the published reference DNA sequence (RefSeq BRCA1: U14680; BRCA2: NM00059) and are not classifiable as either deleterious or as neutral based on available evidence.</p> <p>The association with <i>clinical phenotype</i> at <u>this time</u> is <i>unknown</i> and cannot be used to inform clinical decisions.</p>
None	<ul style="list-style-type: none"> Frequent mutation Common variant Polymorphism Variants of low clinical significance Variants of no clinical significance Neutral variants Benign variants 	<p>Changes in the DNA sequence that do not disrupt the normal function of the encoded protein and are not associated with a clinically important increased risk of disease. If present in the population at a frequency of greater than 1% these may be referred to as polymorphisms.</p>
<p>^a for example assays showing partial loss of normal transcript from aberrant splicing or partially reduced function from an amino acid substitution may require additional evidence to classify as class 4 or 5.</p>		

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary Table (2) Examples of software tools available for clinical variant evaluation online

Resource	Description	Website
Polyphen2	POLYmorphism PHENotypes Polyphen2 is freely available, web-based program from Harvard University that predicts whether an amino acid substitution affects protein structure and function.	http://genetics.bwh.harvard.edu/pph2/
SIFT	Sorting Intolerant From Tolerant SIFT software is freely available and predicts whether an amino acid substitution affects protein function.	http://sift.jcvi.org/
Align-GVGD	Align Grantham Variation and Grantham Deviation. Align-GVGD is a freely available, web-based program from IARC (International Agency for Cancer Research) that combines the known chemical nature of amino acids and the alignments of DNA sequences across species to predict the likely pathogenicity of missense substitutions.	http://agvgd.iarc.fr/about.php
HCI database of prior probabilities	The Huntsman Cancer Institute Database of Prior Probabilities of Pathogenicity for Single Nucleotide Substitutions provides the prior probability of pathogenicity estimate which is the starting point for the multifactorial likelihood estimate for a novel SNP identified through genetic testing for a BRCA gene.	http://priors.hci.utah.edu/PRIORS/
GeneSplicer	GeneSplicer is a freely available flexible system for detecting splice sites in the genomic DNA of various eukaryotes.	http://www.ncbi.nlm.nih.gov/pmc/articles/PMC29713/
Human Splicing Finder	This tool is freely available and aimed at studying pre-mRNA splicing and identifying variants that may disrupt normal splicing.	http://www.umd.be/HSF3/
Commercial software e.g. Alamut	Alamut is commercially available software that provides a single interface to bring together multiple data sources including many of those listed above in a simplified and streamlined tool for rapid assembly of data about any one variant aimed at busy diagnostic laboratories.	http://www.interactivebiosoftware.com/software/alamut/overview
Entrez-Pubmed Google Scholar	Web-based search engines to look for research publications including a specific mutation.	http://www.ncbi.nlm.nih.gov/pubmed http://scholar.google.com
UMD-BRCA1 UMD-BRCA2	The UMD-BRCA1/BRCA2 databases have been set up in a joined national effort through the network of the 16 French diagnostic laboratories. A classification is proposed based on the classical parameters including splicing algorithms. A strong effort is done to perform cosegregation studies.	http://www.umd.be/BRCA1/ http://www.umd.be/BRCA2/

Supplementary Table 3. 5-tier classification *BRCA1/2* VUS modified from [15]

Class	Description	Probability of being pathogenic	Clinical predictive testing of at risk relatives	Management recommendations if at-risk relative has the variant	Research testing of family members
5	Definitely pathogenic	>0.99	Yes	Full high-risk guidelines	Not indicated
4	Likely pathogenic	0.95-0.99	Yes	Full high-risk guidelines	May be helpful to further classify variant
3	Uncertain	0.05-0.949	No	Presence of variant is irrelevant to risk assessment, manage risk based on family history only	May be helpful to further classify variant
2	Likely not pathogenic or of no clinical significance	0.001-0.049	No	Manage risk based on family history only	May be helpful to further classify variant
1	Not pathogenic or of no clinical significance	<0.001	No	Manage risk based on family history only	Not indicated

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary Table 4: Commonly used online database resources that provide some interpretation of *BRCA* sequence variants although

Resource	Description	Website
IARC/LOVD – (Leiden Open Variation Database)	Separate <i>BRCA1</i> and <i>BRCA2</i> variant databases curated by experts at the University of Leiden and IARC. Includes only those VUS with literature references and records the published literature associated with each variant. Each variant is given an IARC class[15] and links to published source data. Submissions of variants are from registered submitters largely from the research community. Curation of submissions for each gene is undertaken by a named expert curator on a voluntary basis.	http://brca.iarc.fr/LOVD/home.php soon to be replaced by http://hci-exlovd.hci.utah.edu/home.php
The Breast Cancer Information Core (BIC)	A database that acts as a central repository for <i>BRCA1</i> and <i>BRCA2</i> variants (deleterious, neutral or VUS) deposited by submitters from research and clinical sites internationally. Recently, a central curation process working to classify all variants according to the IARC 5 tier classification[15] scheme has been introduced.	http://research.nhgri.nih.gov/bic
ClinVar	ClinVar is a freely accessible, public archive of reports of the relationships between human variations and phenotypes presented with supporting evidence and an indication of likely clinical significance. Submissions are from research and some diagnostic laboratories and submitters include a classification for submitted variants, submissions are not currently curated.	http://www.ncbi.nlm.nih.gov/clinvar/
Human Variome Project	The Human Variome Project, under the auspices of UNESCO, has been created as a unified reporting portal and lists 4 separate databases for <i>BRCA1</i> and 4 for <i>BRCA2</i> . As yet it offers no formal curation or attempt to classify variants	http://www.humanvariomeproject.org

Supplementary Table 4: Commonly used online database resources that provide some interpretation of *BRCA* sequence variants

Resource	Description	Website
IARC/LOVD – (Leiden Open (source) Variation Database)	Separate <i>BRCA1</i> and <i>BRCA2</i> variant databases curated by experts at the University of Leiden and IARC. Includes only those VUS with literature references and records the published literature associated with each variant. Each variant is given an IARC class[15] and links to published source data. <u>Submissions of variants are from registered submitters largely from the research community. Curation of submissions for each gene is undertaken by a named expert curator on a voluntary basis.</u>	http://brca.iarc.fr/LOVD/home.php soon to be replaced by http://hci-exlovd.hci.utah.edu/home.php
The Breast Cancer Information Core (BIC)	<u>A database that acts as a central repository for <i>BRCA1</i> and <i>BRCA2</i> variants (deleterious, neutral or VUS) deposited by submitters from research and clinical sites internationally. Recently, a central curation process working to classify all variants according to the IARC 5 tier classification[15] scheme has been introduced. An international collaboration that acts as a central repository for all reported <i>BRCA1</i> and <i>BRCA2</i> deleterious, neutral and VUS variants. The vast majority of classifications are self-reported by the submitting laboratories. Recently, a central curation process working to classify all variants according to the IARC classification[15] scheme has been introduced.</u>	http://research.nhgri.nih.gov/bic
ClinVar	ClinVar is a freely accessible, public archive of reports of the relationships between human variations and phenotypes presented with supporting evidence and an indication of likely clinical significance. <u>Submissions are from research and some diagnostic laboratories and submitters include a classification for submitted variants, submissions are not currently curated.</u>	http://www.ncbi.nlm.nih.gov/clinvar/
Human Variome Project	The Human Variome Project, under the auspices of UNESCO, has been created as a unified reporting portal and lists 4 separate databases for <i>BRCA1</i> and 4 for <i>BRCA2</i> . As yet it offers no formal curation or attempt to classify variants	http://www.humanvariomeproject.org

Figure 1: Important facts about genetic variability


Variation in DNA is common	
Facts and terms	Further information
When a gene is tested for mutations, the sequence of that gene in the individual is compared to an accepted reference sequence for the gene and variation from the reference sequence is often observed.	The human genome contains ~ 6 billion bases of genetic sequence in any diploid cell and on average there is a change in the reference sequence approximately every 500 bases, much of this variation does not obviously impact on function but occasionally it does.
<i>BRCA1</i> and <i>BRCA2</i> are very large genes	Random variation in the reference sequence occurs frequently across the population, not associated with any clinical effect. This is true whether the individual has had cancer or not.
Non-pathogenic variation in sequence is as frequent in the general population as in the cancer affected patient	It is tempting to believe that because a person with cancer has been tested for a mutation in the <i>BRCA1/2</i> gene, that any deviation from the expected code is causative but in reality, in the absence of clear evidence of a loss of function in the mutated copy of the gene, most of the genetic variation is of little or no significance in relation to disease causation
Different ethnic groups have different frequencies of variants	For ethnic populations where relatively little genetic testing has been undertaken, common polymorphisms in that ethnic minority group may be unrecognised in the testing laboratory and are therefore more likely to be reported as of uncertain significance (VUS).
Terms used in describing genetic variants	
Polymorphisms (meaning “many forms”) are common variations observed in more than 1% of the normal population.	When observed frequently in population controls (unaffected by cancer), this type of sequence change is generally not associated with a high cancer risk.
A change in a single base (T,G,A or C) is a single nucleotide variant and if the particular variant is relatively frequently observed in a particular population, it is called a single nucleotide polymorphism (SNP)	Single nucleotide variants usually have no major clinical consequence especially if frequent (i.e. SNPs) but rarely they may affect the function of the protein being coded. Synonymous: a single base change in the exon which results in no change to the expected amino acid at that position so is usually of no consequence but can create or destroy a native donor or acceptor splice site. Non-synonymous: also known as missense mutations lead to a change in one amino acid, these may affect a critical functional domain, although most do not. Variants within an intron (ie non coding part of the gene) can also rarely destroy or create false donor or acceptor splice sites.
In-frame deletions or insertions	An insertion or deletion of three bases starting with the first base of an amino acid code is an in frame deletion or insertion and leads to the insertion or deletion of one or sometimes more amino acids but the full length protein is otherwise predicted to be complete. These may have no effect on protein function since most of the protein is translated correctly.
Variants of uncertain significance (VUS)	Where a change to the expected sequence is observed that has not been observed with any frequency in the testing laboratory and has not been classed as non-pathogenic in literature or databases, genetic testing laboratories will usually report these as a VUS. Gathering multiple lines of additional information may help

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	to clarify pathogenicity [15]. In reality most of these are unlikely to be pathogenic mutations but a few will be.
--	--

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Recommendations for referral of cases to a specialist genetic clinic