

Duncan Street, BM
Christopher A.
Halfpenny, PhD
Ian Galea, PhD

CNS INFLAMMATION OTHER THAN MULTIPLE SCLEROSIS: HOW LIKELY IS DIAGNOSIS?

OPEN

The incidence, diagnostic landscape, and workload impact of CNS inflammatory diseases other than multiple sclerosis (MS) (CIDOMS) in a tertiary setting is unknown. We describe a retrospective case series of 64 patients identified over a 2-year period (2009–2010) at the Wessex Neurological Centre in the United Kingdom, accounting for 4% of all patients seen at the center. As expected, neurosarcoidosis and neuromyelitis optica (NMO) were the most common diagnoses reached (14% each); other diagnoses singly accounted for <10%. However, the likeliest diagnostic outcome (strikingly, in 25%) was nondiagnosis, despite intensive investigation and a mean follow-up period of 3 years. Undiagnosed patients with CIDOMS represented the largest workload of the neurology center.

The Wessex Neurological Centre is a typical regional neurologic service with a catchment population of ~3 million in southern England and on-site expertise in neurology, neurosurgery, neuropathology, neurophysiology, neuroradiology, neuropsychology, neurorehabilitation, and neurophysiotherapy, across which specialist interest in neuroinflammatory disease is adequately represented. The study was approved by the University of Southampton's Ethics and Research Governance. A diagnosis of CIDOMS was made when there was unequivocal evidence of CNS inflammation (with certainty of an inflammatory etiology based on clinical, radiologic, CSF, and other laboratory findings) in the absence of MS or clinically isolated syndrome (CIS). Published diagnostic criteria (or, in their absence, published consensus opinions) were utilized to establish specific CIDOMS diagnoses (tables e-1 and e-2 on the *Neurology*[®] Web site at Neurology.org). MS and CIS were diagnosed according to established criteria.^{1,2} Detailed evaluation was conducted regarding resource utilization, including inpatient episode duration, imaging, invasive procedures, neurophysiology, non-neurology specialist reviews, and laboratory tests. In order to provide a sense of proportion to figures, a similar analysis was performed in patients admitted

during the same period whose final diagnosis was MS. Further method details are available online (e-Methods).

From a total of 1,525 patients admitted to the tertiary neurology center, 81 had a working diagnosis of CIDOMS, which was maintained in 64 cases as a final diagnosis. This represents an incidence rate of 11 cases of CIDOMS per million person-years. Seventeen patients with an initial working diagnosis of CIDOMS received a diagnosis of MS by the end of the follow up-period. A breakdown of individual CIDOMS diagnoses is given in table 1 (for more detail, see tables e-1 and e-2).

Despite their small number ($n = 64$) in comparison with MS patients ($n = 167$), patients with CIDOMS required disproportionately longer inpatient stays, more intensive care, and larger numbers of investigations (see table 1 and figure e-1, $p < 0.001$ across all categories). Naturally, this resulted in higher costs (£520,409.18 vs £259,941.51, i.e., twice as much, $p < 0.0001$, see table e-3 and figure e-2). Among patients with CIDOMS, those without a diagnosis represented the largest workload of the neurology center, since they collectively needed the longest inpatient stay and the greatest number of investigations (see table 1).

The definition of CIDOMS did not include CIS, and it may be argued that cases of CIS with low risk of conversion to MS (with normal MRI and CSF¹) may turn out to have CIDOMS. None of the patients with a diagnosis of CIS converted to a diagnosis of CIDOMS in this study, but this may need longer follow-up, as illustrated by a recent case series.³ However, categorization of low-risk CIS with CIDOMS and high-risk CIS with MS maintains nondiagnosis as the most common outcome in patients with CIDOMS (16 out of 78, i.e., 21%), and maintains patients with CIDOMS as still more resource-intensive than patients with MS (data not shown).

Two undiagnosed patients with CIDOMS had a clinical phenotype that resembled NMO spectrum disorder, though not typical (last 2 cases in table e-2). Restriction of anti-aquaporin antibodies to the CSF,⁴ anti-aquaporin-4 antibody assay sensitivity,^{5,6} and antimyelin oligodendrocyte glycoprotein

Supplemental data
at Neurology.org

Table 1 Number, percentage, and biennial total resource use of patients with CIDOMS by their final diagnostic categories

Diagnostic category	Number	%	Total days in hospital	Days on ICU	All imaging	MRIs	Invasive procedures	Neurophysiology	Specialist reviews	Laboratory tests
Unclassified	16 ^a	25 ^a	295.5 ^a	0	75 ^a	30 ^a	31 ^a	13 ^a	10 ^a	994
Neurosarcoidosis	9	14	278	13.5	59	18	19	4	5	1,021 ^a
Neuromyelitis optica	9	14	117	0	18	6	4	2	1	420
Cerebral vasculitis	4	6	246.5	2	52	10	16	2	9	707
Anti-VGKC encephalitis	4	6	93.5	0	19	3	6	4	1	227
ADEM	3	5	75.5	9	15	5	4	1	0	327
Postinfective	3	5	32	0	9	3	3	3	0	150
Anti-NMDA receptor encephalitis	2	3	216.5	39 ^a	18	3	6	4	2	584
Paraneoplasia in lymphoma	2	3	87.5	0	27	8	5	4	0	326
Opsoclonus-myoclonus	2	3	53	0	6	2	2	0	0	134
Neuro-Behçet	2	3	31.5	0	7	3	3	2	2	92
Atypical MS	2	3	13.5	0	4	3	3	2	1	95
CRION	1	2	10.5	0	3	2	3	2	1	81
Anti-Hu syndrome	1	2	17	0	4	0	0	0	2	38
Lymphocytic hypophysitis	1	2	7	0	2	0	3	0	0	61
Neuro-Sjögren	1	2	4	0	3	3	1	2	1	30
Stiff person syndrome	1	2	9	0	0	0	0	0	0	15
Radiologically isolated syndrome	1	2	0.5	0	1	1	1	0	0	15
Total	64	100	1,588	63.5	322	100	110	45	35	5,317
MS (for comparison)	167	100	689.5	0	57	109	106	43	5	3,735
CIS: low and high risk (for comparison)	25	100	122	0	25	11	21	12	1	496

Abbreviations: ADEM = acute disseminated encephalomyelitis; CIDOMS = CNS inflammatory diseases other than multiple sclerosis; CIS = clinically isolated syndrome; CRION = chronic relapsing inflammatory optic neuritis; ICU = intensive care unit; MS = multiple sclerosis; VGKC = voltage-gated potassium channel. Invasive procedures included catheter cerebral angiogram; lumbar puncture; biopsy of brain, nerve, skin, pinna, or bone marrow; gastroscopy; bronchoscopy; or flexisigmoidoscopy.

^a Highest numbers.

antibodies⁷ may be explanations. Yet again, a sensitivity analysis excluding these patients did not change conclusions.

Collectively CIDOMS are common and consist of up to 25% of the neurologic practice pertaining to inflammatory CNS disorders. The data highlight the importance of education regarding the diagnosis and treatment of these disorders. The striking finding that one-quarter of CIDOMS remained undiagnosed means that, in the absence of a diagnosis of neurosarcoidosis or NMO, an unclassified inflammatory disease is eventually more likely than other rarer diagnoses, which singly accounted for fewer than 10%. Broader serologic testing for individually rare antibodies and discovery of novel biomarkers should facilitate more rapid diagnoses in unclassified cases. Meanwhile, in the absence of a diagnosis, consensus guidelines to help recognition of antibody vs T-cell-mediated neuroinflammatory

disorders will enable a rational empirical approach to treatment based on the likely underlying pathophysiologic mechanism.

From Clinical Neurosciences (D.S., I.G.), Clinical & Experimental Sciences, Faculty of Medicine, University of Southampton; and Wessex Neurological Centre (C.A.H., I.G.), University Hospital Southampton NHS Foundation Trust, UK.

Author contributions: Dr. Street: data acquisition, data analysis, drafting of the manuscript, and review of manuscript for important intellectual content. Dr. Halfpenny: review of the manuscript for important intellectual content. Dr. Galea: data acquisition, data analysis, drafting of the manuscript, review of the manuscript for important intellectual content, and study supervision.

Study funding: Supported by the Wellcome Trust/Academy of Medical Sciences (starter grant to I.G.) and the University of Southampton's Faculty of Medicine (D.S.).

Disclosure: D. Street reports no disclosures relevant to the manuscript. C. Halfpenny has received expenses to attend meetings from Biogen-Idex, Merck-Serono, and Teva-UK. I. Galea has received expenses to attend meetings from Teva-UK, has received commercial research grant support from Merck-Serono and IQ products, and holds research grants from the MS Society, Association of British Neurologists, Wessex

Medical Research, Medical Research Council, and the Engineering, and Physical Sciences Research Council, UK. Go to Neurology.org for full disclosures.

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received October 1, 2013. Accepted in final form December 13, 2013.

Correspondence to Dr. Galea: I.Galea@soton.ac.uk

© 2014 American Academy of Neurology

1. Miller DH, Chard DT, Ciccarelli O. Clinically isolated syndromes. *Lancet Neurol* 2012;11:157–169.
2. Polman CH, Reingold SC, Banwell B, et al. Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald criteria. *Ann Neurol* 2011;69:292–302.
3. Kurne A, Isikay IC, Karlioguz K, et al. A clinically isolated syndrome: a challenging entity: multiple sclerosis or collagen tissue disorders: clues for differentiation. *J Neurol* 2008;255:1625–1635.
4. McKeon A, Pittock SJ, Lennon VA. CSF complements serum for evaluating paraneoplastic antibodies and NMO-IgG. *Neurology* 2011;76:1108–1110.
5. Fujihara K, Leite MI. Seronegative NMO: a sensitive AQP4 antibody test clarifies clinical features and next challenges. *Neurology* 2013;80:2176–2177.
6. Marignier R, Bernard-Valnet R, Giraudon P, et al. Aquaporin-4 antibody-negative neuromyelitis optica: distinct assay sensitivity-dependent entity. *Neurology* 2013;80:2194–2200.
7. Kitley J, Woodhall M, Waters P, et al. Myelin-oligodendrocyte glycoprotein antibodies in adults with a neuromyelitis optica phenotype. *Neurology* 2012;79:1273–1277.

Complete the AAN 2014 *Neurology Compensation and Productivity Survey* by May 9

The AAN launched its second annual *Neurology Compensation and Productivity Survey* in March and needs practicing US members and their practices to contribute their data. It is critical that all US neurologists and practice managers participate in the survey to ensure the most accurate and authoritative data representing the US neurology landscape. Visit AAN.com/view/2014NeuroSurvey to review preparation documents, including an FAQ and Quick Start Guide. Complete the survey by May 9 and get *free access* to the online results and the *Neurology Compensation and Productivity Report*, available in early July 2014. The cost to access the data and report for nonparticipants is \$600 for AAN members and \$1200 for nonmembers.

WriteClick: Rapid Online Correspondence

The editors encourage comments about recent articles through WriteClick:

Go to www.neurology.org and click on the “WriteClick” tab at the top of the page. Responses will be posted within 72 hours of submission.

Before using WriteClick, remember the following:

- WriteClick is restricted to comments about studies published in *Neurology* within the last eight weeks
- Read previously posted comments; redundant comments will not be posted
- Your submission must be 200 words or less and have a maximum of five references; reference one must be the article on which you are commenting
- You can include a maximum of five authors (including yourself)

Neurology®

CNS inflammation other than multiple sclerosis: How likely is diagnosis?

Duncan Street, Christopher A. Halfpenny and Ian Galea

Neurology 2014;82;1187-1189 Published Online before print February 28, 2014

DOI 10.1212/WNL.0000000000000273

This information is current as of February 28, 2014

Updated Information & Services	including high resolution figures, can be found at: http://www.neurology.org/content/82/13/1187.full.html
Supplementary Material	Supplementary material can be found at: http://www.neurology.org/content/suppl/2014/02/28/WNL.0000000000000273.DC1.html
References	This article cites 7 articles, 4 of which you can access for free at: http://www.neurology.org/content/82/13/1187.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Demyelinating disease (CNS) http://www.neurology.org/cgi/collection/all_demyelinating_disease_cn s All Health Services Research http://www.neurology.org/cgi/collection/all_health_services_research Incidence studies http://www.neurology.org/cgi/collection/incidence_studies
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://www.neurology.org/misc/addir.xhtml#reprintsus

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2014 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

