



Cite this: *Dalton Trans.*, 2014, **43**, 14600

Phosphine complexes of aluminium(III) halides – preparation and structural and spectroscopic systematics†

Jennifer Burt, William Levason,* Mark E. Light and Gillian Reid

Six-coordinate pseudo-octahedral complexes $trans-[AlX_2(L-L)_2][AlX_4]$ ($X = Cl, Br$ or I ; $L-L = o-C_6H_4(PMe_2)_2, Me_2P(CH_2)_2PMe_2$) are produced from reaction of AlX_3 with the diphosphine in CH_2Cl_2 ($X = Cl$) or toluene ($X = Br$ or I) solution. Four-coordinate dimers $[Cl_3Al(\mu-L'-L')AlCl_3]$ ($L'-L' = Me_2P(CH_2)_2PMe_2, Cy_2P(CH_2)_2PCy_2$), and the tetrahedral cation $[AlCl_2(o-C_6H_4(PPh_2)_2)][AlCl_4]$ were also obtained. Both four- and five-coordinate complexes $[AlX_3(PMe_3)]$ and $[AlX_3(PMe_3)_2]$ could be isolated with PMe_3 depending upon the ratio of reagents used. These extremely moisture sensitive complexes have been characterised by microanalysis, IR and multinuclear NMR ($^1H, ^{31}P\{^1H\}$ and ^{27}Al) spectroscopy. X-ray crystal structures are reported for $[AlCl_2(o-C_6H_4(PMe_2)_2)_2][AlCl_4]$, $[AlCl_2(Me_2P(CH_2)_2PMe_2)_2][AlCl_4]$, $[Cl_3Al(\mu-Me_2P(CH_2)_2PMe_2)AlCl_3]$, $[Cl_3Al(\mu-Cy_2P(CH_2)_2PCy_2)AlCl_3]$, $[AlCl_3(PMe_3)]$, $[AlCl_3(PMe_3)_2]$, and for the six-coordinate cation complex $[AlCl_2(o-C_6H_4(PPh_2)_2)_2][AlCl_4]$, although a bulk sample of the last could not be isolated. Tertiary arsines ($AsPh_3$ or $AsEt_3$) form only 1:1 complexes even with excess arsine present. The unstable $[AlCl_2(o-C_6H_4(AsMe_2)_2)][AlCl_4]$ is also described, and shown to decompose rapidly in CH_2Cl_2 solution to form the diquaternised diarsine cation $[o-C_6H_4(AsMe_2)_2(CH_2)_2][AlCl_4]_2$, which was fully characterised. Comparisons are drawn with the corresponding gallium(III) systems (Cheng *et al.*, *Inorg. Chem.*, 2007, **46**, 7215–7223) and with AlX_3 complexes of Group 16 ligands (George *et al.*, *Dalton Trans.*, 2014, **43**, 3637–3648), and it is concluded that the differences between the Al and Ga systems reflect the higher Lewis acidity of aluminium(III) towards soft donor ligands.

Received 7th July 2014,
Accepted 8th August 2014

DOI: 10.1039/c4dt02051k

www.rsc.org/dalton

Introduction

Aluminium chloride is an archetypal Lewis acid, widely used in industry and in the laboratory as a catalyst for condensation, polymerisation and isomerisation reactions. It is perhaps best known as a Friedel–Crafts catalyst for alkylations or acylations where the strong affinity for chloride generates incipient carbocations in combination with $[AlCl_4]^-$ anions.^{1,2} Typically, $AlBr_3$ and AlI_3 are weaker Lewis acids, but chemically similar. Solid $AlCl_3$ contains six-coordinate aluminium, but in the melt or when dissolved in inert solvents, dimeric tetrahedral molecules $[Cl_2Al(\mu-Cl)_2AlCl_2]$ are present. Solid $AlBr_3$ is also dimeric, $[Br_2Al(\mu-Br)_2AlBr_2]$, whereas AlI_3 is a chain polymer $\{[I_2Al(\mu-I)]_n\}$.³ The extensive coordination chemistry of these

three halides, mostly with hard N- or O-donor ligands, continues to attract much effort.³ In contrast, AlF_3 is an inert polymer containing six-coordinate aluminium; few complexes are known.⁴ Complexes of aluminium halides with neutral soft-donor ligands have received much less attention and current knowledge is not systematic.³ We recently reported⁵ complexes with thio-, seleno- and telluro-ether ligands, including examples of four-coordinate $[X_3Al(ER_2)]$ ($X = Cl, Br$ or I ; $E = S, Se$ or Te ; $R = alkyl$) and $[(AlCl_3)_2\{o-C_6H_4(CH_2SEt)_2\}]$, as well as six-coordinate $[AlX_2\{MeE(CH_2)_2EME\}_2][AlX_4]$ ($E = S$ or Se). The isolation of the latter was unexpected since gallium(III) halides gave only four-coordinate complexes with the bidentate chalcogenoethers.⁶ Both Al and Ga can achieve six-coordination with thia- and seleno-macrocycles.^{6,7} There are a substantial number of phosphine and arsine complexes of trialkylaluminiums, and even some examples with stibine and bismuthine ligands; all contain four-coordinate aluminium with a single Group 15 donor per Al centre, even when polydentate ligands are used.⁸ In contrast, complexes with AlX_3 are few, and almost all are of the type $[X_3Al(E'R_3)]$ ($E'R_3 = PPh_3$,⁹ $AsPh_3$,¹⁰ $AsEt_3$,¹⁰ $AsMe_3$,¹⁰ $P(mesityl)_3$ ^{11a} or $P(SiMe_3)_3$ ^{11b}) containing tetrahedral aluminium centres.⁸ A rare example of a higher coordination number (five) is found in the trigonal

School of Chemistry, University of Southampton, Southampton SO17 1BJ, UK.

E-mail: wxl@soton.ac.uk

† Electronic supplementary information (ESI) available: The X-ray structural data for $[o-C_6H_4(PMe_2)(PPhMe_2)]_2[AlX_4][X]$ ($X = Cl$ or Br) and $[Cy_2P(H)(CH_2)_2PCy_2(H)]_2[AlCl_4]_2$ and an analysis of M–X bond lengths in $[MX_4]^-$ ($M = Al, Ga, I$) based upon data taken from the Cambridge Structural Database, with derived histograms, are also given. CCDC 1005860–1005870. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4dt02051k



bipyramidal $[\text{AlI}_3(\text{PET}_3)_2]$.¹² Surprisingly, no examples with bidentate phosphine or arsine ligands have been prepared. In a previous study with gallium(III) halides,¹³ four-coordination was dominant, either as neutral dimers $[\text{X}_3\text{Ga}\{\mu\text{-Et}_2\text{-P}(\text{CH}_2)_2\text{PET}_2\}\text{GaX}_3]$ or $[\text{X}_3\text{Ga}\{\mu\text{-Ph}_2\text{As}(\text{CH}_2)_2\text{AsPh}_2\}\text{GaX}_3]$, or tetrahedral cations $[\text{GaX}_2(\text{L-L})]^+$ ($\text{L-L} = o\text{-C}_6\text{H}_4(\text{PPh}_2)_2$ or $o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2$). Octahedral cations, $[\text{GaX}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2]^+$, are only known with $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$, which is a very strong σ -donor ligand, has small steric demands and is pre-organised for chelation.¹³ The larger In(III) centre forms both four-coordinate $[\text{InX}_2(\text{L-L})]^+$ and six-coordinate $[\text{InX}_2(\text{L-L})_2]^+$ cations, as well as neutral six-coordinate dimers, $[\text{In}_2\text{X}_6(\text{L-L})_2]$, and very rare six-coordinate iodoanions $[\text{InI}_4(\text{L-L})]^-$ ($\text{L-L} = o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$, $o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2$).^{8,14}

There has also been much recent interest in exploring the relative Lewis acidity among the Group 13 halides. DFT calculations have predicted that Lewis acidity generally falls $\text{AlCl}_3 > \text{AlBr}_3 \gg \text{AlI}_3$ and $\text{AlCl}_3 > \text{AlBr}_3 > \text{GaCl}_3 > \text{GaBr}_3$ (gas phase) (ref. 8,9,15 and references therein) and this is supported by experimental data, although in some cases intermolecular interactions or solvate molecules can mask these trends in the solid state.⁹ Here we report systematic studies of AlX_3 ($\text{X} = \text{Cl}$, Br or I) complexes with some diphosphine and diarsine ligands and detailed comparisons with the gallium(III) analogues.

Experimental section

All preparations were carried out under rigorously anhydrous conditions *via* a dry dinitrogen atmosphere and standard Schlenk and glove-box techniques. Anhydrous grade aluminium trihalides were obtained commercially (Aldrich) and used as received. The ligands were obtained commercially (Strem or Aldrich), apart from $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$ and $o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2$ which were made by the literature methods.¹⁶ Solvents were dried by distillation from CaH_2 (CH_2Cl_2 , MeCN) or sodium benzophenone ketyl (hexane, toluene). IR spectra were recorded as Nujol mulls between CsI plates using a Perkin Elmer Spectrum 100 spectrometer over the range $4000\text{--}200\text{ cm}^{-1}$. ^1H and $^{31}\text{P}\{^1\text{H}\}$ NMR spectra were recorded using a Bruker AV300 or DPX400 spectrometer and referenced to the residual solvent resonance and external $85\% \text{H}_3\text{PO}_4$ respectively. ^{27}Al NMR spectra were recorded with a Bruker DPX400 spectrometer and referenced to external $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$. Microanalytical measurements were performed by Medac Ltd or London Metropolitan University.

trans- $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlCl}_4]$

To a suspension of AlCl_3 (0.067 g, 0.51 mmol) in CH_2Cl_2 (5 mL) was added $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$ (0.100 g, 0.51 mmol) in CH_2Cl_2 (5 mL) to form a colourless solution, which was allowed to stir for 1.5 h. The solvent volume was reduced to about 3 mL *in vacuo*. The resulting white precipitate was isolated by filtration and dried *in vacuo* to give a white powder. Yield: 0.071 g (43%). Small colourless blocks suitable for

single crystal X-ray diffraction study were grown from a CH_2Cl_2 solution containing AlCl_3 and $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$ in a 2:1 mol ratio kept at $-18\text{ }^\circ\text{C}$. Anal. Calc. for $\text{C}_{20}\text{H}_{32}\text{Al}_2\text{Cl}_6\text{P}_4$: C, 36.2; H, 4.9. Found: C, 36.0; H, 4.8%. ^1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.74$ (s, [24H], CH_3), 7.67–7.77 (m, [8H], C_6H_4). $^{31}\text{P}\{^1\text{H}\}$ NMR ($\text{CH}_2\text{Cl}_2\text{-CD}_2\text{Cl}_2$, 295 K): $\delta = -42.8$ (sextet, $^1J_{\text{PAI}} = 155$ Hz); (253 K): -42.6 (sextet). ^{27}Al NMR ($\text{CH}_2\text{Cl}_2\text{-CD}_2\text{Cl}_2$, 295 K): $\delta = 103.3$ (s, AlCl_4^-), 0.7 (quintet, $^1J_{\text{AIP}} = 155$ Hz); (253 K): 103.4 (s), 0.4 (quintet). IR (Nujol): $\nu = 488$ (vs, AlCl_4^-), 410 (s, AlCl) cm^{-1} .

trans- $[\text{AlBr}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlBr}_4]$

AlBr_3 (0.135 g, 0.51 mmol) was dissolved in toluene (5 mL) to form a yellow solution. To this was added $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$ (0.100 g, 0.51 mmol) in toluene (5 mL) which immediately led to the solution turning colourless, with evidence of a large amount of white precipitate. The reaction was stirred for 2 h, then the white powder was isolated by filtration and dried *in vacuo*. Yield: 0.213 g (91%). Anal. Calc. for $\text{C}_{20}\text{H}_{32}\text{Al}_2\text{Br}_6\text{P}_4$: C, 25.8; H, 3.5. Found: C, 26.0; H, 3.3%. ^1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.80$ (s, [24H], CH_3), 7.71–7.78 (m, [8H], C_6H_4). $^{31}\text{P}\{^1\text{H}\}$ NMR ($\text{CH}_2\text{Cl}_2\text{-CD}_2\text{Cl}_2$, 295 K): $\delta = -42.6$ (br s); (223 K): -42.6 (br s). ^{27}Al NMR ($\text{CH}_2\text{Cl}_2\text{-CD}_2\text{Cl}_2$, 295 K): $\delta = 80.5$ (s, AlBr_4^-), -12.0 (br s); (253 K): 80.5 (s), -11.4 (br s). IR (Nujol): $\nu = 398$ (vs, AlBr_4^-), 326 (m, AlBr) cm^{-1} .

trans- $[\text{AlI}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlI}_4]$

This was made similarly from AlI_3 (0.166 g, 0.41 mmol) and $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$ (0.080 g, 0.41 mmol) in toluene (10 mL). Yield: 0.231 g (94%). Anal. Calc. for $\text{C}_{20}\text{H}_{32}\text{Al}_2\text{I}_6\text{P}_4$: C, 19.8; H, 2.7. Found: C, 20.0; H, 2.8%. ^1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.90$ (s, [24H], CH_3), 7.79–7.89 (m, [8H], C_6H_4). $^{31}\text{P}\{^1\text{H}\}$ NMR ($\text{CH}_2\text{Cl}_2\text{-CD}_2\text{Cl}_2$, 295 K): $\delta = -42.3$ (br s); (223 K): -42.2 (br s). ^{27}Al NMR ($\text{CH}_2\text{Cl}_2\text{-CD}_2\text{Cl}_2$, 295 K): $\delta = -26.5$ (s, AlI_4^-); (223 K): -23.9 (s). IR (Nujol): $\nu = 334$ (vs, AlI_4^-) 281 (m, AlI) cm^{-1} .

trans- $[\text{AlCl}_2\{\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2\}_2][\text{AlCl}_4]$

To a suspension of AlCl_3 (0.090 g, 0.67 mmol) in CH_2Cl_2 (5 mL) was added $\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2$ (0.100 g, 0.67 mmol) in CH_2Cl_2 (5 mL). The resulting solution was stirred for 2 h, then the solvent volume was reduced to about 2 mL *in vacuo* and layered with hexane (2.5 mL), whereupon small colourless blocks suitable for single crystal X-ray diffraction study grew. The crystalline material was then isolated by filtration and dried *in vacuo* to give a white solid. Yield: 0.145 g (77%). Anal. Calc. for $\text{C}_{12}\text{H}_{32}\text{Al}_2\text{Cl}_6\text{P}_4$: C, 25.4; H, 5.7. Found: C, 25.2; H, 5.9%. ^1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.44$ (br s, [24H], CH_3), 2.00 (br s, [8H], CH_2). $^{31}\text{P}\{^1\text{H}\}$ NMR (CD_2Cl_2 , 295 K): $\delta = -40.6$ (sextet, $^1J_{\text{PAI}} = 164$ Hz). ^{27}Al NMR ($\text{CH}_2\text{Cl}_2\text{-CD}_2\text{Cl}_2$, 295 K): $\delta = 103.3$ (s, AlCl_4^-), 1.2 (quintet, $^1J_{\text{AIP}} = 164$ Hz). IR (Nujol): $\nu = 478$ (vs, br, AlCl_4^-), 377 (s, AlCl) cm^{-1} .

trans- $[\text{AlBr}_2\{\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2\}_2][\text{AlBr}_4]$

AlBr_3 (0.133 g, 0.50 mmol) was dissolved in toluene (5 mL) to form a yellow solution. To this was added $\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2$ (0.075 g, 0.50 mmol) in toluene (5 mL) which caused the immediate precipitation of a white solid. The reaction was



stirred for 1 h, then the white powder was isolated by filtration and dried *in vacuo*. Yield: 0.176 g (85%). Anal. Calc. for $C_{12}H_{32}Al_2Br_6P_4$: C, 17.3; H, 3.9. Found: C, 17.2; H, 3.9%. 1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.52$ (br s, [24H], CH_3), 2.04 (br s, [8H], CH_2). $^{31}P\{^1H\}$ NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = -38.0$ (br s). ^{27}Al NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = 80.4$ (s, $AlBr_4^-$), -12.3 (br s). IR (Nujol): $\nu = 394$ (vs, $AlBr_4^-$), 326 (m, $AlBr$) cm^{-1} .

trans-[$AlI_2\{Me_2P(CH_2)_2Me_2\}_2][AlI_4]$

AlI_3 (0.204 g, 0.50 mmol) was dissolved in toluene (5 mL) to form a yellow solution. To this was added $Me_2P(CH_2)_2PMe_2$ (0.078 g, 0.51 mmol) in toluene (5 mL) which caused the immediate precipitation of a white solid. The reaction was stirred for 2 h, then the white powder was isolated by filtration and dried *in vacuo*. Yield: 0.270 g (97%). Anal. Calc. for $C_{12}H_{32}Al_2I_6P_4$: C, 12.9; H, 2.9. Found: C, 13.0; H, 3.0%. 1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.63$ (br s, [24H], CH_3), 2.09 (br s, [8H], CH_2). $^{31}P\{^1H\}$ NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = -39.1$ (br s); (203 K): -37.7 (br s). ^{27}Al NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = -27.1$ (s, AlI_4^-), -33.9 (br s); (203 K): -26.1 (s). IR (Nujol): $\nu = 337$ (vs, AlI_4^-), 285 (s, AlI) cm^{-1} .

$[(AlCl_3)_2\{\mu-Me_2P(CH_2)_2PMe_2\}]$

To a suspension of $AlCl_3$ (0.178 g, 1.33 mmol) in CH_2Cl_2 (5 mL) was added $Me_2P(CH_2)_2PMe_2$ (0.099 g, 0.66 mmol) in CH_2Cl_2 (5 mL). The resulting solution was stirred for 1 h, then the solvent volume was reduced to about 4 mL *in vacuo* whereupon a white solid precipitated out. The solid was isolated by filtration and dried *in vacuo*. Yield: 0.055 g. The solid is a mixture of *trans*-[$AlCl_2\{Me_2P(CH_2)_2Me_2\}_2][AlCl_4]$ and $[(AlCl_3)_2\{\mu-Me_2P(CH_2)_2PMe_2\}]$. Small colourless blocks suitable for single crystal X-ray diffraction study were grown from a CH_2Cl_2 solution containing $AlCl_3$ and $Me_2P(CH_2)_2PMe_2$ in a 1.5 : 1 ratio kept at -18 °C. *Spectroscopic data for this complex obtained from the mixture, the resonances of the trans-[$AlCl_2\{Me_2P(CH_2)_2Me_2\}_2][AlCl_4]$ present were identical to those listed above:* 1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.90$ (m, [12H], CH_3), 2.17 (d, [4H], CH_2). $^{31}P\{^1H\}$ NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = -31.6$ (br s). ^{27}Al NMR (CH_2Cl_2 - CD_2Cl_2 , 298 K): $\delta = 109.3$ (br s); (253 K): 108.9 (s).

$[AlCl_2\{o-C_6H_4(PPh_2)_2\}][AlCl_4]$

To a solution of *o*- $C_6H_4(PPh_2)_2$ (0.075 g, 0.17 mmol) in CH_2Cl_2 (10 mL) was added $AlCl_3$ (0.045 g, 0.34 mmol). The resulting solution was stirred for 1.5 h, then the solvent volume was reduced to about 3 mL *in vacuo* and hexane (3 mL) was added. The resulting white precipitate was isolated by filtration and dried *in vacuo*. Yield = 0.035 g (29%). Anal. Calc. for $C_{30}H_{24}Al_2Cl_6P_2$: C, 50.5; H, 3.4. Found: C, 50.5; H, 3.7%. 1H NMR (CD_2Cl_2 , 295 K): $\delta = 7.22$ - 7.53 (aromatics). $^{31}P\{^1H\}$ NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = -8.3$ (v br); (180 K): -12.4 (v br). ^{27}Al NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = 103.3$ (s, $AlCl_4^-$). IR (Nujol): $\nu = 483$ (br, s, $AlCl_4^-$) cm^{-1} . Some small crystals of $[AlCl_2\{o-C_6H_4(PPh_2)_2\}][AlCl_4]$ were obtained from a similar preparation by careful hexane layering.

$[(AlCl_3)_2\{\mu-Cy_2P(CH_2)_2PCy_2\}]$

To a suspension of $AlCl_3$ (0.063 g, 0.47 mmol) in CH_2Cl_2 (5 mL) was added a solution of $Cy_2P(CH_2)_2PCy_2$ (0.101 g, 0.24 mmol) in CH_2Cl_2 (5 mL). The resulting solution was stirred for 1 h, then the solvent volume was reduced to about 3 mL *in vacuo*. Small colourless blocks suitable for single crystal X-ray diffraction study grew upon cooling the solution to -18 °C. Anal. Calc. for $C_{26}H_{48}Al_2Cl_6P_2$: C, 45.3; H, 7.0. Found: C, 45.1; H, 7.0%. 1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.32$ - 1.89 (m [44H]), 2.53 (m, [4H]), $^2J_{PH} = 13$ Hz). $^{31}P\{^1H\}$ NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = -0.5$ (s); (213 K): -1.4 (br). ^{27}Al NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = 110.6$ (s); (213 K): 111.3 (s). IR (Nujol): $\nu = 478$ (vs, br), 375 (w, br, $AlCl$) cm^{-1} .

$[AlCl_3(PMe_3)]$

To a suspension of $AlCl_3$ (0.177 g, 1.33 mmol) in CH_2Cl_2 (5 mL) was added PMe_3 (0.100 g, 1.31 mmol) in CH_2Cl_2 (5 mL). The resulting colourless solution was stored at -18 °C and over a few days large, colourless, temperature-sensitive crystals formed. These were separated and dried. The solvent was then removed *in vacuo* and the resulting white solid was washed with hexane (5 mL), and dried *in vacuo*. Yield: 0.174 g (63%). Anal. Calc. for $C_3H_9AlCl_3P$: C, 17.2; H, 4.3. Found: C, 17.2; H, 4.4%. 1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.50$ (d, $^2J_{HP} = 10$ Hz, CH_3). $^{31}P\{^1H\}$ NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = -42.6$ (br s); (248 K): -42.3 (sextet); (223 K): -42.3 (sextet, $^1J_{PAI} = 275$ Hz). ^{27}Al NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = 108.8$ (s); (223 K): 111.4 (d, $^1J_{AIP} = 275$ Hz). IR (Nujol): $\nu = 482$ (vs, br, $AlCl$) cm^{-1} .

trans-[$AlCl_3(PMe_3)_2]$

To a suspension of $AlCl_3$ (0.133 g, 1.00 mmol) in CH_2Cl_2 (5 mL) was added PMe_3 (0.154 g, 2.02 mmol) in CH_2Cl_2 (5 mL). The resulting solution was stirred for 1.5 h, then the solvent was removed *in vacuo* and the resulting white solid was washed with hexane (6 mL). The hexane was decanted off and the white powder was dried *in vacuo*. Yield: 0.191 g (67%). Small colourless crystals suitable for single crystal X-ray diffraction study were grown from a concentrated CH_2Cl_2 reaction solution kept at -18 °C. 1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.28$ (d, $^2J_{HP} = 8.0$ Hz, CH_3). $^{31}P\{^1H\}$ NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = -37.9$ (s); (253 K): -38.6 (br s); (183 K): -39.7 (broad, ill-defined coupling). ^{27}Al NMR (CH_2Cl_2 - CD_2Cl_2 , 295 K): $\delta = 66.0$ (s); (253 K): 61.3 (s). IR (Nujol): $\nu = 488$ (vs, br, $AlCl$) cm^{-1} .

$[AlBr_3(PMe_3)]$

To a yellow solution of $AlBr_3$ (0.175 g, 0.66 mmol) in toluene (3 mL) was added PMe_3 (0.050 g, 0.66 mmol) in toluene (3 mL). The resulting colourless solution was stirred for 1.5 h, then the solvent was removed *in vacuo* to yield a white powder, which was dried *in vacuo*. Yield: 0.186 g (83%). Anal. Calc. for $C_3H_9AlBr_3P$: C, 10.5; H, 2.6. Found: C, 10.6; H, 2.6%. 1H NMR (CD_2Cl_2 , 295 K): $\delta = 1.39$ (d, $^2J_{HP} = 9.6$ Hz, CH_3). $^{31}P\{^1H\}$ NMR (toluene- d_8 toluene, 295 K): $\delta = -40.9$ (sextet, $^1J_{AIP} = 242$ Hz); (CH_2Cl_2 - CD_2Cl_2 , 193 K): $\delta = -37.1$ (sextet, $^1J_{PAI} = 259$ Hz). ^{27}Al NMR (toluene- d_8 toluene, 295 K): $\delta = 101.5$ (d, $^1J_{AIP} = 242$ Hz);



(CH₂Cl₂-CD₂Cl₂, 193 K): $\delta = 102.7$ (d, $^1J_{\text{AlP}} = 258$ Hz). IR (Nujol): $\nu = 414$ (m), 393 (s, AlBr) cm⁻¹.

[AlI₃(PMe₃)]

This was made similarly from AlI₃ (0.282 g, 0.69 mmol) and PMe₃ (0.053 g, 0.69 mmol) in toluene (8 mL). Yield: 0.264 g (78%). Anal. Calc. for C₃H₉AlI₃P: C, 7.4; H, 1.9. Found: C, 7.5; H, 2.0%. ¹H NMR (CD₂Cl₂, 295 K): $\delta = 1.45$ (d, $^2J_{\text{HP}} = 10.0$ Hz, CH₃). ³¹P{¹H} NMR (CH₂Cl₂-CD₂Cl₂, 295 K): $\delta = -42.8$ (br s); (193 K): -40.8 (sextet, $^1J_{\text{PAI}} = 205$ Hz). ²⁷Al NMR (CH₂Cl₂-CD₂Cl₂, 295 K): $\delta = 50.2$ (br s); (193 K): 52.6 (d, $^1J_{\text{AlP}} = 209$ Hz). IR (Nujol): $\nu = 367$ (s), 340 (s, AlI) cm⁻¹.

[AlBr₃(PMe₃)₂]

To a yellow solution of AlBr₃ (0.173 g, 0.65 mmol) in toluene (2 mL) was added PMe₃ (0.100 g, 1.31 mmol) in toluene (3 mL) which immediately led to the solution turning colourless, with formation of a large amount of white precipitate. The reaction was stirred for 45 min, then the solvent volume was reduced to about 1 mL *in vacuo*. A white powder was isolated by filtration and dried *in vacuo* for 15 min. Yield: 0.182 g (66%). Anal. Calc. for C₆H₁₈AlBr₃P₂: C, 17.2; H, 4.3. Found: C, 17.1; H, 4.4%. ¹H NMR (CD₂Cl₂, 295 K): $\delta = 1.25$ (d, $^2J_{\text{HP}} = 7.0$ Hz, CH₃). ³¹P{¹H} NMR (CH₂Cl₂-CD₂Cl₂, 295 K): $\delta = -37.4$ (br s); (223 K): -32.6 (br s); (193 K): -30.3 (s). ²⁷Al NMR (CH₂Cl₂-CD₂Cl₂, 295 K): $\delta = 44.5$ (br s); (193 K): no resonance. IR (Nujol): $\nu = 373$ (s, AlBr) cm⁻¹.

[AlI₃(PMe₃)₂]

This was made similarly from AlI₃ (0.135 g, 0.33 mmol) and PMe₃ (0.050 g, 0.66 mmol) in toluene (10 mL). Yield: 0.090 g (49%). Anal. Calc. for C₆H₁₈AlI₃P₂: C, 12.9; H, 3.2. Found: C, 12.8; H, 3.3%. ¹H NMR (CD₂Cl₂, 295 K): $\delta = 1.33$ (d, $^2J_{\text{HP}} = 9.2$ Hz, CH₃). ³¹P{¹H} NMR (CH₂Cl₂-CD₂Cl₂, 295 K): $\delta = -31.6$ (br s); (223 K): -28.1 (br s); (193 K): -27.2 (s). ²⁷Al NMR (CH₂Cl₂-CD₂Cl₂, 295 K): $\delta = 4.4$ (br s); (193 K): no resonance. IR (Nujol): $\nu = 324$ (s, AlI) cm⁻¹.

The two [AlI₃(AsR₃)] (R = Et or Ph)¹⁰ were made by reaction of a 1 : 1 molar ratio of AlI₃ and AsR₃ in toluene. [AlI₃(AsPh₃)]: ²⁷Al NMR (toluene-toluene-d⁸, 295 K): $\delta = 33.1$ (br s); (203 K): no resonance; [AlI₃(AsEt₃)]: ²⁷Al NMR (toluene-toluene-d⁸, 295 K): $\delta = 31.9$ (s); (243 K): 32.4 (s); (203 K): 31.4 (br s).

[AlCl₂{*o*-C₆H₄(AsMe₂)₂}] [AlCl₄]

To a suspension of AlCl₃ (0.093 g, 0.70 mmol) in toluene (5 mL) was added *o*-C₆H₄(AsMe₂)₂ (0.101 g, 0.35 mmol) in toluene (5 mL), resulting in a colourless solution with a white solid suspended in it. After the reaction was stirred for 1.5 h the solid was isolated by filtration, washed with hexane and dried *in vacuo* to give a sticky white solid. Anal. Calc. for C₁₀H₁₆Al₂As₂Cl₆: C, 21.7; H, 2.9. Found: C, 21.6; H, 3.1%. ¹H NMR (CD₂Cl₂, 295 K): $\delta = 1.76$ (s, [12H], CH₃), 7.62–7.70 (m, [4H], C₆H₄). ²⁷Al NMR (CH₂Cl₂-CD₂Cl₂, 295 K): $\delta = 103.2$ (s, AlCl₄⁻); (203 K): 103.2 (s). IR (Nujol): $\nu = 490$ (s, br, AlCl) cm⁻¹.

[*o*-C₆H₄(AsMe₂)₂(CH₂)] [AlCl₄]₂

To a suspension of AlCl₃ (0.094 g, 0.70 mmol) in CH₂Cl₂ (5 mL) was added *o*-C₆H₄(AsMe₂)₂ (0.100 g, 0.35 mmol) in CH₂Cl₂ (5 mL). Initially a colourless solution formed, with a white solid precipitating out after 15 min stirring. The reaction was stirred for 2.5 h and then allowed to sit overnight, whereupon small colourless crystals suitable for single crystal X-ray diffraction study grew. Over the period of a week the white precipitate slowly redissolved and more colourless crystals grew, which were isolated and dried *in vacuo* to give a white powder. Yield: 0.052 g (23%). Anal. Calc. for C₁₁H₁₈Al₂As₂Cl₈: C, 20.7; H, 2.8. Found: C, 20.6; H, 2.7%. ESI+ (CH₃CN): m/z 191.2 [M + CH₃CN]²⁺. ¹H NMR (CD₃CN, 295 K): $\delta = 2.43$ (s, [12H], CH₃), 3.43 (s, [2H], CH₂), 8.02–8.15 (m, [4H], C₆H₄). ²⁷Al NMR (CH₃CN-CD₃CN, 295 K): $\delta = 103.5$ (s, AlCl₄⁻). IR (Nujol): $\nu = 478$ (vs, br, AlCl₄⁻) cm⁻¹.

X-Ray experimental

Details of the crystallographic data collection and refinement parameters are given in Table 1. Crystals suitable for single crystal X-ray analysis were obtained as described above. Data collections used a Rigaku AFC12 goniometer equipped with an enhanced sensitivity (HG) Saturn724+ detector mounted at the window of an FR-E+ SuperBright molybdenum ($\lambda = 0.71073$ Å) rotating anode generator with VHF Varimax optics (70 μ m focus) with the crystal held at 100 K (N₂ cryostream). Structure solution and refinements were performed with either SHELX (S/L)97 or SHELX(S/L)2013¹⁷ and were straightforward, except where detailed below. H atoms bonded to C were placed in calculated positions using the default C–H distance and refined using a riding model. Distance restraints were used in [AlCl₃(PMe₃)] to stop hydrogen atoms disordering across the mirror plane. [*o*-C₆H₄(AsMe₂)₂(CH₂)] [AlCl₄]₂ was treated as a racemic twin with the batch scale of 0.241.

Results and discussion

All syntheses, manipulations and spectroscopic measurements on the aluminium pnictogen complexes require rigorous exclusion of moisture, which otherwise causes displacement of the phosphorus donor from the aluminium by water, and generation of phosphonium salts, [R₂P(H) \cap PR₂][AlX₄] or [R₂P(H) \cap PR₂(H)][AlX₄]₂. The latter are readily identified by ¹H, ³¹P{¹H} and ³¹P NMR spectroscopy and during the course of this work X-ray structures were obtained on several examples (see ESI†). The diarsine is very easily displaced from the aluminium by moisture, but protonates less readily. We also found that while CH₂Cl₂ can be used to prepare the complexes with AlCl₃, attempts to use CH₂Cl₂ for the synthesis of AlBr₃ or AlI₃ complexes result in fast Cl/X exchange and incorporation of substantial amounts of chloride. This arises from the ability of the aluminium halides to promote reaction with CH₂Cl₂, as seen in other systems.^{5,18} In these cases toluene was used for the syntheses. Once isolated, the *pure* complexes with AlBr₃ and AlI₃ react only slowly with CH₂Cl₂, and it remains the NMR




Table 1 X-Ray crystallographic data^a

Compound	[AlCl ₃ (PMe ₃) ₂]	[AlCl ₃ (PMe ₃) ₂] ₂	[AlCl ₃ { <i>o</i> -C ₆ H ₄ (PMe ₂) ₂ }][AlCl ₄]	[AlCl ₂ { <i>o</i> -C ₆ H ₄ (PPh ₂) ₂ }][AlCl ₄]
Formula	C ₃ H ₉ AlCl ₃ P	C ₆ H ₁₈ AlCl ₃ P ₂	C ₂₀ H ₃₂ Al ₂ Cl ₆ P ₄	C ₆₀ H ₁₀₈ Al ₂ Cl ₆ P ₄
<i>M</i>	209.40	285.47	663.00	1159.52
Crystal system	Monoclinic	Orthorhombic	Monoclinic	Triclinic
Space group	<i>P</i> 2 ₁ / <i>m</i> (no. 11)	<i>Pnma</i> (no. 62)	<i>C</i> 2/ <i>c</i> (no. 15)	<i>P</i> $\bar{1}$ (no. 2)
<i>a</i> /Å	6.414(3)	9.970(3)	13.278(4)	10.024(3)
<i>b</i> /Å	10.292(4)	10.599(4)	14.545(4)	11.752(4)
<i>c</i> /Å	7.649(4)	13.562(4)	16.554(5)	24.438(8)
<i>a</i> /°	90	90	90	81.926(14)
<i>β</i> /°	113.698(12)	90	104.126(6)	85.52(2)
<i>γ</i> /°	90	90	90	82.25(2)
<i>U</i> /Å ³	462.4(4)	1433.1(8)	3100.2(16)	2818.9(16)
<i>Z</i>	2	4	4	2
<i>μ</i> (Mo-K α)/mm ⁻¹	1.174	0.883	0.828	0.489
<i>F</i> (000)	212	592	1360	1192
Total no. reflns	4220	14433	14366	24430
Unique reflns	1113	1486	3544	12194
<i>R</i> _{int}	0.056	0.140	0.032	0.076
No. of params, restraints	44, 9	66, 0	151, 0	652, 0
<i>R</i> ₁ , <i>wR</i> ₂ [<i>I</i> > 2 σ (<i>I</i>)] ^b	0.052, 0.124	0.065, 0.087	0.030, 0.065	0.079, 0.159
<i>R</i> ₁ , <i>wR</i> ₂ (all data)	0.055, 0.125	0.088, 0.091	0.035, 0.067	0.135, 0.193
Compound	[(AlCl ₃) ₂ { <i>μ</i> -Me ₂ P(CH ₂) ₂ PMe ₂ }]	[AlCl ₃] ₂ { <i>μ</i> -C ₆ H ₄ (PMe ₂) ₂ }][AlCl ₄]	[AlCl ₃] ₂ { <i>μ</i> -C ₆ H ₄ (PMe ₂) ₂ }][AlCl ₄]	[<i>o</i> -C ₆ H ₄ (AsMe ₂) ₂ (CH ₂)][AlCl ₄] ₂
Formula	C ₆ H ₁₆ Al ₂ Cl ₆ P ₂	C ₁₂ H ₃₂ Al ₂ Cl ₆ P ₄	C ₂₆ H ₄₈ Al ₂ Cl ₆ P ₂	C ₁₁ H ₁₈ Al ₂ As ₂ Cl ₈
<i>M</i>	416.82	566.92	689.24	637.65
Crystal system	Monoclinic	Orthorhombic	Triclinic	Orthorhombic
Space group	<i>P</i> 2 ₁ / <i>c</i> (no. 14)	<i>Pbca</i> (no. 61)	<i>P</i> $\bar{1}$ (no. 2)	<i>Pnca</i> ₂₁ (no. 33)
<i>a</i> /Å	10.738(6)	19.631(5)	8.251(2)	14.7537(10)
<i>b</i> /Å	7.948(4)	16.157(4)	9.830(2)	14.9361(10)
<i>c</i> /Å	11.772(6)	34.831(8)	11.921(3)	10.9633(8)
<i>a</i> /°	90	90	113.905(8)	90
<i>β</i> /°	113.938(9)	90	91.329(6)	90
<i>γ</i> /°	90	90	103.390(7)	90
<i>U</i> /Å ³	918.2(8)	11048(5)	852.4(4)	2415.9(3)
<i>Z</i>	2	16	1	4
<i>μ</i> (Mo-K α)/mm ⁻¹	1.182	0.916	0.666	3.719
<i>F</i> (000)	420	4672	362	1248
Total no. reflns	10169	92232	8428	19941
Unique reflns	2093	10858	3884	4813
<i>R</i> _{int}	0.062	0.1169	0.114	0.049
No. of params, restraints	73, 0	449, 0	163, 0	213, 1
<i>R</i> ₁ , <i>wR</i> ₂ [<i>I</i> > 2 σ (<i>I</i>)] ^b	0.051, 0.076	0.0503, 0.0985	0.0859, 0.1751	0.0409, 0.0954
<i>R</i> ₁ , <i>wR</i> ₂ (all data)	0.065, 0.081	0.0871, 0.1148	0.1743, 0.2276	0.0499, 0.1002

^a Common items: *T* = 100 K; wavelength (Mo-K α) = 0.71073 Å; θ (max) = 27.5°; *R*₁ = $\sum ||F_o| - |F_c|| / \sum |F_o|$; *wR*₂ = $[\sum w(F_o - F_c)^2 / \sum wF_o^2]^{1/2}$.





Scheme 1 The types of Al(III) phosphine complexes formed.

solvent of choice since it is very weakly coordinating and freezes at 176 K, with data collected immediately from freshly prepared solutions.

Diphosphine complexes

The reaction of $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$ with AlCl_3 in CH_2Cl_2 , irrespective of the mol ratio of reactants used (1 : 1 or 1 : 2), formed white crystals with a 1 : 1 $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\text{-AlCl}_3$ stoichiometry (see Scheme 1). These were shown by the X-ray structure determination to be $\text{trans-}[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlCl}_4]$, containing a pseudo-octahedral cation (Fig. 1) and the familiar tetrahedral anion. Even with a four-fold excess of AlCl_3 , the $^{31}\text{P}\{^1\text{H}\}$ and ^{27}Al NMR spectra showed no evidence for the presence of a pseudo-tetrahedral cation, $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}]^+$. The crystals are isomorphous with $\text{trans-}[\text{GaCl}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{GaCl}_4]$,¹³ and the cell dimensions, $d(\text{M-P})$ ($\text{M} = \text{Al}$ or Ga), $\langle \text{Cl-M-P} \rangle$ and $\langle \text{P-M-P} \rangle$ are very similar, while $d(\text{M-Cl})$ is ~ 0.05 Å longer for $\text{M} = \text{Ga}$. We return to these comparisons later. The corresponding reactions with AlBr_3 and AlI_3 also formed only $[\text{AlX}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlX}_4]$. They precipitate as fine powders from toluene and slow reaction with CH_2Cl_2 (see above) precludes using this solvent for crystallization for the bromo and iodo complexes, although it was used satisfactorily in the less reactive gallium systems.¹³ The complexes are also decomposed by MeCN with partial displacement of the phosphine. The spectroscopy of the three complexes show that all contain




Fig. 1 The structure of the centrosymmetric cation $\text{trans-}[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2]^+$ showing the atom numbering scheme. Ellipsoids are drawn at the 50% probability level and H atoms are omitted for clarity. Symmetry operation: $a = -x, -y, 1 - z$. Selected bond lengths (Å) and angles ($^\circ$): $\text{Al1-Cl1} = 2.2692(7)$, $\text{Al1-P1} = 2.4745(8)$, $\text{Al1-P2} = 2.4832(7)$, $\text{Cl1-Al1-P1} = 87.703(14)$, $\text{Cl1-Al1-P2} = 87.46(3)$, $\text{P1-Al1-P2} = 81.415(13)$.

six-coordinate trans isomers; the first examples of six-coordination in aluminium phosphine complexes.⁸

The far-IR spectra contain strong peaks due to the t_2 mode of $[\text{AlX}_4]^-$ at 488 ($\text{X} = \text{Cl}$), 398 ($\text{X} = \text{Br}$) or 334 ($\text{X} = \text{I}$) cm^{-1} (ref. 19) and weaker features at lower energy assigned to the a_{2u} modes in the cations (with local D_{4h} symmetry) at 410 ($\text{X} = \text{Cl}$), 326 ($\text{X} = \text{Br}$) or 281 ($\text{X} = \text{I}$) cm^{-1} .



Table 2 Selected NMR spectroscopic data

Complex	$\delta(^{31}\text{P}\{^1\text{H}\})$ 295 K ^a ($^1J_{\text{Al-P}}/\text{Hz}$)	$\Delta = \delta(^{31}\text{P}_{\text{complex}} - \delta(^{31}\text{P}_{\text{ligand}})^a$ (295 K)	$\delta(^{27}\text{Al})^{a,b}$ (295 K)
$[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlCl}_4]$	-42.8 (sextet, 155)	+12.2	0.7 (quintet)
$[\text{AlBr}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlBr}_4]$	-42.6 (s)	+12.4	-12.0 (s)
$[\text{AlI}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlI}_4]$	-42.3 (s)	+12.7	—
$[\text{AlCl}_2\{\text{Me}_2\text{P}(\text{CH}_2)_2\text{Me}_2\}_2][\text{AlCl}_4]$	-40.6 (sextet, 164)	+6.4	1.2 (quintet)
$[\text{AlBr}_2\{\text{Me}_2\text{P}(\text{CH}_2)_2\text{Me}_2\}_2][\text{AlBr}_4]$	-38.0 (s)	+9	-12.3 (s)
$[\text{AlI}_2\{\text{Me}_2\text{P}(\text{CH}_2)_2\text{Me}_2\}_2][\text{AlI}_4]$	-39.1 (br s)	+8	-33.9 (br s)
$[(\text{AlCl}_3)_2\{\mu\text{-Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2\}_2]$	-31.6 (br s)	+15.4	109.3 (s)
$[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PPh}_2)_2\}_2][\text{AlCl}_4]$	-8.3 (v br)	~+5	—
$[(\text{AlCl}_3)_2\{\mu\text{-Cy}_2\text{P}(\text{CH}_2)_2\text{PCy}_2\}_2]$	-0.5 (br, s)	-2.7	110.6 (s)
$[\text{AlCl}_3(\text{PMe}_3)]$	-42.6 (br s)	+19.4	108.8 (s)
$[\text{AlCl}_3(\text{PMe}_3)_2]$	-37.9 (s)	+24	66.0 (s)
$[\text{AlBr}_3(\text{PMe}_3)]$	-40.9 (sextet, 240) ^c	+21	101.5 (d) ^c
$[\text{AlBr}_3(\text{PMe}_3)_2]$	-37.4 (s)	+24.5	44.5 (s)
$[\text{AlI}_3(\text{PMe}_3)]$	-42.8 (br s)	+19	50.2 (s)
$[\text{AlI}_3(\text{PMe}_3)_2]$	-31.6 (br s);	+30.5	4.4 (s)
$[\text{AlI}_3(\text{AsEt}_3)]$	—	—	31.9 (s) ^c
$[\text{AlI}_3(\text{AsPh}_3)]$	—	—	33.1 (s) ^c

^a $\text{CH}_2\text{Cl}_2\text{-CD}_2\text{Cl}_2$ except c., s = singlet, br s = broad singlet, v br = very broad. ^b ²⁷Al resonances of $[\text{AlX}_4]^-$ anions not listed; $\delta([\text{AlCl}_4]^-) = 103$, $\delta([\text{AlBr}_4]^-) = 80.5$, $\delta([\text{AlI}_4]^-) = -26.5$. ^c Toluene/d⁸ toluene.




Fig. 2 (a) ²⁷Al NMR of $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2]^+$ and (b) ³¹P{¹H} NMR of $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2]^+$ in CH_2Cl_2 at 295 K.

Their ¹H NMR spectra in CD_2Cl_2 at 295 K each show single broad $\delta(\text{Me})$ resonances without resolved $^2J_{\text{PH}}$ couplings, confirming that only the *trans* forms are present. Similarly, the ³¹P{¹H} NMR spectra of all three complexes show resonances with modest high frequency coordination shifts which differ very little with X (Table 2). For X = Cl, the ³¹P{¹H} resonance shows a well-defined six line pattern due to coupling to the quadrupolar ²⁷Al nucleus (²⁷Al, *I* = 5/2, 100%), but for X = Br the coupling is not clearly resolved, and for X = I only a broad singlet ($w_{1/2} \sim 500$ Hz) is present. The ²⁷Al NMR spectrum of *trans*- $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlCl}_4]$ shows a very sharp singlet at $\delta = 103.4$, due to $[\text{AlCl}_4]^-$ (ref. 20) and a binomial quintet at $\delta = +0.7$ from coupling to four equivalent P atoms in the cation (Fig. 2).

The NMR spectra are essentially invariant over the temperature range 295–233 K, in contrast to those of the corresponding thioether or selenoether complexes, which show fast dissociative ligand exchange is present at ambient temperatures.⁵ *trans*- $[\text{AlBr}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlBr}_4]$ similarly shows ²⁷Al resonances at $\delta = 80.5$ (s) assigned to $[\text{AlBr}_4]^-$ and a broad singlet at $\delta = -12.0$ (s) from the cation, whereas for *trans*- $[\text{AlI}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}_2][\text{AlI}_4]$ only the ²⁷Al resonance of the anion ($\delta = -26.5$ (s)) is observed. The loss of coupling and then the loss of the cation resonance along the series X = Cl → Br → I can be ascribed to increasing quadrupolar relaxation rates as the electric field gradients around the Al nucleus increase with the changing donor sets.



The reaction of AlCl_3 with the sterically bulkier $o\text{-C}_6\text{H}_4(\text{PPh}_2)_2$ in either toluene or CH_2Cl_2 using a variety of conditions and molar ratios, invariably produced white powders with an $\text{AlCl}_3\text{-}o\text{-C}_6\text{H}_4(\text{PPh}_2)_2$ stoichiometry of 2:1. These showed a strong band in the IR spectrum at 483 cm^{-1} indicative of $[\text{AlCl}_4]^-$, leading to the proposed composition, $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PPh}_2)_2\}][\text{AlCl}_4]$, containing a pseudo-tetrahedral cation. The ^{27}Al NMR spectrum shows only the sharp resonance for the anion over the range 295 to 180 K, whilst the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum (CH_2Cl_2 solution) at ambient temperatures is a very broad feature at $\delta \sim -8$. On cooling the solution, the latter drifts to low frequency, reaching $\delta \sim -12$ at 180 K, but remaining broad. Addition of $o\text{-C}_6\text{H}_4(\text{PPh}_2)_2$ to the solution shows a second resonance at $\delta = -13.3$ (295 K) which is the value for the free diphosphine, showing intermolecular exchange is slow on the NMR timescale at room temperature. Gallium(III) halides form only the $[\text{GaX}_2\{o\text{-C}_6\text{H}_4(\text{PPh}_2)_2\}][\text{GaX}_4]$ complexes with this ligand, irrespective of the reaction conditions or ratio of reagents used.^{13,21} The reason for the broad $^{31}\text{P}\{^1\text{H}\}$ NMR resonance in the aluminium complex is not entirely clear; the X-ray crystal structures of $[\text{SnCl}_2\{o\text{-C}_6\text{H}_4(\text{PPh}_2)_2\}]^{22}$ and $[\text{GeCl}_2\{o\text{-C}_6\text{H}_4(\text{PPh}_2)_2\}]^{23}$ show the diphosphine coordinated in a very asymmetric manner, probably best described as κ^1 , and these complexes also exhibit broad $^{31}\text{P}\{^1\text{H}\}$ NMR resonances at all temperatures 295–190 K, which were attributed to intramolecular exchange between the 'free' and coordinated phosphorus donor groups in the diphosphine. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of the aluminium complex also rules out other than occasional traces of the phosphonium species $[o\text{-C}_6\text{H}_4(\text{PPh}_2)(\text{PPh}_2\text{H})]^+$ or $[o\text{-C}_6\text{H}_4(\text{PPh}_2\text{H})_2]^{2+}$.²⁴ The preparation of $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PPh}_2)_2\}][\text{AlCl}_4]$ produced a small number of colourless crystals, which were found to be *trans*- $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PPh}_2)_2\}][\text{AlCl}_4]$, containing a pseudo-octahedral cation, although the bulk product was reproducibly $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PPh}_2)_2\}][\text{AlCl}_4]$, based upon microanalysis and spectroscopic data. We assume that the *trans*- $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PPh}_2)_2\}][\text{AlCl}_4]$ crystallised preferentially from a solution in which it was a minor constituent. The structure of the cation is shown in Fig. 3. The $d(\text{Al-P})$ are $\sim 0.1\text{ \AA}$ longer than those in *trans*- $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}]^+$, reflecting the relatively weaker donor power of the diphenylphosphino groups, and possible steric hindrance. All attempts to isolate the six-coordinate complex as a bulk product have failed, but its existence was very surprising given the presence of two very bulky ligands chelating to the small aluminium centre, and the absence of similar gallium complexes.

The reaction of AlX_3 ($\text{X} = \text{Cl}, \text{Br}$ or I) with $\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2$ in a 1:1 mol ratio produces *trans*- $[\text{AlX}_2\{\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2\}][\text{AlX}_4]$, which contrasts with the corresponding GaX_3 complexes of diphosphinoethanes which produce only the dimers $[\text{X}_2\text{Ga}\{\mu\text{-R}_2\text{P}(\text{CH}_2)_2\text{PR}_2\}]\text{GaX}_3$.¹³ The presence of six-coordinate cations and $[\text{AlX}_4]^-$ was confirmed by the X-ray structure of *trans*- $[\text{AlCl}_2\{\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2\}][\text{AlCl}_4]$ (Fig. 4).

The multinuclear NMR data obtained for the three *trans*- $[\text{AlX}_2\{\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2\}][\text{AlX}_4]$ (Table 2 and Experimental section) show these to be the only forms present in CH_2Cl_2




Fig. 3 The structure of the centrosymmetric Al1 centred cation in *trans*- $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{PPh}_2)_2\}_2][\text{AlCl}_4]$ showing the atom labelling scheme. The C_6 rings are numbered cyclically with only the *ipso*-carbon atoms labelled and H atoms are excluded for clarity. Ellipsoids are drawn at the 50% probability level. The other cation is similar. Symmetry operation: $a = 2 - x, 1 - y, 1 - z$. Selected bond lengths (\AA) and angles ($^\circ$): $\text{Al1-Cl1} = 2.2222(13)$, $\text{Al1-P1} = 2.5574(15)$, $\text{Al1-P2} = 2.5843(15)$, $\text{P1-Al1-P2} = 78.69(4)$, $\text{Cl1-Al1-P1} = 87.80(5)$, $\text{Cl1a-Al1-P1} = 92.20(5)$, $\text{Cl1-Al1-P2} = 87.13(5)$, $\text{Cl1a-Al1-P2} = 92.87(5)$.




Fig. 4 The structure of the cation in *trans*- $[\text{AlCl}_2\{\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2\}_2][\text{AlCl}_4]$ showing the atom labelling scheme and with H atoms excluded for clarity. Ellipsoids are drawn at the 50% probability level. The other cation is similar. Selected bond lengths (\AA) and angles ($^\circ$): $\text{Cl1-Al1} = 2.2911(16)$, $\text{Cl2-Al1} = 2.2959(16)$, $\text{P1-Al1} = 2.4929(16)$, $\text{P2-Al1} = 2.4773(16)$, $\text{P3-Al1} = 2.4863(16)$, $\text{P4-Al1} = 2.4899(16)$, $\text{P3-Al1-P4} = 83.56(5)$, $\text{P2-Al1-P1} = 83.51(5)$, $\text{Cl1-Al1-Cl2} = 179.83(8)$, $\text{Cl1-Al1-P2} = 91.96(6)$, $\text{Cl2-Al1-P2} = 87.88(5)$, $\text{Cl1-Al1-P1} = 89.50(5)$, $\text{Cl2-Al1-P1} = 90.44(5)$, $\text{Cl1-Al1-P3} = 89.03(5)$, $\text{Cl1-Al1-P4} = 91.35(5)$, $\text{Cl2-Al1-P4} = 88.81(5)$, $\text{Cl2-Al1-P3} = 91.02(5)$.

solution, with the chemical shifts and coupling patterns similar to those discussed above for the complexes of $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$. Notably, for *trans*- $[\text{AlI}_2\{\text{Me}_2\text{P}(\text{CH}_2)_2\text{PMe}_2\}_2][\text{AlI}_4]$



[AlI₄], the ²⁷Al NMR resonance of the cation, $\delta = -33.9$, was observed as a broad singlet, illustrating how the presence or absence of a resonance due to the quadrupolar nucleus varies with very small changes in the coordination environment, which clearly affect the electric field gradient and hence the relaxation rate. The solution NMR data showed no evidence for *cis*-isomers, whereas [AlX₂{MeS(CH₂)₂SMe}₂}⁺ show mixtures of *cis* and *trans* forms in solution at low temperatures, both of which were crystallographically characterised in *trans*-[AlCl₂{MeS(CH₂)₂SMe}₂][AlCl₄] and *cis*-[AlI₂{MeS(CH₂)₂SMe}₂][AlI₄].⁵

Reaction of Me₂P(CH₂)₂PMe₂ with AlCl₃ in a 1 : ≤2 mol ratio in CH₂Cl₂ or toluene resulted in a mixture containing varying amounts of *trans*-[AlCl₂{Me₂P(CH₂)₂PMe₂}_2][AlCl₄] and a second complex identified by a combination of ¹H, ³¹P{¹H} and ²⁷Al NMR spectroscopy as [Cl₃Al{μ-Me₂P(CH₂)₂PMe₂}-AlCl₃]. Despite many attempts, we have been unable to isolate a pure bulk sample of the diphosphine-bridged complex, and similar studies with AlBr₃ or AlI₃ showed that only the [AlX₂{Me₂P(CH₂)₂PMe₂}_2][AlX₄] type formed, even with excess AlX₃, although the solids containing excess aluminium halide were extremely moisture sensitive, and prone to forming [Me₂P(CH₂)₂PMe₂H][AlX₄]. A few colourless crystals formed from one reaction mixture containing AlCl₃-Me₂P(CH₂)₂PMe₂ in a 1.5 : 1 mol ratio were found to be of the bridged complex, thus confirming its identity (Fig. 5).

The molecule is centrosymmetric with rather shorter *d*(Al-P) and *d*(Al-Cl) than in the corresponding six-coordinate complexes described above, due to the decreased coordination number. Although not isomorphous, it is also generally similar to the structures reported¹³ for [(GaX₃)₂{Et₂P(CH₂)₂-PEt₂}] (X = Br or I). The formation of the ligand-bridged dimer in the chloride system alone shows that steric factors from the increased radius of the halide co-ligands are not the driving force, but increasing the cone angle at phosphorus would be expected to favour a lower coordination number at Al.




Fig. 5 The structure of centrosymmetric [(AlCl₃)₂{μ-Me₂P(CH₂)₂PMe₂}] showing the atom labelling scheme. Ellipsoids are drawn at the 50% probability level and H atoms are omitted for clarity. Symmetry operation: $a = -x, 1 - y, -z$. Selected bond lengths (Å) and angles (°): Al1–Cl2 = 2.1166(14), Al1–Cl3 = 2.1203(15), Al1–Cl1 = 2.1284(15), Al1–P1 = 2.4059(14), Cl2–Al1–Cl3 = 112.15(6), Cl2–Al1–Cl1 = 112.54(6), Cl3–Al1–Cl1 = 110.78(5), Cl2–Al1–P1 = 106.34(5), Cl3–Al1–P1 = 109.97(6), Cl1–Al1–P1 = 104.68(5).




Fig. 6 The structure of centrosymmetric [(AlCl₃)₂{μ-Cy₂P(CH₂)₂PCy₂}] showing the atom labelling scheme. Ellipsoids are drawn at the 50% probability level and H atoms are omitted for clarity. Symmetry operation: $a = 2 - x, 1 - y, 1 - z$. Selected bond lengths (Å) and angles (°): Al1–Cl3 = 2.114(3), Al1–Cl1 = 2.123(3), Al1–Cl2 = 2.129(3), Al1–P1 = 2.407(3), Cl3–Al1–Cl1 = 112.98(12), Cl3–Al1–Cl2 = 113.23(12), Cl1–Al1–Cl2 = 112.00(12), Cl3–Al1–P1 = 106.20(11), Cl1–Al1–P1 = 108.76(11), Cl2–Al1–P1 = 102.90(10).

To explore this we examined the AlCl₃ complexes of Et₂P(CH₂)₂PEt₂ and Cy₂P(CH₂)₂PCy₂. The former gave a similar mixture of six- and four-coordinate species to that described and these complexes were not studied further, but the more bulky cyclohexyl-substituted ligand gave only the dimer, [(AlCl₃)₂{μ-Cy₂P(CH₂)₂PCy₂}].

The [(AlCl₃)₂{μ-Cy₂P(CH₂)₂PCy₂}] is extremely sensitive and very readily decomposed by trace moisture, and quite quickly by standing in CH₂Cl₂ solution at room temperature, generating [AlCl₄][−] and phosphonium cations. The X-ray structure (Fig. 6) was obtained from a very small crystal, and confirms the centrosymmetric dimer. The *d*(Al-P), *d*(Al-Cl), <Cl–Al–Cl and <Cl–Al–P are similar to those in the dimer shown in Fig. 5, showing the bulky cyclohexyl groups do not exert any great steric effects at the four-coordinate aluminium centre.

The ²⁷Al NMR (CH₂Cl₂-CD₂Cl₂, 295 K) of this complex shows a singlet at $\delta = 110.6$, consistent with a tetrahedral aluminium centre, whilst the ³¹P{¹H} NMR is a singlet at $\delta = -0.5$. The coordination shift (Δ) for this complex is very small and negative (−0.3) which contrasts with the small positive values of Δ seen for the other diphosphine complexes. This observation parallels those observed in trimethylaluminium–phosphine complexes, where small phosphines exhibit small positive coordination shifts, but as the steric bulk of the phosphine increases the Δ values become negative, attributed to steric effects.²⁵

Monophosphine complexes

The reaction of AlCl₃ with PMe₃ in a 1 : 1 molar ratio in anhydrous CH₂Cl₂ forms four-coordinate [AlCl₃(PMe₃)], whilst using a 1 : ≥2 molar ratio produced five-coordinate *trans*-[AlCl₃(PMe₃)₂]. The X-ray structures of both were determined (Fig. 7 and 8) and reveal the expected C_{3v} and D_{3h} geometries respectively. The *d*(Al–Cl) and *d*(Al–P) distances are ~0.07 Å






Fig. 7 The structure of $[\text{AlCl}_3(\text{PMe}_3)]$ showing the atom labelling scheme. The molecule has a mirror plane. Ellipsoids are drawn at the 50% probability level and H atoms are omitted for clarity. Symmetry operation: $a = x, 1/2 - y, z$. Selected bond lengths (Å) and angles ($^\circ$): Al1–Cl1 = 2.1291(15), Al1–Cl2 = 2.130(2), Al1–P1 = 2.392(2), Cl1–Al1–Cl1a = 111.15(10), Cl1–Al1–Cl2 = 111.06(6), Cl1–Al1–P1 = 107.22(6), Cl2–Al1–P1 = 108.94(9).




Fig. 8 The structure of *trans*- $[\text{AlCl}_3(\text{PMe}_3)_2]$ showing the atom numbering scheme. The molecule lies on a mirror plane going through C4, P2, Al1, Cl2, P1, C1. Ellipsoids are drawn at the 50% probability level and H atoms are omitted for clarity. Symmetry operation: $a = x, 1/2 - y, z$. Selected bond lengths (Å) and angles ($^\circ$): Al1–Cl2 = 2.203(2), Al1–Cl1 = 2.2029(14), Al1–P1 = 2.452(2), Al1–P2 = 2.459(2), Cl2–Al1–Cl1 = 121.22(5), Cl1–Al1–Cl1 = 117.56(9), P1–Al1–P2 = 177.31(9), Cl2–Al1–P1 = 88.79(8), Cl1–Al1–P1 = 90.44(6), Cl2–Al1–P2 = 88.52(8), Cl1–Al1–P2 = 90.96(6).

longer in the five-coordinate complex, attributed to the increased coordination number. The $d(\text{Al}–\text{P})$ in *trans*- $[\text{AlCl}_3(\text{PMe}_3)_2]$ is also shorter (by 0.06 Å) than that in *trans*- $[\text{AlI}_3(\text{PET}_3)_2]$,¹² consistent with the rather weaker Lewis acidity expected for aluminium iodide.

The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of $[\text{AlCl}_3(\text{PMe}_3)]$ at 295 K in CH_2Cl_2 – CD_2Cl_2 exhibits a broad resonance at $\delta = -42.6$, which on cooling the solution, resolves into a six-line coupling pattern with $^1J_{\text{PAI}} = 275$ Hz. The corresponding ^{27}Al NMR spectrum at 295 K is a singlet at $\delta = 108.8$, which splits into a doublet with the same $^1J_{\text{PAI}}$ on cooling. In contrast, the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of $[\text{AlCl}_3(\text{PMe}_3)_2]$ shows ill-defined coupling

even at low temperatures, whilst the ^{27}Al NMR spectrum is a singlet at $\delta = 66.0$ (295 K). Examination of the $^{31}\text{P}\{^1\text{H}\}$ spectra of either complex in CH_2Cl_2 solution containing added PMe_3 , shows singlets at room temperature due to fast exchange between the ‘free’ and coordinated phosphine, but on cooling the exchange slows and separate resonances for the complex and PMe_3 are observed. Even with a large excess of PMe_3 there was no evidence for the formation of a six-coordinate species such as $[\text{AlCl}_3(\text{PMe}_3)_3]$ or $[\text{AlCl}_2(\text{PMe}_3)_4]^+$.

The corresponding reactions of PMe_3 with AlX_3 ($X = \text{Br}$ or I) were carried out in toluene, since in CH_2Cl_2 rapid formation of aluminium chloro-species was observed, as discussed. Depending on the $\text{PMe}_3 : \text{AlX}_3$ ratio used, either $[\text{AlX}_3(\text{PMe}_3)]$ or $[\text{AlX}_3(\text{PMe}_3)_2]$ were isolated as very moisture sensitive white powders. The pure complexes dissolve in CD_2Cl_2 and NMR spectra can be obtained from the freshly prepared solutions without significant Cl/X exchange. However, if some AlX_3 is present (or added to the solution) rapid halide scrambling occurs. The spectroscopic properties of the bromide and iodide complexes (Experimental section) are much as expected and the NMR spectra (Table 2) show systematic chemical shift changes with the aluminium coordination number and the halogen present.

Diarsine complexes

A series of arsine complexes $[\text{AlX}_3(\text{AsR}_3)]$ ($X = \text{Cl}, \text{Br}$ or I ; $R = \text{Me}, \text{Et}$ or Ph) have been described, along with the X-ray structures of $[\text{AlX}_3(\text{AsPh}_3)]$ ($X = \text{Cl}$ or I).¹⁰ These show the presence of the expected pseudo-tetrahedral molecules. In passing we note that ^{27}Al NMR spectra reported (obtained from *in situ* reaction mixtures in CH_2Cl_2) for $X = \text{Br}$ or I , show resonances in the regions typical of chloro-species ($\delta(^{27}\text{Al}) = 100$ – 112), indicating Cl/X exchange with the solvent had most likely occurred. To confirm this we obtained ^{27}Al NMR spectra for $[\text{AlI}_3(\text{AsR}_3)]$ prepared and recorded in toluene solution, which showed $\delta = 33.1$ ($R = \text{Ph}$) or $\delta = 31.9$ (s) ($R = \text{Et}$), similar to the values reported for the phosphine complexes (above). We also found that addition of AsEt_3 to a toluene solution of $[\text{AlI}_3(\text{AsEt}_3)]$ failed to generate any new resonance, indicating that a five-coordinate complex did not form, which is in contrast to the phosphine systems.

The reaction of AlCl_3 with the diarsine, $o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2$, in toluene solution gave a slightly sticky white solid with an $\text{AlCl}_3 : o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2$ composition of 2 : 1, which we formulate as $[\text{AlCl}_2\{o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2\}][\text{AlCl}_4]$ based upon IR and ^{27}Al NMR spectroscopic evidence for the presence of the anion. Storing the CH_2Cl_2 solution produced from reacting AlCl_3 with $o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2$ in a 2 : 1 molar ratio for a few days formed a modest yield (23%) of colourless crystals. These were identified as $[o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2(\text{CH}_2)][\text{AlCl}_4]_2$, in which the diarsine has been diquaternized by the solvent, yielding a dication with a CH_2 bridge between the arsenic centres. Spectroscopic data for the cation are reported in the Experimental section and the structure is shown in Fig. 9. The dimensions are unexceptional, but the formation of this species again shows that the reactivity of the aluminium halide systems is not limited to ligand






Fig. 9 The structure of the cation in $[o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2(\text{CH}_2)]^+$ showing the atom numbering scheme. Ellipsoids are drawn at the 50% probability level and H atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): As1–C2 = 1.907(6), As1–C5 = 1.910(5), As1–C1 = 1.910(7), As1–C11 = 1.968(7), As2–C3 = 1.906(6), As2–C10 = 1.907(5), As2–C4 = 1.910(7), As2–C11 = 1.969(7), As1–C11–As2 = 104.2(3).

coordination. The diphosphine analogue, $[o\text{-C}_6\text{H}_4(\text{PMe}_2)_2(\text{CH}_2)]^{2+}$ was obtained in (failed) attempts to make phosphine complexes of SnF_2 , and may be synthesised from the diphosphine and CH_2I_2 in toluene.²² A related diarsine cation with an oxido bridge, $[o\text{-C}_6\text{H}_4(\text{AsMe}_2\text{Cl})(\mu\text{-O})\text{AsMe}_2]^+$ is formed by hydrolysis of $[\text{VOCl}_2\{o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2\}_x]$ ($x = 1$ or 2) or $[\text{VOCl}_2\{o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2\}]$ in thf solution.²⁶

Comparisons with gallium systems

Gallium(III) halides form four-coordinate $[\text{GaX}_3(\text{PR}_3)_3]$, $[\text{X}_3\text{Ga}(\mu\text{-L-L})\text{GaX}_3]$ or $[\text{GaX}_2(\text{L-L})]^+$ with most phosphines and diphosphines (L-L), with six-coordination observed only with $o\text{-C}_6\text{H}_4(\text{PMe}_2)_2$, in the cations $[\text{GaX}_2\{o\text{-C}_6\text{H}_4(\text{PMe}_2)_2\}]^+$.^{8,13} In contrast, as described above, aluminium(III) halides can produce four-, five- or six-coordinate complexes depending upon the phosphine present. In p-block chemistry coordination numbers and geometries are controlled by a combination of steric and electronic effects, and the structural and composition data reported above, combined with literature data,^{8–13,21} should allow some elucidation of the factors involved. Considering steric effects first, the single bond covalent radii of Al(III) and Ga(III) are quoted in standard texts as very similar (~ 1.25 Å), although the ionic radius of Ga^{3+} is ~ 0.07 Å larger than that of Al^{3+} .²⁷ In a study of complexes with Group 16 donor ligands, we found that for comparable complexes, $d(\text{M-X})$ and $d(\text{M-E})$ ($\text{E} = \text{S}, \text{Se}, \text{Te}$) were very similar for $\text{M} = \text{Al}$ or Ga and typically ~ 0.2 Å longer for $\text{M} = \text{In}$.⁵ DFT calculations predict that the bond lengths involving neutral ligands for a fixed metal (Al, Ga or In) generally increase (for gas phase molecules) $\text{Cl} < \text{Br} < \text{I}$, reflecting decreasing Lewis acidity down the series.^{5,7–9} However, in some cases solid state effects (packing effects, intermolecular interactions, hydrogen

Table 3 Summary of the mean M–X bond lengths ($d(\text{M-X})$ in Å), with associated errors, in $[\text{MX}_4]^-$

$[\text{MX}_4]^-$	X = Cl	X = Br	X = I
M = Al	2.126 ± 0.018	2.289 ± 0.015	2.527 ± 0.012
M = Ga	2.166 ± 0.018	2.321 ± 0.016	2.544 ± 0.018
M = In	2.337 ± 0.022	2.484 ± 0.018	2.702 ± 0.014

bonding or interaction with solvent molecules) can mask this trend.⁹ In order to explore the issue of size between Al and Ga centres, we compared $d(\text{M-X})$ ($\text{X} = \text{Cl}, \text{Br}$ or I) for the four-coordinate $[\text{MX}_4]^-$ ions, with data taken from the Cambridge Structural Database.²⁸ The results are shown in Table 3 and as histograms in the ESI† and show that the $d(\text{M-X})$ are essentially the same for gallium and aluminium. The fact that aluminium(III) has the same covalent radius to gallium(III) allows us to rule out steric effects as an explanation for the ability of aluminium to achieve a higher coordination number than gallium in many of the phosphine systems described in the present work. The similar size of Al and Ga is explained as being due to the “3d block contraction”, *i.e.* the increased nuclear charge resulting from insertion of the 3d transition metals, not completely offset by the screening effect of the 3d electrons.

The trend to higher coordination numbers found for aluminium(III) is thus an electronic effect, resulting from higher Lewis acidity towards phosphine ligands, as predicted by the DFT studies.^{7,15}

Conclusions

The synthesis, structures and spectroscopic characteristics of a range of four-, five- and six-coordinate aluminium complexes with phosphine donor ligands have been described. The contrasts with gallium(III) phosphine chemistry have been highlighted and shown to be due to greater Lewis acidity of the aluminium centre in these systems, consistent with computational predictions.^{9,15} In contrast to many d-block metals where analogous phosphine and arsine ligands produce very similar chemistries, the affinity of AlX_3 for arsines is markedly less, and only four-coordinate complexes were identified. This discrimination is also seen in complexes of other light p-block elements; SiX_4 ($\text{X} = \text{Cl}, \text{Br}$ or I) form phosphine complexes but do react with AsMe_3 or $o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2$,²⁹ whilst GeCl_4 produces the structurally authenticated *trans*- $[\text{GeCl}_4(\text{AsR}_3)_2]$ ($\text{R} = \text{Me}$ or Et), but does not form a complex with $o\text{-C}_6\text{H}_4(\text{AsMe}_2)_2$.³⁰ The high reactivity of AlX_3 also contributes to the synthetic challenges in this area, with their ability to activate chloro-carbon solvents toward reaction at the P- or As-centres.

Acknowledgements

We thank EPSRC for support (EP/I033394/1) and Dr M. Webster for helpful discussions. The SCFED Project (<http://www.scfed.net>) is a multidisciplinary collaboration of



British universities investigating the fundamental and applied aspects of supercritical fluids.

References

- (a) S. Dagherne and S. Bellemin-Laponnaz, in *The group 13 metals aluminium, gallium, indium and thallium*, ed. S. Aldridge and A. J. Downs, Wiley, NY, 2011, p. 654; (b) J. A. Miller, in *The chemistry of aluminium, gallium, indium and thallium*, ed. A. J. Downs, Blackie, London, 1993, p. 372.
- (a) *Comprehensive organic synthesis*, ed. B. M. Tost, Pergamon, Oxford, 1991, ch. 3, vol. 2, p. 702; (b) T. Ooi and K. Maruoka, in *Lewis acids in organic synthesis*, ed. H. Yamamoto, Wiley VCH, Weinheim, Germany, 2000, p. 191.
- (a) G. H. Robinson, in *Comprehensive Coordination Chemistry II*, ed. J. A. McCleverty and T. J. Meyer, Elsevier, Oxford, 2004, vol. 3, p. 347; (b) S. Aldridge, in *The group 13 metals aluminium, gallium, indium and thallium*, ed. S. Aldridge and A. J. Downs, Wiley, NY, 2011, p. 75.
- S. L. Benjamin, W. Levason and G. Reid, *Chem. Soc. Rev.*, 2013, **42**, 1460.
- K. George, M. Jura, W. Levason, M. E. Light and G. Reid, *Dalton Trans.*, 2014, **43**, 3637.
- C. Gurnani, M. Jura, W. Levason, R. Ratnani, G. Reid and M. Webster, *Dalton Trans.*, 2008, 6274.
- K. George, M. Jura, W. Levason, M. E. Light, L. P. Ollivere and G. Reid, *Inorg. Chem.*, 2012, **51**, 2231.
- J. Burt, W. Levason and G. Reid, *Coord. Chem. Rev.*, 2014, **260**, 65.
- A. Y. Timoshkin, M. Bodensteiner, T. N. Sevastianova, A. S. Lisovenko, E. I. Davydova, M. Scheer, C. Grassi and A. V. Butlak, *Inorg. Chem.*, 2012, **51**, 11602.
- E. Conrad, J. Pickup, N. Burford, R. McDonald and M. J. Ferguson, *Can. J. Chem.*, 2010, **88**, 797.
- (a) G. Menard and D. W. Stephan, *J. Am. Chem. Soc.*, 2010, **132**, 1796; (b) R. L. Wells, A. T. McPhail, J. A. Laske and P. S. White, *Polyhedron*, 1994, **13**, 2737.
- A. Ecker and H. Schnoekel, *Z. Anorg. Allg. Chem.*, 1998, **624**, 813.
- F. Cheng, A. L. Hector, W. Levason, G. Reid, M. Webster and W. Zhang, *Inorg. Chem.*, 2007, **46**, 7215.
- F. Cheng, S. I. Friend, A. L. Hector, W. Levason, G. Reid, M. Webster and W. Zhang, *Inorg. Chem.*, 2008, **47**, 9691.
- F. Bessac and G. Frenking, *Inorg. Chem.*, 2006, **45**, 6956.
- (a) R. D. Feltham, A. Kasenally and R. S. Nyholm, *J. Organomet. Chem.*, 1967, **7**, 285; (b) E. P. Kyba, S. T. Liu and R. L. Harris, *Organometallics*, 1983, **2**, 1877.
- G. M. Sheldrick, *Acta Crystallogr., Sect. A: Fundam. Crystallogr.*, 2008, **64**, 112.
- (a) R. T. Tjahjanto, M. F. Peintinger, T. Bredow and J. Beck, *Eur. J. Inorg. Chem.*, 2112, 3625; (b) N. Burford, B. W. Royan, R. E. vH. Spence and T. S. Cameron, *J. Chem. Soc., Dalton Trans.*, 1990, 1521.
- K. Nakamoto, *Infrared and Raman spectra of inorganic and coordination compounds*, Wiley, NY, 4th edn, 1986.
- J. Mason, *Multinuclear NMR*, Plenum, NY, 1987.
- M. Sigl, A. Schier and H. Schmidbaur, *Eur. J. Inorg. Chem.*, 1998, 203.
- C. Gurnani, A. L. Hector, E. Jager, W. Levason, D. Pugh and G. Reid, *Dalton Trans.*, 2013, **42**, 8364.
- F. Cheng, A. L. Hector, W. Levason, G. Reid, M. Webster and W. Zhang, *Inorg. Chem.*, 2010, **49**, 752.
- (a) M. Sigl, A. Schier and H. Schmidbaur, *Z. Naturforsch., B: Chem. Sci.*, 1998, **53**, 1313; (b) M. Jura, W. Levason, E. Petts, G. Reid, M. Webster and W. Zhang, *Dalton Trans.*, 2010, 10264.
- A. R. Barron, *J. Chem. Soc., Dalton Trans.*, 1988, 3047.
- B. M. Gray, A. L. Hector, W. Levason, G. Reid, M. Webster, W. Zhang and M. Jura, *Polyhedron*, 2010, **29**, 1630.
- (a) M. J. Taylor and P. J. Brothers, in *The chemistry of aluminium, gallium, indium and thallium*, ed. A. J. Downs, Blackie, London, 1993, p. 111; (b) B. Cordero, V. Gomez, A. E. Platero-Prats, M. Reves, J. Echeverria, E. Cremades, F. Barragan and S. Alvarez, *Dalton Trans.*, 2008, 2832.
- Cambridge Structural Database (CSD version 5.34; latest update = May 2013), accessed on the 21st January 2014, <http://www.ccdc.cam.ac.uk>.
- W. Levason, D. Pugh and G. Reid, *Inorg. Chem.*, 2013, **52**, 5185.
- (a) M. F. Davis, W. Levason, G. Reid, M. Webster and W. Zhang, *Dalton Trans.*, 2008, 353; (b) S. M. Godfrey, I. Mushtaq and R. Pritchard, *Dalton Trans.*, 1999, 1319.

