

Author's Accepted Manuscript

Effects of acute ocean acidification on spatially-diverse polar pelagic foodwebs: insights from on-deck microcosms

GA Tarling, V Peck, P Ward, NS Ensor, E Achterberg, E Tynan, AJ Poulton, E Mitchell, MV Zubkov



www.elsevier.com/locate/dsr2

PII: S0967-0645(16)30017-0
DOI: <http://dx.doi.org/10.1016/j.dsr2.2016.02.008>
Reference: DSRII4006

To appear in: *Deep-Sea Research Part II*

Received date: 21 May 2015
Revised date: 7 January 2016
Accepted date: 5 February 2016

Cite this article as: GA Tarling, V Peck, P Ward, NS Ensor, E Achterberg, E Tynan, AJ Poulton, E Mitchell and MV Zubkov, Effects of acute ocean acidification on spatially-diverse polar pelagic foodwebs: insights from on-deck microcosms, *Deep-Sea Research Part II* <http://dx.doi.org/10.1016/j.dsr2.2016.02.008>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and a review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

Effects of acute ocean acidification on spatially-diverse polar pelagic foodwebs: insights from on-deck microcosms

Tarling GA₁, Peck V₁, Ward P₁, Ensor NS_{1,2}, Achterberg E_{3,4}, Tynan E₃, Poulton AJ₅, Mitchell E₆, Zubkov MV₅

1. British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Rd, Cambridge, CB3 0ET, UK
2. Marine Scotland Science, Marine Laboratory, 375 Victoria Road, Aberdeen, AB11 9DB, UK
3. Ocean and Earth Science, National Oceanography Centre Southampton, University of Southampton, Southampton SO14 3ZH, UK
4. GEOMAR Helmholtz Centre for Ocean Research, Kiel, 24148, Kiel, Germany
5. National Oceanography Centre, Waterfront Campus, European Way, Southampton, SO14 3ZH, UK
6. Scottish Association for Marine Science, Dunstaffnage Marine Laboratory, Oban, Argyll, PA37 1QA, UK

Abstract

The polar oceans are experiencing some of the largest levels of ocean acidification (OA) resulting from the uptake of anthropogenic carbon dioxide (CO₂). Our understanding of the impacts this is having on polar marine communities is mainly derived from studies of single species in laboratory conditions, while the consequences for food web interactions remain largely unknown. This study carried out experimental manipulations of natural pelagic communities at different high latitude sites in both the northern (Nordic Seas) and southern hemispheres (Scotia and Weddell Seas). The aim of this study was to identify more generic responses and greater experimental reproducibility through implementing a series of short term (4 day), multilevel (3 treatment) carbonate chemistry manipulation experiments on unfiltered natural surface ocean communities, including grazing copepods. The experiments were successfully executed at six different sites, covering a diverse range of environmental conditions and differing plankton community compositions. The study identified the interaction between copepods and dinoflagellate cell abundance to be significantly altered by elevated levels of dissolved CO₂ (pCO₂), with dinoflagellates decreasing relative to ambient conditions across all six experiments. A similar pattern was not observed in any other major

phytoplankton group. The patterns indicate that copepods show a stronger preference for dinoflagellates when in elevated $p\text{CO}_2$ conditions, demonstrating that changes in food quality and altered grazing selectivity may be a major consequence of ocean acidification. The study also found that transparent exopolymeric particles (TEP) generally increased when $p\text{CO}_2$ levels were elevated, but the response was dependent on the exact set of environmental conditions. Bacteria and nanoplankton showed a neutral response to elevated $p\text{CO}_2$ and there was no significant relationship between changes in bacterial or nanoplankton abundance and that of TEP concentrations. Overall, the study illustrated that, although some similar responses exist, these contrasting high latitude surface ocean communities are likely to show different responses to the onset of elevated $p\text{CO}_2$.

Key Words: Arctic, Southern Ocean, copepod, phytoplankton, dinoflagellates, bacteria, nanoplankton, transparent exopolymeric particles, $p\text{CO}_2$

Introduction

The on-going shift in seawater carbonate chemistry and related reduction of ocean pH (termed ocean acidification, OA) caused by increasing atmospheric carbon dioxide (CO_2) levels has the potential to affect a wide range of marine organisms (Fabry et al. 2009, Cooley et al. 2009). Taxa that inhabit oceanic surface layers are particularly at risk since that is where the greatest changes are being observed. Some taxa respond sensitively to elevated CO_2 levels, whereas the response in others is mixed and often highly species-specific (Engel et al. 2008, de Nooijer et al. 2009, Doney et al. 2012, Beaufort et al. 2011, Whiteley 2011). Our present understanding of OA impacts is largely limited to single species responses while the consequences for food web interactions remain relatively unknown (Rossoll et al. 2012). For instance, the response at the community level may dampen the overall impact to elevated CO_2 levels through altered physiological performance, prey switching and species replacement (Rossoll et al. 2013).

The solubility of CO₂ increases with decreasing temperatures, such that polar oceans have naturally higher CO₂ and correspondingly lower carbonate ion concentrations (Sabine et al. 2004). With a low buffer capacity and a relatively low starting baseline, polar oceans are particularly susceptible to ocean acidification and are expected to be the first to experience carbonate mineral undersaturation and large decreases in pH (McNeil and Matear 2008, Steinacher et al., 2009). As a result, the polar regions, and particularly the Arctic, have been a focus for studies into the effects of projected OA on the marine environment (Riebesell et al. 2013).

Most studies of the effect of OA on pelagic organisms have involved laboratory and mesocosm experiments. Laboratory monoculture experiments allow experimental conditions to be tightly controlled and are valuable at identifying possible mechanisms and impacts. However, the outcomes of these experiments cannot be fully realistic because species are studied in isolation from the natural ecosystem in which there may be shifts from more pH-sensitive to more pH-tolerant species. Furthermore, the experimental conditions themselves (e.g. nutrient levels) may not be representative of typical oceanic conditions. Some of these shortcomings are overcome in carrying out mesocosm experiments, where large volumes (50 to 75 m³) of natural planktonic communities are incubated in situ and manipulated with regards artificially altering pH, carbonate chemistry and nutrient levels (Riebesell et al. 2010). The logistical demands of such operations have, nevertheless, restricted their deployment to inshore regions which may not adequately cover the range of conditions and communities found in the open ocean.

To date, the main approach to considering planktonic community responses to OA in open-ocean environments has been through short-term (1 to 14 d) incubations of natural communities on scientific vessels using on-deck microcosms (1 to 50 L; Riebesell et al. 2000, Tortell et al. 2002, Hutchins et al. 2007, Richier et al. 2014). Although the incubation volumes are relatively small compared to mesocosms, this approach does allow identical experiments to be repeated across natural gradients in carbonate chemistry, temperature and nutrients. The approach has the

advantage of examining a range of natural communities adapted to their particular environmental ranges. Comparisons can be made on how each community responds to the same stressor (i.e. changes in carbonate chemistry). The disadvantage of this approach is in the sudden induced change in pH and carbonate chemistry to which the community is given little time to adapt. Nevertheless, alternative longer-term, multigenerational approaches (Cripps et al. 2014) are best suited to studies of single species, and over timescales that are unfeasible for spatially extensive, vessel-based research. We used the short-term on-deck microcosm approach to consider the response of planktonic foodwebs, from a range of polar and sub-polar environments, to acute OA perturbations.

OA may affect food-webs in a number of ways, each with their own feedbacks to other parts of the system. Short-term acute-change experiments of this sort cannot characterise all such changes.

However, in performing identical experiments across a range of environmental starting conditions, they can identify where there are common responses to acute change, or otherwise, how responses vary according to the initial environmental conditions and community composition. In particular, we investigate two separate, but not mutually exclusive, hypotheses:

- i) that acute OA change will lead to the accumulation of transparent exopolymeric particles (TEP) which will stimulate the microbial food web;
 - ii) that acute OA change will alter the nutritional quality of individual phytoplankton species and lead to a change in zooplankton grazing preferences, so altering phytoplankton community composition and potentially zooplankton body condition.
- i) Stimulation of TEP production TEP are gel-like particles that form through coagulation of the polysaccharide fraction of dissolved organic matter (Passow, 2002). It has been proposed that increasing levels of $p\text{CO}_2$ in the surface ocean leads to more partitioning of the organic carbon fixed by marine primary production into the dissolved rather than the particulate fraction. This enhances the accumulation of dissolved organic carbon (DOC) in the surface ocean, and the concurrent accumulation of TEP. Such an accumulation was first noted by Engel (2002) in on-board experiments

under increased $p\text{CO}_2$, although whether the accumulation was due to increased production or modification of TEP structure remains uncertain (Weinbauer et al. 2011). Enhanced TEP has also been associated with nutrient stress (Engel 2002, Engel et al. 2004, Bellerby et al. 2008). Increased TEP aggregation can result in higher abundances and production of bacteria by attracting cells to a microbial hotspot (Simon et al. 2002). Nevertheless, contrasting relationships between TEP and bacteria have been found across diverse ocean areas (Corzo et al. 2005, Passow et al. 2001, Hung et al. 2003, Santschi et al. 2003, Passow and Alldredge 1994, Baskar and Bhosle 2006). We investigated if TEP production is affected by acute pH change and whether there is evidence of an enhancement of the microbial community as a result. Furthermore, we also considered whether any such patterns are altered by an additional trophic level (i.e. copepod grazers), given that copepods can directly feed on clusters of TEP (Passow and Alldredge 1999, Ling and Alldredge 2003).

ii) Changes to grazing preferences There is growing evidence that enhanced $p\text{CO}_2$ levels alter the nutritional quality of individual phytoplankton species available to grazers via species specific changes in their biochemistry (Riebesell and Tortell 2011). One major factor is taxonomic differences in the kinetic properties of carbon acquisition. Diatoms, for instance, have a much higher enzymatic specificity for carbon than dinoflagellates (Tortell 2000), which affects the rate of carbon enrichment per unit $p\text{CO}_2$. $p\text{CO}_2$ increase may also alter carbohydrate and fatty acid content in certain phytoplankton (Wynn-Edwards et al. 2014). Zooplankton have specific stoichiometric dietary requirements (Nobili et al. 2013) and an ability to select the prey species that suit their overall needs (Kleppel 1993). Therefore, this study will consider how zooplankton grazing preferences may alter between different acute OA treatments, and whether zooplankton body condition is affected.

We conducted short-term bioassay CO_2 perturbation experiments using natural seawater collected from the surface ocean and brought into a controlled laboratory on the deck of the research ship. This natural unfiltered seawater was manipulated to raise $p\text{CO}_2$ to future predicted levels, if at an acute rate. Locally-caught herbivorous copepods were subsequently introduced into certain incubation chambers to simulate natural food-web scenarios. The procedure was repeated in a

number of Arctic and Southern Ocean oceanic settings, with contrasting environmental characteristics. In so doing, the study examines both short-term physiological responses, through acute exposure to raised pCO₂ levels, alongside adaptive community variability, given that the communities being incubated were extracted directly from their natural setting. This means that responses will be dependent on initial community structure alongside community changes resulting from competition between the multiple genotypes present within the incubation. Overall, this study examines the effects of acute changes to CO₂ levels on real-world planktonic food-webs found in both polar oceans.

1. Methods

2.1 Sampling stations

This study was undertaken using on-board incubations of unfiltered seawater with and without additions of net-caught zooplankton as part of two open-ocean research campaigns. The research cruises were carried out on board RRS James Clark Ross in June-July 2012 in the sub-Arctic and Arctic Ocean (Cruise JR271) and January to February 2013 in the Southern Ocean (Cruise JR274). As part of these cruises, successful experimental incubations were performed in a total of six high-latitude environments (Table 1) with a diverse set of starting conditions. Two of the incubations were performed in Arctic waters, one in the open Greenland Sea (E03 271) and the other at the Greenland sea-ice margin (E04 271). A further two incubations were performed in the Southern Ocean, the first towards its northern limits close to South Georgia (E03 274), and the other, further south in the Weddell Sea, close to the South Sandwich Islands (E04 274). The remaining two incubations were carried out outside of true polar waters, one in the North Sea (E01 271) and the other in the sub-Arctic Barents Sea (E05 271).

2.2 Water collection

The incubations were set up with unfiltered seawater taken from the upper-mixed layer at the station location. On the day of the experimental setup, vertical CTD profiles were performed to characterise water column variables (temperature, salinity, chlorophyll fluorescence) and determine water column structure. The depth of water collection for the experimental setup was then decided upon based on these initial profiles and was typically from the middle of the upper mixed layer. The unfiltered water was collected from dedicated CTD casts using a titanium frame CTD and acid-cleaned 24 x 10 L Rosette OTE (Ocean Test Equipment) bottles. On deck, the bottles were immediately transferred to a class-100 filtered air environment within a trace-metal clean container to avoid contamination. The unfiltered water was transferred using acid-cleaned silicon tubing to 1 L acid-cleaned clear Duran bottles and the bottles checked for any extraneous copepods, which were removed before the bottles were sealed, pending carbonate chemistry manipulation and the addition of hand-picked live copepods from net-catches (see below). Subsamples were collected simultaneously for time-zero (T0) measurements of each of the variables to be measured over the subsequent time course (Table 2).

Further subsamples at time zero (T0) were taken directly from the CTD and immediately measured for total alkalinity (TA) and dissolved inorganic carbon (DIC) to characterise the carbonate chemistry system in seawater; details for the analytical methods can be found below. The remaining variables of the carbonate system were calculated with the CO2SYS programme (version 1.05, Lewis and Wallace, 1998; van Heuven et al., 2011), using the constants of Mehrbach et al. (1973) refitted by Dickson and Millero (1987). Carbonate chemistry in the experimental bottles was subsequently manipulated using equimolar additions of strong acid (HCl, 1 mol L⁻¹) and HCO₃⁻ (1 mol L⁻¹), as recommended by Gattuso et al. (2010) for increasing DIC at constant TA. The volumes of HCl and HCO₃⁻ required to adjust pCO₂ to the chosen target values (750 and 1000 µatm) were calculated from the measured ambient state of the carbonate system in seawater using CO2SYS.

2.3 Zooplankton collection

Immediately after the collection of water, a motion-compensated Bongo net (61 cm diameter, 100 μm and 200 μm mesh) was deployed between 0 and 200 m to capture live zooplankton. Sub-samples of the net catches were immediately analysed to determine the approximate species composition and decide on the most appropriate calanoid species (and developmental stage) to use for the incubation experiments. The choice of species varied between experimental stations, with *Calanus finmarchicus* being used in the North Sea (E01 271), *C. hyperboreus* in the Arctic (E03 271 and E04 271) and *Calanoides acutus* in the Southern Ocean (E03 274 and E04 274). The exact species identity of the calanoid used in the Barents Sea was unclear and classified as *Calanus finmarchicus*/*glacialis*. The chosen developmental stage was mainly copepodite stage V (CV), although, in E04 274, it was necessary to combine this stage with females because numbers were relatively low. Individual copepods were picked out from sub-samples analysed under a light microscope using titanium forceps (to minimise iron contamination) and acid washed pipettes and transferred to 10 mL glass vials containing filtered seawater and maintained at ambient sea-surface temperature. Each glass vial contained either 5 or 10 specimens, the number differing between stations to account for individual specimen size and in situ concentrations. A total of 18 glass vials were populated with copepods in readiness for their subsequent introduction into the incubation experimental bottles.

2.4 Set-up and running of experiments

In the trace-metal clean container, the 1 L Duran bottles containing the unfiltered sea-water were modified with varying aliquots of equimolar HCl and HCO_3^- to achieve nominal pCO_2 levels of either 750 μatm or 1000 μatm . Eight individual Duran bottles per pCO_2 level were prepared, while a further ten Duran bottles were left unmodified. This number allowed for two sets of triplicate bottles containing copepods in ambient conditions, 750 μatm pCO_2 and 1000 μatm pCO_2 , as well as two sets of single bottles without copepods for each pCO_2 treatment and two sets of double bottles without copepods in ambient conditions. Accordingly, the copepods in the glass vials were gently poured into

six of the eight modified bottles for each pCO₂ treatment, and six of the ten unmodified bottles. Each of the bottles was topped up to the rim and the lids sealed. The 26 bottles (i.e. 2 identical sets of 13 bottles) were placed on 3 plankton wheels within a controlled temperature container and maintained at ambient surface-water temperature (Fig. 2). Irradiance (100 μmol m⁻² s⁻¹) was provided by daylight simulation LED panels (Powerpax, UK) over a 24 h light cycle which approximated the ambient photoperiod and light intensity at 5 m (Richier et al. 2014).

The first set of 13 bottles was removed from the plankton wheels after 24 h and inspected to determine if there were any dead copepods in the incubation. The bottles were then immediately subsampled for the concentration of nutrients (including ammonium), TA and DIC and transparent exopolymeric particles (TEP) as well as the abundance of bacteria, nanoplankton, and major phytoplankton taxa. It was not possible to make all measurements on all bottles because of the volumes required for each analytical method. However, each measurement was made on at least one bottle with copepods and one without copepods at the three different pCO₂ levels (ambient, 750 μatm and 1000 μatm). The copepods were filtered out of the remaining water, rinsed in deionised water (Milli-Q, Millipore) and snap-frozen in liquid nitrogen for subsequent elemental composition analysis.

The remaining set of 13 bottles were checked for copepod mortalities at 48, 72 and 96 h after the start of the incubation. At the 96 h timepoint, the take-down procedure was repeated, with the exception that samples for bacteria, nanoplankton and phytoplankton abundance were not taken, given that they would most likely be subject to exponential growth and overgrazing effects respectively after this length of time in volumes of 1L.

2.5 Measurements of biological and chemical variables

Nutrients Samples for macronutrients (nitrate (NO₃⁻) + nitrite (NO₂⁻)), silicic acid (dSi) and phosphate (PO₃⁻) were collected directly from the incubation bottles into a 25 mL polystyrene container and

stored at 4°C pending analysis within 12 h. The samples were run on a Skalar San+Segmented Flow Autoanalyzer using colorimetric techniques (Kirkwood, 1996) with the exception that the flow rate of the sample through the phosphate channel was increased to improve reproducibility and peak shape.

For ammonium analysis, 10 mL samples were collected in polypropylene vials and reagent added, following Kerouel and Aminot (1997). Fluorometric analysis was performed 24 h after the addition of reagent. Concentrations were estimated to an accuracy of nanmoles L⁻¹.

Carbonate chemistry Samples for dissolved inorganic carbon (DIC) and total alkalinity (TA) were collected from the incubation bottles in 250 mL borosilicate glass bottles. TA and DIC measurements were performed using either a VINDTA 3C system (Marianda, Germany), an Apollo SciTech DIC analyser (AS-C3) or an Apollo SciTech Alkalinity Titrator (AS-ALK2), depending on availability of the instruments during the cruises. On the VINDTA 3C system, DIC was measured by coulometric titration and TA by potentiometric titration and calculated using a modified Gran plot approach (Bradshaw et al., 1981). The Apollo uses a CO₂ infrared detector (LICOR 7000) to measure DIC and an open cell titration to measure TA. We have routinely used the different instruments in our lab and have found no systematic differences between them, therefore the data is inter-comparable with no significant difference between the methods. Certified Reference Materials (CRMs) (A.G. Dickson, Scripps Institution of Oceanography, batch 117 in the Arctic, batches 119 and 120 in the Southern Ocean) were used at the beginning, middle and end of each run on all instruments as a control for both DIC and TA. Precision was taken as the standard deviation of repeated analysis of CRMs. For the VINDTA 3C instrument, this was $\pm 2 \mu\text{mol kg}^{-1}$ for TA and $\pm 3 \mu\text{mol kg}^{-1}$ for DIC. For the Apollo instrument, TA was $\pm 1.5 \mu\text{mol kg}^{-1}$ and DIC $\pm 3 \mu\text{mol kg}^{-1}$. The remaining variables of the carbonate system were calculated with the CO2SYS program (MATLAB version 1.1; Lewis and Wallace, 1998; van Heuven et al., 2011) using the carbonate equilibrium constants of Mehrbach et al. (1973)

refitted by Dickson and Millero (1987), the boric acid dissociation constants of Dickson (1990) and the ratio of total boron to salinity recommended by Lee et al. (2010).

TEP Samples of TEP were collected by filtration of 200 mL of seawater through 25 mm diameter polycarbonate filters (0.45 μm pore-size, Sterlitech) at constant vacuum (200 mBar). Three separate filtrations were carried out on each 1 L bottle that was analysed. The particles retained on the filters were stained with 500 μL of 0.02% aqueous solution of Alcian blue in 0.06% acetic acid (pH 2.5). The dye was pre-filtered using a polycarbonate filter (0.2 μm pore-size; Sterlitech) before use. Stained filters were rinsed once with deionised water (Milli-Q, Millipore) and then transferred into 15 mL polypropylene centrifuge tubes (Fisher Scientific) and stored at -20°C . Onshore, the amount of Alcian Blue adsorbed onto the filters was determined following a soak in 6 mL of 80% sulphuric acid for 2 h and determination of the absorbance of the resulting solution at 787 nm (absorption maximum) using a spectrophotometer (U-1800, Hitachi). The amount of Alcian blue in the solution was directly related to the weight of the polysaccharide that was retained on the filter (Passow and Alldredge, 1995).

Bacteria and nanoplankton For each sample, 1.6 mL of seawater was removed and placed into a 2 mL polypropylene screw cap vial containing 80 μL of paraformaldehyde (PFA, 1% final concentration). The vials were then placed in a fridge and left for no longer than 12 hours before being analysed by flow cytometry (BD FACSort instrument) to enumerate bacterioplankton and protist abundance (Marie et al. 1997, Zubkov et al. 2007). Samples were stained with SYBR green I nucleic acid dye for a minimum of 30 minutes, before being analysed. Samples were run at a low flow rate of 0.02 mL min^{-1} for 1 minute to determine bacterioplankton concentration and at a flow rate of 0.2 mL min^{-1} for 3 minutes to determine concentrations of phototrophic and heterotrophic protists. For each sample, a known volume of bead solution was added as an internal standard. The bead solution consisted of Fluoresbrite multicolour 0.5 μm and 1.0 μm beads dispersed in 400 mL of

sterile Milli-Q (Millipore) water. The bead concentration was calculated before use using the FACS caliber flow cytometer and a syringe pump (Zubkov and Burkhill 2006).

Phytoplankton composition and abundance 100 mL of seawater was taken from the incubation bottles and immediately preserved in a 2% final concentration of acidic Lugol's solution stored in 250 mL amber glass bottles. Phytoplankton (>10 μm) was identified and enumerated in 50 mL Hydro-Bios settling chambers on a SP-95-I inverted microscope, following Poulton et al. (2007).

Elemental analysis of copepods Total carbon, hydrogen and nitrogen of either single copepods or batches of 5 or 10 copepods were determined using a CE-440 Elemental Analyser (Exeter Analytical, Inc.). The copepods were either taken immediately from the net catch (T0 population) or post-incubation, rinsed in deionised water and placed in tin capsules before snap freezing in liquid nitrogen and maintained at -80°C for analysis on-shore. On-shore, the capsules were dried overnight at 50°C and then combusted at 1800°C in the Elemental Analyser to determine total carbon, hydrogen and nitrogen through analysis of the products of combustion by gas chromatography.

Elemental analysis of copepods from experiment E01 271 was not performed because of a processing error.

2.6 Statistical treatment

Differences between pCO_2 treatments (treatment effects) were analysed using a 1-way ANOVA provided that they first passed tests for normality (Kolmogorov-Smirnov test) and equal variance (variability about group means), else they were analysed non-parametrically with a 1-way ANOVA on ranks. Paired t-tests were used to determine the impact of the addition of copepods to incubations, subject to the same tests for normality and equal variance. Attempts to fit linear regressions were made to the relationship between the relative changes in TEP and either bacteria or nanoplankton, using the Sigmaplot 12.5 package (Sysstat Software Inc.) using a least-squared fitting method. Fits where the slope of the curve were not significantly different from 0 were otherwise considered as

trends of which the consistency was compared between treatments. The influence of copepods on the abundance of bacteria or nanoplankton was assessed through comparison of 95% confidence intervals between incubations with- and without copepods (Zar 1996).

2. Results

3.1 Starting conditions

Physical conditions Experiments at 4 of the 6 stations were carried out under true polar conditions where surface mixed-layer temperatures were between 2.2°C and -1.6°C (E03 271, E04 271, E03 274 and E04 274). Experiment E05 271 was initiated in the sub-Arctic Barents Sea, with surface mixed-layer temperatures at around 6.5°C. Experiment E01 271 was sited in the North Sea, where the temperature was 10.7°C (Table 2). Nutrient concentrations varied considerably, with phosphate and nitrate being at non-limiting concentrations in both of the Southern Ocean stations (E03 274 and E04 274) in contrast to the very low levels observed at the North Sea station (E01 271). This is reflected in the N:P atomic ratio, which was close to the Redfield ratio of 16 in the Southern Ocean, but as low as 0.6 in the North Sea. The Arctic and sub-Arctic stations showed varying degrees of nutrient limitation. Silicate was also abundant in the Southern Ocean stations compared to the other stations, with the lowest concentrations being observed in the North Sea and Barents Sea. pH was lowest at the South Georgia Southern Ocean station (E03 274) at 8.050, with a corresponding pCO₂ level of 384 µatm. All other stations had pH values ranging between 8.1 and 8.2 and pCO₂ levels in the region of 300 µatm.

Biological conditions Chl-a concentrations in the mixed layer reached as high as 4.2 µg L⁻¹ at the Weddell Sea station (E04 271), and 3.03 µg L⁻¹ at the Greenland sea-ice margin station (E04 271) but only 0.31 µg L⁻¹ in the North Sea (E01 271). There appeared to be little correspondence between levels of Chl-a and TEP across stations, with relatively high TEP levels occurring at stations where Chl-a was both low (E01 271) and moderately high (E05 271). TEP concentrations were comparatively

low in both Southern Ocean stations (E03 274, E04 274). The highest abundance of bacteria was observed in the open Greenland Sea (E03 271) and the Barents Sea (E05 271), at above 1 million cells mL⁻¹, while the Greenland sea-ice margin station (E04 271) contained the lowest numbers, at less than half of those levels of abundance. Nannoplankton abundance similarly was highest at stations E03 271 and E05 271, and were almost an order of magnitude lower at the Weddell Sea station (E04 274).

The phytoplankton community analysed from the Lugol's samples was numerically dominated by diatoms at stations in the North Sea (E01 271), the Greenland sea-ice margin station (E04 271) and in the Weddell Sea (E04 274, Fig. 2). Dinoflagellates were the dominant taxa in the Barents Sea (E05 271) while there was more of an even mix of phytoplankton taxa in the open Greenland Sea (E03 271) and near South Georgia (E03 274). Those stations dominated by diatoms had relatively high cell counts, while cell counts were low where dinoflagellates made up a large fraction of the phytoplankton community.

The choice of calanoid species for the incubations differed between stations, based on their relative availability in net-catch samples (Table 1). There was as much variability in the C:N mass ratio within as between species, indicating that body condition varied according to locality (Table 3, Fig. 2). Higher C:N mass ratios are considered to reflect better body condition in that such individuals contain greater amount of body reserves. *Calanus hyperboreus* CV in the open Greenland Sea (E03 271) had a median C:N mass ratio that was almost 30% higher than the same species and stage found at the Greenland sea-ice margin (E04 274). Similarly, *Calanoides acutus* (CV and female combined) in the Weddell Sea had C:N mass ratios that were at least double of those collected at South Georgia (CV only, E03 274). Therefore, the physiological states of the populations chosen as grazers in each set of incubations were diverse, reflecting different life-history and feeding trajectories up to the point of capture.

3.2 Outcome of manipulations on water chemistry

The influence of the equimolar additions of acid and base on nutrient and silicic acid levels and the carbonate chemistry is fully detailed in Appendices 1, 2 and 3. The manipulations were successful in increasing DIC levels whilst maintaining relatively constant levels of TA. However, in some instances, the $p\text{CO}_2$ target levels were not attained, with the measured levels of $p\text{CO}_2$ after the manipulations being below those intended. The effects of the additions on nutrient and silicic acid concentrations was negligible given the minute volumes that were added.

3.3 Response of TEP and the microbial community

TEP response After 96 hours, TEP concentrations increased in all of the control (ambient conditions, no copepods) incubations across all six sets of experiments (bottle effects, Table 4). The rate of increase varied between experiments, with the highest rate observed at the Barents Sea station (E05 271, $5.71 \mu\text{g xanthan equivalent L}^{-1} \text{h}^{-1}$) where the corresponding N:P atomic ratio was relatively low (5.5). The lowest rates occurred in the Southern Ocean stations (E03 274, E04 274, 0.1 to $0.4 \mu\text{g xanthan equivalent L}^{-1} \text{h}^{-1}$), where N:P was close to the Redfield ratio of 16 (Redfield 1958).

Relative to this tendency for TEP accumulation over time, the only experiment to show an increase above that otherwise expected after 24 h was E04 274 for the $1000 \mu\text{atm}$ treatment (treatment effects, Table 4). However, by 96 h, significant increases were also found in E01 271, E04 271 and E05 271. Only experiments E01 271 and E04 274 showed significant increases in both the $750 \mu\text{atm}$ and $1000 \mu\text{atm}$ treatments after 96 h, the others being limited to significant increases in the $1000 \mu\text{atm}$ treatments only.

The effects of the presence of copepods on TEP were mixed (copepod effects, Table 4). None of the experiments showed any significant increase or decrease in TEP above background increases by 24 h. By 96 h, four of the six experiments showed significant increases in TEP (E01 271, E03 271, E05 271 and E03 274), while one experiment showed a significant decrease (E04 271) and one no significant change (E04 274).

Bacterial response Compared to the variance in bacterial cell counts for the starting conditions (T0 samples), there was no significant change in the abundance of bacteria after 24 h either at ambient conditions or in the two pCO₂ treatments (Fig. 3). The presence or absence of copepods in the incubations had a significant impact on bacterial abundance in certain instances. In ambient conditions in E04 271 and in the 750 µatm treatment in E04 274, the presence of copepods significantly reduced the number of bacterial cells. However, bacterial numbers were significantly enhanced by the presence of copepods in the 1000 µatm treatments in experiments E05 271 and E03 274. For the majority of experiments however, the presence or absence of copepods had very little impact on bacterial numbers under ambient or increased pCO₂ levels.

Nannoplankton response There was similarly little pattern in the response of the nannoplankton community to incubation conditions (Fig. 4). No significant change in the abundance of nannoplankton cells after 24 h was detected in either ambient conditions or the two pCO₂ treatments compared to the variance observed in the T0 samples. The presence or absence of copepods in the incubations did not make a significant impact in the majority of instances. However in the 1000 µatm treatment of E01 271, and at ambient conditions in E04 274, there was a significant enhancement in nannoplankton abundance when copepods were present.

Relationship between TEP and microbial abundance In the control incubations (ambient conditions without the presence of copepods), there was a general negative trend between relative changes in TEP concentration and bacterial cell abundance, although the relationship was not significant (ANOVA, $F = 4.237$, $P = 0.109$, 5 df, Fig. 5). This trend was mainly driven by the two Southern Ocean experiments (E03 274, E04 274), where there was a relatively large increase in TEP in E03 274 associated with a fall in bacterial abundance and a similarly large decrease in TEP in E04 274, where bacterial abundance increased. The relative changes in TEP were less marked in incubations that contained copepods. Overall, the relationship between TEP and bacteria was weak but potentially

negative, with neither increases in pCO₂ levels nor the presence of copepods having a strong impact on the relationship.

A similar pattern was observed with regards the relationship between the relative change in TEP concentration and nanoplankton abundance (Fig. 6). As with the bacteria, some of the highest rates of change in nanoplankton were observed in the E04 274 experiment, where numbers increased 2 fold when incubated in the control incubations and 1.5 fold in all incubations containing copepods. Correspondingly, there was a drop in TEP concentrations by almost 50% over 24 h. By contrast, in E03 274, nanoplankton abundance almost halved in the control incubations where there was a concomitant doubling in TEP concentration. These two extremities were the main drivers of a negative relationship between TEP concentration and nanoplankton abundance that was particularly evident in the control incubations. Nevertheless, the trend did not prove to be significant ($F = 6.346$, $P = 0.065$, 5 df). This trend was weaker in incubations containing copepods either in ambient conditions or at raised pCO₂ levels.

Phytoplankton community response In the majority of control incubations (i.e. ambient conditions, no copepods), the abundance of phytoplankton cells increased over 24 h, many by between 1 and 5 fold (Bottle effects, Table 5). These increases were observed across all four of the major phytoplankton taxa counted (diatoms, dinoflagellates, ciliates and flagellates). The exception was experiment E04 274, where cell abundance of diatoms, dinoflagellates and ciliates in control incubations decreased.

The impact of copepods on phytoplankton cell abundance was mixed, with cell numbers increasing in some instances and decreasing in others (Copepod effects, Table 5). No major phytoplankton taxon either universally increased or decreased across all experiments when incubated in ambient conditions with copepods, rather, responses varied according to taxon in each incubation. For example, in experiment E01 271, all taxa continued to increase in cell numbers when copepods were present while, in E05 271, all but the flagellates decreased.

Similarly mixed responses in phytoplankton cell abundance were observed in the pCO₂ treatments (Treatment effects, Table 5, Fig. 7). Compared to the responses observed in ambient conditions, some taxa responded negatively and other positively to increased pCO₂ within the same experiment (e.g. E03 271, E03 274). However, in two experiments (E01 271 and E04 274), all 4 major phytoplankton taxa responded negatively to both the 750 µatm and 1000 µatm treatments when copepods were present. Of particular interest however was the response of dinoflagellates, in which there was a negative response within the incubations containing copepods at both 750 µatm and 1000 µatm pCO₂ across all experiments while, in the corresponding incubations without copepods, the response was almost wholly positive. The difference in the dinoflagellate response between incubations containing copepods to those without copepods was significant in both the 750 µatm and 1000 µatm treatments (paired t-test: 750 µatm $t = -3.38$, $P = 0.02$, 5 df; 1000 µatm $t = -3.38$, $P = 0.02$). Such a consistent response to the presence or absence of copepods at raised pCO₂ levels was not observed in the other major phytoplankton taxa.

Copepod response Compared to the variability observed in animals that were frozen immediately after capture (T0 population), there was no significant increase in the carbon content of copepods incubated for 24 h or 96 h in any of the treatments (Table 4). Over the course of the experiments therefore, the incubated copepods did not appear to increase or decrease in body mass to a detectable level.

C:N mass ratios of incubated copepods also did not appear to differ significantly from the T0 populations in the majority of instances (Fig. 8), indicating that copepods neither added to nor used up body reserves during the course of the experiments. However, significant differences were found in experiment E05 271, where the C:N mass ratio increased relative to the T0 population in all but the 24 h 1000 µatm incubation (Kruskall-Wallis 1-way ANOVA, $H = 55.45$, $P < 0.001$, 6 df).

Furthermore, there was a significant increase in the C:N mass ratio of E03 271 copepods incubated for 96 h in ambient conditions (Kruskall-Wallis 1-way ANOVA, $H = 18.07$, $P = 0.006$, 6 df).

Nevertheless, on the whole, neither the length of the incubation nor the addition of pCO₂ had an acute influence on copepod body mass or body condition.

3. Discussion

The present study carried out short-term (up to 4 d) incubation experiments of natural open-ocean communities with and without the presence of grazing copepods in microcosms in which pCO₂ levels had been manipulated. The major advantage of the approach was in the repetition of the same experimental technique on a diverse range of polar and sub-polar natural communities. The addition of a mesozooplankton grazer to these microcosms added to the complexity of the experimental system and made the interpretation of results more challenging. Nevertheless, we found that this approach revealed responses to acute OA change. In particular, the impact of pCO₂ treatments on dinoflagellate cell numbers was consistently and significantly affected by the presence or absence of copepods in the microcosms across each of the six diverse environments in which the experiments were performed. By comparison, the other phytoplankton taxa did not show any consistent response between treatments or experiments.

This study also set out to determine whether the widely reported observation of increased TEP as a result of CO₂ perturbation was prominent within a multi-trophic level experimental setup, and whether this had any impact on the microbial community. We found that TEP increased naturally at varying rates within control microcosms (i.e. ambient conditions without copepods). However, elevated pCO₂ treatments resulted in significant increases in TEP concentrations in some instances, but no change or significant decreases in others. Furthermore, there was no significant relationship between any observed change in TEP concentrations and the abundance of either bacteria or nanoplankton. Indeed, although weak, the general trend was that both sets of microorganisms were less abundant in those experiments where TEP was increasing most rapidly, which is contrary to the expectation that these organisms thrive on increased TEP production (Bellerby et al. 2008).

4.1 Sources of error

One of the main methodological errors was in the under achievement of pCO₂ targets in certain experiments. Although the manipulations were successful in elevating DIC levels while maintaining a constant TA, the measured pCO₂ were sometimes several hundred μatm below their target levels. Discrepancies between measured pCO₂ values and those calculated from DIC and TA have been previously identified, particularly at high pCO₂ levels (Hoppe et al. 2010, Hoppe et al. 2012). Amongst the sources of error are uncertainties in calculation of the carbonate system, miscalculations of the perturbation amounts, and handling errors when adding the chemicals to the incubation bottles. There is also the potential for measurement error, which we estimated to be up to 60 μatm pCO₂ when also including errors propagated in the calculation of the carbonate system. pCO₂ levels within the bottles were also subject to drift during the period of incubation, by an average of 36 μatm after 24 h and 60 μatm after 96 h. For this reason, the following discussion does not emphasise the precise differences in the response to the two different pCO₂ treatments but focuses more on the consistency of responses between all incubations where pCO₂ was manipulated.

4.2 Dinoflagellate response to acute pCO₂ elevation

Our study found a significant difference in the response of dinoflagellates to acute pCO₂ elevation, depending on the presence or absence of grazing copepods. Across all six experiments, there was a decline in dinoflagellate cell numbers in raised pCO₂ levels when copepods were present, whereas dinoflagellate numbers increased when copepods were absent. No significant change was observed in any of the other major phytoplankton taxa. One explanation for this pattern is the direct impact of copepods on dinoflagellates in that the feeding behaviour and preferences of the copepods alter under enhanced pCO₂ conditions, resulting in them selecting dinoflagellates to a greater extent than under ambient conditions. Alternatively, the presence of copepods may have had an indirect negative effect on dinoflagellate abundance when pCO₂ levels were raised.

Support for the direct impact of copepods on dinoflagellates assumes that dinoflagellates become an increasingly suitable food item under elevated $p\text{CO}_2$. Kleppel et al. (1991) was one of the first to argue that dinoflagellates are more nutritious than diatoms to zooplankton grazers. Dinoflagellates contain 1.1-3.5 times more carbohydrate and 1.8-6 times more lipid than diatoms of equivalent volume (Hitchcock 1982). Laboratory experiments have also shown that diets containing certain dinoflagellates enhance copepod egg production rates (Stoecker and Egloff 1987) whereas diatoms could suppress these rates (Miralto et al. 1999, Ianora et al. 2004, although see Irigoien et al. 2002). Nevertheless, to explain the pattern in the present study, dinoflagellates must not only be the prey item of choice, but become increasingly preferred when $p\text{CO}_2$ is raised. Under high $p\text{CO}_2$ conditions, increasing carbon availability has the potential to change the stoichiometry of nutrients to primary producers. Carbon to nutrient ratios, which are already comparatively high in diatoms, will increase under such scenarios, making them of inferior quality to herbivorous consumers (Sterner and Elser 2002, Boersma et al. 2008, Malzahn et al. 2007, Schoo et al. 2013, Nobili et al. 2013). The lower enzymatic specificity for CO_2 in dinoflagellates makes them less responsive to CO_2 enrichment and their carbon to nutrient ratios are less affected than diatoms (Tortell 2000), and they retain their quality as food items. Consumers, such as calanoid copepods, are more constrained with respect to their body nutrient composition than primary producers and thus have to cope with excess amounts of carbon relative to essential nutrients since they aim to maintain homeostasis (Sterner and Elser 2002). Selection of the most stoichiometrically suitable food items is likely to play a major role in maintaining this homeostasis. Nevertheless, this selectivity may alter during different life-stages, and carbon-rich foods may be preferable during phases when the copepods are actively storing reserves as fatty acids, particularly long chain polyunsaturated fatty acids. Such fatty acids may also be altered to varying extents in different phytoplankton taxa under enhanced $p\text{CO}_2$ (Wynn-Edwards et al. 2014).

Furthermore, a direct impact of copepods on dinoflagellates means that these grazers must have the capacity to select between different prey items (Kleppel 1993). There has been a long history of study and debate on the capacity of copepods to feed selectively, dating back to Esterly (1916). Copepods are generally thought of as filtration feeders, meaning that they principally capture particles no smaller than the inter-setal distance on their feeding appendages. However, Koehl and Strickler (1981) and Paffenhöfer et al. (1982) demonstrated that, in the low Reynolds number environments in which they exist, copepods are suspension feeders where the second maxillae push parcels of water containing particles towards the mouth rather than filtering particles from the water. There is also abundant evidence that copepods can select based on food quality. For instance, Mullin (1963) demonstrated that *Calanus helgolandicus* ingests more exponentially growing than senescent diatoms per unit time. Cowles et al. (1988) showed that *Acartia tonsa* distinguishes nutritional quality in mixtures of nutritionally rich and poor phytoplankton species. The same finding was made by Kiørboe (1989) who further found a relationship between the quality of ingested food and egg production rates. In summary, direct impacts of copepods on dinoflagellates at acutely enhanced $p\text{CO}_2$ levels is supported by other findings that enhanced $p\text{CO}_2$ is likely to decrease the nutritional value of dinoflagellates less than diatoms, and that copepods are capable of selecting phytoplankton cells based on their quality as food items.

It is more difficult to find supporting evidence for the possibility that copepods have an indirect negative impact on dinoflagellate abundance under enhanced $p\text{CO}_2$ conditions. One means in which this could occur is that the copepods may alter the chemical environment, through respiration and excretion, making it less suited to dinoflagellate growth when $p\text{CO}_2$ is raised. Flynn et al. (2015) found that altering nutrient conditions changes the competitive balance between different phytoplankton species exposed to OA. However, there was little evidence that nutrient levels were significantly altered after 24 h and CO_2 becomes more, rather than less, available as a result of the experimental treatments and should be non-limiting to cell division. Another possibility is that the copepods grazing on other prey items may cause a trophic cascade, releasing predatory pressure on

other dinoflagellates grazers. However, dinoflagellates are relatively large cells and are unlikely to be major prey items of any other heterotrophic organisms within the incubations other than copepods. Therefore, the direct impact of copepods on dinoflagellates through selective grazing is the most likely explanation for the observed pattern.

4.3 Copepod body condition

We found there to be no detectable change in copepod body condition in CV and females stages of *Calanus finmarchicus*/*glacialis*, *C. hyperboreus*, and *Calanoides acutus*, assessed in terms of the C:N mass ratio, as a result of raised pCO₂ treatments either after 24 h or 96 h of incubation. Neither was there a concomitant rise or fall in the mass of carbon and nitrogen, which would indicate a change in body mass over the incubation period. The exception was experiment E05 271, where the C:N mass ratio significantly increased. However, the same level of increase was observed in both ambient and raised pCO₂ conditions, indicating that this pattern was more likely to be a bottle effect than a treatment effect.

The fact that there was little response to pCO₂ treatments in any of the copepod species is consistent with the majority of other studies on the sensitivity of copepods to raised CO₂. In terms of direct physiological responses, most studies have found no direct effects on copepods at CO₂ levels within the range expected by the end of this century (Kurihara et al. 2004; Watanabe et al. 2006, Kurihara and Ishimatsu 2008, Zhang et al. 2011, Vehmaa et al. 2012, Weydmann et al., 2012, McConville et al. 2013, Niehoff et al. 2013). It is to be noted that the majority of these studies focussed on adult females whereas only a few have concentrated on other life-history stages which are likely to show greater sensitivity to elevated pCO₂ (Kurihara et al. 2004, Fitzer et al. 2012, Lewis et al. 2013, Cripps et al. 2014).

There has been less attention on how copepod body condition responds to changes in the quality of their food as a result of increased levels of pCO₂. For instance, Urabe et al. (2003) found lower

growth rates of the freshwater species *Daphnia pulicaria* at high $p\text{CO}_2$ due to altered algal C:P ratios. Schoo et al. (2013) found that the copepod *Acartia tonsa* showed decreased development rate when fed the algae *Rhodomonas salina* grown under increased $p\text{CO}_2$ levels. In a culture containing the diatom *Thalassiosira pseudonana* and the copepod *Acartia tonsa*, Rossoll et al. (2012) found that exposure to $p\text{CO}_2$ levels of $750 \mu\text{atm}$ led to a decrease in polyunsaturated fatty acids in the diatoms and a corresponding decrease in copepod egg production (Rossoll et al. 2012). Therefore, even though it is likely that copepods have the ability to select food items based on their quality, their stoichiometric homeostasis and growth potential will be affected when there is a lack of good quality food items available. The diversity of food items in the incubation chambers in the present study may have allowed the copepods in the present study to prevent decreases in body condition, at least for the short term duration of the incubations.

4.4 Net change in TEP

TEP increased in all of the control incubations (i.e. ambient conditions, without copepods) to varying degrees, with some of the highest rates of increase occurring in the most nutrient stressed sites (North Sea, Barents Sea). These increases took some days to be detectable, with no significant increases being found after 24 h. When compared to the changes documented in the control incubation, 4 out of the 6 experiments showed significant increases in the $1000 \mu\text{atm } p\text{CO}_2$ treatment, with the remaining 2 experiments either showing a neutral or negative change. Only 2 of the experiments showed a significant increase in the $750 \mu\text{atm } p\text{CO}_2$ treatment, the remaining 4 not showing any significant change. The impact of copepods on TEP concentrations was mixed, leading to increased concentrations in some experiments but neutral or negative changes in others.

Initial studies on the influence of raised $p\text{CO}_2$ on TEP formation found a relatively consistent pattern of greater TEP formation when $p\text{CO}_2$ levels were raised acutely (Bellerby et al. 2008, Weinbauer et al. 2011). The prevailing hypothesis is that, under high $p\text{CO}_2$ conditions, more of the organic carbon fixed by photosynthesis is channelled into the dissolved fraction and released from the cells. This

leads to a greater standing stock of TEPs as the released matter coagulates into particulates. Higher concentration of TEP in surface waters is therefore believed to be one of the most predictable consequences of OA (Weinbauer et al. 2011).

Nevertheless, there has recently been some doubt regarding the universality of the response of TEP to increased $p\text{CO}_2$ levels. In rolling tank experiments, where TEP was either in equilibrium or non-equilibrium with its precursors, Passow (2012) found that OA as expected in the future oceans had no impact on the process of TEP formation from its precursors. In field observations, MacGilchrist et al. (2014) found a significant but highly variable effect of acute $p\text{CO}_2$ increase on TEP production, with increased TEP production being closely linked to phytoplankton growth in some instances and as a result of greater amounts of TEP being produced per cell in others. The MacGilchrist et al. (2014) study used similar incubation techniques to the present study but was carried out in a variety of environmental settings in European shelf seas. Their findings match those of the present study in that the TEP response to raised $p\text{CO}_2$ levels is more likely to be dependent on the specific environmental setting and cannot be assumed to be universal.

As well as being conducted in high latitude environments, the present study differed from that of MacGilchrist et al. (2014) in introducing a further trophic level through the addition of grazing copepods. Zooplankton have been found to feed effectively on clusters of TEP (Passow and Alldredge 1999, Ling and Alldredge 2003). Nevertheless, there was no consistent impact of copepods on TEP in the present study, suggesting that it is unlikely to be a preferred food item. This may be particularly the case given that TEP is carbon rich (Passow 2012) and of low quality relative to nutrient rich phytoplankton taxa such as dinoflagellates (Kleppel 1993).

4.5 Response of bacteria and nanoplankton

We found that there was no significant change in the abundance of either bacteria or nanoplankton in response to raised $p\text{CO}_2$ levels. This is consistent with the results of mesocosm

studies where no or only a small effect of elevated $p\text{CO}_2$ levels was found on bacterial abundance (Rochelle-Newall et al. 2004, Allgaier et al. 2008, Paulino et al 2008). Nevertheless, according to Weinbauer et al. (2011), no study has shown a clear negative effect of elevated $p\text{CO}_2$ on prokaryotic production at pH levels relevant in the context of OA. Thus, our study supports the wider body of data so far collected that suggests a neutral or positive effect of OA on the growth rate of microorganism populations.

The relationship between the relative change in TEP concentrations and bacterial abundance in the present study was not significant, although there was a trend of bacterial numbers decreasing as TEP increased. The relationship between TEP concentration and nanoplankton abundance showed a similar trend. Corzo et al. (2005), Passow et al. (2001), Hung et al. (2003) and Santschi et al. (2003) have reported positive relationships between TEP and bacteria, whereas Passow and Alldredge (1994) and Bhaskar and Bhosle (2006) found negative or no correlation at all. In the Southern Ocean, Ortega-Retuerta et al. (2009) found a positive relationship between TEP and bacterial abundance within the upper mixed layer, although the slope was relatively shallow. There was a much stronger relationship below the mixed layer. It was proposed that the relationship was mediated by phytoplankton and may not be a direct relationship between the two components. The present study only considered the upper mixed layer community which may have constrained our ability to identify any such relationship between TEP and microorganisms. Alternatively, it is possible that the increase in TEP we consistently observed over the course of the incubation experiments was the result of bottle effects to which the bacterial- and nanoplankton communities could not respond. In summary, we found no evidence to support the view that the microbial community is stimulated by the greater levels of TEP production that may result from acute increases in $p\text{CO}_2$.

4. Concluding remarks

This study completed six controlled microcosm experiments on natural communities extracted from a diverse set of high latitude open-ocean sites and exposed to acute increases in pCO₂. The spatial diversity of the experiments allowed the determination of whether different communities showed similar or differing responses to a common acute perturbation in pCO₂. Although the short-term and acute nature of the perturbation was not a true simulation of the changes likely to take place over the coming decades in the natural environment, it did provide an insight into whether these differing communities are equally placed in adapting to the predicted environmental changes. Across a range of high latitude environments, we identified a similar trophic response in copepods to their phytoplankton prey, which was induced by elevated pCO₂. We also demonstrated that the response of TEP to acute pCO₂ perturbation depends on the precise set of environmental conditions and cannot be assumed to be highly predictable. Trophic impacts on the body condition of the grazing copepod community were not resolved and are better suited to longer-term experimental approaches.

Acknowledgements

This work was funded under the UK Ocean Acidification thematic programme (UKOA) via the UK Natural Environment Research Council (NERC) grants NE/H017267/1 to GA Tarling, NE/H017348/1 to T Tyrrell, EP Achterberg and CM Moore, NE/H017097/1 to AJ Poulton and MV Zubkov and NE/H016988/1 to RJG Leakey. The UK Department of Environment, Food and Rural Affairs (Defra) and the UK Department of Energy and Climate Change (DECC) also contributed to funding UKOA. Our travel and work in Greenland, Iceland and Svalbard coastal and offshore waters was granted thanks to permissions provided by the Danish, Icelandic and Norwegian diplomatic authorities. We also thank the captains and crew of the RRS James Clark Ross during cruises JR271 and JR274, the technical staff from the British Antarctic Survey, National Marine Facilities, and the Scottish Association for Marine Science, Oban. Further thanks to P Enderlein for Bongo net maintenance, CM

Moore for assistance with acid-base manipulations, M Esposito and R Sanders for macronutrient data, H Lawson for Lugols count data, C Georges for JR274 flow cytometry data, P Geissler for elemental analyses and M Ribas-Ribas (NOC) and R Thomas (BODC) for data management. We are grateful to R Leakey for leading cruise JR271 and for critical comments on this work.

References

- Allgaier, M., Riebesell, U., Vogt, M., Thyrrhaug, R., Grossart, H.-P., 2008. Coupling of heterotrophic bacteria to phytoplankton bloom development at different pCO₂ levels: a mesocosm study. *Biogeosciences Discuss.* 5, 317-359.
- Beaufort, L., Probert, I., de Garidel-Thoron, T., Bendif, E., Ruiz-Pino, D., Metzl, N., Goyet, C., Buchet, N., Coupel, P., Grelaud, M., 2011. Sensitivity of coccolithophores to carbonate chemistry and ocean acidification. *Nature* 476, 80-83.
- Bellerby, R.G.J., Schulz, K.G., Riebesell, U., Neill, C., Nondal, G., Heegaard, E., Johannessen, T., Brown, K.R., 2008. Marine ecosystem community carbon and nutrient uptake stoichiometry under varying ocean acidification during the PeECE III experiment. *Biogeosciences* 5, 1517-1527.
- Bhaskar, P., Bhosle, N.B., 2006. Dynamics of transparent exopolymeric particles (TEP) and particle-associated carbohydrates in the Dona Paula bay, west coast of India. *J. Earth Syst. Sci.* 115, 403-413.
- Boersma, M., Aberle, N., Hantzsche, F.M., Schoo, K.L., Wiltshire, K.H., Malzahn, A.M., 2008. Nutritional limitation travels up the food chain. *Int. Rev. Hydrobiol.* 93, 479-488.
- Bradshaw, A.L., Brewer, P.G., Shafer, D.K., Williams, R.T., 1981. Measurements of total carbon dioxide and alkalinity by potentiometric titration in the GEOSECS program. *Earth Planet. Sci. Lett.* 55, 99115, doi:99110.91016/90012-99821X(99181)90090-X.
- Cooley, S.R., Kite-Powell, H.L., Doney, S.C., 2009. Ocean Acidification's potential to alter global marine ecosystem services. *Oceanography* 22, 172-181.
- Corzo, A., Rodríguez-Gálvez, S., Lubian, L., Sobrino, C., Sangrá, P., Martínez, A., 2005. Antarctic marine bacterioplankton subpopulations discriminated by their apparent content of nucleic acids

- differ in their response to ecological factors. *Polar Biol.* 29, 27-39.
- Cowles, T., Olson, R., Chisholm, S., 1988. Food selection by copepods: discrimination on the basis of food quality. *Mar. Biol.* 100, 41-49.
- Cripps, G., Lindeque, P., Flynn, K.J., 2014. Have we been underestimating the effects of ocean acidification in zooplankton? *Global Change Biol.* 20, 3377-3385.
- de Nooijer, L.J., Toyofuku, T., Kitazato, H., 2009. Foraminifera promote calcification by elevating their intracellular pH. *P. Natl. Acad. Sci. USA* 106, 15374-15378.
- Dickson, A.G., 1990. Standard potential of the reaction: $\text{AgCl(s)} + 12\text{H}_2\text{(g)} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *J. Chem. Thermodyn.* 22, 13127, doi:13110.11016/10021-19614(13190)90074-Z.
- Dickson, A.G., Millero, F.J., 1987. A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep-Sea Res.* 34, 1733-1743.
- Dickson, A.G., Sabine, C.L., Christian, J.R., 2007. Guide to best practices for ocean CO_2 measurements. N. Pac. Mar. Sci. Org, Sidney, BC, Canada, p. 191
http://cdiac.ornl.gov/oceans/Handbook_2007.html.
- Doney, S.C., Ruckelshaus, M., Emmett Duffy, J., Barry, J.P., Chan, F., English, C.A., Galindo, H.M., Grebmeier, J.M., Hollowed, A.B., Knowlton, N., 2012. Climate change impacts on marine ecosystems. *Ann. Rev. Mar. Sci.* 4, 11-37.
- Engel, A., 2002. Direct relationship between CO_2 uptake and transparent exopolymer particles production in natural phytoplankton. *J. Plankt. Res.* 24, 49-53.
- Engel, A., Schulz, K., Riebesell, U., Bellerby, R., Delille, B., Schartau, M., 2008. Effects of CO_2 on particle size distribution and phytoplankton abundance during a mesocosm bloom experiment (PeECE II). *Biogeosciences Discuss.* 5, 509-521.
- Engel, A., Thoms, S., Riebesell, U., Rochelle-Newall, E., Zondervan, I., 2004. Polysaccharide aggregation as a potential sink of marine dissolved organic carbon. *Nature* 428, 929-932.
- Esterly, C.O., 1916. The feeding habits and food of pelagic copepods and the question of nutrition by

- organic substances in solution in the water. *Unlv. Calif. Publ. Zool.* 16, 171-184.
- Fabry, V.J., McClintock, J.B., Mathis, J.T., Grebmeier, J.M., 2009. Ocean Acidification at high latitudes: the bellwether. *Oceanography* 22, 160-171.
- Fitzer, S.C., Caldwell, G.S., Close, A.J., Clare, A.S., Upstill-Goddard, R.C., Bentley, M.G., 2012. Ocean acidification induces multi-generational decline in copepod naupliar production with possible conflict for reproductive resource allocation. *J. Exp. Mar. Biol. Ecol.* 418, 30-36.
- Flynn KJ, Clark DR, Mitra A, Fabian H, Hansen PJ, Glibert PM, Wheeler GL, Stoecker DK, Blackford JC, Brownlee C (2015) Ocean acidification with (de) eutrophication will alter future phytoplankton growth and succession. *Proc. R. Soc. London B* 282, 20142604.
- Gattuso, J.-P., Gao, K., Lee, K., Rost, B., Schulz, K.G., 2010. Approaches and tools to manipulate the carbonate chemistry, in: Riebesell, U., Fabry, V.J., Hansson, L., Gattuso, J.P. (Eds.), *Guide to Best Practices for Ocean Acidification Research and Data Reporting*. Publications Office of the European Union, Luxembourg, p. 263 pp.
- Hitchcock, G.L., 1982. A comparative study of the size-dependent organic composition of marine diatoms and dinoflagellates. *J. Plankt. Res.* 4, 363-377.
- Hoppe, C., Langer, G., Rokitta, S., Wolf-Gladrow, D., Rost, B., 2010. On CO₂ perturbation experiments: over-determination of carbonate chemistry reveals inconsistencies. *Biogeosciences Discuss.* 7, 1707-1726.
- Hoppe, C.J.M., Langer, G., Rokitta, S.D., Wolf-Gladrow, D.A., Rost, B., 2012. Implications of observed inconsistencies in carbonate chemistry measurements for ocean acidification studies. *Biogeosciences* 9, 2401-2405.
- Hung, C.C., Guo, L., Schultz, G.E., Pinckney, J.L., Santschi, P.H., 2003. Production and flux of carbohydrate species in the Gulf of Mexico. *Global Biogeochem. Cy.* 17, 1055, doi:10.1029/2002GB001988.
- Hutchins, D., Fu, F.-X., Zhang, Y., Warner, M., Feng, Y., Portune, K., Bernhardt, P., Mulholland, M., 2007. CO₂ control of *Trichodesmium* N₂ fixation, photosynthesis, growth rates, and elemental

- ratios: implications for past, present, and future ocean biogeochemistry. *Limnol. Oceanogr.* 52, 1293-1304.
- Ianora, A., Miralto, A., Poulet, S.A., Carotenuto, Y., Buttino, I., Romano, G., Casotti, R., Pohnert, G., Wichard, T., Colucci-D'Amato, L., 2004. Aldehyde suppression of copepod recruitment in blooms of a ubiquitous planktonic diatom. *Nature* 429, 403-407.
- Irigoien, X., Harris, R.P., Verheye, H.M., Joly, P., Runge, J., Starr, M., Pond, D., Campbell, R., Shreeve, R., Ward, P., 2002. Copepod hatching success in marine ecosystems with high diatom concentrations. *Nature* 419, 387-389.
- K erouel, R., Aminot, A., 1997. Fluorometric determination of ammonia in sea and estuarine waters by direct segmented flow analysis. *Mar. Chem.* 57, 265-275.
- Kj rboe, T., 1989. Phytoplankton growth rate and nitrogen content: implications for feeding and fecundity in a herbivorous copepod. *Mar. Ecol. Prog. Ser.* 55, 229-234.
- Kirkwood, D.S., 1996. Nutrients: a practical note on their determination in seawater. ICES, Copenhagen, Denmark.
- Kleppel, G., 1993. On the diets of calanoid copepods. *Mar. Ecol. Prog. Ser.* 99, 183-183.
- Kleppel, G.S., Holliday, D., Pieper, R., 1991. Trophic interactions between copepods and microplankton: a question about the role of diatoms. *Limnol. Oceanogr.* 36, 172-178.
- Koehl, M., Strickler, J.R., 1981. Copepod feeding currents: Food capture at low Reynolds numbers. *Limnol. Oceanogr.* 26, 1062-1073.
- Kurihara, H., Ishimatsu, A., 2008. Effects of high CO₂ seawater on the copepod (*Acartia tsuensis*) through all life stages and subsequent generations. *Mar. Pollut. Bull.* 56, 1086-1090.
- Kurihara, H., Shimode, S., Shirayama, Y., 2004. Effects of raised CO₂ concentration on the egg production rate and early development of two marine copepods (*Acartia steueri* and *Acartia erythraea*). *Mar. Pollut. Bull.* 49, 721-727.
- Lee, K., Kim, T.-W., Byrne, R.H., Millero, F.J., Feely, R.A., Liu, Y.-M., 2010. The universal ratio of boron to chlorinity for the North Pacific and North Atlantic oceans. *Geochim. Cosmochim. Ac.* 74, 1801-

1811.

- Lewis, C.N., Brown, K.A., Edwards, L.A., Cooper, G., Findlay, H.S., 2013. Sensitivity to ocean acidification parallels natural pCO₂ gradients experienced by Arctic copepods under winter sea ice. *Proc. Natl. Acad. Sci. USA* 110, E4960-E4967.
- Lewis, E., Wallace, D.W.R., 1998. Program Developed for CO₂ System Calculations. ORNL/CDIAC-105. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee.
- Ling, S.C., Alldredge, A.L., 2003. Does the marine copepod *Calanus pacificus* consume transparent exopolymer particles (TEP)? *J. Plankt. Res.* 25, 507-515.
- MacGilchrist, G., Shi, T., Tyrrell, T., Richier, S., Moore, C., Dumousseaud, C., Achterberg, E.P., 2014. Effect of enhanced pCO₂ levels on the production of dissolved organic carbon and transparent exopolymer particles in short-term bioassay experiments. *Biogeosciences* 11, 3695-3706.
- Malzahn, A.M., Aberle, N., Clemmesen, C., Boersma, M., 2007. Nutrient limitation of primary producers affects planktivorous fish condition. *Limnol. Oceanogr.* 52, 2062-2071.
- Marie, D., Partensky, F., Jacquet, S., Vaulot, D., 1997. Enumeration and cell cycle analysis of natural populations of marine picoplankton by flow cytometry using the nucleic acid stain SYBR Green I. *Appl. Environ. Microbiol.* 63, 186-193.
- McConville, K., Halsband, C., Fileman, E.S., Somerfield, P.J., Findlay, H.S., Spicer, J.I., 2013. Effects of elevated CO₂ on the reproduction of two calanoid copepods. *Mar. Pollut. Bull.* 73, 428-434.
- McNeil, B.I., Matear, R.J., 2008. Southern Ocean acidification: A tipping point at 450-ppm atmospheric CO₂. *Proc. Natl. Acad. Sci. U.S.A.* 105, 18860-18864.
- Mehrbach, C., Culberson, C.H., Hawley, J.E., Pytkowicz, R.M., 1973. Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnol. Oceanogr.* 18, 897-907.
- Miralto, A., Barone, G., Romano, G., Poulet, S., Ianora, A., Russo, G., Buttino, I., Mazzarella, G., Laabir, M., Cabrini, M., 1999. The insidious effect of diatoms on copepod reproduction. *Nature*

- 402, 173-176.
- Mullin, M.M., 1963. Some factors affecting the feeding of marine copepods in the genus *Calanus*.
Limnol. Oceanogr. 8, 239-250.
- Niehoff, B., Knüppel, N., Daase, M., Czerny, J., Boxhammer, T., 2013. Mesozooplankton community development at elevated CO₂ concentrations: results from a mesocosm experiment in an Arctic fjord. *Biogeosciences* 10, 1391-1406.
- Nobili, R., Robinson, C., Buitenhuis, E., Castellani, C., 2013. Food quality regulates the metabolism and reproduction of *Temora longicornis*. *Biogeosciences Discuss.* 10, 3203-3239.
- Ortega-Retuerta, E., Frazer, T.K., Duarte, C.M., Ruiz-Halpern, S., Tovar-Sánchez, A., Arrieta, J.M., Rechea, I., 2009. Biogeneration of chromophoric dissolved organic matter by bacteria and krill in the Southern Ocean. *Limnol. Oceanogr.* 54, 1941-1950.
- Paffenhöfer, G.-A., Strickler, J., Alcaraz, M., 1982. Suspension-feeding by herbivorous calanoid copepods: a cinematographic study. *Mar. Biol.* 67, 193-199.
- Passow, U., 2002. Transparent exopolymer particles (TEP) in aquatic environments. *Prog. Oceanogr.* 55, 287-333.
- Passow, U., 2012. The abiotic formation of TEP under different ocean acidification scenarios. *Mar. Chem.* 128, 72-80.
- Passow, U., Alldredge, A., 1995. A dye binding assay for the spectrophotometric measurement of transparent exopolymer particles (TEP). *Limnol. Oceanogr.* 40, 1326-1335.
- Passow, U., Alldredge, A.L., 1994. Distribution, size and bacterial-colonization of transparent exopolymer particles (TEP) in the ocean. *Mar. Ecol. Prog. Ser.* 113, 185–198.
- Passow, U., Alldredge, A.L., 1999. Do transparent exopolymer particles (TEP) inhibit grazing by the euphausiid *Euphausia pacifica*? *J. Plankt. Res.* 21, 2203-2217.
- Passow, U., Shipe, R., Murray, A., Pak, D., Brzezinski, M., Alldredge, A., 2001. The origin of transparent exopolymer particles (TEP) and their role in the sedimentation of particulate matter. *Cont. Shelf Res.* 21, 327-346.

- Paulino, A., Egge, J., Larsen, A., 2008. Effects of increased atmospheric CO₂ on small and intermediate sized osmotrophs during a nutrient induced phytoplankton bloom. *Biogeosciences* 5, 739-748.
- Poulton, A.J., Moore, C.M., Seeyave, S., Lucas, M.I., Fielding, S., Ward, P., 2007. Phytoplankton community composition around the Crozet Plateau, with emphasis on diatoms and *Phaeocystis*. *Deep-Sea Res. II* 54, 2085-2105.
- Redfield, A., 1958. The biological control of chemical factors in the environment. *Amer. Sci*, 205-221.
- Richier, S., Achterberg, E., Dumousseaud, C., Poulton, A., Suggett, D., Tyrrell, T., Zubkov, M., Moore, C., 2014. Carbon cycling and phytoplankton responses within highly-replicated shipboard carbonate chemistry manipulation experiments conducted around Northwest European Shelf Seas. *Biogeosciences Discuss* 11, 3489-3534.
- Riebesell, U., Fabry, V.J., Hansson, L., Gattuso, J.-P., 2010. Guide to best practices for ocean acidification research and data reporting. Publications Office of the European Union Luxembourg.
- Riebesell, U., Gattuso, J.-P., Thingstad, T., Middelburg, J., 2013. Preface "Arctic ocean acidification: pelagic ecosystem and biogeochemical responses during a mesocosm study". *Biogeosciences* 10, 5619-5626.
- Riebesell, U., Tortell, P., 2011. Effects of ocean acidification on pelagic organisms and ecosystems, in: Gattuso, J., Hanson, L. (Eds.), *Ocean Acidification*. Oxford University Press, Oxford, pp. 99-121.
- Riebesell, U., Zondervan, I., Rost, B., Tortell, P.D., Zeebe, R.E., Morel, F.M., 2000. Reduced calcification of marine plankton in response to increased atmospheric CO₂. *Nature* 407, 364-367.
- Rochelle-Newall, E., Delille, B., Frankignoulle, M., Gattuso, J.-P., Jacquet, S., Riebesell, U., Terbrüggen, A., Zondervan, I., 2004. Chromophoric dissolved organic matter in experimental mesocosms maintained under different pCO₂ levels. *Mar. Ecol. Prog. Ser.* 272, 25-31.
- Rossoll, D., Bermúdez, R., Hauss, H., Schulz, K.G., Riebesell, U., Sommer, U., Winder, M., 2012. Ocean acidification-induced food quality deterioration constrains trophic transfer. *Plos One* 7, e34737.
- Rossoll, D., Sommer, U., Winder, M., 2013. Community interactions dampen acidification effects in a

- coastal plankton system. *Mar. Ecol. Prog. Ser.* 486, 37-46.
- Sabine, C.L., Feely, R.A., Gruber, N., Key, R.M., Lee, K., Bullister, J.L., Wanninkhof, R., Wong, C.S., Wallace, D.W.R., Tilbrook, B., Millero, F.J., Peng, T.H., Kozyr, A., Ono, T., Rios, A.F., 2004. The oceanic sink for anthropogenic CO₂. *Science* 305, 367-371.
- Santschi, P.H., Hung, C.C., Schultz, G., Alvarado-Quiroz, N., Guo, L., Pinckney, J., Walsh, I., 2003. Control of acid polysaccharide production and ²³⁴Th and POC export fluxes by marine organisms. *Geophys. Res. Lett.* 30.
- Schoo, K.L., Malzahn, A.M., Krause, E., Boersma, M., 2013. Increased carbon dioxide availability alters phytoplankton stoichiometry and affects carbon cycling and growth of a marine planktonic herbivore. *Mar. Biol.* 160, 2145-2155.
- Simon, M., Grossart, H.-P., Schweitzer, B., Ploug, H., 2002. Microbial ecology of organic aggregates in aquatic ecosystems. *Aquat. Microb. Ecol.* 28, 175-211.
- Steinacher, M., Joos, F., Frolicher, T., Plattner, G.-K., Doney, S.C., 2009. Imminent ocean acidification in the Arctic projected with the NCAR global coupled carbon cycle-climate model. *Biogeosciences* 6, 515–533.
- Sterner, R.W., Elser, J.J., 2002. *Ecological stoichiometry: the biology of elements from molecules to the biosphere.* Princeton University Press.
- Stoecker, D.K., Egloff, D.A., 1987. Predation by *Acartia tonsa* Dana on planktonic ciliates and rotifers. *J. Exp. Mar. Biol. Ecol.* 110, 53-68.
- Tortell, P.D., 2000. Evolutionary and ecological perspectives on carbon acquisition in phytoplankton. *Limnol. Oceanogr.* 45, 744-750.
- Tortell, P.D., DiTullio, G.R., Sigman, D.M., Morel, F., 2002. CO₂ effects on taxonomic composition and nutrient utilization in an Equatorial Pacific phytoplankton assemblage. *Mar. Ecol. Prog. Ser.* 236, 37-43.
- Urabe, J., Togari, J., Elser, J.J., 2003. Stoichiometric impacts of increased carbon dioxide on a planktonic herbivore. *Global Change Biol.* 9, 818-825.

- van Heuven, S., Pierrot, D., Rae, J.W.B., E., L., Wallace, D.W.R., 2011. MATLAB Program Developed for CO₂ System Calculations. ORNL/CDIAC-105b. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee.
- Vehmaa, A., Brutemark, A., Engström-Öst, J., 2012. Maternal effects may act as an adaptation mechanism for copepods facing pH and temperature changes. *Plos One* 7, e48538.
- Watanabe, Y., Yamaguchi, A., Ishida, H., Harimoto, T., Suzuki, S., Sekido, Y., Ikeda, T., Shirayama, Y., Mac Takahashi, M., Ohsumi, T., 2006. Lethality of increasing CO₂ levels on deep-sea copepods in the western North Pacific. *J. Oceanogr.* 62, 185-196.
- Weinbauer, M.G., Mari, X., Gattuso, J.-P., 2011. Effect of ocean acidification on the diversity and activity of heterotrophic marine microorganisms, in: Gattuso, J.P., Hansson, L. (Eds.), *Ocean acidification*. Oxford University Press, Oxford, pp. 83-98.
- Weydmann, A., Søreide, J.E., Kwasniewski, S., Widdicombe, S., 2012. Influence of CO₂-induced acidification on the reproduction of a key Arctic copepod *Calanus glacialis*. *J. Exp. Mar. Biol. Ecol.* 428, 39-42.
- Whiteley, N., 2011. Physiological and ecological responses of crustaceans to ocean acidification. *Mar. Ecol. Prog. Ser.* 430, 257-271.
- Wynn-Edwards, C., King, R., Davidson, A., Wright, S., Nichols, P.D., Wotherspoon, S., Kawaguchi, S., Virtue, P., 2014. Species-specific variations in the nutritional quality of Southern Ocean phytoplankton in response to elevated pCO₂. *Water* 6, 1840-1859.
- Zar, J.H., 1996. *Biostatistical analysis*. Prentice Hall International, Englewood Cliffs, New Jersey.
- Zhang, D., Li, S., Wang, G., Guo, D., 2011. Impacts of CO₂-driven seawater acidification on survival, egg production rate and hatching success of four marine copepods. *Acta Oceanol. Sin.* 30, 86-94.
- Zubkov, M.V., Burkill, P.H., 2006. Syringe pumped high speed flow cytometry of oceanic phytoplankton. *Cytom. Part A* 69A, 1010-1019.
- Zubkov, M.V., Burkill, P.H., Topping, J.N., 2007. Flow cytometric enumeration of DNA-stained oceanic planktonic protists. *J. Plankton Res.* 29, 79-86.

Tables

Bioassay	Date	Location	Latitude (°)	Longitude (°)	Water depth (m)	Mixed-layer depth (m)	Incubated taxa
E01 271	3/6/12	North Sea	56.26664	2.63319	74	13.600 (2.408)	Calanus finmarchicus CV
E03 271	13/6/12	Arctic	76.17524	2.54953	3758	29.667 (7.506)	Calanus hyperboreus CV
E04 271	18/6/12	Arctic	78.28063	4.28433	2257	11.000 (1.414)	Calanus hyperboreus CV
E05 271	24/6/12	Arctic	72.89080	26.00273	362	44.000 (4.301)	Calanus finmarchicus/glacialis CV
E03 274	25/1/13	Southern Ocean	-52.68930	-36.62303	2445	62.600 (10.569)	Calanoides acutus CV
E04 274	1/2/13	Southern Ocean	-58.08701	-25.92901	2899	35.828 (5.824)	Calanoides acutus CV and female

Table 1: Locations and attributes of the six experimental stations, including the species and stage of copepod added to the experimental incubation bottles

Bioassay	Mixed layer T (°C)	Mixed layer salinity	NO ₃ +NO ₂ μmol L ⁻¹	Si μM L ⁻¹	PO ₄ μmol L ⁻¹	[NH ₄ ⁺] nmol L ⁻¹	N:P (atomic ratio)	TA (μmol kg ⁻¹)	DIC (μmol kg ⁻¹)	pH	pCO ₂ (μatm)	TEP (μg L ⁻¹ xanthan equiv.)	Total bacteria (cell mL ⁻¹)	Total nannoplankt. (cells mL ⁻¹)
E01 271	10.700 (0.071)	35.100 (0.000)	0.043 (0.006)	1.351 (0.006)	0.077 (0.012)	2.974 (1.030)	0.604	2325.300 (0.436)	2082.267 (0.503)	8.153 (0.000)	300.452 (0.365)	176.570 (16.717)	806803.712 (41963.039)	5444.164 (341.646)
E03 271	1.880 (0.795)	34.920 (0.045)	9.253 (0.051)	5.821 (0.025)	0.650 (0.000)	111.437 (134.775)	14.407	2305.267 (6.529)	2130.000 (3.205)	8.159 (0.014)	289.287 (9.705)	99.037 (10.046)	117395.0914 (14221.856)	41275.690 (2242.764)
E04 271	-1.575 (0.050)	32.625 (0.050)	4.200 (0.156)	12.214 (0.023)	0.790 (0.000)	145.211	5.500	2234.567 (2.901)	2106.400 (5.671)	8.129 (0.009)	304.682 (7.219)	90.973 (6.804)	460466.414 (85179.903)	3839.254 (629.680)
E05 271	6.450 (0.058)	35.000 (0.000)	5.380 (0.544)	3.867 (0.151)	0.437 (0.025)	162.527 (162.836)	12.693	2314.050 (4.031)	2108.700 (2.022)	8.146 (0.003)	304.295 (2.154)	111.855 (11.460)	113533.3889 (15287.2784)	12071.861 (1273.412)
E03 274	2.224 (0.021)	33.932 (0.002)	24.080 (0.252)	19.393 (0.095)	1.597 (0.006)	812.623 (70.177)	15.590	2287.800	2152.342 (6.112)	8.050	384	3.293 (1.678)	653401.452 (26549.817)	4358.355 (623.205)
E04 274	0.400 (0.037)	33.706 (0.005)	18.537 (0.605)	71.310 (0.473)	1.190 (0.046)	519.415 (308.750)	16.014	2293.000	2129.500 (4.355)	8.167	283	65.473 (2.887)	804153.919 (79884.942)	1476.402 (404.310)

Table 2: Starting conditions for experimental incubations. Values represent averages, while (SD) is included where more than 3 measurements were made.

Experiment/incubation		C (μg)	H (μg)	N (μg)	C:N (mass ratio)
E03 271					
T0 population		2228.53 (1033.07)	343.91 (154.36)	290.24 (97.60)	7.38 (1.44)
Ambient	24 h	2370.11 (718.46)	378.04 (104.09)	284.68 (50.08)	8.19 (1.51)
	96h	2944.47 (603.86)	464.80 (121.29)	330.63 (54.75)	8.86 (0.56)
750 μatm	24 h	2460.75 (775.92)	358.65 (123.53)	296.81 (60.40)	8.15 (1.06)
	96 h	2563.23 (595.72)	425.49 (99.41)	287.04 (51.66)	8.86 (0.59)
1000 μatm	24 h	1814.57 (623.17)	268.11 (62.74)	259.23 (88.34)	7.07 (1.40)
	96 h	2513.16 (738.55)	393.35 (120.84)	288.09 (63.67)	8.60 (0.85)
E04 271					
T0 population		1000.20 (520.57)	161.43 (83.04)	167.72 (49.15)	5.70 (1.47)
Ambient	24 h	1211.67 (346.37)	197.27 (55.64)	176.69 (30.51)	6.75 (1.04)
	96h	1454.91 (968.46)	190.75 (86.75)	195.59 (94.16)	7.06 (2.09)
750 μatm	24 h	908.51 (343.28)	149.38 (56.26)	161.69 (39.02)	5.55 (1.25)
	96 h	1330.19 (389.65)	216.62 (63.42)	191.71 (40.46)	6.85 (1.08)
1000 μatm	24 h	872.04 (237.25)	140.82 (38.32)	150.84 (32.41)	5.75 (0.90)
	96 h	1066.25 (882.90)	177.15 (144.76)	184.06 (73.86)	5.17 (2.05)
E05 271					
T0 population		74.06 (18.73)	10.83 (3.01)	17.92 (3.70)	4.15 (0.68)
Ambient	24 h	65.28 (22.50)	7.04 (3.14)	9.63 (5.01)	7.59 (2.07)
	96h	66.01 (26.04)	7.25 (3.74)	11.76 (5.33)	6.37 (2.77)
750 μatm	24 h	71.28 (35.02)	8.25 (4.41)	11.15 (4.74)	6.97 (3.13)
	96 h	68.60 (36.65)	7.40 (4.92)	12.31 (5.93)	5.57 (1.68)
1000 μatm	24 h	69.61 (35.78)	7.75 (4.68)	13.74 (5.14)	4.97 (1.21)
	96 h	66.98 (25.27)	7.30 (3.52)	11.92 (3.50)	5.66 (1.43)
E03 274					
T0 population		54.80 (12.20)	9.61 (2.21)	17.85 (3.26)	3.05 (0.21)
Ambient	24 h	66.75 (23.67)	9.76 (3.38)	18.32 (4.56)	3.57 (0.69)
	96h	50.75 (29.74)	6.90 (4.30)	14.56 (5.74)	3.32 (0.59)
750 μatm	24 h	45.33 (17.59)	6.63 (2.53)	14.59 (3.49)	3.03 (0.58)
	96 h	37.66 (14.32)	5.35 (2.14)	12.43 (4.48)	3.29 (1.58)
1000 μatm	24 h	52.16 (21.87)	7.33 (3.28)	15.55 (4.96)	3.30 (0.50)
	96 h	44.49 (21.68)	6.23 (3.04)	13.79 (4.76)	3.16 (0.69)
E04 274					
T0 population		256.57 (128.63)	46.51 (21.76)	45.02 (14.77)	5.45 (1.06)
Ambient	24 h	286.53 (188.66)	45.50 (30.36)	47.34 (20.72)	5.63 (1.63)
	96h	222.33 (158.99)	35.08 (28.00)	38.39 (18.41)	5.38 (1.61)
750 μatm	24 h	250.58 (152.42)	43.55 (29.73)	41.98 (18.13)	5.61 (1.59)
	96 h	218.71 (160.44)	34.64 (30.97)	35.56 (21.39)	6.09 (1.78)
1000 μatm	24 h	260.74 (153.65)	43.36 (30.60)	44.10 (16.74)	5.78 (1.81)
	96 h	218.09 (192.69)	37.14 (40.93)	38.43 (18.88)	4.98 (1.98)

Table 3: Average (SD) elemental composition of copepods either at time zero (T0) or after 24 h or 96 h of incubation at ambient or elevated pCO₂ conditions. All copepods were stage CV with the exception of EO4 274 where a combination of CV and females were used.

Bioassay	T0 TEP (μg xanthan equiv. L ⁻¹) [SD]	Bottle effects $\Delta\text{TEP h}^{-1}$	Treatment effects				Copepod effects	
			24 h (750)	24 h (1000)	96 h (750)	96 h (1000)	24 h	96 h
JR271 (E01)	176.6 [16.7]	0.6677	↔	↔	↑	↑	↔	↑
JR271 (E03)	99.0 [10.0]	1.0553	↔	↔	↔	↔	↔	↑
JR271 (E04)	91.0 [6.98]	5.8465	↔	↔	↔	↑	↔	↓
JR271 (E05)	111.9 [11.5]	1.6261	↔	↔	↔	↑	↔	↑
JR274 (E03)	3.3 [1.7]	0.1004	↔	↔	↔	↓	↔	↑
JR274 (E04)	65.5 [2.9]	0.4496	↔	↑	↑	↑	↔	↔

Table 4: Average concentrations [SD] of transparent exopolymeric particles (TEP) at the start of the incubation experiments (T0). Bottle effects were calculated as $(\text{TEP}_{96\text{h, ambient, non-copepod}} - \text{TEP}_{\text{T0}})/96$ and represent the rate of increase of TEP over the maximum duration of the experiments (96 h) in the control incubations (ambient, non-copepod). Treatment effects were determined through performing a 1-way ANOVA comparing concentrations of TEP in incubations with elevated levels of pCO₂ with those of ambient conditions. All of incubations in this test contained copepods. Copepod effects were determined through performing a t-test comparing concentrations of TEP in incubations containing copepods with those without copepods. ↓ denotes significantly lower concentrations of TEP (P<0.05), ↑ significantly higher TEP (P<0.05) and ↔ no significant difference in TEP.

Bioassay Phytoplankton group (Phy _x)	T0 Cells mL ⁻¹	Bottle effects	Copepo d effects	Treatment effects			
				With copepods		Without copepods	
				750	1000	750	1000
E01 271							
Diatoms	282.65	+	+	-	-	+	+
Dinoflagellates	7.37	+	+	-	-	+	+
Ciliates	0.35	+	+	-	-	+	+
Flagellates	0.27	+++	+	-	-	+	+
E03 271							
Diatoms	5.19	-	-	++	++	+	+
Dinoflagellates	13.01	+	+	-	-	+	+
Ciliates	3.84	-	+	+	-	-	+
Flagellates	0.84	+++	+	+	+	-	-
E04 271							
Diatoms	249.96	++	-	-	+	-	-
Dinoflagellates	9.54	++	+	-	-	+	0
Ciliates	1.35	0	-	-	-	+	+
Flagellates	0.68	++	+	-	-	+	+
E05 271							
Diatoms	0.77	++	-	-	-	0	0
Dinoflagellates	27.4	+	-	-	-	-	-
Ciliates	0.08	+	--	++	+	+	-
Flagellates	0	0	+	-	-	-	+
E03 274							
Diatoms	14.03	++	-	+	+	-	-
Dinoflagellates	4.41	-	+	-	-	+++	+++
Ciliates	0.85	++	+	+	+	-	+
Flagellates	0.22	+	+	-	-	-	+
E04 274							
Diatoms	1084.44	-	+	-	0	+	-
Dinoflagellates	8.22	-	-	-	-	0	+
Ciliates	28.25	-	+	-	-	+	-
Flagellates	1.64	+	--	-	-	-	-

Table 5: Concentrations (cells mL⁻¹) of the four major phytoplankton groups (Phy_x) in each of the experimental incubations. Bottle effects were calculated as $(Phy_{x,T0} - Phy_{x,ambient, non-copepods})/Phy_{x,T0}$, Copepods effects as $(Phy_{x,ambient, copepods} - Phy_{x,treatment, copepods})/Phy_{x,ambient, copepods}$ and Treatment effects as $(Phy_{x,copepods} - Phy_{x,non-copepods})/Phy_{x,copepods}$. '0' represents fold changes of between -0.05 to +0.05; '+' or '-', up to a 1 fold increase; '++' or '--', a 1 to 5 fold increase; '+++' or '---', a more than 5 fold increase.

Figure Legends

Fig. 1: Proportion and total cell concentration (cells mL⁻¹) of the four major phytoplankton taxa at the start of each of the six incubation experiments (T₀).

Fig. 2: Carbon to nitrogen mass ratio (C:N) of a subset of the selected grazing copepods at the start of the incubation experiments (T₀). The species/stages of the selected copepod grazers for each experiment are listed in Table 1. Specimens from Station E01 271 were not analysed as a result of a processing error.

Fig. 3: Concentration of total bacterial cells (cell mL⁻¹) at the start of each incubation experiment (T₀) and in each of the three treatments (ambient, 750 μatm pCO₂ and 1000 μatm pCO₂). Open circle denotes average concentration within incubations with copepods, filled circles, without copepods. Bars represent 95% confidence intervals of incubations containing copepods. Instances where the filled circles fell outside these confidence intervals indicate significant differences between concentrations in copepod versus non-copepod incubations.

Fig. 4: Concentration of total nanoplankton cells (cell mL⁻¹) at the start of each incubation experiment (T₀) and in each of the three treatments (ambient, 750 μatm pCO₂ and 1000 μatm pCO₂). Open circle denotes average concentration of incubations with copepods, filled circles, without copepods. Bars represent 95% confidence intervals of incubations containing copepods. Instances where the filled circles fell outside of these confidence intervals indicate significant differences between concentrations in copepod versus non-copepod incubations.

Fig. 5: Relative change in bacterial cell concentrations in relation to the relative change in the concentration of transparent exopolymeric particles (TEP) across all six incubation experiments. Dashed lines represent trends fitted through least-squared regression. None of the trends were statistically significant.

Fig. 6: Relative change in nanoplankton cell concentrations in relation to the relative change in the concentration of transparent exopolymeric particles (TEP) across all six incubation experiments. Dashed lines represent trends fitted through least-squared regression. None of the trends were statistically significant.

Fig. 7: Relative response level of the four major phytoplankton taxa to treatments relative to that observed in incubations in ambient conditions. Response levels were graded as between 0 and 3 for each incubation experiment, as detailed in Table 1. * indicates a significant difference between incubations with copepods and those without copepods (paired t-test, $P < 0.05$).

Fig. 8: Carbon to nitrogen mass ratio (C:N) of the selected copepod grazers either at the start of the incubation experiments (T0) or at their completion. The species/stages of the selected copepod grazers for each experiment are listed in Table 1. Specimens from Station E01 271 were not analysed as a result of a processing error. * indicates a significant difference to the starting population (1-way ANOVA, $P < 0.05$).

Appendix 1: Parameter measurements on completion of the experimental incubations in ambient conditions. Starting conditions are given in Table 2.

Bioassay	Duration	Copepods	N:P (atomic ratio)	TA (μmol kg^{-1})	DIC (μmol kg^{-1})	pH	pCO ₂ (μatm)	Average (SD) TEP ($\mu\text{g L}^{-1}$ xanthan equiv.)	Average (SD) total bacteria (cells mL^{-1})	Average (SD) total nannoflag. (cells mL^{-1})
E01 271	24	Yes	7.0	2322.5	2080.0	8.1	305.2	149.5 (7.0)	922122.7 (126373.0)	6054.3 (1548.3)
	24	No	1.215	2316.8	2078.1	8.1	309.8	143.3 (9.8)	827276.1	5912.2
	96	Yes	2.1	2309.1	2099.4	8.1	364.3	135.2 (10.6)		
	96	No	1.843	2306.1	2121.6	8.0	422.7	244.4 (37.0)		
E03 271	24	Yes	18.2	2299.9	2146.9	8.1	348.4	142.7 (11.2)	924927.7 (26989.6)	43954.0 (1680.8)
	24	No	14.403	2297.6	2132.2	8.1	322.0	128.4 (7.4)	872248.4	40346.4
	96	Yes	28.6	2299.4	2154.6	8.1	368.0	223.9 (11.2)		
	96	No	17.778	2301.4	2125.3	8.1	302.0	200.3 (2.8)		
E04 271	24	Yes	5.9	2229.3	2120.9	8.0	386.7	116.6 (19.9)	423270.3 (5039.0)	4470.2 (114.5)
	24	No	3.923	2225.5	2102.3	8.1	344.1	142.0 (15.8)	515806.7	4656.7
	96	Yes	4.6	2228.9	2112.3	8.1	363.6	192.9 (26.1)		
	96	No	0.236	2228.4	2028.8	8.3	195.2	651.3 (7.4)		
E05 271	24	Yes	12.5	2309.1	2121.0	8.1	348.7	145.1 (8.1)	1380493.1 (69973.4)	12950.0 (2351.8)
	24	No	12.689	2311.5	2123.9	8.1	352.9	160.0 (9.8)	1430876.3	14696.8
	96	Yes	10.1	2313.5	2096.0	8.2	296.9	145.1 (3.7)		
	96	No	8.582	2307.7	2088.0	8.2	292.8	268.0 (35.4)		
E03 274	24	Yes	15.4	2291.1	2142.7	8.1	342.0	5.7 (5.0)	673354.5 (232254.2)	4476.0 (217.1)
	24	No	15.300	2288.1	2138.1	8.1	337.0	7.6 (1.2)	361558.2	1527.6
	96	Yes	16.0	2262.5	2130.9	8.1	374.0	26.6 (1.9)		
	96	No	15.808	2286.7	2138.4	8.1	339.0	12.9 (3.6)		
E04 274	24	Yes	15.7	2291.9	2113.1	8.2	259.0	51.0 (11.8)	1043846.7 (28753.2)	3727.4 (470.1)
	24	No	15.256	2293.7	2119.3	8.2	267.0	37.9 (21.4)	1155703.6	4354.8
	96	Yes	15.9	2296.5	2085.2	8.3	215.0	82.1 (12.2)		
	96	No	16.494	NA	2164.3	NA	NA	108.6 (25.6)		

Appendix 2: Parameter measurements on completion of the experimental incubations in the 750 μatm pCO_2 treatments. Starting conditions are given in Table 2.

Bioassay	Duration	Copepods	N:P (atomic ratio)	TA (μmol kg^{-1})	DIC (μmol kg^{-1})	pH	pCO_2 (μatm)	Average (SD) TEP (μg L^{-1} gum xanthan equiv.)	Average (SD) total bacteria (cells mL^{-1})	Average (SD) total nannoplankt. (cells mL^{-1})
E01 271	24	Yes	2.9	2322.0	2199.3	7.9	625.3	155.1 (8.4)	851978.2 (253396.5)	4011.9 (3242.0)
	24	No	2.0	2331.5	2207.0	7.9	620.6		756796.9	5064.8
	96	Yes	1.9	2333.5	2246.8	7.8	823.3	153.2 (7.5)		
	96	No	0.5	2328.8	2231.6	7.8	759.6			
E03 271	24	Yes	15.9	2314.0	2258.4	7.8	722.9	147.0 (6.7)	946909.8 (41416.8)	45213.7 (5527.9)
	24	No	15.4	2305.6	2235.6	7.8	640.9		950633.5	43088.2
	96	Yes	26.7	2308.1	2252.7	7.8	737.2	259.9 (20.7)		
	96	No	17.6	2306.2	2223.3	7.9	578.8			
E04 271	24	Yes	7.0	2231.7	2206.3	7.7	781.8	128.4 (20.1)	425599.1 (10102.0)	4869.8 (949.0)
	24	No	4.1	2232.1	2189.3	7.8	667.9		474527.5	5157.9
	96	Yes	5.5	2236.9	2197.9	7.8	706.5	183.6 (15.6)		
	96	No	0.1	NA	NA	NA	NA			
E05 271	24	Yes	12.4	2319.1	2226.3	7.8	664.2	140.2 (12.4)	1364566.6 (29206.0)	12495.6 (1920.1)
	24	No	12.7	2309.3	2215.1	7.8	653.1		1380145.0	15539.9
	96	Yes	9.9	2316.4	2188.5	7.9	531.4	170.0 (12.1)		
	96	No	9.1	2319.1	2195.6	7.9	535.8			
E03 274	24	Yes	15.2	2289.6	2180.9	8.0	451.0	3.3 (0.7)	836115.7 (19018.9)	2876.5 (1717.3)
	24	No	15.1	2291.8	2178.6	8.0	437.0		830448.8	3355.0
	96	Yes	15.9	2292.4	2174.3	8.0	420.0	27.5 (2.7)		
	96	No	15.8	2289.2	2177.6	8.0	440.0			
E04 274	24	Yes	15.4	2293.4	2178.3	8.0	398.0	49.0 (5.7)	1042692.9 (39800.6)	3492.6 (333.9)
	24	No	15.7	2296.0	2175.8	8.1	385.0		1248353.9	3576.1
	96	Yes	16.0	2298.0	2160.9	8.1	342.0	155.6 (9.5)		
	96	No	15.4	2296.4	2168.9	8.1	365.0	15.4		

Appendix 3: Parameter measurements on completion of the experimental incubations in the 1000 μatm pCO_2 treatments. Starting conditions are given in Table 2.

Bioassay	Duration	Copepods	N:P (atomic ratio)	TA (μmol kg^{-1})	DIC (μmol kg^{-1})	pH	pCO_2 (μatm)	Average (SD) TEP ($\mu\text{g L}^{-1}$ xanthan equiv.)	Average (SD) total bacteria (cells mL^{-1})	Average (SD) total nannoplankt. (cells mL^{-1})
E01 271	24	Yes	2.0	2334.3	2224.6	7.8	688.8	173.1 (21.0)	866416.6 (180754.5)	6313.0 (545.4)
	24	No	1.1	2342.3	2224.9	7.9	659.0		534542.3	3207.0
	96	Yes	2.8	2334.2	2263.9	7.7	943.8	173.7 (6.0)		
	96	No	1.0	2333.7	2253.1	7.7	866.5			
E03 271	24	Yes	17.2	2312.2	2272.8	7.7	827.1	130.9 (4.3)	950025.8 (47509.0)	47774.8 (2671.6)
	24	No	15.5	2310.8	2269.0	7.8	805.3		921361.4	40506.9
	96	Yes	23.0	2311.4	2299.2	7.7	1030.4	208.4 (19.7)		
	96	No	17.2	2314.9	2269.6	7.8	793.0			
E04 271	24	Yes	7.2	2240.1	2231.5	7.7	910.3	138.3 (7.5)	449130.9 (49422.8)	4965.5 (470.7)
	24	No	4.1	2233.4	2212.6	7.7	810.1		454930.6	4513.3
	96	Yes	3.0	2236.1	2211.4	7.8	774.9	377.7 (33.1)		
	96	No	0.1	2240.1	2148.4	8.0	417.8			
E05 271	24	Yes	12.1	2320.5	2261.9	7.7	870.0	151.3 (13.5)	1363161.4 (21832.2)	12451.0 (1685.0)
	24	No	12.9	2314.3	2247.2	7.8	807.1		1243892.1	11405.5
	96	Yes	10.1	2318.3	2234.5	7.8	716.7	179.3 (18.1)		
	96	No	1.0	2325.3	2209.9	7.9	577.5			
E03 274	24	Yes	15.4	2295.2	2206.9	7.9	528.0	5.0 (1.1)	824785.6 (18588.5)	2981.2 (163.9)
	24	No	15.2	2296.9	2211.3	7.92	540		481463.0	2800.4
	96	Yes	16.0	2297.8	2210.6	8.0	493.0	18.9 (2.4)		
	96	No	16.1	2292.2	2201.4	7.93	516			
E04 274	24	Yes	15.6	2294.1	2208.6	7.9	538.0	70.5 (2.9)	1060930.4 (125832.4)	3279.4 (228.2)
	24	No	15.3	2298.4	2203	7.98	463		1043688.3	3384.3
	96	Yes	16.1	2297.2	2199.0	8.0	453.0	120.0 (0.8)		
	96	No	16.3	2303.3	2184.4	8.04	390			





Fig. 1






Fig. 2

Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8