



OPEN ACCESS

ORIGINAL ARTICLE

Exome-wide analysis of rare coding variation identifies novel associations with COPD and airflow limitation in *MOCS3*, *IFIT3* and *SERPINA12*

Victoria E Jackson,¹ Ioanna Ntalla,^{1,2} Ian Sayers,³ Richard Morris,^{4,5} Peter Whincup,⁶ Juan-Pablo Casas,^{7,8} Antoinette Amuzu,⁹ Minkyung Choi,⁹ Caroline Dale,⁹ Meena Kumari,^{10,11} Jorgen Engmann,¹² Noor Kalsheker,¹³ Sally Chappell,¹³ Tamar Guetta-Baranes,¹³ Tricia M McKeever,¹⁴ Colin N A Palmer,¹⁵ Roger Tavendale,¹⁵ John W Holloway,^{16,17} Avan A Sayer,^{18,19} Elaine M Dennison,^{18,20} Cyrus Cooper,^{18,19} Mona Bafadhel,²¹ Bethan Barker,^{22,23} Chris Brightling,^{22,23} Charlotte E Bolton,²⁴ Michelle E John,²⁴ Stuart G Parker,²⁵ Miriam F Moffat,²⁶ Andrew J Wardlaw,^{22,23} Martin J Connolly,²⁷ David J Porteous,²⁸ Blair H Smith,²⁹ Sandosh Padmanabhan,³⁰ Lynne Hocking,³¹ Kathleen E Stirrups,^{2,32} Panos Deloukas,^{2,33} David P Strachan,⁶ Ian P Hall,³ Martin D Tobin,^{1,23} Louise V Wain¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/thoraxjnl-2015-207876>).

For numbered affiliations see end of article.

Correspondence to

Victoria E Jackson, Department of Health Sciences, University of Leicester, University Road, Leicester LE1 7RH, UK; vej3@le.ac.uk

Received 24 September 2015

Revised 5 January 2016

Accepted 29 January 2016

ABSTRACT

Background Several regions of the genome have shown to be associated with COPD in genome-wide association studies of common variants.

Objective To determine rare and potentially functional single nucleotide polymorphisms (SNPs) associated with the risk of COPD and severity of airflow limitation.

Methods 3226 current or former smokers of European ancestry with lung function measures indicative of Global Initiative for Chronic Obstructive Lung Disease (GOLD) 2 COPD or worse were genotyped using an exome array. An analysis of risk of COPD was carried out using ever smoking controls (n=4784). Associations with % predicted FEV₁ were tested in cases. We followed-up signals of interest ($p < 10^{-5}$) in independent samples from a subset of the UK Biobank population and also undertook a more powerful discovery study by meta-analysing the exome array data and UK Biobank data for variants represented on both arrays.

Results Among the associated variants were two in regions previously unreported for COPD; a low frequency non-synonymous SNP in *MOCS3* (rs7269297, $p_{\text{discovery}} = 3.08 \times 10^{-6}$, $p_{\text{replication}} = 0.019$) and a rare SNP in *IFIT3*, which emerged in the meta-analysis (rs140549288, $p_{\text{meta}} = 8.56 \times 10^{-6}$). In the meta-analysis of % predicted FEV₁ in cases, the strongest association was shown for a splice variant in a previously unreported region, *SERPINA12* (rs140198372, $p_{\text{meta}} = 5.72 \times 10^{-6}$). We also confirmed previously reported associations with COPD risk at *MMP12*, *HHIP*, *GPR126* and *CHRNA5*. No associations in novel regions reached a stringent exome-wide significance threshold ($p < 3.7 \times 10^{-7}$).

Conclusions This study identified several associations with the risk of COPD and severity of airflow limitation, including novel regions *MOCS3*, *IFIT3* and *SERPINA12*, which warrant further study.

Key messages

What is the key question?

- Do low frequency exonic variants influence susceptibility to COPD, and severity of airflow limitation?

What is the bottom line?

- Low frequency single nucleotide polymorphisms (SNPs) in *MOCS3* and *IFIT3* were associated with risk of COPD and a rare splice variant in *SERPINA12* was associated with severity of airflow limitation.

Why read on?

- These genomic regions have not previously been implicated in lung function or COPD and these findings could therefore provide further insight into COPD susceptibility and severity.

INTRODUCTION

COPD is a major public health concern, being a leading cause of morbidity and mortality worldwide.¹ The Global Initiative for Chronic Obstructive Lung Disease (GOLD) recommends that the impact of COPD on an individual patient should be assessed by considering breathlessness, symptoms and exacerbation risk, in combination with the severity of airflow limitation, which can be graded using %predicted FEV₁.² Approximately 1%–2% of COPD cases can be attributed to α 1-antitrypsin (AAT) deficiency, a rare inherited disorder, caused by mutations within the *SERPINA1* gene.^{3,4} For the remainder of COPD cases, cigarette smoking is recognised as the most

To cite: Jackson VE, Ntalla I, Sayers I, et al. Thorax Published Online First: [please include Day Month Year] doi:10.1136/thoraxjnl-2015-207876

Chronic obstructive pulmonary disease

significant risk factor⁵; however, there is also a genetic component, with several genomic regions showing association with COPD risk or airflow limitation to date, including *CHRNA3/5*, *HHIP*,³ *HTR4*, *GSTCD*, *TNS1*,⁶ *MMP12*^{7 8} and *FAM13A*.⁹ COPD diagnosis is confirmed using measures of lung function, so it is likely that the genetic determinants of COPD and lung function will overlap. Indeed, many loci identified in large genome-wide association studies (GWAS) of FEV₁ and the ratio of FEV₁ to forced vital capacity (FEV₁/FVC) in general population samples^{10–13} have subsequently been shown to be associated with COPD or airflow limitation.^{6 9 14 15}

Despite the successes in identifying genes associated with lung function and COPD, these known loci only explain a small proportion of the expected heritability.¹³ Large GWAS undertaken to date have generally focused on common variants (typically >5% minor allele frequency (MAF))^{3 9–14}; one hypothesis is that some of the so-called ‘missing heritability’ might be accounted for by variants of lower frequencies. In this study, we set out to investigate the role of low frequency, functional variants in COPD, and to confirm the role of single nucleotide polymorphisms (SNPs) previously showing association with lung function. It is hypothesised that rare variants are more likely than common variants to have deleterious effects; identifying such SNPs could lead to greater understanding of the pathways and biological mechanisms underlying airflow obstruction and COPD, and could translate to novel targets for treatment.

We genotyped cases with a history of smoking and airflow limitation, indicative of GOLD 2 COPD or worse, and control samples using an exome chip array to which we had added custom content comprising 2585 SNPs tagging regions which had shown suggestive association ($p < 2.21 \times 10^{-3}$) with lung function in a previous large genome-wide HapMap-imputed study.¹³ The exome chip genotyping array design contains mostly non-synonymous, splice or stop codon altering variants that are likely to affect protein structure and function, with the majority of variants being low frequency (MAF 1%–5%) or rare (MAF < 1%).

In this study, we carried out discovery case–control analyses (COPD cases vs controls) and analyses of %predicted FEV₁ in cases, as a measure of severity of airflow limitation. Replication was undertaken using a subset of the UK Biobank Lung Exome Variant Evaluation (BiLEVE) study, a collection of 48 931 individuals from UK Biobank with high-quality lung function and smoking data who were genotyped on an array that includes substantial overlap with the exome chip.¹⁶ We also adopted a more powerful discovery strategy for COPD risk and severity of airflow limitation, by meta-analysing data for the subset of exome chip variants that were measured in both the COPD exome chip consortium and the UK BiLEVE study.

METHODS

Study participants and phenotypes

A total of 3487 ever smokers with airflow limitation indicative of GOLD 2 COPD or worse were identified from 12 UK collections as cases (case collections described in online supplementary table S1). Individuals met case criteria if they had FEV₁/FVC ≤ 0.7 and %predicted FEV₁ $\leq 80\%$ (according to the National Health and Nutrition Examination Survey (NHANES) III spirometric reference equations¹⁷), did not have a doctor diagnosis of asthma and had reported current, or former smoking. Five of the sample collections (n=1398 samples, table 1) were COPD cohorts, with all individuals having irreversible airflow limitation, and meeting GOLD 2 criteria based on post-bronchodilator spirometry. The remaining cases were taken

from general population cohorts; for these samples, only pre-bronchodilator spirometry measures were available. We used general population controls with exome chip data, from Generation Scotland: Scottish Family Health Study (GS:SFHS), British 1958 Birth Cohort (1958BC), Oxford Biobank and GoDARTS (Genetics of Diabetes and Audit Research Tayside Study), listed in table 1 with clinical characteristics. All controls were current or former smokers and were free of lung disease, according to available spirometry and phenotype information.

We used a subset of the UK BiLEVE study¹⁶ for replication of novel signals, and for a larger discovery meta-analysis. A total of 24 457 heavy smokers (mean 35 pack-years) were genotyped as part of the UK BiLEVE study, selected such that 9748 individuals formed a low FEV₁ group (based on %predicted FEV₁), 4906 individuals formed a high FEV₁ group and 9803 had average FEV₁. We selected 4231 samples from the low FEV₁ group, with airflow limitation consistent with GOLD 2 or worse as cases and 8979 samples from the high and average FEV₁ groups with FEV₁/FVC > 0.7, %predicted FEV₁ > 80% and no doctor diagnosis of COPD for use as controls. All spirometry measures were prebronchodilator, all samples were heavy smokers and individuals with a doctor diagnosis of asthma or other lung diseases were excluded. The %predicted FEV₁ was estimated using NHANES III spirometric reference equations.¹⁷

An overview of the full study design is shown in figure 1.

Genotyping

All 3487 cases and 1032 GS:SFHS controls were genotyped together using the Illumina Human Exome BeadChip with additional custom content for regions which have previously shown modest association with lung function (description of custom content design in online supplementary methods). The remaining discovery analyses control samples were genotyped separately using the Illumina Human Exome BeadChip.

The UK BiLEVE samples were genotyped using the Affymetrix UK BiLEVE array, which includes rare variants selected from the same sequencing project as the Illumina Human Exome BeadChip alongside additional content.¹⁶ Of the 807 411 SNPs included on the Affymetrix UK BiLEVE array, 74 891 were also present on the Illumina Human Exome BeadChip; this subset of SNPs, which were directly genotyped on both arrays, was selected for the discovery meta-analysis.

Quality control of genotype data

Discovery exome analysis

Genotypes were called using Illumina’s Gencall algorithm in Genomestudio¹⁸ with refinement of rare variants with missing calls undertaken using zCall.¹⁹ Standard quality control (QC) filters were applied, in accordance with the Exome-chip Quality Control SOP V.5, as developed within the UK exome chip consortium²⁰ and are fully described in online supplementary methods. In brief, SNPs were excluded if they had low call rate (<99%) or deviated from Hardy Weinberg Equilibrium ($p < 10^{-4}$) and samples were excluded if they were duplicates, sex mismatches, heterozygosity outliers (>3 SD from mean), had an excess of singleton SNPs, or were ancestral outliers. Clusterplots for all SNPs of interest were inspected, to ensure accuracy of genotype calling.

UK BiLEVE data


The QC procedure of the UK BiLEVE genotype data is described elsewhere.¹⁶

Table 1 Clinical characteristics of samples passing genotype QC

Sample collection	n	Sex	Age	%Predicted FEV ₁	FEV ₁ /FVC	Pack-years	
		Male, n (%)	Mean (SD)	Mean (SD)	Mean (SD)	Samples with data (n)	Mean (SD)
Discovery analyses airflow limitation cases (total n=3226, with pack-years n=2517)							
GS:SFHS	508	224 (44.1%)	58.9 (8.94)	64.84 (12.64)	0.580 (0.108)	482	29.32 (24.96)
British Regional Heart Study	425	425 (100%)	70.1 (5.46)	59.41 (14.66)	0.597 (0.084)	0	—
British Women's Heart and Health Study	254	0 (0%)	69.3 (5.46)	64.26 (12.40)	0.603 (0.074)	203	28.1 (18.36)
UK COPD cohort*	209	129 (61.7%)	68.7 (8.11)	37.94 (15.29)	0.447 (0.119)	199	50.07 (27.79)
Hertfordshire Cohort Study	317	203 (64.0%)	66.1 (2.79)	62.89 (13.57)	0.589 (0.101)	312	32.25 (23.37)
COPDBeAT*	87	62 (71.3%)	67.6 (8.77)	45.19 (16.24)	0.480 (0.115)	86	38.69 (21.24)
Nottingham COPD study*	76	48 (63.2%)	67.2 (8.97)	50.29 (15.04)	0.482 (0.111)	74	49.02 (26.86)
Nottingham smokers	125	78 (62.4%)	63.1 (8.60)	46.27 (17.65)	0.503 (0.125)	124	41.75 (20.61)
Gedling study	33	26 (78.8%)	69.0 (8.23)	59.67 (16.81)	0.593 (0.103)	31	45.47 (33.40)
English Longitudinal Study of Aging	166	75 (45.2%)	66.0 (8.17)	54.84 (17.24)	0.526 (0.149)	0	—
EU COPD Gene Scan*	277	155 (56.0%)	67.0 (8.68)	38.51 (14.74)	0.467 (0.120)	277	46.43 (20.56)
GoTARDIS Study*	749	412 (55.0%)	68.8 (8.97)	52.16 (14.14)	0.509 (0.110)	729	43.26 (21.59)
Discovery analyses controls (total n=4784, with pack-years n=3889)							
GS:SFHS	961	552 (57.4%)	54.5 (8.41)	98.18 (10.92)	0.783 (0.051)	961	28.92 (16.86)
British 1958 Birth Cohort	1429	888 (62.1%)	44 (0)	100.90 (13.46)	0.809 (0.060)	1046	14.74 (10.07)
Oxford Biobank	1770	832 (47.0%)	41.6 (5.77)	—	—	1682	9.09 (9.34)
GoDARTS	624	402 (64.4%)	59.0 (10.75)	—	—	200	35.46 (25.89)
UK Biobank Lung Exome Variant Evaluation samples (meta-analysis and replication)							
Airflow limitation cases	4231	2379 (56.2%)	59.54 (6.86)	61.76 (11.8)	0.607 (0.076)	4231	42.41 (21.10)
Controls	8979	4260 (47.4%)	56.19 (7.92)	101.40 (8.1)	0.773 (0.038)	8979	30.43 (14.41)

*Sample collection is COPD case cohort.

GS:SFHS, Generation Scotland: Scottish Family Health Study; GoTARDIS, Tayside Allergy and Respiratory Disease Information System; QC, quality control.

† Case-control analysis of COPD cases with FEV₁/FVC ≤ 0.70 and %predicted FEV₁ ≤ 80% vs general population controls‡ Analysis of percent predicted FEV₁ in COPD cases**Figure 1** Two-stage study design. Stage 1: exome discovery analyses. Stage 2: Follow-up in UK BiLEVE: A. Replication of signals; B. meta-analysis of UK COPD exome chip consortium and UK BiLEVE.

Chronic obstructive pulmonary disease

Statistical analyses

SNP associations with COPD risk were carried out using a logistic regression model, adjusting for age, sex and pack-years and assuming an additive genetic model. Associations with untransformed %predicted FEV₁ in cases were tested, using a linear regression model, with adjustment for pack-years (analysis of severity of airflow limitation). Since not all samples had pack-years data available, secondary analyses were carried out without adjustment for pack-years, for both the COPD risk and severity of airflow limitation analyses, allowing the inclusion of all samples. Single variant analyses were carried out using PLINK V1.07.²¹ Using a Bonferroni correction for the number of tests undertaken, a significance level of $p < 3.7 \times 10^{-7}$ would be required in the exome single variant analysis to retain a type 1 error of 5%. We defined SNPs of interest as those with $p < 10^{-5}$ in the discovery exome analysis; for these SNPs, we undertook replication analyses in the UK BiLEVE study to corroborate findings (see online supplementary methods). We set a Bonferroni corrected significance level for replication, for the number of SNPs in novel loci taken forward to replication ($p < 0.017$ for analysis of COPD risk). Gene-based analyses using SKAT-O were additionally undertaken; the methods and results of these analyses are described in the online supplementary information.

Custom content single variant analyses

Custom content comprising 2585 SNPs tagging regions which had shown suggestive association ($p < 2.21 \times 10^{-3}$) with lung function in a previous large genome-wide HapMap-imputed study¹³ were also included on the array for cases and GS:SFHS controls. Additional controls from 1958BC and Busselton Health Study (BHS) with genome-wide data were also used; full methods and results of this analysis are given in the supplementary information.

Meta-analysis with UK BiLEVE data

Single variant associations with COPD risk and severity of airflow limitation in the UK BiLEVE samples were carried out using PLINK v1.07,²¹ identically to the corresponding discovery analysis with pack-years adjustment. We carried out an inverse-variance-weighted meta-analysis of the union of SNPs included in the discovery exome and UK BiLEVE analyses (described in online supplementary methods).

RESULTS

Discovery exome analysis

3226 cases and 4784 controls passed all sample and SNP genotype QC and were used in the exome analysis (exclusions in online supplementary table S1). Clinical characteristics of these samples are summarised in table 1. Of the SNPs which passed all QC criteria in both cases and controls, 135 818 were polymorphic, of which 101 308 (74.6%) had a MAF < 1%.

Analyses of COPD risk

We carried out pack-years adjusted analysis of COPD risk, including 2517 cases and 3889 controls, in addition to an unadjusted analysis, using all 3226 cases and 4784 controls (quantile-quantile plots shown in online supplementary figure S1). A total of four SNPs in three regions met the $p < 10^{-5}$ significance threshold in the pack-years adjusted analysis, with five SNPs in four regions showing $p < 10^{-5}$ in the unadjusted analysis (figure 2).

In the pack-years adjusted analysis (table 2A and figure 2A), the most significant association was for the previously reported COPD/smoking region 15q25 (sentinel SNP rs8034191 OR: 1.38, MAF=34.8%, $p = 2.42 \times 10^{-7}$). This signal was replicated in the UK BiLEVE study. Two novel signals of association with COPD risk ($p < 10^{-5}$) were rs3813803 within *SMPDL3B* (OR: 1.37, MAF=29.2%, $p = 1.04 \times 10^{-6}$) and low frequency SNP rs7269297 within *MOCS3* (OR: 0.25, MAF=1.1%, $p = 3.08 \times 10^{-6}$). There was evidence of replication, just above the Bonferroni corrected level of significance ($p < 0.017$) for rs7269297 in the UK BiLEVE study ($p = 7.27 \times 10^{-5}$ for meta-analysis of discovery and UK BiLEVE results, table 2A).

A further two loci were associated with COPD risk in the analysis unadjusted for pack-years: rs3827522 within *PRICKLE1* (OR: 0.12, MAF=0.4%, $p = 1.03 \times 10^{-7}$) and rs17368582 within *MMP12* (OR: 0.712, MAF=12.2%, $p = 5.01 \times 10^{-6}$, table 2A and figure 2B); however, there was no evidence of replication of these associations with COPD risk in UK BiLEVE. rs2276109, another SNP within *MMP12*, (MAF=5.6%) which is strongly correlated with rs17368582 ($r^2 = 0.84$), has previously been associated with COPD risk in smokers.⁷ Overall, no associations in novel regions met exome-wide significance ($p < 3.7 \times 10^{-7}$).

Analyses of severity of airflow limitation

Although no SNPs reached the $p < 10^{-5}$ significance level in either the pack-years adjusted, or the unadjusted analysis (see online supplementary figures S2 and S3), six SNPs showed some evidence of association ($p < 10^{-4}$) in one or both analyses (see online supplementary table S2). Of note, rs28929474, the z-allele within the *SERPINA1* gene, showed modest association in the unadjusted analysis ($\beta = -6.17\%$, MAF=2.0%, $p = 2.83 \times 10^{-5}$).

UK BiLEVE meta-analysis results

Analyses of COPD risk

For the 57 234 polymorphic SNPs common to both the COPD exome chip consortium samples and the UK BiLEVE study, a meta-analysis of discovery and UK BiLEVE study results was undertaken in which three regions showed association with risk of COPD ($p < 10^{-5}$, figure 3, online supplementary figure S4 and table 2B). The *GYPB/HHIP* and *GPR126* regions have previously been reported as showing association with lung function and COPD or airflow limitation risk.^{3 10 14} The *IFIT3* region signal (rs140549288 p.Val352Leu in *IFIT3*, OR: 1.92, MAF=0.7%, $p = 7.49 \times 10^{-6}$) represents a novel rare variant signal of association with COPD.

Analyses of severity of airflow limitation

A total of 54 168 SNPs were included in the meta-analysis of severity of airflow limitation (see online supplementary figures S5 and S6). One SNP showed association with $p < 10^{-5}$: rs140198372, a variant which alters the sequence at a site where the splicing of an intron takes place (splice site) in *SERPINA12* ($\beta = -33.51\%$, MAF=0.03%, $p = 5.72 \times 10^{-6}$, table 3).

Sensitivity analyses to assess COPD case criteria

Of our 3226 COPD cases defined as described above, 1398 also had a GOLD 2 or worse COPD based on postbronchodilator spirometry. We carried out a sensitivity analysis for all SNPs identified in our discovery or meta-analyses of COPD risk, by repeating the discovery analyses including only those 1398 COPD cases which underwent reversibility testing. This analysis




Figure 2 (A) Analysis of COPD risk, with pack-years adjustment (single nucleotide polymorphisms (SNPs) with minor allele frequency (MAF) $>0.05\%$ only; SNPs with $p < 10^{-5}$ highlighted). (B) Analysis of COPD risk, without pack-years adjustment (SNPs with MAF $>0.05\%$ only; SNPs with $p < 10^{-5}$ highlighted).

showed consistent estimated effect sizes (see online supplementary table S3 and figure S7), and in particular, the ORs were not substantially attenuated for rs7269297 in *MOCS3* (sensitivity analysis OR: 0.276; original discovery OR: 0.251), nor rs140549288 in *IFIT3* (sensitivity analysis OR: 2.554; original discovery OR: 2.156).

Association of novel loci with smoking behaviour

Given the disparity of smoking behaviour in our cases and control samples (table 1), we further investigated whether either of the two novel COPD risk loci were associated with smoking behaviour, to ascertain whether the associations with COPD may be explained by differences in smoking. Neither of the sentinel SNPs showed significant association with heavy versus never smoking within UK BiLEVE ($p=0.956$ for rs7269297 and $p=0.945$ for rs140549288) study. We further undertook a look-up in the publically available results of a GWAS from the Tobacco and Genetics consortium²² for associations with rs7269297 in *MOCS3* (rs140549288 was not available in data) and a number of smoking traits; however, no evidence for association with smoking behaviour was found (cigarettes per day

$p=0.610$; ever vs never smoking $p=0.172$; current vs former smoking $p=0.699$).

DISCUSSION

We carried out analyses of exome chip variants with COPD risk and %predicted FEV₁ among cases, through which we identified a number of SNPs in both known COPD regions and at novel loci that showed suggestive association ($p < 10^{-5}$) with risk of COPD. These novel regions (region plots: online supplementary figure S8) warrant further investigation as they may provide insight into the underlying biological mechanisms of COPD and airflow limitation in smokers and could provide novel therapeutic targets. The most significant associations in both the discovery exome analysis and the meta-analysis were with SNPs in the 15q25 region, previously identified through GWAS as being associated with smoking behaviour,^{22–24} lung cancer,²⁵ COPD³ and airflow obstruction.¹⁴ In addition, we independently replicated previously reported associations of *HHIP*,^{3 10} *GPR126*¹⁴ and *MMP12*^{7 8} with COPD risk.

We identified novel associations between COPD risk and low frequency or rare coding SNPs in two genes: *MOCS3*

Table 2 Top associations in exome discovery analyses and meta-analysis of COPD risk(A) SNPs with $p < 10^{-5}$ in either the pack-years adjusted or unadjusted discovery analyses

rs no.	CHR	Position	Coded allele	Gene	Discovery pack-years adjusted analysis (2517 cases, 3889 controls)				Discovery unadjusted analysis (3226 cases, 4784 controls)				UK BiLEVE pack-years adjusted analysis (4231 cases, 8979 controls)				Meta-analysis of discovery and UK BiLEVE pack-year adjusted analyses	
					MAF (MAC)		Association result		MAF (MAC)		Association result		MAF (MAC)		Association result		Association result	
					Cases	Controls	OR (95% CI)	p Value*	Cases	Controls	OR (95% CI)	p Value*	Cases	Controls	OR (95% CI)	p Value*	OR (95% CI)	p Value*
rs3813803	1	28282292	C	<i>SMPDL3B</i> (non-synonymous)	30.6% (1541)	28.3% (2203)	1.370 (1.207 to 1.554)	2.41×10^{-6}	30.3% (1956)	28.5% (2722)	1.288 (1.160 to 1.430)	2.11×10^{-6}	28.7% (2418)	29.4% (5269)	0.968 (0.911 to 1.029)	0.298	1.033 (0.978 to 1.092)	0.241
rs17368582	11	102738075	C	<i>MMP12</i> (synonymous)	11.1% (561)	12.9% (1001)	0.767 (0.642 to 0.915)	3.22×10^{-3}	11.1% (719)	12.8% (1229)	0.712 (0.615 to 0.824)	5.01×10^{-6}	12.0% (1015)	12.2% (2198)	0.982 (0.902 to 1.069)	0.676	0.938 (0.868 to 1.013)	0.101
rs3827522	12	42853871	A	<i>PRICKLE1</i> (non-synonymous)	0.2% (11)	0.4% (27)	0.184 (0.065 to 0.519)	1.39×10^{-3}	0.2% (14)	0.5% (46)	0.123 (0.057 to 0.266)	1.03×10^{-7}	0.3% (21)	0.3% (45)	0.907 (0.518 to 1.585)	0.731	0.633 (0.386 to 1.039)	0.071
rs8034191	15	78806023	C	near <i>AGPHD1</i> (intergenic)	38.0% (1912)	32.7% (2546)	1.374 (1.218 to 1.550)	2.42×10^{-7}	37.7% (2432)	32.9% (3144)	1.364 (1.234 to 1.507)	1.18×10^{-9}	39.2% (3315)	35.2% (6320)	1.156 (1.092 to 1.224)	6.85×10^{-7}	1.193 (1.133 to 1.257)	2.79×10^{-11}
rs7269297	20	49576664	G	<i>MOC53</i> (non-synonymous)	0.7% (37)	1.4% (110)	0.251 (0.140 to 0.448)	3.08×10^{-6}	0.8% (54)	1.5% (139)	0.423 (0.262 to 0.680)	3.98×10^{-4}	1.2% (98)	1.4% (252)	0.742 (0.578 to 0.953)	0.019	0.626 (0.497 to 0.789)	7.27×10^{-5}

(B) SNPs with $p < 10^{-5}$ in the meta-analysis (only most statically significant SNP in each region shown)

rs no.	CHR	Position	Coded allele	Gene	Discovery pack-years adjusted analysis (2517 cases, 3889 controls)				UK BiLEVE pack-years adjusted analysis (4231 cases, 8979 controls)				Meta-analysis of discovery and UK BiLEVE pack-year adjusted analyses	
					MAF (MAC)		Association result		MAF (MAC)		Association result		Association result	
					Cases	Controls	OR (95% CI)	p Value*	Cases	Controls	OR (95% CI)	p Value*	OR (95% CI)	p Value*
rs1828591	4	145480780	A	GYPA/HHIP (intergenic)	35.6% (1794)	39.1% (3042)	0.9167 (0.814, 1.032)	0.153	36.6% (3088)	40.0% (771)	0.867 (0.819, 0.918)	9.88×10^{-7}	0.876 (0.832, 0.922)	5.75×10^{-7}
rs4896582	6	142703877	A	GPR126 (intronic)	29.3% (1473)	31.7% (2468)	0.8594 (0.757, 0.974)	0.018	28.0% (2349)	30.2% (5344)	0.879 (0.826, 0.934)	3.87×10^{-5}	0.875 (0.827, 0.925)	2.53×10^{-6}
rs140549288	10	91099466	C	IFIT3 (exonic), LIPA (intronic)	0.8% (38)	0.6% (44)	2.156 (1.046, 4.445)	0.037	0.9% (79)	0.6% (100)	1.880 (1.378, 2.565)	6.87×10^{-5}	1.924 (1.441, 2.560)	8.56×10^{-6}

*p Values in bold significant at $p < 10^{-5}$ level.

BiLEVE, Biobank Lung Exome Variant Evaluation; MAC, minor allele count; MAF, minor allele frequency; SNPs, single nucleotide polymorphisms.




Figure 3 Meta-analysis of COPD risk in discovery exome analysis and UK Biobank Lung Exome Variant Evaluation samples.

(rs7269297, serine to alanine, MAF=1.3%, $p_{\text{discovery}}=3.08 \times 10^{-6}$, PolyPhen prediction: benign) and *IFIT3* (rs140549288, valine to leucine, MAF=0.7%, $p_{\text{meta}}=8.56 \times 10^{-6}$, PolyPhen prediction: benign). The protein encoded by *MOCS3* adenylates and activates molybdopterin synthase, an enzyme required to synthesise molybdenum cofactor²⁶ and is expressed in bronchial epithelium and smooth muscle layer of the bronchus.²⁷ *IFIT3* is associated with interferon- α antiviral activity and has been found to be up-regulated in respiratory syncytial virus infection²⁸ and in human lung epithelial cells infected with dengue virus.²⁹ The SNP rs140549288 is also located within an intron of *LIPA*; the product of this gene is involved in the hydrolysis of cholesterol esters and triglycerides and other SNPs within this gene have previously been associated with coronary artery disease.³⁰

The z-allele within the *SERPINA1* gene was associated with a lower %predicted FEV₁ in cases (unadjusted analysis: $p_{\text{discovery}}=2.83 \times 10^{-5}$); as well as being a well-established cause of AAT deficiency,^{3 4} this SNP has also previously been associated with an increased annual decline in FEV₁ in a general population sample³¹ and increased airflow limitation in COPD cases.³² In the present study, the z-allele was associated with an increased risk of COPD, although this was not statistically significant (OR: 1.27, $p=0.252$). The likely reason for the lack of a significant association with this known COPD locus is that some of the case collections excluded individuals with AAT deficiency, resulting in selection bias. In the meta-analysis of severity of

airflow limitation, we identified a very rare SNP within another serine protease inhibitor gene, *SERPINA12*, not previously associated with COPD (rs140198372, MAF=0.03%, $p_{\text{meta}}=5.72 \times 10^{-6}$). *SERPINA12* and *SERPINA1* lie 96.6 kb apart on chromosome 14 (rs140198372 and the z-allele in *SERPINA1* are not in linkage disequilibrium ($r^2=9.0 \times 10^{-6}$)). *SERPINA12* has been associated with cardiovascular diseases, being implicated in obesity and type 2 diabetes.³³

One of the primary challenges associated with identifying low frequency variants associated with disease is limited statistical power, and this could explain our lack of strong statistically significant findings. Indeed, none of the reported associations in novel regions met a stringent exome-wide significance level ($p < 3.8 \times 10^{-7}$) overall. In the present study, we would have just 54% power to detect an association with an SNP associated with COPD risk with a MAF of 1% and an OR of 2, at the $p < 3.8 \times 10^{-7}$ level. Furthermore, recent analyses undertaken by the UK10K Consortium found no evidence of low frequency SNPs having large effects, upon a series of traits.³⁴ Due to the limited power to detect single variant associations of rare variants with modest effect sizes, we additionally adopted gene-based analyses using SKAT-O, a method which combines information from several rare variants (see online supplementary information). In these analyses, we only identified one gene meeting our elected significance level ($p < 10^{-5}$); this gene-based signal in *PRICKLE1* was found however, to be driven by a single SNP, which was identified as being associated with COPD risk in

Table 3 Top associations ($p < 10^{-5}$) in meta-analysis of severity of airflow limitation

rs no.	CHR	Position	Coded allele	Gene	Severity of airflow limitation, adjusted for pack-years (n=2517)			UK BiLEVE pack-years adjusted analysis (n=4231)			Meta-analysis of discovery and UK BiLEVE pack-year adjusted analyses	
					MAF (MAC)	Beta (95% CI)	p Value	MAF (MAC)	Beta (95% CI)	p Value	Beta (95% CI)	p Value
rs140198372	14	94953832	A	<i>SERPINA12</i> (splice site)	0.059% (3)	−29.23 (−49.50 to −8.96)	2.59×10^{-5}	0.012% (1)	−38.35 (−59.88 to −16.82)	4.11×10^{-4}	−33.51 (−48.27 to −18.76)	5.72×10^{-6}

*p Values in bold significant at $p < 10^{-5}$ level.

BiLEVE, Biobank Lung Exome Variant Evaluation; MAC, minor allele count; MAF, minor allele frequency.

Chronic obstructive pulmonary disease

the single variant discovery analysis, but which was not replicated in the UK BiLEVE data.

Another limitation of this study is that a number of our cases had only prebronchodilator spirometry; for these samples, it could not be determined whether their airflow limitation was reversible, and so a proportion of these cases may not have met the clinical definition of COPD. We undertook case-control sensitivity analyses using our discovery samples, restricting cases to the subset of 1398 individuals taken from COPD cohorts and who had known irreversible airflow limitation. The effect estimates of our top hits did not substantially change in this sensitivity analysis, suggesting that our broader case definition, including samples that did not undergo reversibility testing, did not result in substantial misclassification bias.

A further potential source of bias in this study was the heavier smoking history in our cases compared with the control samples. For the two SNPs identified through the analyses of COPD risk, we found no evidence of association with smoking in data from the UK BiLEVE study, suggesting that the associations with COPD risk were not driven by the imbalances in smoking behaviour.

Finally, it was not possible to validate the findings of this study through additional genotyping; however for the three reported loci, consistent results were observed in both the discovery and the UK BiLEVE samples. It would not be expected to see the same false positive result in these two independent samples, therefore, strengthening the evidence for these being true associations.

In summary, we have identified potentially interesting associations with low frequency and rare SNPs and COPD risk in two regions not previously implicated in COPD or lung function. We further identified an association of %predicted FEV₁ in individuals with COPD with a very rare SNP in *SERPINA12*. Further confirmation of these associations in larger independent collections of COPD cases and controls is needed. This study also provides further evidence that the z-allele within *SERPINA1* may be related to severity of airflow limitation in COPD. While large sample sizes may be required to definitively identify novel loci, we present evidence to support the notion that the genetic contribution to COPD risk comprises polygenic contributions of rare, low frequency and common genetic variants. Future studies, alone or in combination, should aim to target the full allele frequency range to unravel the genetic architecture of COPD.

Author affiliations

- ¹Department of Health Sciences, University of Leicester, Leicester, UK
- ²William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK
- ³Division of Respiratory Medicine, University of Nottingham, Queen's Medical Centre, Nottingham, UK
- ⁴School of Social & Community Medicine, University of Bristol, Bristol, UK
- ⁵Department of Primary Care & Population Health, UCL, London, UK
- ⁶Population Health Research Institute, St George's, University of London, London, UK
- ⁷University College London, Farr Institute of Health Informatics, London, UK
- ⁸Cochrane Heart Group, London, UK
- ⁹Department of Non-communicable Disease Epidemiology, Faculty of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, London, UK
- ¹⁰SER, University of Essex, Colchester, Essex, UK
- ¹¹Department of Epidemiology and Public Health, UCL, London, UK
- ¹²Institute of Cardiovascular Science, UCL, London, UK
- ¹³School of Life Sciences, University of Nottingham, Nottingham, UK
- ¹⁴Division of Epidemiology and Public Health, Nottingham City Hospital, University of Nottingham, Nottingham, UK
- ¹⁵Cardiovascular and Diabetes Medicine, School of Medicine, University of Dundee, Dundee, UK
- ¹⁶Human Development & Health, Faculty of Medicine, University of Southampton, Southampton General Hospital, Southampton, UK

- ¹⁷NIHR Southampton Respiratory Biomedical Research Unit, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton General Hospital, Southampton, UK
- ¹⁸MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton General Hospital, Southampton, UK
- ¹⁹NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton General Hospital, Southampton, UK
- ²⁰Victoria University, Wellington, New Zealand
- ²¹Respiratory Medicine Unit, Nuffield Department of Medicine, University of Oxford, Oxford, UK
- ²²Institute for Lung Health, Department of Infection, Immunity and Inflammation, University of Leicester, Leicester, UK
- ²³National Institute for Health Research Respiratory Biomedical Research Unit, Glenfield Hospital, Leicester, UK
- ²⁴Nottingham Respiratory Research Unit, University of Nottingham, City Hospital Campus, Nottingham, UK
- ²⁵Institute for Ageing and Health, Newcastle University, Campus for Ageing and Vitality, Newcastle upon Tyne, UK
- ²⁶Department of Molecular Genetics and Genomics, National Heart and Lung Institute, Imperial College London, London, UK
- ²⁷Freemasons' Department of Geriatric Medicine, University of Auckland, New Zealand
- ²⁸Generation Scotland, Centre for Genomic and Experimental Medicine, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK
- ²⁹Division of Population Health Sciences, University of Dundee, Dundee, UK
- ³⁰Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, UK
- ³¹Institute of Medical Sciences, University of Aberdeen, Aberdeen, UK
- ³²Department of Haematology, University of Cambridge, Cambridge, UK
- ³³Princess Al-Jawhara Al-Brahim Centre of Excellence in Research of Hereditary Disorders, King Abdulaziz University, Jeddah, Saudi Arabia

Acknowledgements This research used the ALICE and SPECTRE High Performance Computing Facilities at the University of Leicester and was supported by the National Institute for Health Research (NIHR) Leicester Respiratory Biomedical Research Unit. This article/paper/report presents independent research funded partially by the NIHR. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. This research has been conducted using the UK Biobank Resource.

Contributors Case collection study concept, or data acquisition and quality control: IS, IPH, DPS, RM, PW, JPC, AA, MC, CD, MK, JE, NK, SC, TGB, TMM, CNAP, RT, JWH, AAS, EMD, CC, MB, BB, CB, CEB, MEJ, SGP, MFM, AJW, MJC, BJP, BHS, SP and LH. Genotype data acquisition and QC: KES and PD. Central study design, analysis and writing of manuscript: VEJ, IN, LVW, MDT, IPH and IS.

Funding British Women's Heart and Health Study is funded by the Department of Health grant no. 90049 and the British Heart Foundation grant no. PG/09/022. British Regional Heart Study is supported by the British Heart Foundation (grant RG/13/16/30528). CB (COPDBeAT) received funding from the Medical Research Council UK (grant no. G0601369), CB (COPDBeAT) and AJW (UKCOPD) were supported by the National Institute for Health Research (NIHR Leicester Biomedical Research Unit). MB (COPDBeAT) received funding from the NIHR (grant no. PDF-2013-06-052). Hertfordshire Cohort Study received support from the Medical Research Council, Arthritis Research UK, the International Osteoporosis Foundation and the British Heart Foundation; NIHR Biomedical Research Centre in Nutrition, University of Southampton; NIHR Musculoskeletal Biomedical Research Unit, University of Oxford. Generation Scotland: Scottish Family Health Study is funded by the Chief Scientist Office, Scottish Government Health Directorates, grant number CZD/16/6 and the Scottish Funding Council grant HR03006. EU COPD Gene Scan is funded by the European Union, grant no. QLG1-CT-2001-01012. English Longitudinal Study of Aging is funded by the Institute of Aging, NIH grant No. AG176440651. GoDARTs is funded by the Wellcome Trust grants 072960, 084726 and 104970. MDT has been supported by MRC fellowship G0902313. UK Biobank Lung Exome Variant Evaluation study was funded by a Medical Research Council strategic award to MDT, IPH, DPS and LVW (MC_PC_12010).

Competing interests None declared.

Ethics approval Several (meta-analysis design).

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- Rabe KF, Hurd S, Anzueto A, *et al.* Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2007;176:532–55.
- Global Strategy for the Diagnosis, Management and Prevention of COPD. 2015. http://www.goldcopd.org/uploads/users/files/GOLD_Report_2015_Apr2.pdf
- Pillai SG, Ge D, Zhu G, *et al.* A genome-wide association study in chronic obstructive pulmonary disease (COPD): identification of two major susceptibility loci. *PLoS Genet* 2009;5:e1000421.
- Thun GA, Imboden M, Ferrarotti I, *et al.* Causal and synthetic associations of variants in the *SERPINA* gene cluster with alpha1-antitrypsin serum levels. *PLoS Genet* 2013;9:e1003585.
- Mayer AS, Newman LS. Genetic and environmental modulation of chronic obstructive pulmonary disease. *Respir Physiol* 2001;128:3–11.
- Soler Artigas M, Wain LV, Repapi E, *et al.* Effect of five genetic variants associated with lung function on the risk of chronic obstructive lung disease, and their joint effects on lung function. *Am J Respir Crit Care Med* 2011;184:786–95.
- Hunninghake GM, Cho MH, Tesfaigzi Y, *et al.* MMP12, lung function, and COPD in high-risk populations. *N Engl J Med* 2009;361:2599–608.
- Haq I, Chappell S, Johnson SR, *et al.* Association of MMP-12 polymorphisms with severe and very severe COPD: a case control study of MMPs-1, 9 and 12 in a European population. *BMC Med Genet* 2010;11:7.
- Cho MH, Boutaoui N, Klanderman BJ, *et al.* Variants in *FAM13A* are associated with chronic obstructive pulmonary disease. *Nat Genet* 2010;42:200–2.
- Wilk JB, Chen TH, Gottlieb DJ, *et al.* A Genome-wide association study of pulmonary function measures in the Framingham Heart Study. *PLoS Genet* 2009;5:e1000429.
- Repapi E, Sayers I, Wain LV, *et al.* Genome-wide association study identifies five loci associated with lung function. *Nat Genet* 2010;42:36–44.
- Hancock DB, Eijgelsheim M, Wilk JB, *et al.* Meta-analyses of genome-wide association studies identify multiple loci associated with pulmonary function. *Nat Genet* 2010;42:45–52.
- Soler Artigas M, Loth DW, Wain LV, *et al.* Genome-wide association and large-scale follow up identifies 16 new loci influencing lung function. *Nat Genet* 2011;43:1082–90.
- Wilk JB, Shrine NRG, Loehr LR, *et al.* Genome-wide association studies identify *CHRNA5/3* and *HTR4* in the development of airflow obstruction. *Am J Respir Crit Care Med* 2012;186:622–32.
- Castaldi PJ, Cho MH, Litonjua AA, *et al.* The association of genome-wide significant spirometric loci with chronic obstructive pulmonary disease susceptibility. *Am J Respir Cell Mol Biol* 2011;45:1147–53.
- Wain L, Shrine N, Miller S, *et al.* Novel insights into the genetics of smoking behaviour, lung function, and chronic obstructive pulmonary disease (UK BiLEVE): a genetic association study in UK Biobank. *Lancet Respir Med* 2015;3:769–81.
- Hankinson JL, Odencrantz JR, Fedan KB. Spirometric reference values from a sample of the general U.S. population. *Am J Respir Crit Care Med* 1999;159:179–87.
- Illumina Inc. Illumina GenCall Data Analysis Software. 2005.
- Goldstein JI, Crenshaw A, Carey J, *et al.* zCall: a rare variant caller for array-based genotyping: Genetics and population analysis. *Bioinformatics* 2012;28:2543–5.
- Mahajan A, Robertson N, Rayner W. Exome-Chip Quality Control SOP. Version 5, 2012-11-20. 2012.
- Purcell S, Neale B, Todd-Brown K, *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;81:559–75.
- The Tobacco and Genetics Consortium. Genome-wide meta-analyses identify multiple loci associated with smoking behavior. *Nat Genet* 2010;42:441–7.
- Thorgerirsson TE, Gudbjartsson DF, Surakka I, *et al.* Sequence variants at *CHRNA3-CHRNA6* and *CYP2A6* affect smoking behavior. *Nat Genet* 2010;42:448–53.
- Liu JZ, Tozzi F, Waterworth DM, *et al.* Meta-analysis and imputation refines the association of 15q25 with smoking quantity. *Nat Genet* 2010;42:436–40.
- McKay JD, Hung RJ, Gaborieau V, *et al.* Lung cancer susceptibility locus at 5p15.33. *Nat Genet* 2008;40:1404–6.
- Matthies A, Nimtz M, Leimkühler S. Molybdenum cofactor biosynthesis in humans: identification of a persulfide group in the rhodanese-like domain of MOCS3 by mass spectrometry. *Biochemistry* 2005;44:7912–20.
- Uhlen M, Oksvold P, Fagerberg L, *et al.* Towards a knowledge-based Human Protein Atlas. *Nat Biotech* 2010;28:1248–50.
- Ternette N, Wright C, Kramer HB, *et al.* Label-free quantitative proteomics reveals regulation of interferon-induced protein with tetratricopeptide repeats 3 (IFIT3) and 5'-3'-exoribonuclease 2 (XRN2) during respiratory syncytial virus infection. *Virology* 2011;8:442.
- Hsu YL, Shi SF, Wu WL, *et al.* Protective roles of interferon-induced protein with tetratricopeptide repeats 3 (IFIT3) in dengue virus infection of human lung epithelial cells. *PLoS ONE* 2013;8:e79518.
- Wild PS, Zeller T, Schillert A, *et al.* A genome-wide association study identifies *LIPA* as a susceptibility gene for coronary artery disease. *Circ Cardiovasc Genet* 2011;4:403–12.
- Dahl M, Tybjaerg-Hansen A, Lange P, *et al.* Change in lung function and morbidity from chronic obstructive pulmonary disease in α 1-antitrypsin MZ heterozygotes: a longitudinal study of the general population. *Ann Intern Med* 2002;136:270–9.
- Molloy K, Hersh CP, Morris VB, *et al.* Clarification of the risk of chronic obstructive pulmonary disease in α 1-antitrypsin deficiency PiMZ heterozygotes. *Am J Respir Crit Care Med* 2014;189:419–27.
- Kim DS, Burt AA, Crosslin DR, *et al.* Novel common and rare genetic determinants of paraoxonase activity: *FTO*, *SERPINA12*, and *ITGAL*. *J Lipid Res* 2013;54:552–60.
- Walter K, Min JL, Huang J, *et al.* UK10K Consortium. The UK10K project identifies rare variants in health and disease. *Nature* 2015;526:82–90.

THORAX

Exome-wide analysis of rare coding variation identifies novel associations with COPD and airflow limitation in *MOCS3*, *IFIT3* and *SERPINA12*

Victoria E Jackson, Ioanna Ntalla, Ian Sayers, Richard Morris, Peter Whincup, Juan-Pablo Casas, Antoinette Amuzu, Minkyung Choi, Caroline Dale, Meena Kumari, Jorgen Engmann, Noor Kalsheker, Sally Chappell, Tamar Guetta-Baranes, Tricia M McKeever, Colin N A Palmer, Roger Tavendale, John W Holloway, Avan A Sayer, Elaine M Dennison, Cyrus Cooper, Mona Bafadhel, Bethan Barker, Chris Brightling, Charlotte E Bolton, Michelle E John, Stuart G Parker, Miriam F Moffat, Andrew J Wardlaw, Martin J Connolly, David J Porteous, Blair H Smith, Sandosh Padmanabhan, Lynne Hocking, Kathleen E Stirrups, Panos Deloukas, David P Strachan, Ian P Hall, Martin D Tobin and Louise V Wain

Thorax published online February 25, 2016

Updated information and services can be found at:

<http://thorax.bmj.com/content/early/2016/02/25/thoraxjnl-2015-207876>

These include:

References

This article cites 31 articles, 3 of which you can access for free at:
<http://thorax.bmj.com/content/early/2016/02/25/thoraxjnl-2015-207876#BIBL>

Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See:
<http://creativecommons.org/licenses/by/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (177)
[Health education](#) (1122)
[Smoking](#) (952)
[Tobacco use](#) (954)

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>