

Ground State Electroluminescence

Mauro Cirio,¹ Simone De Liberato,² Neill Lambert,³ and Franco Nori^{3,4}

¹*Interdisciplinary Theoretical Science Research Group (iTHES), RIKEN, Wako-shi, Saitama 351-0198, Japan*

²*School of Physics and Astronomy, University of Southampton, Southampton SO17 1BJ, United Kingdom*

³*CEMS, RIKEN, Saitama, 351-0198, Japan*

⁴*Department of Physics, University of Michigan, Ann Arbor, MI 48109-1040, USA*

(Dated: August 25, 2015)

Electroluminescence, the emission of light in the presence of an electric current, provides information on the allowed electronic transitions of a given system. It is commonly used to investigate the physics of strongly-coupled light-matter systems, whose eigenfrequencies are split by the strong coupling with the photonic field of a cavity. Here we show that, together with the usual electroluminescence, systems in the ultrastrong light-matter coupling regime emit a uniquely quantum radiation when a flow of current is driven through them. While standard electroluminescence relies on the population of excited states followed by spontaneous emission, the process we describe herein extracts bound photons by the dressed ground state and it has peculiar features that unequivocally distinguish it from usual electroluminescence.

PACS numbers: 42.50.Pq, 78.60.Fi, 71.36.+c

The proper dimensionless parameter to study perturbatively the resonant interaction between light and matter in cavity quantum electrodynamics (QED) is the ratio between vacuum Rabi frequency, Ω_R , and the bare frequency of the excitation, ω_C . When such a quantity, usually referred to as normalised coupling, $\eta \equiv \frac{\Omega_R}{\omega_C}$, becomes non-negligible, higher-order perturbative phenomena start to become observable. This is the so-called ultrastrong-coupling regime, that has been experimentally achieved in a number of solid-state systems [1–11]. The non-perturbative nature of the light-matter coupling in this regime leads to a range of very rich (and as yet not fully understood) phenomenology, including quantum phase transitions [12–14], modification of energy transport [15, 16] and light emission properties [17–19], or the appearance of cavity-assisted chemical and thermodynamic effects [20–23]. One of the main consequences of the ultrastrong-coupling regime is to modify the ground state of the system, such that it becomes a multimode squeezed state, containing a finite population of bound excitations that can only be observed if the system parameters are nonadiabatically modulated in time [24–31]. This quantum vacuum radiation, that has strong similarities with the dynamical Casimir effect [32–37], is a hallmark of the quantum nature of the light-matter interaction, and it can provide insights into the quantum nature of the ground state [38].

Here we investigate a unique quantum electrodynamical effect wherein one can observe the photonic excitations bound in the ground state when an electrical current is driven through the system. We call this effect *ground state electroluminescence* (GSE), in order to distinguish it from the standard polaritonic electroluminescence observable when a current passes in a strongly-coupled light-matter system [9, 39–49]. We will see that such a GSE is not only intense enough to be observable




FIG. 1: (a) Energy diagram for the uncoupled system ($\Omega_R = 0$). Photons can leave the cavity at a rate Γ_{cav} and electrons can populate/leave the system at rates $\Gamma_{\text{in/out}}$. (b) Diagram levels for the strongly coupled regime at resonance $\omega_C = \omega_e$. (c) Schematic showing standard electroluminescent emission. (d) Schematic showing the dominant GSE process.

in some near-future solid-state cavity QED experiments, but it has also unique features that clearly distinguish it from the usual polaritonic electroluminescence.

Our aim is to present a novel QED effect, a priori relevant for many of the systems in which the USC regime has been observed. For this reason we will keep the discussion as generic and system-independent as possible, investigating the simplest model presenting GSE effects. A discussion on possible implementations will be discussed later. We will thus consider a toy model exhibiting GSE: a two-level electronic system ultrastrongly coupled

to a single mode of a photonic resonator. When a single electron is present, this is just the Rabi model, one of the simplest models used to investigate the physics of strong light-matter coupling. The important element we add here is that, given that we are interested in electroluminescence, we will consider the system to be electronically open: i.e., we will account for the possibility of electrons to tunnel in and out of the structure. While it is not necessary for our theory, in order to simplify the already rather heavy notation, we will neglect the possibility to have a doubly-occupied state. Depending on the specific implementation, this assumption can be physically justified assuming the system to be in a Coulomb-blockade regime. Thus, for this electronic system we only consider a 3D Hilbert space spanned by the empty state $|s\rangle$, and the two singly-occupied states $|g\rangle$ and $|e\rangle$, with Hamiltonian

$$H = \hbar\omega_C a^\dagger a + \hbar\omega_e |e\rangle\langle e| - \hbar\omega_s |s\rangle\langle s| + \Omega_R(a + a^\dagger)(|e\rangle\langle g| + |g\rangle\langle e|),$$

where a is the annihilation operator of the photonic mode of energy $\hbar\omega_C$, and $\hbar\omega_e$ and $\hbar\omega_s$ the energies of the $g \rightarrow e$ and $s \rightarrow g$ transitions. For definiteness, hereafter we will limit ourselves to the resonant case $\omega_e = \omega_C$. As the Hamiltonian in Eq. (1) conserves the number of electrons, it can be diagonalised separately in the zero and one electron sectors. In the first sector the Hamiltonian is diagonal in the n -photons states $|s, n\rangle$, of energy $n\hbar\omega_C - \hbar\omega_s$; while in the second sector we recover exactly the Hamiltonian of a quantum Rabi model [50]. We will call respectively $|G\rangle$ and $|\pm\rangle$ its ground state and first polaritonic doublet, with respective energies $\hbar\omega_G$ and $\hbar\omega_G + \hbar\omega_\pm$. For the sake of clarity, we will neglect higher-lying states and temperature effects, as both are inessential to GSE. In order to be able to investigate electroluminescence, we couple such a system to two external electronic reservoirs. Fixing the chemical potential of one of them to $-\hbar\omega_s$, so that it acts only as a sink for electrons, the voltage across the system is described by the chemical potential μ of the other reservoir, leading to the effective electronic injection and extraction rates $\Gamma_{\text{in/out}}$. We also couple the cavity itself to a reservoir, describing the extra-cavity photonic modes in which the luminescence is emitted, leading to a photonic lifetime Γ_{cav} . A diagram of the relevant levels and transition rates for vanishing couplings is in Fig. 1(a).

Note that we will consider strongly-coupled systems, whose eigenstates are linear superpositions of different electronic and photonic excitations. Therefore, the bare electronic and photonic lifetimes will give rise to different transition rates between different pairs of dressed states, that can be calculated using the different methods that have been developed to deal with strongly-coupled open systems [27, 51–53]. The levels and transition rates for strong couplings is in Fig. 1(b). Limiting ourselves to

the low-lying states that we will consider here, we thus define $\Gamma_{\text{in}}^{n \rightarrow j}$ and $\Gamma_{\text{out}}^{j \rightarrow n}$ as the electronic injection and extraction rates to and from the states $|s, n\rangle$ and $|j\rangle$, $j \in \{G, \pm\}$, and Γ_{cav}^\pm as the photonic transition rates from $|\pm\rangle$ to $|G\rangle$. Notice that in the zero temperature case we are considering here, electron tunnelling can couple two states only if μ is larger than their energy mismatch, that is

$$\begin{aligned} \Gamma_{\text{in}}^{n \rightarrow G} &\propto \Theta(\mu + n\hbar\omega_C - \hbar\omega_G) \\ \Gamma_{\text{in}}^{n \rightarrow \pm} &\propto \Theta(\mu + n\hbar\omega_C - \hbar\omega_G - \hbar\omega_\pm), \end{aligned} \quad (1)$$

where Θ is the Heaviside function.

In order to introduce our discussion of GSE, we will start by reviewing standard electroluminescence in polaritonic systems for $\eta \ll 1$ [48, 49]. In this regime, the Rabi Hamiltonian reduces to a Jaynes-Cummings model

$$|G\rangle \simeq |g, 0\rangle, \quad |\pm\rangle \simeq \frac{|g, 1\rangle \pm |e, 0\rangle}{\sqrt{2}}, \quad (2)$$

leading to $\hbar\omega_G = 0$ and $\Gamma_{\text{out}}^{G \rightarrow n} = \Gamma_{\text{out}}\delta_{n,0}$, $\Gamma_{\text{in}}^{n \rightarrow G} = \Gamma_{\text{in}}\delta_{n,0}$: as the ground state $|G\rangle$ contains no photons, when we extract the electron we cannot end in a state with a photonic component and vice versa. A current can thus pass through the structure for any $\mu \geq 0$, without any photonic emission, through the path

$$|s, 0\rangle \xrightarrow{\Gamma_{\text{in}}} |G\rangle \xrightarrow{\Gamma_{\text{out}}} |s, 0\rangle. \quad (3)$$

Only for $\mu \geq \hbar\omega_G + \hbar\omega_\pm$, the electrons can excite states with a photonic component, corresponding to $\Gamma_{\text{in}}^{0 \rightarrow \pm} > 0$, then decaying to the ground state before being extracted. This leads to the usual polaritonic electroluminescence at energy $\hbar\omega_\pm$, through the processes

$$|s, 0\rangle \xrightarrow{\Gamma_{\text{in}}^{0 \rightarrow \pm}} |\pm\rangle \xrightarrow[\downarrow \hbar\omega_\pm]{\Gamma_{\text{cav}}^\pm} |G\rangle \xrightarrow{\Gamma_{\text{out}}} |s, 0\rangle, \quad (4)$$

where we highlighted the transition (and its energy) expected to emit light.

The situation radically changes when instead $\eta \simeq 1$. In this case Eq. 2 is not valid anymore, and the ground state has a non-negligible photonic component, $\langle G|a^\dagger a|G\rangle \neq 0$ [24]. As $|G\rangle$ is now a linear superposition of states with one electron and n photons, when the electron is extracted, it has a finite probability to project the system in the state $|s, n\rangle$, with no electrons and n photons ($\Gamma_{\text{out}}^{G \rightarrow n} > 0$), that will eventually escape out of the cavity. This process is what we term GSE: emission of photons when a current passes through the ground state. The fundamental GSE process, emitting at the cavity frequency $\hbar\omega_C$, has the form

$$|s, 0\rangle \xrightarrow{\Gamma_{\text{in}}^{0 \rightarrow G}} |G\rangle \xrightarrow[\downarrow \hbar\omega_C]{\Gamma_{\text{out}}^{G \rightarrow 1}} |s, 1\rangle \xrightarrow{\Gamma_{\text{cav}}} |s, 0\rangle. \quad (5)$$

If the photonic lifetime is longer than the electron injection and extraction times, GSE photons can accumulate in the cavity, leading to the occupation of higher-lying states of the Rabi Hamiltonian and luminescence




FIG. 2: Emission spectrum $S(\omega)$ for GSE alone (blue solid line, $\mu = \hbar\omega_G$) and GSE plus standard electroluminescence (yellow solid line, $\mu = \hbar\omega_+$). The peaks corresponding to the processes emitting at the bare cavity frequency $\hbar\omega_C$ and at the polaritonic frequencies $\hbar\omega_{\pm}$ are explicitly shown. The dashed line represents the black-body radiation spectrum for $k_B T \sim 0.1 \hbar\omega_C$. Parameters: $\eta = 0.1$, $\Gamma = 0.5 \times 10^{-6} \omega_C$, $\Gamma_{\text{cav}} = 7 \times 10^{-4} \omega_C$.

at the respective frequencies. Under the hypothesis of large enough cavity losses, we will limit ourselves to processes of first order in $\frac{\Gamma_{\text{in/out}}}{\Gamma_{\text{cav}}}$

$$|s, 0\rangle \xrightarrow{\Gamma_{\text{in}}^{0 \rightarrow G}} |G\rangle \xrightarrow{\Gamma_{\text{out}}^{G \rightarrow 1}} |s, 1\rangle \xrightarrow{\Gamma_{\text{in}}^{1 \rightarrow \pm}} |\pm\rangle \xrightarrow{\Gamma_{\text{cav}}^{\pm}} |G\rangle \downarrow \hbar\omega_{\pm}$$

$$\xrightarrow{\Gamma_{\text{out}}^{G \rightarrow 0}} |s, 0\rangle, \quad (6)$$

that clearly emit at the polaritonic frequencies $\hbar\omega_{\pm}$. We emphasize that the processes in Eqs. (5, 6) do not depend upon $\Gamma_{\text{in}}^{0 \rightarrow \pm}$, and thus from Eq. (1) they happen for values of the applied bias $\mu < \hbar\omega_G + \hbar\omega_-$, such that the standard electroluminescence in Eq. (4) is forbidden.

The GSE emission rate can be calculated writing a rate equation for the populations P_j of the respective states $|j\rangle$. Limiting ourselves to the first satellite peaks, and to the regime $\mu < \hbar\omega_G + \hbar\omega_-$, in which no standard electroluminescence is possible, we have

$$\begin{aligned} \dot{P}_{s,0} &= -P_{s,0}\Gamma_{\text{in}}^{s,0} + P_G\Gamma_{\text{out}}^{G \rightarrow 0} + P_{\pm}\Gamma_{\text{out}}^{\pm \rightarrow 0} + P_{s,1}\Gamma_{\text{cav}} \\ \dot{P}_{s,1} &= -P_{s,1}(\Gamma_{\text{cav}} + \Gamma_{\text{in}}^{s,1}) + P_G\Gamma_{\text{out}}^{G \rightarrow 1} + P_{\pm}\Gamma_{\text{out}}^{\pm \rightarrow 1} \\ \dot{P}_G &= -P_G\Gamma_{\text{out}}^G + P_{s,0}\Gamma_{\text{in}}^{0 \rightarrow G} + P_{s,1}\Gamma_{\text{in}}^{1 \rightarrow G} + P_{\pm}\Gamma_{\text{cav}}^{\pm} \\ \dot{P}_+ &= -P_+(\Gamma_{\text{cav}}^+ + \Gamma_{\text{out}}^+) + P_{s,1}\Gamma_{\text{in}}^{1 \rightarrow +} \\ \dot{P}_- &= -P_-(\Gamma_{\text{cav}}^- + \Gamma_{\text{out}}^-) + P_{s,1}\Gamma_{\text{in}}^{1 \rightarrow -}, \end{aligned} \quad (7)$$

where repeated \pm stands for sum over its repeated constituents. In the steady state, this system can be solved for the populations and, consequently, the emission rates for the fundamental GSE in Eq. (5), f_C , and for the first satellites in Eq. (6), f_{\pm} , can be calculated as

$$f_C = P_{s,1}\Gamma_{\text{cav}}, \quad f_+ = P_+\Gamma_{\text{cav}}^+, \quad f_- = P_-\Gamma_{\text{cav}}^-. \quad (8)$$

Developing perturbatively the coupled eigenstates of the Rabi Hamiltonian $|G\rangle$ and $|\pm\rangle$ over the uncoupled states $|g, n\rangle$ and $|e, n\rangle$, and considering $\Gamma_{\text{in}} = \Gamma_{\text{out}} \equiv \Gamma$, we can obtain estimates for the radiation intensity of the main GSE peak in Eq. (5) and for the satellites from Eq. (6). To the lowest nontrivial order in η and to the first order in $\frac{\Gamma}{\Gamma_{\text{cav}}}$ they read

$$f_C = \frac{\eta^2 \Gamma}{8} \left(1 - \frac{\Gamma}{\Gamma_{\text{cav}}}\right), \quad f_{\pm} = \frac{\eta^2 \Gamma}{16} \frac{\Gamma}{\Gamma_{\text{cav}}}, \quad (9)$$

where, as expected, the emission at $\hbar\omega_{\pm}$, that relies upon a photon buildup in the cavity as from Eq. (6), is weighted down by a factor $\frac{\Gamma}{\Gamma_{\text{cav}}} \ll 1$.

For comparison we also calculated, to the same order, the standard electroluminescence intensity for the case $\mu \geq \hbar\omega_G + \hbar\omega_+$, obtaining

$$f'_C = \frac{\Gamma}{6} \left(\frac{2\Gamma}{\Gamma_{\text{cav}}} + \eta^2 \right), \quad f'_{\pm} = \frac{\Gamma}{6} \left(1 \pm \frac{\eta}{2} \right) \left(1 - \frac{2\Gamma}{\Gamma_{\text{cav}}} \right) \quad (10)$$

To the dominant order in both variables we obtain, as expected, a term not dependent on η emitting at $\hbar\omega_{\pm}$, that is the standard electroluminescence described in Eq. (4). Note that also in this case we recover higher-order processes allowed by the buildup of photonic population in the cavity (and thus of higher order in $\frac{\Gamma}{\Gamma_{\text{cav}}}$); in this case the first one emitting at $\hbar\omega_C$

$$|s, 0\rangle \xrightarrow{\Gamma_{\text{in}}^{0 \rightarrow \pm}} |\pm\rangle \xrightarrow{\Gamma_{\text{out}}^{\pm \rightarrow 1}} |s, 1\rangle \xrightarrow{\Gamma_{\text{cav}}} |s, 0\rangle \downarrow \hbar\omega_C. \quad (11)$$

In order to test such an intuitive understanding of GSE, we studied the system numerically solving [54, 55] the master equation

$$\dot{\rho} = -i[H, \rho] + \mathcal{L}_{\text{in}}(\rho) + \mathcal{L}_{\text{out}}(\rho) + \mathcal{L}_{\text{cav}}(\rho), \quad (12)$$

where ρ is the density operator of the system with a converged cutoff over the maximum number of photons and the Lindblad operators $\mathcal{L}_{\text{in/out/cav}}$ are projected over the dressed basis of the system [27, 51]. Exploiting the input-output formalism compatible with the dressed state analysis developed in Ref. [53], we can write the extra-cavity emission spectrum as

$$S(\omega) = \frac{\Gamma_{\text{cav}}}{2\pi} \int_{-\infty}^{\infty} d\tau e^{-i\omega\tau} \langle X^+(\tau) X^-(0) \rangle, \quad (13)$$

where $X = (a + a^\dagger)$, $X^- = \sum_{j>i} \langle i| X |j\rangle |i\rangle \langle j|$, and X^+ is its hermitian conjugate. In Fig. 2 we plot the emitted spectrum from Eq. (13) for $\mu = \hbar\omega_G$ (blue solid line) in which we can clearly see the central GSE peak at $\hbar\omega_C$ and the satellites around $\hbar\omega_{\pm} \simeq \hbar\omega_C \pm \Omega$. Figure 2 also plots (yellow solid line) the spectrum for $\mu = \hbar\omega_G + \hbar\omega_+$. In this case, while the central peak remains substantially unchanged, standard electroluminescence becomes dominant over the satellite frequencies. There we plot, for




FIG. 3: (a) Integrated emission spectrum $S(\omega)$ (as in Fig. 2) for the central peak and one of the satellites channels for $\mu = \hbar\omega_G$, obtained by integrating the numerical spectrum (solid lines) and from the analytical estimates in Eq. (9) (hollow dots). In the inset we illustrate the GSE processes from Eqs. (5,6). (b) Integrated emissions for $\mu = \hbar\omega_G + \hbar\omega_+$, obtained by integrating the numerical spectrum (solid lines), and from the analytical estimates in Eq. (10) (hollow dots). Inset: electroluminescence from Eqs. (4,11).

comparison, the black body emission from the cavity for $k_B T = 0.1\hbar\omega_C$. In Fig. 3 we plot (solid lines) the integrated emission of the central peak and of the two satellites respectively for $\mu = \hbar\omega_G$ (a) and $\mu = \hbar\omega_G + \hbar\omega_+$ (b), as a function of the perturbative parameter η . The hollow dots in the same Figures are obtained by the analytical estimates from Eq. (9) and Eq. (10), respectively. Schematics of the involved processes are shown as insets. We clearly obtain a very good quantitative agreement between our theory and the numerical simulations, proving the correctness of our theoretical analysis.

Various developing technological platforms seem possible candidates to experimentally observe GSE and some of them, like hybrid QED [40, 41, 56–70] and circuit QED [1, 2, 39, 71–77], in which electronic transitions are coupled to microwave superconducting resonators, seem close to achieve all the needed requirements. In particular, a double quantum dot coupled to a superconducting

resonator seems to us the system better described by the simple model studied here, where $|s\rangle$ is the empty state, $|g\rangle$ and $|e\rangle$ are the hybridised states of the two dots, and the doubly-occupied state is Coulomb blocked. Remarkably, electroluminescence has already been observed in those systems [40, 41]. Given that the ratio $\frac{\Gamma}{\Gamma_{\text{cav}}}$ can be engineered in those systems, from Eq. (9) we obtain a rough estimate of the photon population, due to GSE, of $P \simeq 0.1\eta^2$. Considering a conservative lower bound to the detectable photon population in microwave resonators of $P_{\text{min}} = 10^{-3}$ [41], $\eta > 0.1$ should thus suffice to observe GSE. This normalised coupling has been achieved in circuit QED [1, 2]. In hybrid systems, experiments are moving toward higher couplings and the appropriate regime may become accessible in the near future.

In conclusion, we have highlighted a novel cavity QED phenomenon, leading to the purely quantum emission of photons out of the ground state when an electric current passes through it. Such a novel form of quantum vacuum emission could be observed in various near-future cavity QED setups, leading to a further step forward in the understanding and engineering of the quantum vacuum in the ultrastrong light-matter coupling regime.

Acknowledgments.— This work is partially supported by the RIKEN iTHES Project, the MURI Center for Dynamic Magneto-Optics via the AFOSR award number FA9550-14-1-0040, the IMPACT program of JST, and a Grant-in-Aid for Scientific Research (A). MC is supported by the Canon Foundation in Europe and the RIKEN iTHES program. NL is partially supported by the FY2015 Incentive Research Project. SDL is Royal Society Research Fellow. SDL acknowledges support from the Engineering and Physical Sciences Research Council (EPSRC), research grant EP/L020335/1.

-
- [1] T. Niemczyk, F. Deppe, H. Huebl, E. P. Menzel, F. Hocke, M. J. Schwarz, J. J. Garcia-Ripoll, D. Zueco, T. Hummer, E. Solano, A. Marx, and R. Gross, *Circuit quantum electrodynamics in the ultrastrong-coupling regime*, Nat. Phys. **6**, 772 (2010).
 - [2] A. Baust, E. Hoffmann, M. Haeberlein, M. J. Schwarz, P. Eder, J. Goetz, F. Wulschner, E. Xie, L. Zhong, F. Quijandria, D. Zueco, J.-J. Garcia Ripoll, L. Garcia-Alvarez, G. Romero, E. Solano, K. G. Fedorov, E. P. Menzel, F. Deppe, A. Marx, R. Gross, *Ultrastrong coupling in two-resonator circuit QED*, arXiv:1412.7372 (2015).
 - [3] V. M. Muravev, I. V. Andreev, I. V. Kukushkin, S. Schmult, and W. Dietsche, *Observation of hybrid plasmon-photon modes in microwave transmission of coplanar microresonators*, Phys. Rev. B **83**, 075309 (2011).
 - [4] A. A. Anappara, S. De Liberato, A. Tredicucci, C. Ciuti, G. Biasiol, L. Sorba and F. Beltram, *Signatures of the ultrastrong light-matter coupling regime*, Phys. Rev. B **79**,

- 201303 (2009).
- [5] M. Geiser, F. Castellano, G. Scalari, M. Beck, L. Nevou, and J. Faist, *Ultrastrong Coupling Regime and Plasmon Polaritons in Parabolic Semiconductor Quantum Wells*, Phys. Rev. Lett. **108**, 106402 (2012).
 - [6] T. Schwartz, J. A. Hutchison, C. Genet and T. W. Ebbesen, *Reversible Switching of Ultrastrong Light-Molecule Coupling*, Phys. Rev. Lett. **106**, 196405 (2011).
 - [7] M. Porer, J.-M. Menard, A. Leitenstorfer, R. Huber, R. Degl'Innocenti, S. Zanotto, G. Biasiol, L. Sorba, and A. Tredicucci, *Nonadiabatic switching of a photonic band structure: Ultrastrong light-matter coupling and slow-down of light*, Phys. Rev. B **85**, 081302 (2012).
 - [8] B. Askenazi, A. Vasanelli, A. Delteil, Y. Todorov, L. C. Andreani, G. Beaudoin, I. Sagnes, and C. Sirtori, *Ultra-strong lightmatter coupling for designer Reststrahlen band*, New. J. Phys. **16**, 043029 (2014).
 - [9] C. R. Gubbin, S. A. Maier, and S. Kéna-Cohen, *Low-voltage polariton electroluminescence from an ultrastrongly coupled organic light-emitting diode*, App. Phys. Lett. **104**, 233302 (2014).
 - [10] S. Gambino, M. Mazzeo, A. Genco, O. Di Stefano, S. Savasta, S. Patané, D. Ballarini, F. Mangione, G. Lerario, D. Sanvitto, and G. Gigli, *Exploring LightMatter Interaction Phenomena under Ultrastrong Coupling Regime*, ACS Phot. **1**, 1042 (2014).
 - [11] C. Maissen, G. Scalari, F. Valmorra, M. Beck, J. Faist, S. Cibella, R. Leoni, C. Reichl, C. Charpentier, and W. Wegscheider, *Ultrastrong coupling in the near field of complementary split-ring resonators*, Phys. Rev. B **90**, 205309 (2014).
 - [12] N. Lambert, C. Emary and T. Brandes, *Entanglement and the Phase Transition in Single-Mode Superradiance*, Phys. Rev. Lett. **92**, 073602 (2004).
 - [13] P. Nataf and C. Ciuti, *No-go theorem for superradiant quantum phase transitions in cavity QED and counter-example in circuit QED*, Nat. Comm. **1**, 72 (2010).
 - [14] A. Baksic, P. Nataf, and C. Ciuti, *Superradiant phase transitions with three-level systems*, Phys. Rev. A **87**, 023813 (2013).
 - [15] J. Feist, F. J. Garcia-Vidal, *Extraordinary Exciton Conductance Induced by Strong Coupling*, Phys. Rev. Lett. **114**, 196402 (2015).
 - [16] E. Orgiu, J. George, J. A. Hutchison, E. Devaux, J.-F. Dayen, B. Doudin, F. Stellacci, C. Genet, P. Samori, and T. Ebbesen, *Conductivity in organic semiconductors hybridized with the vacuum field*, arXiv:1409.1900 (2014).
 - [17] S. De Liberato, *Light-Matter Decoupling in the Deep Strong Coupling Regime: The Breakdown of the Purcell Effect*, Phys. Rev. Lett. **112**, 016401 (2014).
 - [18] J. J. García-Ripoll, B. Peropadre, and S. De Liberato, *Light-matter decoupling and A^2 term detection in superconducting circuits*, arXiv:1410.7785 (2014).
 - [19] M. Bamba and T. Ogawa, *Laser under ultrastrong electromagnetic interaction with matter*, arXiv:1410.3912 (2014).
 - [20] J. A. Hutchison, T. Schwartz, C. Genet, E. Devaux, and T. W. Ebbesen, *Modifying Chemical Landscapes by Coupling to Vacuum Fields*, Angew. Chemie **124**, 1624 (2012).
 - [21] J. A. Hutchison, A. Liscio, T. Schwartz, A. Canaguier-Durand, C. Genet, V. Palermo, P. Samori, and T. W. Ebbesen, *Tuning the work-function via strong coupling*, Adv. Mater. **25**, 2481 (2013).
 - [22] J. Galego, F. J. Garcia-Vidal, and J. Feist, *Cavity-induced modifications of molecular structure in the strong coupling regime*, arXiv:1506.03331.
 - [23] J. A. Cwik, P. Kirton, S. De Liberato, and J. Keeling, *Self-consistent molecular adaptation induced by strong coupling*, arXiv:1506.08974.
 - [24] C. Ciuti, G. Bastard and I. Carusotto, *Quantum vacuum properties of the intersubband cavity polariton field*, Phys. Rev. B **72**, 115303 (2005).
 - [25] V. V. Dodonov and A. V. Dodonov, *QED effects in a cavity with a time-dependent thin semiconductor slab excited by laser pulses*, J. Phys. B **39**, 1 (2006).
 - [26] S. De Liberato, C. Ciuti and I. Carusotto, *Quantum Vacuum Radiation Spectra from a Semiconductor Microcavity with a Time-Modulated Vacuum Rabi Frequency*, Phys. Rev. Lett. **98**, 103602 (2007).
 - [27] S. De Liberato, D. Gerace, I. Carusotto, and C. Ciuti, *Extracavity quantum vacuum radiation from a single qubit*, Phys. Rev. A **80**, 053810 (2009).
 - [28] A. Agnesi, C. Braggio, G. Bressi, G. Carugno, F. Della Valle, G. Galeazzi, G. Messineo, F. Pirzio, G. Reali, G. Ruoso, D. Scarpa, and D. Zanello, *MIR: An experiment for the measurement of the dynamical Casimir effect*, J. Phys. Conf. **161**, 012028 (2009).
 - [29] D. Faccio and I. Carusotto, *Dynamical Casimir Effect in optically modulated cavities*, Europ. Phys. Lett. **96**, 24006 (2011).
 - [30] I. Carusotto, S. De Liberato, D. Gerace, and C. Ciuti, *Back-reaction effects of quantum vacuum in cavity quantum electrodynamics*, Phys. Rev. A **85**, 023805 (2012).
 - [31] A. Auer and G. Burkard, *Entangled photons from the polariton vacuum in a switchable optical cavity*, Phys. Rev. B **85**, 235140 (2012).
 - [32] P. D. Nation, J. R. Johansson, M. P. Blencowe, and F. Nori, *Stimulating uncertainty: Amplifying the quantum vacuum with superconducting circuits*, Rev. Mod. Phys. **84**, 1 (2012).
 - [33] A. Lambrecht, M.-T. Jaekel, and S. Reynaud, *Motion Induced Radiation from a Vibrating Cavity*, Phys. Rev. Lett. **77**, 615 (1996).
 - [34] J. R. Johansson, G. Johansson, C.M. Wilson, F. Nori, *Dynamical Casimir Effect in a Superconducting Coplanar Waveguide*, Phys. Rev. Lett. **103**, 147003 (2009).
 - [35] J. R. Johansson, G. Johansson, C.M. Wilson, F. Nori, *Dynamical Casimir effect in superconducting microwave circuits*, Phys. Rev. A **82**, 052509 (2010).
 - [36] J.R. Johansson, G. Johansson, C. M. Wilson, P. Delsing, F. Nori, *Nonclassical microwave radiation from the dynamical Casimir effect*, Phys. Rev. A **87**, 043804 (2013).
 - [37] C. M. Wilson, G. Johansson, A. Pourkabirian, M. Simoen, J. R. Johansson, T. Duty, F. Nori, and P. Delsing, *Observation of the dynamical Casimir effect in a superconducting circuit*, Nature **479**, 376 (2011).
 - [38] J. Lolli, A. Baksic, D. Nagy, V. E. Manucharyan, C. Ciuti, *Ancillary Qubit Spectroscopy of Vacua in Cavity and Circuit Quantum Electrodynamics*, Phys. Rev. Lett. **114**, 183601 (2015).
 - [39] O. Astafiev, K. Inomata, A. O. Niskanen, T. Yamamoto, Yu. A. Pashkin, Y. Nakamura and J. S. Tsai, *Single artificial-atom lasing*, Nature **449**, 588 (2007).
 - [40] Y.-Y. Liu, K. D. Petersson, J. Stehlik, J. M. Taylor and J. R. Petta, *Photon Emission from a Cavity-Coupled Double Quantum Dot*, Phys. Rev. Lett. **113**, 036801 (2014).
 - [41] A. Stockklauser, V. F. Maisi, J. Basset, K. Cujia, C. Re-

- ichl, W. Wegscheider, T. Ihn, A. Wallraff and K. Ensslin, *Microwave Emission from Hybridized States in a Semiconductor Charge Qubit*, Phys. Rev. Lett. **115**, 046802 (2015).
- [42] A. A. Khalifa, A. P. Love, D. N. Krizhanovskii, M. S. Skolnick, and J. S. Roberts, *Electroluminescence emission from polariton states in GaAs-based semiconductor microcavities*, Appl. Phys. Lett. **92**, 061107 (2008).
- [43] S. I. Tsintzos, N. T. Pelekanos, G. Konstantinidis, Z. Hatzopoulos, and P. G. Savvidis, *A GaAs polariton light-emitting diode operating near room temperature*, Nature **453**, 372 (2008).
- [44] L. Sapienza, A. Vasanelli, R. Colombelli, C. Ciuti, Y. Chassagneux, C. Manquest, U. Gennser, and C. Sirtori, *Electrically Injected Cavity Polaritons*, Phys. Rev. Lett. **100**, 136806 (2008).
- [45] G. H. Lodden and R. J. Holmes, *Polarization splitting in polariton electroluminescence from an organic semiconductor microcavity with metallic reflectors*, Appl. Phys. Lett. **98**, 233301 (2011).
- [46] P. Jouy, A. Vasanelli, Y. Todorov, L. Sapienza, R. Colombelli, U. Gennser, and C. Sirtori, *Intersubband electroluminescent devices operating in the strong-coupling regime*, Phys. Rev. B **82**, 045322 (2010).
- [47] M. Geiser, G. Scalari, F. Castellano, M. Beck, and J. Faist, *Room temperature terahertz polariton emitter*, Appl. Phys. Lett. **101**, 141118 (2012).
- [48] S. De Liberato and C. Ciuti, *Quantum model of microcavity intersubband electroluminescent devices*, Phys. Rev. B **77**, 155321 (2008).
- [49] S. De Liberato and C. Ciuti, *Quantum theory of electron tunneling into intersubband cavity polariton states*, Phys. Rev. B **79**, 075317 (2009).
- [50] D. Braak, *Integrability of the Rabi Model*, Phys. Rev. Lett. **107**, 100401 (2011).
- [51] F. Beaudoin, J. M. Gambetta, and A. Blais, *Dissipation and ultrastrong coupling in circuit QED*, Phys. Rev. A **84**, 043832 (2011).
- [52] S. De Liberato, *Comment on System-environment coupling derived by Maxwell's boundary conditions from the weak to the ultrastrong light-matter-coupling regime*, Phys. Rev. A **89**, 017801 (2014).
- [53] A. Ridolfo, M. Leib, S. Savasta, and M. J. Hartmann, *Photon Blockade in the Ultrastrong Coupling Regime*, Phys. Rev. Lett. **109**, 193602 (2012).
- [54] J. R. Johansson, P. D. Nation, and F. Nori, *QuTiP: An open-source Python framework for the dynamics of open quantum systems*, Comput. Phys. Commun. **183**, 1760 (2012).
- [55] J. R. Johansson, P. D. Nation, and F. Nori, *QuTiP 2: A Python framework for the dynamics of open quantum systems*, Comput. Phys. Commun. **184**, 1234 (2013).
- [56] Z.-L. Xiang, S. Ashhab, J. Q. You, and F. Nori, *Hybrid quantum circuits: Superconducting circuits interacting with other quantum systems*, Rev. Mod. Phys. **85**, 623 (2013).
- [57] M. R. Delbecq, V. Schmitt, F. D. Parmentier, N. Roch, J. J. Viennot, G. Fève, B. Huard, C. Mora, A. Cottet and T. Kontos, *Coupling a Quantum Dot, Fermionic Leads, and a Microwave Cavity on a Chip*, Phys. Rev. Lett. **107**, 256804 (2011).
- [58] M. R. Delbecq, L. E. Bruhat, J. J. Viennot, S. Datta, A. Cottet and T. Kontos, *Photon-mediated interaction between distant quantum dot circuits*, Nature Commun. **4**, 1400 (2013).
- [59] T. Frey, P. J. Leek, M. Beck, A. Blais, T. Ihn, K. Ensslin and A. Wallraff, *Dipole Coupling of a Double Quantum Dot to a Microwave Resonator*, Phys. Rev. Lett. **108**, 046807 (2012).
- [60] T. Frey, P. J. Leek, M. Beck, J. Faist, A. Wallraff, K. Ensslin, T. Ihn and M. Büttiker, *Quantum dot admittance probed at microwave frequencies with an on-chip resonator*, Phys. Rev. B **86**, 115303 (2012).
- [61] K. D. Petersson, L. W. McFaul, M. D. Schroer, M. Jung, J. M. Taylor, A. A. Houck and J. R. Petta, *Circuit quantum electrodynamics with a spin qubit*, Nature **490**, 380 (2012).
- [62] H. Toida, T. Nakajima and S. Komiyama, *Vacuum Rabi Splitting in a Semiconductor Circuit QED System*, Phys. Rev. Lett. **110**, 066802 (2013).
- [63] A. Wallraff, A. Stockklauser, T. Ihn, J. R. Petta and A. Blais, *Comment on Vacuum Rabi Splitting in a Semiconductor Circuit QED System*, Phys. Rev. Lett. **111**, 249701 (2013).
- [64] J. J. Viennot, M. R. Delbecq, M. C. Dartiailh, A. Cottet and T. Kontos, *Out-of-equilibrium charge dynamics in a hybrid circuit quantum electrodynamics architecture*, Phys. Rev. B **89**, 165404 (2014).
- [65] Y. Mu and C. M. Savage, *One-atom lasers*, Phys. Rev. A **46**, 5944 (1992).
- [66] J. McKeever, A. Boca, A. D. Boozer, J. R. Buck and H. J. Kimble, *A One-Atom Laser in a Regime of Strong Coupling*, Nature **425**, 268 (2003).
- [67] Y.-Y. Liu, J. Stehlik, C. Eichler, M. J. Gullans, J. M. Taylor J. R. Petta, *Semiconductor double quantum dot micromaser*, Science **347**, 285 (2015).
- [68] G.-W. Deng, D. Wei, S.-X. Li, J. R. Johansson, W.-C. Kong, H.-O. Li, G. Cao, M. Xiao, G.-C. Guo, F. Nori, H.-W. Jiang and G.-P. Guo, *Coupling two distant double quantum dots to a microwave resonator*, arXiv:1409.4980 (2014).
- [69] N. Lambert, C. Flindt, F. Nori, *Photon-mediated electron transport in hybrid circuit-QED*, EPL **103**, 17005 (2013).
- [70] N. Lambert, F. Nori, C. Flindt, *Bistable Photon Emission from a Solid-State Single-Atom Laser*, arXiv:1505.07991 (2015).
- [71] J.Q. You, F. Nori, *Superconducting circuits and quantum information*, Physics Today **58** (11), 42-47 (2005).
- [72] J.Q. You, F. Nori, *Atomic physics and quantum optics using superconducting circuits*, Nature **474**, 589 (2011).
- [73] S. Ashhab, J. R. Johansson, A. M. Zagoskin and F. Nori, *Single-artificial-atom lasing using a voltage-biased superconducting charge qubit*, New J. Phys. **11**, 023030 (2009).
- [74] M. Devoret, S. Girvin, and R. Schoelkopf, *Circuit-QED: How strong can the coupling between a Josephson junction atom and a transmission line resonator be?*, Ann. Phys. (Leipzig) **16**, 767 (2007).
- [75] D. A. Rodrigues, J. Imbers and A. D. Armour, *Quantum Dynamics of a Resonator Driven by a Superconducting Single-Electron Transistor: A Solid-State Analogue of the Micromaser*, Phys. Rev. Lett. **98**, 067204 (2007).
- [76] J. Hauss, A. Fedorov, C. Hutter, A. Shnirman and G. Schön, *Single-Qubit Lasing and Cooling at the Rabi Frequency*, Phys. Rev. Lett. **100**, 037003 (2008).
- [77] J. Q. You, Y. X. Liu, C. P. Sun and F. Nori, *Persistent single-photon production by tunable on-chip micromaser with a superconducting quantum circuit*, Phys. Rev. B **75**, 104516 (2007).