

**Running title: C9orf72 neurons recapitulate features of ALS/FTD**

**Title: C9orf72 hexanucleotide expansions are associated with altered ER calcium homeostasis and stress granule formation in iPSC-derived neurons from patients with amyotrophic lateral sclerosis and frontotemporal dementia**

Ruxandra Dafinca<sup>1</sup>, Jakub Scaber<sup>1</sup>, Nida'a Ababneh<sup>1</sup>, Tatjana Lalic<sup>1</sup>, Gregory Weir<sup>1</sup>, Helen Christian<sup>2</sup>, Jane Vowles<sup>3</sup>, Andrew G. L. Douglas<sup>2</sup>, Alexandra Fletcher-Jones<sup>1</sup>, Cathy Browne<sup>3</sup>, Mahito Nakanishi<sup>4</sup>, Martin R. Turner<sup>1</sup>, Richard Wade-Martins<sup>2</sup>, Sally A Cowley<sup>3\*</sup>, Kevin Talbot<sup>1\*</sup>

<sup>1</sup>Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford UK

<sup>2</sup>Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, UK

<sup>3</sup>James Martin Stem Cell Facility, Sir William Dunn School of Pathology, University of Oxford, Oxford, UK

<sup>4</sup>Research Center for Stem Cell Engineering, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Ibaraki, Japan

\*Corresponding authors:

Prof. Kevin Talbot

Nuffield Department of Clinical Neurosciences, John Radcliffe Hospital, West Wing Level 6, Headley Way Oxford, OX3 9DU, UK

E-mail: [kevin.talbot@ndcn.ox.ac.uk](mailto:kevin.talbot@ndcn.ox.ac.uk)

Telephone : 01865 223380

Dr Sally A. Cowley

James Martin Stem Cell Facility, Sir William Dunn School of Pathology, University of Oxford, South Parks Road, Oxford, OX1 3RE, UK

E-mail: [sally.cowley@path.ox.ac.uk](mailto:sally.cowley@path.ox.ac.uk)

Telephone: 01865 275600

Author contributions:

Ruxandra Dafinca: Conception and design, collection and assembly of data, data analysis and interpretation, manuscript writing, final approval of manuscript

Jakub Scaber: MN differentiation, RNA fluorescence *in situ* hybridisation

Nidaa Ababneh: Stem cell reprogramming, iPS expansion and characterization

Tatjana Lalic: Electrophysiology experiments

Gregory Weir: Electrophysiology experiments

Helen Christian: Electron Microscopy data collection and interpretation

Jane Vowles: Stem cell reprogramming, iPS expansion

Andrew G. L. Douglas: Southern blotting

Alexandra Fletcher-Jones: Partial collection of survival data

Cathy Browne: Stem cell reprogramming

Mahito Nakanishi: Provision of reprogramming factors

Martin R. Turner: Provision of patients

Richard Wade-Martins: supervision of ER Ca<sup>2+</sup> study

Sally A Cowley: Collection and assembly of data, final approval of manuscript

Kevin Talbot: Conception and design, generation of funding, provision of study patients, manuscript writing, editing and final approval of manuscript

Key words:

Amyotrophic lateral sclerosis, frontotemporal dementia, induced pluripotent stem cells, C9orf72, calcium dysregulation, motor neurons

## Abstract

An expanded hexanucleotide repeat in a noncoding region of the *C9orf72* gene is a major cause of amyotrophic lateral sclerosis (ALS), accounting for up to 40% of familial cases and 7% of sporadic ALS in European populations. We have generated induced pluripotent stem cells (iPSCs) from fibroblasts of patients carrying *C9orf72* hexanucleotide expansions, differentiated these to functional motor and cortical neurons and performed an extensive phenotypic characterization. In *C9orf72* iPSC-derived motor neurons, decreased cell survival is correlated with dysfunction in  $\text{Ca}^{2+}$  homeostasis, reduced levels of the anti-apoptotic protein Bcl-2, increased endoplasmic reticulum (ER) stress and reduced mitochondrial membrane potential. Furthermore, *C9orf72* motor neurons, and also cortical neurons, show evidence of abnormal protein aggregation and stress granule formation. This study is an extensive characterization of iPSC-derived motor neurons as cellular models of ALS carrying *C9orf72* hexanucleotide repeats, which describes a novel pathogenic link between *C9orf72* mutations, dysregulation of calcium signalling and altered proteostasis and provides a potential pharmacological target for the treatment of ALS and the related neurodegenerative disease frontotemporal dementia (FTD).

## Introduction

A hexanucleotide (GGGGCC) expansion in the first intron of the *C9orf72* gene accounts for approximately 40% of cases of familial amyotrophic lateral sclerosis (fALS), up to 7% of sporadic ALS, and approximately 20% of familial frontotemporal lobar degeneration (FTLD), establishing a firm genetic link between ALS and FTD [1-3]. The expansion is located in an intronic or promoter region upstream of the *C9orf72* coding sequence, and the number of (GGGGCC)<sub>n</sub> hexanucleotide repeats ranges between ~100 and ~4000 repeats in patients [1, 2, 4]. *C9orf72*-associated cases exhibit distinctive neuropathological changes characterized by abundant cytoplasmic and intranuclear inclusions immunopositive for ubiquitin and sequestosome-1/p62, suggesting disrupted protein degradation [5-7].

While the function of the *C9orf72* gene and the pathogenic mechanisms of the hexanucleotide expansion are currently unknown, several hypotheses have been proposed. A toxic gain of function mechanism, mediated by the accumulation of (GGGGCC)<sub>n</sub>-rich RNA transcripts within nuclear foci observed in the frontal cortex and spinal cord of patients, might lead to sequestration of RNA binding proteins and disruption of the translation of diverse mRNAs [1], or increased nucleolar stress [8]. Alternatively, reduced transcription of *C9orf72* could lead to neurodegeneration by interfering with the constitutive function of the protein [1, 9]. Lastly, repeat-associated non-ATG (RAN) translation, occurring in the absence of an initiating ATG codon, over the GGGGCC repeat expansion has been shown to produce homopolymeric proteins prone to aggregation [10, 11].

The generation of human motor neurons in culture from induced pluripotent stem cells, reprogrammed from skin fibroblasts of patients with neurodegenerative diseases such as ALS, offers a potentially powerful tool with which to study the key pathological processes in motor neuron degeneration and for screening drugs of potential therapeutic benefit. Previous studies have demonstrated that RNA foci and RAN-translation products can be detected in iPSC-derived motor neurons from ALS/FTD patients with *C9orf72* hexanucleotide expansions [12-14]. Evidence of defects in autophagy, sequestration of RNA-binding proteins by the expanded repeat, changes in gene transcription and altered neuronal excitability suggest that these models can display disease-relevant phenotypes which can be corrected by targeting the expanded RNA with antisense oligonucleotides [13, 14].

Calcium ( $\text{Ca}^{2+}$ ) dysregulation is believed to play an important role in the pathophysiology of ALS [15] and  $\text{Ca}^{2+}$  overload in the cytoplasm of neurons is a potential mechanism that may link excitotoxicity to neuronal death [16]. The endoplasmic reticulum (ER) is the largest intracellular  $\text{Ca}^{2+}$  store and high ER  $\text{Ca}^{2+}$  concentration plays an essential role in the activity of protein synthesis and processing. Disturbances in ER  $\text{Ca}^{2+}$  homeostasis have been linked to chronic activation of the ER stress response and downstream compensatory mechanisms to protect the cell, such as the unfolded protein response (UPR) and autophagy [17]. ER-stress induced cell death can proceed in both  $\text{Ca}^{2+}$ -dependent and  $\text{Ca}^{2+}$ -independent ways. The anti-apoptotic proteins Bcl-2 and Bcl-XL are essential modulators of  $\text{Ca}^{2+}$  signalling in the cell, mediating the decrease and release of  $\text{Ca}^{2+}$  in the ER thereby protecting mitochondria from  $\text{Ca}^{2+}$  overload [18, 19].

In order to explore whether altered  $\text{Ca}^{2+}$  homeostasis and ER stress are cellular phenotypes associated with *C9orf72*-related ALS/FTD, we generated induced Pluripotent Stem Cell (iPSC) lines from skin fibroblasts from patients carrying *C9orf72* hexanucleotide expansions and differentiated them to motor and cortical neurons. Here we show that in *C9orf72* iPSC-derived motor neurons loss of  $\text{Ca}^{2+}$  homeostasis is associated with a decrease in mitochondrial potential and increase in ER stress. Increased susceptibility to cell death was accompanied by reduced levels of the anti-apoptotic Bcl-2 and Bcl-X<sub>L</sub> proteins and upregulation of the pro-apoptotic protein BAK. Furthermore, both *C9orf72* motor and cortical neurons showed elevated levels of SQST-1/p62 and the formation of poly-A-binding protein (PABP) positive stress granules. Our results indicate disease-specific alterations in motor and cortical neurons of ALS/FTD patients carrying *C9orf72* repeat expansions and reveal a novel pathogenic link between *C9orf72*,  $\text{Ca}^{2+}$  dysregulation, ER stress and loss of proteostasis.

## Results

### Reprogramming of patient fibroblasts to generate induced pluripotent stem cells

Two lines from each of three patients (C9-T2, C9-7245, C9-02) were generated, along with lines from control donors. Fig 1 shows the representative characterisation data for one of the patient iPSC lines generated for this study, iPSC C9-T2-7 (data for the other previously unpublished iPSC lines, including control lines, is given in Fig S1). SNP analysis confirmed the identity of the iPSC in relation to their parental fibroblasts and that they had not acquired large-scale copy number variations, although a small number of small-scale indels (well below the level that would be detected by G-banding) were noted, particularly in the C9-7245 lines (Fig 1 A and S1 A). SNP datasets for these lines have been deposited in GEO under the accession number GSE64582.

iPSC lines were positive for pluripotency markers (TRA-1-60 and Nanog Fig 1 B; also Fig S1 B) and showed human pluripotent stem cell-like morphology (cell-cell-contact-dependent clusters, with high nucleus to cytoplasm ratio, Fig 1 C and S1 C). PluriTest is a quantitative assessment of pluripotency based on transcriptome data from iPSCs in comparison to a large reference set of genome-wide profiles from multiple cell and tissue types. It measures a ‘pluripotency score’ (expression of pluripotency genes, with high scores correlating with high pluripotency, y axis) and a ‘novelty’ score (expression of non-pluripotent stem cell genes, therefore this score inversely correlates with pluripotency, x axis). All the patient iPSC lines were deemed to be fully reprogrammed and pluripotent by this analysis (Fig 1 D). The Illumina HT12v4 transcriptome array results have been deposited in the Gene Expression Omnibus (GEO) under accession number GSE64583. Finally, all banked iPSC line

stocks had undetectable levels of residual Sendai virus RNA by qRT-PCR (Fig 1 E and Fig S1 D).

### ***C9orf72* iPSCs differentiate to functional motor neurons**

*In vitro* cell differentiation protocols recapitulate signalling events that occur during embryogenesis. The differentiation of iPSCs to spinal motor neurons (MNs) is a three-stage process that involves neuralization, caudalization and ventralization. The iPSC lines were differentiated to MNs according to three previously published protocols, with some modifications [20-22] (Fig S2 A-C).

The fully differentiated motor neurons were identified by immunostaining with the MN marker HB9 and neuronal marker  $\beta$ -III tubulin (Tuj1), which indicated successful differentiation of iPSCs to motor neurons (Fig S2 D). The iPSC-derived motor neurons showed expression of synaptophysin, a synaptic vesicle protein essential for the formation of synapses, as well as the expression of choline acetyltransferase (ChAT), which demonstrates an advanced degree of cholinergic neuron maturation (Fig S2 D,E).

Two iPSC lines were reprogrammed from each patient's fibroblasts and differentiated to MNs for characterization. To assess differentiation efficiency, motor neurons positive for Islet1/Tuj1 were counted in control lines AH017-13, OX3-9, OX1-19, and in *C9orf72* patient lines C9-T2-6, C9-T2-7, C9-7245-1, C9-7245-3, C902-2 and C902-10 (Fig 2 A). Differentiation efficiency was assessed after 8 weeks of neural culture, with a highly neuralized population indicated by a mean of 82% of Tuj1-positive cells across all lines (Fig 2 B). Approximately half of the Tuj1-positive neural cells in these cultures were also positive for the postmitotic MN marker Islet1

(42.7%-46.4% of total cells) (Fig 2 C). Cortical neurons were differentiated in parallel and functional experiments were carried out after 70-100 days in culture [23] (Fig S2 E). The sizes of the expansions were determined by Southern blotting in the mature MNs and estimated at 510-690 repeats for C9-T2-6, 420-640 repeats for C9-T2-7, 1210 for C9-7245-1, 1380 for C9-7245-3, 1000 for C902 (Fig S3 A). The presence of the repeats was also confirmed by repeat-primed (rp)-PCR in C9-02-2 and C9-7245-3 MNs and the absence of repeats was confirmed in the control lines OX1-19 and AH017-13 (Fig S3 B).

To assess the functional maturation of the iPSC-derived MNs, we performed whole-cell patch clamp recordings to determine their electrophysiological properties. Trains of action potentials were elicited in both control and patient MNs in response to depolarizing current injection in current-clamp recordings (Fig 2 D-E). Inward and outward ionic currents were recorded by voltage clamp in both controls and patients (Fig 2 F-G).  $\text{Ca}^{2+}$  dynamics were investigated using the calcium-sensitive dye Fluo-4AM. In the absence of external stimulation, spontaneous calcium transients were detectable in the cell bodies and processes of the neurons (Fig 2 H). The neurons were then stimulated with kainate (KA), to activate ionotropic receptors, and with KCl to depolarize the membrane and activate voltage-gated  $\text{Ca}^{2+}$  channels. Exposure to KCl induced  $\text{Ca}^{2+}$  transients in Islet1+ MNs (Fig 2 I). Similarly, exposure to kainate induced characteristic increases in intracellular  $\text{Ca}^{2+}$  levels in 80% of the cells (Fig 2 J).

### **RNA foci and RAN dipeptides are present in *C9orf72* MNs**

In the MNs derived from *C9orf72* patients, we identified a significant increase in intranuclear GGGGCC-repeat containing RNA foci in all lines (Fig 3 A,B). The specificity of the probe was confirmed by treatment with RNase, which reduced the number of foci substantially (Fig S4 A), and with DNase I, after which the number of foci was maintained. To investigate evidence for the RAN translation hypothesis, we performed a dot blot analysis using anti-GA, GP, GR and PR antibodies and we could detect traces of all the dipeptides in the MNs from *C9orf72* patients (Fig 3 C).

### **Elevated ER Ca<sup>2+</sup> levels, altered mitochondrial morphology and reduced Bcl-2 levels in *C9orf72* motor neurons**

Although a direct link between calcium signalling and *C9orf72* expansions has not yet been reported, calcium dysregulation is an established feature of the pathogenesis of ALS. The levels of Ca<sup>2+</sup> stores in the ER were analysed in fully differentiated MNs from 8 separate cultures of three independent differentiations for each cell line (OX3-9, C9-T2-6, C9-T2-7, C9-7245-1 and C9-7245-3). To assess the total content of ER Ca<sup>2+</sup> stores, 10  $\mu$ M thapsigargin (TG) was applied to the motor neurons and the peak increase in fluorescence intensity resulting from ER Ca<sup>2+</sup> depletion into the cytoplasm was measured (Fig 4 A, B). TG acts as a potent inhibitor of the SERCA pump on the ER membrane, blocking the reuptake of Ca<sup>2+</sup> ions from the cytoplasm. In our cultures, we detected significantly higher levels of ER Ca<sup>2+</sup> content in the *C9orf72* iPSC-derived MNs, which showed up to 2-fold increase in cytoplasmic fluorescence intensity compared to the control MNs ( $p < 0.05$ , One-Way ANOVA) (Fig 4 C).

Abnormal elevations in ER  $\text{Ca}^{2+}$  levels can cause ER stress and can activate the unfolded protein response (UPR). We investigated levels of proteins associated with the UPR and ER stress responses and we found significantly elevated levels of GRP78/BiP in C9-7245-1 and C9-7245-3 iPSC-derived MNs compared to AH017-13 and OX1-61 MNs ( $p < 0.05$ , One-Way ANOVA) (Fig 4 D, E). We did not detect differences in the splicing of XBP-1 between patients and controls (Fig S5 A).

The Bcl-2 family of proteins are key components involved in the modulation of calcium homeostasis in the ER, controlling the release of  $\text{Ca}^{2+}$  ions through the  $\text{IP}_3\text{R}$  and through channels formed by its own dimerization [19]. The balance between pro-apoptotic and anti-apoptotic members of the Bcl-2 family is essential for controlling ER stress-induced apoptosis and mitochondrial membrane permeability [17]. In *C9orf72* MNs, we detected an imbalance between pro-apoptotic and anti-apoptotic members of the Bcl-2 family, which showed up to 50% downregulation of anti-apoptotic Bcl-2 in C9-7245-1, C9-7245-3 and C902-2 patient lines (Fig 4 F, G) and 75% reduction of Bcl- $\text{X}_\text{L}$  ( $p < 0.05$ , One-Way ANOVA) (Fig 4 H). Significant upregulation of pro-apoptotic BAK was detected in the MNs with large expansions, which showed up to 2.5 fold increase in expression ( $p < 0.05$ , One-Way ANOVA) (Fig 4 I). A significant upregulation of the pro-apoptotic BIM was also detected in some of the MN lines (Fig S5 B).

The Bcl-2 family of proteins is distributed throughout the ER and mitochondrial membranes, where they play critical roles in regulating intracellular  $\text{Ca}^{2+}$  homeostasis and exchange of  $\text{Ca}^{2+}$  between the 2 organelles. Following dysfunction of these proteins and  $\text{Ca}^{2+}$  overload, the mitochondria undergo typical morphological changes, such as swelling and perturbation of the outer membrane, accompanied by the release

of apoptotic factors [24]. To assess these alterations, we used the potentiometric dye MitoTracker Red CMXRos, accumulation of which is dependent on mitochondrial potential (Fig 4 J). We found up to 30% reduction in mitochondrial membrane potential in C9 MNs compared to controls ( $p < 0.01$ , One-Way ANOVA) (Fig 4 K).

In line with these findings, analysis of morphological alterations of the mitochondria using electron microscopy (EM) also revealed swollen mitochondria in the neurons from *C9orf72* patients compared to the healthy control (Fig 4 L). Alterations of the mitochondrial morphology and abnormalities in the cristae structures were observed in up to 88% of *C9orf72* neurons compared to only 2% normal healthy neurons (Fig 4 M).

These mitochondrial alterations are consistent with the release of apoptotic factors, such as cytochrome *c*, which was found to be elevated in all patient lines, specifically in the C9-7245-1 compared to healthy controls ( $p < 0.05$ , One-Way ANOVA) (Fig 4 N, O).

These results indicate an association between *C9orf72* expansions and elevated ER  $\text{Ca}^{2+}$  concentrations and stress, as well as mitochondrial alterations which may be the drivers for downstream apoptosis.

### ***C9orf72* iPSC-derived MNs and CNs have increased susceptibility to apoptosis**

To investigate whether MNs and CNs carrying pathogenic hexanucleotide expansions in *C9orf72* develop an apoptotic phenotype under normal culture conditions, we assessed markers of cellular stress and cell death.

Caspases are present in the cell as inactive pro-enzymes which can be activated by proteolytic cleavage, the cleavage of caspase-3 being an essential process that initiates

the apoptotic disassembly of the cell. In neurons from *C9orf72* patients we found significantly increased immunostaining for cleaved caspase-3 compared to the control neurons from OX1-19, OX3-9 and AH017-13 (Fig 5 A). We detected between 8.1% and 10.5% of neurons positive for cleaved caspase-3 in the *C9orf72* patient lines, compared to a maximum of 2.9% in OX1-19 and 1.9% in OX3-9 ( $p < 0.05$ , One-Way ANOVA) (Fig 5 B). In line with this finding, the ALS/FTD-derived MNs also showed significantly increased staining with propidium iodide (PI) in up to 32% of *C9orf72* MNs compared to healthy MNs, indicating reduced survival compared to control MNs ( $p < 0.01$ , One-Way ANOVA) (Fig 5 C).

In cortical neurons cultures, we found a significantly high frequency of cleaved caspase-3 in the C902-2 and C902-10 lines, with up to 35% of neurons positive for apoptotic markers compared to controls ( $p < 0.05$ , One Way ANOVA) (Fig 5 D-E). These results also correlated with high PI staining, which was found in up to 40% of the neurons analysed in the C902 patient lines ( $p < 0.001$ , Student's *t* test) (Fig 5 F).

Taken together, these results demonstrate that the presence of the *C9orf72* hexanucleotide expansions activates caspase-3 dependent apoptosis in both motor and cortical neurons of patients.

### **Stress granules and protein aggregates are present in *C9orf72* iPSC-derived MNs and CNs**

SQST1/p62, a marker of the UPR and autophagy flux is found in neuronal aggregates in autopsy samples from *C9orf72* ALS/FTD cases. We found significantly elevated levels of SQST1/p62 in MNs derived from each *C9orf72* patient, and the patient with

the highest number of repeats showed up to 2-fold higher expression of SQST1/p62 compared to control MNs at baseline ( $p < 0.05$ , One-Way ANOVA) (Fig 6 A-C). No differences were detected in the autophagic marker LC3-II/LC3-I in MNs (Fig S5 C, D). In cortical neurons carrying the *C9orf72* expansions we also found approximately 40% neurons positive for p62 aggregates (Fig 6 D-F). In contrast to motor neurons, these results correlated with an increase in levels of LC3-II in C902-2 and C902-10 compared to controls OX3-9 and AH017-13 ( $p < 0.05$ , One-Way ANOVA) (Fig 6 G).

Stress granules are formed when RNA binding proteins aggregate through their glycine rich domains. These granules function to sequester, silence or degrade specific RNA transcripts as part of a mechanism that adapts patterns of local RNA translation to facilitate the response to cellular stress [25]. The poly-A binding protein (PABP) is a major component of stress granules and serves as a robust marker of the activation of the stress response. In our neuronal culture, we found up to 5-fold increase in the frequency of neurons with PABP-positive granules under resting conditions in all five *C9orf72* patient MN lines compared to controls ( $p < 0.05$ , One-Way ANOVA) (Fig 6 H-I). The *C9orf72* MNs were also immunopositive for the early stress granule marker, TIA-1 (Fig 6 H). PABP positive stress granules were detected in up to 40% of *C9orf72* cortical neurons compared to a maximum of 20% control cortical neurons (Fig 6 J-K).

## **Discussion**

The inability to directly study neurons in patients suffering from neurodegenerative disorders has been a significant impediment to the understanding of basic

mechanisms, particularly those occurring early in the disease process. Recent developments in stem cell technology have substantially expanded the range of cellular models available in ALS by allowing the direct observation of pathomechanisms in MNs derived from induced pluripotent stem cells obtained from fibroblasts harbouring the genetic background conferring disease susceptibility.

The mechanisms underlying *C9orf72* toxicity in motor neurons remain to be elucidated, but possibilities include haploinsufficiency and toxic gain-of-function, mediated either by RNA foci or toxic aggregation-prone polypeptides resulting from RAN translation [1, 10, 26, 27]. These potential mechanisms are not mutually exclusive and may each play a role. In this study, we identified several converging pathways that are altered in MNs and CNs from *C9orf72* patients.

RNA foci have been consistently detected in the cortical neurons of patients with hexanucleotide repeats [1, 14] and we detected discrete RNA foci in our motor neuron cultures from all *C9orf72* patient lines. Since the presence and number of RNA foci did not correlate with other phenotypes, the relationship of RNA foci to the pathology we observe in this cellular model requires further investigation. A dot blot analysis using high levels of proteins revealed the presence of the GA, GP, GR and PR dipeptides in the patient lines, but given their low abundance and difficulty of detection, it is challenging to analyse their contribution to neuronal degeneration in our cellular model. A recent study has shown the formation of stress granules in neuronal cells transfected with poly-GA, poly-GP and poly-PA, suggesting that RAN translation may be associated with stress granule formation and neurodegeneration [28]. In our cultures of *C9orf72* MNs and CNs, we detected the frequent presence of

stress granules under baseline conditions. Stress granules are conserved cytoplasmic aggregates of nontranslating messenger ribonucleoprotein complexes (mRNPs) and they are dynamic structures which dissociate when conditions return to normal or they are cleared by autophagy [29]. The accumulation of PABP<sup>+</sup> stress granules suggests impairments in autophagy and proteasomal degradation and the convergence of these two dysfunctional pathways may partially explain the vulnerability of *C9orf72* MNs and CNs to degeneration.

Calcium is known to exert a complex regulatory role in the ER-mitochondrial crosstalk, playing an essential part in cell death signalling [30]. Bcl-2 together with Bcl-X<sub>L</sub> are located on the ER membrane where they lower the [Ca<sup>2+</sup>]<sub>ER</sub> by forming Ca<sup>2+</sup> permeable leakage pores [31]. Reduced levels of Bcl-2 and Bcl-X<sub>L</sub> contribute to non-physiological accumulation of Ca<sup>2+</sup> ions into the ER due to reduced leakage, and we confirmed elevated levels of ER Ca<sup>2+</sup> stores in the *C9orf72* MNs. We have recently shown that TDP-43 pathogenic mutations in mouse motor neurons and in a human cellular model also interfere with ER Ca<sup>2+</sup>, but with the opposite effect, showing reduced ER Ca<sup>2+</sup> levels and decreased ER Ca<sup>2+</sup> release [32].

The tight connection between the ER Ca<sup>2+</sup> stores and mitochondria allows cells to respond promptly and efficiently to various Ca<sup>2+</sup> stimuli. However, under pathological conditions, the high levels of Ca<sup>2+</sup> from the ER will overload the mitochondria and consequently induce mitochondrial swelling, rupture of the outer membrane and release of apoptotic factors from the mitochondria [24]. The anti-apoptotic protein Bcl-2 is located on the mitochondrial membrane where it prevents pro-apoptotic BAK/BAX to oligomerize and to form channels that allow release of

the apoptotic factor cytochrome *c* into the cytosol. Low levels of Bcl-2 and increased levels of BAK promote the release of cytochrome *c* from the mitochondria triggering the activation of caspases that regulate the induction and execution of apoptosis [33]. Consistent with this, we detected high levels of cleaved caspase-3 and cytochrome *c* in the patient-derived MNs and CNs, indicating the C9 neurons are undergoing caspase-dependent apoptosis.

Disruption of ER Ca<sup>2+</sup> homeostasis also plays a direct role in the induction of ER stress by interfering with the function of Ca<sup>2+</sup>-sensitive ER chaperones [34]. Constitutive activation of ER stress in *C9orf72* MNs, which interestingly appears to be greater in cells with a higher number of hexanucleotide repeats, may contribute to the induction of ER-stress associated apoptosis. This is consistent with a previous study of *C9orf72*-patient iPSC-derived MNs which found upregulation of transcripts involved in UPR activation and ER stress [35]. While all MNs carrying repeats showed a survival deficit and an active apoptotic pathway, only the patient-derived MNs with large hexanucleotide expansions showed ER stress under baseline conditions. These results suggest that ER stress is activated by high numbers of pathogenic repeats, but is not the exclusive pathway in the induction of neuronal death. In accordance with our observations, a similar study using SOD1<sup>+A4V</sup> iPSC-derived MNs found elevated ER stress in patient MNs, but treatment with salubrinal or knockdown of XBP-1 did not rescue the survival deficit, suggesting that ER stress is only one of many components that may be contributing to MN death [35]. Selective neuronal vulnerability may involve specific combinations of mutually reinforcing stressors and altered pathways that ultimately converge on neuronal death [36]. Consequently, the gradual increase of stress in an age-dependent fashion in vulnerable

MNs may underlie the progression of neurodegeneration in ALS. Taken together, our results highlight the importance of ER  $\text{Ca}^{2+}$  regulation in different *in vivo* and *in vitro* models and supports the hypothesis that MN degeneration in ALS is mediated by cell-type specific impairments in intracellular  $\text{Ca}^{2+}$  handling [37-39].

Motor neurons and cortical neurons from *C9orf72* patients also showed high levels of SQST1/p62, which is an adaptor molecule for selective autophagosomal degradation of ubiquitinated targets and its accumulation reflects the inefficiency of protein aggregate removal by the autophagic machinery [40]. SQST1/p62 is a common component of protein aggregates found in neurodegenerative diseases, such as Lewy bodies in Parkinson's disease, neurofibrillary tangles in Alzheimer's disease, and in huntingtin aggregates [41-43]. Elevated p62 levels have previously been reported in cortical neurons derived from iPSCs of FTD patients carrying hexanucleotide expansions in *C9orf72* and it is a pathological marker in post-mortem analysis from *C9orf72* carriers [5, 7]. We detected high SQST1/p62 levels in motor neurons under basal conditions without changes in LC3-II/LC3-I turnover, indicating a reduced autophagic flux in *C9orf72* motor neurons compared to healthy controls. Interestingly, elevated SQST1/p62 levels were correlated with increased LC3-II in cortical neurons from an ALS/FTD patient, suggesting a potentially distinct accumulation of autophagosomes in these cells and impairment in the autophagic process, which is specific to the cortical neurons and is not recapitulated in motor neurons.

## **Conclusions**

Our study identified several cellular phenotypes that associate with the repeat expansions in *C9orf72* in MNs and CNs and molecular changes that may account for their vulnerability in pathological conditions. We have identified an apoptotic pathway that is activated in the *C9orf72* neurons, which involves ER calcium dysregulation and stress, followed by mitochondrial alterations, the release of apoptotic factors from the mitochondria and the formation of stress granules. It is unclear where this chain of phenotypes is initiated, whether the presence of toxic RAN dipeptides induces ER stress and dysregulation of calcium, or whether the presence of the RNA foci leads to the recruitment of essential  $\text{Ca}^{2+}$  regulators transcripts. Further work is currently underway to analyse the molecular mechanism linking the hexanucleotide repeats in *C9orf72* and the neuronal phenotypes. In the current study, a high number of repeat expansions (>1000) correlated with a more severe phenotype in our patient MN cultures, but a more extensive study is required to establish whether there is a correlation between the number of repeats and severity of the cellular phenotypes. Although our observations suggest that the cellular phenotypes observed are robust enough to be used to as a tool to screen potential therapies, further work is required to establish the exact mechanistic relationship between the pathological pathways activated in *C9orf72* MNs, CNs and neurodegeneration.

## **Materials and Methods**

## **Ethics Statement**

The human iPSC cell lines derived for this study were derived from human skin biopsy fibroblasts, following signed informed consent, with approval from the South East Wales Research Ethics Committee (REC ref no. 12/WA/0186) and the South Central – Berkshire Research Ethics Committee, UK (REC 10/H0505/71).

## **Generation and culture of iPSC Lines**

In this study, we used 6 iPSC-derived lines from 4 healthy controls: NHDF-1; OX3-9; OX1 clones 19 and 61; AH017 (clones 3 and 13 ) and 6 lines from 3 C9orf72 patients: C9-T2 (clones 6 and 7); C9-7245 (clones 1 and 3); C9-02 (clones 2 and 10). All iPSC lines were derived from skin biopsy fibroblasts in the James Martin Stem Cell Facility, University of Oxford, under standardised protocols to minimise any potential variation attributable to laboratory differences and/or handling. Fibroblasts tested negative for mycoplasma using MycoAlert (Lonza) and the derived iPSC lines also tested negative using the same test. The control iPSC line iPS-NHDF-1 [44 year-old female] [44] and iPS-OX1-19 [36M] [45], derived with Yamanaka retroviruses, have been published previously. C9-T2-6, C9-T2-7 [62M], 7245-1, 7245-3 [48F] and the control lines iPS-OX3-9 [49M] and iPS AH017-13 [67F] (previously published [46]) were derived using the SeVdp(KOSM)302L Sendai virus system [47, 48]. This contains the genes for all four reprogramming transcription factors (Klf4, Oct4, Sox2, c-Myc) expressed from a single transcript, packaged into a single Sendai virus, ensuring consistency of gene dosage ratio. Sendai virus is a negative strand RNA virus that is retained in the cytoplasm. As such, the likelihood of insertion of genetic material into the genome is negligible. This version of the system also contains a target for mir302; mir302 is expressed in pluripotent cells, but not in the originating

fibroblasts, so the viral RNA will be detected and degraded once the cells have reprogrammed to pluripotency, ensuring complete removal of exogenous genetic material within a few passages. AH017-13 [67F], OX1-61 [36M], C902-02, C902-10 [62F] were derived using the commercially available Sendai virus-based reprogramming system Cytotune, which contains the reprogramming genes Klf4, Oct4, Sox2, c-Myc packaged into individual viruses (Life Technologies, used according to the manufacturer's instructions).

Transduced fibroblasts were transferred onto mitotically-inactivated mouse embryonic feeder cells (MEFs; derived in-house from outbred Swiss mice established and maintained at the Department of Pathology, Oxford [49, 50], or MEFs purchased from Merck, derived from CF1 mice) on 0.1% gelatin coated plates on day 2 and were cultured in iPS medium consisting of knock-out DMEM (Invitrogen), 10% knock-out-Serum Replacement (Invitrogen), 2 mM Glutamax-I (Gibco), 100 U/mL penicillin (Invitrogen), 100 µg/mL streptomycin (Invitrogen), 1% non-essential amino acids (Invitrogen), 0.5 mM β-mercaptoethanol (Invitrogen), 10 ng/mL bFGF (R&D). Medium was replaced (50%) daily, and substituted with MEF-conditioned medium from day 10 onwards. Colonies displaying iPSC morphology were picked on ~day 21-28 and transferred onto MEFs by manual dissection every 5–7 days. iPSC lines were adapted to feeder-free conditions onto Matrigel-coated plates (BD Matrigel hESC-qualified Matrix) in mTeSR™1 (100% daily medium change, StemCell Technologies). Bulk and routine passaging was by 0.5mM EDTA[51] to make large-scale, quality-controlled, cryopreserved master-stocks (between ~p10-20). The initial master-stock underwent all the QC tests described, and all subsequent submaster-stocks underwent SNP analysis for tracking and genome integrity. The number of

feeder-free passages was kept to a minimum, to ensure that differentiation experiments were set up within a narrow window of passage numbers. If single-cell dissociation was required for particular applications, medium was supplemented with Rock inhibitor Y27632 (10  $\mu$ M; Calbiochem) on the day of passage.

### **Assessment of Genome Integrity, and Tracking**

Genome integrity was assessed by Illumina Human CytoSNP-12v2.1 beadchip array (~300,000 markers) or Illumina HumanOmniExpress-24 (~700,000 markers) and analysed using KaryoStudio and GenomeStudio software (Illumina). SNP deviations in the iPSC lines were compared to the original pool of fibroblasts. This also confirmed the identity of the iPSC to the original fibroblasts. SNP datasets have been deposited in GEO under the accession number GSE64582.

### **Sendai clearance assay**

Clearance of Sendai virus from reprogrammed cell lines was confirmed by PCR. RNA was extracted using an RNeasy kit (Qiagen) from iPSC, from fibroblasts (negative control) and from fibroblasts infected with Sendai reprogramming viruses 5 days previously (positive control). Reverse transcription was carried out using a RetroScript kit (Ambion), using 2  $\mu$ g template RNA in 20  $\mu$ l reaction volume. 2  $\mu$ l of 1:10 dilution of cDNA product was used in a 25  $\mu$ l PCR reaction. For lines reprogrammed with SeVdp(KOSM)302L, QRT-PCR was carried out on an Applied Biosystems StepOne Plus Real Time PCR machine, with StepOne software, using Applied Biosystems 2xSYBR green PCR mix+ROX and 60 degree Celsius anneal. Target gene transcript levels were compared to actin  $\beta$  control (actin  $\beta$  primers, Eurogentec), and subsequently to the positive control[52].. Sendai-specific primers

were 5'-AGACCCTAAGAGGACGAAGACAGA-3' and 5'-ACTCCCATGGCGTAACTCCATAG-3'. For lines reprogrammed with the Cytotune Sendai reprogramming system, PCR was carried out according to the manufacturers' instructions, with a primer pair for each of the four reprogramming viruses and also for the backbone, using actin  $\beta$  as a positive control for each line (actin  $\beta$  primers, Eurogentec), and products were visualised by running on a 1.5% agarose gel with ethidium bromide, and 2-log ladder or 100bp ladder (New England Biolabs). Primer pairs were: SeV F: GGA TCA CTA GGT GAT ATC GAG C; R: ACC AGA CAA GAG TTT AAG AGA TAT GTA TC; SOX2 F: ATG CAC CGC TAC GAC GTG AGC GC; R: AAT GTA TCG AAG GTG CTC AA; KLF4 F: TTC CTG CAT GCC AGA GGA GCC C; R: AAT GTA TCG AAG GTG CTC AA; c-MYC F: TAA CTG ACT AGC AGG CTT GTC G; R: TCC ACA TAC AGT CCT GGA TGA TGA TG; OCT4 F: CCC GAA AGA GAA AGC GAA CCA G; R: AAT GTA TCG AAG GTG CTC AA.

### **PluriTest**

RNA was extracted from iPSC using an RNeasy kit (Qiagen) for Illumina HT12v4 transcriptome array analysis. The data files were then uploaded to [www.pluritest.org](http://www.pluritest.org) and scored for pluripotency, as previously described. The Illumina HT12v4 transcriptome array results have been deposited in the Gene Expression Omnibus (GEO) under accession number GSE64583.

### **Flow cytometry**

iPSC lines were assessed for expression of pluripotency markers by flow cytometry. Cells were lifted with TrypLE (Life Technologies), fixed with 4% paraformaldehyde

in PBS for 10 minutes then permeabilised in methanol at -20 degrees Celsius.  $0.5-1 \times 10^6$  cells were then washed and stained in flow cytometry buffer consisting of PBS, human IgG (10  $\mu\text{g}/\text{mL}$  Sigma), FCS (1% Hyclone) and sodium azide (0.01%), with an antibody or an isotype-matched control (with same fluorophore, from the same manufacturer, at 1  $\mu\text{g}$  antibody/million cells) for 30 min. After the primary staining, cells were washed three times, and fluorescence was measured using a FACS Calibur (Becton Dickinson) – data from 10,000 cells were collected and analysed using FlowJo software, gated on FSC-SSC dot plots, and plotted as histograms (y axis normalised to mode). The following antibodies (clone, isotype control, supplier) were used: TRA-1-60 (B119983, IgM-488, Biolegend), NANOG (D73G4, IgG-647, Cell Signaling Technology)

### **Differentiation of iPSCs to motor neurons**

Motor neurons were differentiated from iPSCs using 3 different previously published protocols, with several modifications [20, 21]. In one of the differentiation protocols, the differentiation process was initiated by neural induction of embryoid bodies (EBs) and was followed by neural progenitor stages and neural maturation. After 4 days in neural induction medium, the EBs were plated onto Geltrex-coated plates and caudalization of the neural progenitors was obtained by addition of retinoic acid (RA) to the neural differentiation medium at a concentration of 0.1  $\mu\text{M}$ . After 10 days, the colonies displaying neural rosettes structures were isolated and expanded in suspension in the form of neurospheres in the presence of the ventralizing growth factor, sonic hedgehog (SHH) at a concentration of 100 ng/mL. The protocol was modified at this step to enhance cell survival by addition of Rock inhibitor Y27632, which prevents cell detachment and is a widely used supplement in stem cell cultures

[53, 54]. After two weeks of culture as neurospheres, the cells were plated on Geltrex-coated coverslips and terminal differentiation was induced by addition of brain derived neurotrophic factor (BDNF 10 ng/mL), glial-derived neurotrophic factor (GDNF, 10 ng/mL), insulin-like growth factor (IGF-1, 10 ng/mL), cyclic AMP (cAMP) and ascorbic acid. The differentiating MNs were cultured for another 3-4 weeks before functional assays were performed.

Using the second differentiation protocol [21], the iPSCs were cultured as monolayers and neural induction was started by addition of compound c (1  $\mu$ M) in DMEM/F12 and Neurobasal medium, supplemented with N2, B27, Heparin 0.2%. After three days, the medium was supplemented with retinoic acid (1  $\mu$ M) and after another three days, SAG and purmorphamine (1  $\mu$ M) were added and compound c removed. After 9 days in culture, the neural precursors were re-plated at a low density for differentiation and the medium was supplemented with the growth factors BDNF, GDNF and IGF-1 (10 ng/mL), ascorbic acid and cAMP. After 20 days, the motor neurons were re-plated for experiments and allowed to mature for another 3-4 weeks before functional experiments.

Motor neurons were differentiated using a third differentiation protocol [22], with extensive modifications. Monolayers of iPSCs were induced in DMEM/F12/Neurobasal with N2 supplement, B27 supplement, ascorbic acid (0.5  $\mu$ M), 2-mercaptoethanol (1x), compound C (3  $\mu$ M) and Chir99021 (3  $\mu$ M) for 4 days. After 4 days in culture, RA (1  $\mu$ M) and SAG (500 nM) were added to the medium. The next day, Chir99021 and compound C were removed from the medium and the cells were cultured for another 4-5 days. Neural precursors were then dissociated and plated at a low density for 7 days. The medium was supplemented with growth factors BDNF (10 ng/mL), GDNF (10 ng/mL), DAPT (10  $\mu$ M) and laminin (0.5  $\mu$ g/mL).

After 7 days, DAPT and laminin were removed from the medium and the neurons were allowed to mature for another 2-4 weeks before functional experiments.

### **Differentiation of iPSCs to cortical neurons**

The differentiation of iPSCs to cortical neurons was performed following a previously established protocol [23]. Briefly, the iPSCs were grown on Matrigel as monolayers and neural differentiation was induced by changing the medium to basic neural medium DMEM/F12: Neurobasal 1:1, 1 x N2, 1 x B27, Insulin 5 µg/mL (Sigma), non-essential amino acids 100 µM (Life Technologies), 2-Mercaptoethanol 100 µM, Antibiotic-antimycotic (Life Technologies) and supplementing it with 500 ng/mL Noggin (PeproTech) and 10 µM SB431542 (Enzo Life Sciences). Medium was changed every 24 hours until neural rosettes could be observed (12-16 days). The neuroepithelial layer was lifted using Dispase (Life Technologies) and the medium was supplemented with 20 ng/mL FGF2 (PeproTech) for 4-6 days. The neural precursors were split using Accutase (Life Technologies) at a low density. The medium was then supplemented with 10 ng/mL BDNF and 10 ng/mL GDNF, while FGF2 was withdrawn. The neurons were allowed to extend processes and to mature for 60-70 days before functional assays.

### **Immunocytochemistry**

Mature MNs and CNs were fixed with 4% paraformaldehyde-PBS for 15 minutes and incubated with 10% goat serum for blocking for 1 hr at room temperature. The cells were incubated overnight at 4<sup>0</sup>C with rabbit anti-cleaved-caspase-3 (Cell Signalling, 1:350), rabbit anti-PABP (1:500), mouse anti-Islet 1 (DSHB, 1:200), mouse or rabbit

anti-Tuj1 (Covance, 1:1000), mouse anti-p62 (Abcam, 1:500), rabbit anti-GA (ProteinTech, 1:500), rabbit anti-GP (ProteinTech, 1:500), rabbit anti-PR (ProteinTech, 1:500), rabbit anti-GR (ProteinTech, 1:500), goat anti-TIA-1 (Santa Cruz Biotechnology, 1:500). After washing with 0.1% TritonX/PBS 3 times for 10 minutes, the samples were incubated with Alexa Fluor 488 or Alexa Fluor 568 conjugated goat anti-rabbit or goat anti-mouse secondary antibodies (Life Technologies) for 1 hr at room temperature. Nuclei were stained with DAPI. Fluorescence was visualized using a confocal microscope Zeiss LSM.

### **Immunoblotting**

Cells were washed with phosphate-buffered saline (PBS) and then scraped in PBS. Cells were lysed in RIPA buffer (0.5% sodium deoxycholate, 0.1% SDS, 1% NP-40 and 1% protease inhibitor cocktail (Sigma)), sonicated and incubated on ice for 30 minutes. After centrifugation at 5000x g for 10 minutes, the supernatant was retained and protein concentration was quantified using the BCA assay according to the manufacturer's instructions (Sigma). Protein was boiled in Laemmli buffer for 5 minutes at 95°C and 10 µg of protein were loaded on gels. For Western blot analysis, lysates were resolved on a SDS-PAGE (10% or 12% Tris-glycine gel) using a Novex X-Cell electrophoresis system (Life Technologies) under constant current (125 mA for 1hr) and transferred to a methanol activated polyvinylidene difluoride membrane (Immobilon-P, PVDF) using the Novex Blot system under constant voltage (100 mA for 90 mins). Blots were blocked in blocking buffer (PBS, 1% Tween-20, 5% skimmed milk) for 1 hr at room temperature and then incubated in PBS+1% Tween-20+1% milk with primary antibodies overnight at 4°C. Primary antibodies used were mouse anti-Bcl-2 (Dako, 1:200), mouse anti-SQST1/p62 (Abcam, 1:1000), rabbit

anti-GRP78/BiP (Abcam, 1:1000), mouse anti- $\beta$ -actin (Sigma, 1:5000). Horseradish peroxidase (HRP)-conjugated anti-mouse IgG or anti-rabbit IgG (BioRad) were used as secondary antibodies and the signal was visualized using an ECL or ECL Plus detection system (Amersham Pharmacia Biotech). The integrated optical density of each band was measured in ImageJ and expression was normalized to actin levels in the same blot for comparative expression assessment.

For dot blot analysis, a strip of nitrocellulose membrane was used and 10  $\mu$ L containing 10  $\mu$ g of protein were loaded as a dot. The blot was allowed to dry and blocked in 5% skim milk 0.1% Tween/PBS for 1 hr. The blocking buffer was removed and primary antibodies were added to each dot (1:1000) and incubated for 1 hr at room temperature. The membrane was washed 3 times for 10 minutes and HRP-conjugated anti-rabbit secondary antibodies were added for 1 hr at room temperature. The membrane was washed 3 times for 10 minutes in 0.1% Tween/PBS and the signal was detected by ECL (Amersham Pharmacia Biotech).

### **Propidium iodide staining**

Propidium iodide (PI) staining was performed according to the manufacturer's instructions (Sigma). Briefly, the neurons were incubated with 1  $\mu$ g/mL PI stock solution and RNase A for 10 minutes at 37<sup>o</sup>C. The solution was then washed with PBS and cells were fixed in 4% PFA. Immunocytochemistry was then performed as described above.

### **Mitochondrial staining**

Cultures of differentiated MNs were incubated with 100 nM of the potentiometric dye MitoTracker Red CMXRos (Thermo Fisher) for 45 minutes at 37°C. The neurons were then washed with PBS, fixed in 4% PFA and immunostained for Islet1 and SMI-32 according to the immunocytochemistry protocol above.

### **ER calcium imaging**

Mature MNs were loaded with 10 µM Fluo 4-AM dissolved in 0.2% dimethyl sulfoxide (Sigma-Aldrich) with 0.04% Pluronic acid in Hank's Balanced Salts Solution (HBSS) supplemented with CaCl<sub>2</sub> and MgCl<sub>2</sub>. To image Ca<sup>2+</sup> signals, we used wide-field fluorescence microscopy using a Nikon TE 2000 inverted microscope equipped with a 40 x oil-immersion objective and EMCCD camera (Evolve 128, Photometrics) for imaging fluorescence emission (510-600 nm) at a frame rate of 300 s<sup>-1</sup>. Fluorescence was imaged at 25°C from a 12 x 12 µm (128 x 128 pixel) region within 18 mm cover slips, on which MNs were cultured. ER Ca<sup>2+</sup> concentration levels were assessed by fluorescence measurements upon application of 1 µM thapsigargin (Life Technologies) in HBSS depleted of CaCl<sub>2</sub> and MgCl<sub>2</sub>. Fluorescence measurements are expressed as a ratio ( $\Delta F/F_0$ ) of the mean change in fluorescence ( $\Delta F$ ) at a pixel relative to the resting fluorescence at that pixel before stimulation ( $F_0$ ). Image analysis was performed using MetaMorph and ImageJ software.

### **RNA FISH**

Cells grown on 12 mm RNase-free coverslips in 24 well plates were fixed using 4% paraformaldehyde for 20 mins, washed in PBS and permeabilised using 0.2% Triton X-100 (Sigma) in PBS for 30 minutes. Cells were then dehydrated in a graded series of alcohols, air dried and rehydrated in PBS, briefly washed in 2X standard sodium

citrate (SSC) and then incubated in pre-hybridisation solution and hybridisation solution as previously described using Cy3-conjugated 2'-O-methyl RNA probes (Integrated DNA Technologies) [55]. Probe sequences were (CCCCGG)<sub>4</sub> or (CAGG)<sub>6</sub>. Optional treatments using RNase (0.1mg/ml; Life Technologies) or DNase (150U/ml, Qiagen) were performed at 37°C for 1 hour following permeabilisation. Following hybridisation washes, staining with Tuj1 (1:2000, Covance) and Islet1 (1:200, DSHB) were performed as described above.

### **Electrophysiology**

A coverslip containing iPSC-derived MNs was placed in a recording chamber mounted onto the stage of an upright microscope and the somata were targeted for recording using IR-DIC optics. Cells were continuously superfused with oxygenated aCSF (95% O<sub>2</sub>/5% CO<sub>2</sub>) containing 130 mM NaCl, 25 mM NaHCO<sub>3</sub>, 2.5 mM KCl, 1.25 mM NaH<sub>2</sub>PO<sub>4</sub>, 2 mM CaCl<sub>2</sub>, 2 mM MgCl<sub>2</sub>, and 10 mM glucose. Patch-clamp electrodes (4–7 MΩ) were filled with an intracellular solution containing 120 mM K-gluconate, 10 mM KCl, 10 mM HEPES, 4 mM MgATP, 0.3 mM NaGTP, 10 mM Na-phosphocreatine. Recordings were obtained using a Multiclamp 700B amplifier and digitized at 10–20 kHz using Digidata 1550 acquisition board.

### **Southern blotting**

5 µg of DNA were digested with 2 µl each of FastDigest DdeI and FastDigest AluI (Thermo Scientific) for 2 hours at 37°C. Digested DNA underwent electrophoresis in 0.8% agarose gel in 1X TAE and was transferred to a positively charged nylon

membrane (Roche) by capillary blotting. Roche DIG-labelled DNA molecular weight marker II was used as a ladder. DNA was cross-linked to the membrane by UV irradiation. Prehybridisation was carried out in 20 ml DIG Easy Hyb solution (Roche) at 53°C for 2 hours. 10 ng DIG-labelled (GGGGCC)<sub>5</sub> probe (custom synthesised by Integrated DNA Technologies, Inc) was incubated at 95°C for 5 minutes, snap cooled on ice for 30 seconds and immediately added to 7 ml DIG Easy Hyb solution pre-warmed to 60°C. Hybridisation took place overnight at 53°C. Stringency washes were performed at 64°C as follows: 2X standard sodium citrate (SSC), 0.1% sodium dodecyl sulphate (SDS) for 5 minutes, 2X SSC, 0.1% SDS for 10 minutes, 1X SSC, 0.1% SDS for 15 minutes three times, 0.5X SSC, 0.1% SDS for 15 minutes twice.

The membrane was prepared for detection using the DIG Wash and Block buffer set (Roche). The membrane was washed with Wash buffer for 2 minutes then incubated with Blocking solution for 1-2 hours. Anti-digoxigenin-AP Fab fragments (Roche) were incubated with the membrane at room temperature for 20-25 minutes at 1 in 10,000 dilution in Blocking solution. Three washes were carried out for 5, 15 and 15 minutes followed by a 5 minute equilibration step using the supplied Detection buffer. CDP-Star (Roche) was added to the membrane for at least 5 minutes and signal was detected by exposure to X-ray film. Repeat number was estimated by interpolating a plot of log<sub>10</sub> of band size in base-pairs versus distance travelled on the blot, with subtraction of 199 bp from the estimated base-pair number to account for the wild-type allele as performed by Beck et al [56].

## **qRT-PCR**

RNA was isolated from iPSC-derived MNs using the Qiagen RNeasy kit, according to the manufacturer's instructions. 500 µg of total RNA was used for reverse transcription reaction using SuperScript III reverse transcriptase (Life Technologies). Quantitative real-time PCR was performed with 100 µg of RNA reverse transcribed to cDNA and SYBR Green qPCR Supermix (Roche). Primers used for qPCR are listed below in Table 1.

Table 1: Oligos used for qRT-PCR.

Oligo	Sequence
BAK_fwd	5'-GGGTCTATGTTCCCCAGGAT-3'
BAK_rev	5'-GCAGGGGTAGAGTTGAGCA-3'
Bcl-XL_fwd	5'-ATGAACTCTTCCGGGATGG-3'
Bcl-XL_rev	5'-TGGACTCAAGGCTCTAGGTG-3'
BIM_fwd	5'-GGTCCTCCAGTGGGTATTTCTCTT-3'
BIM_rev	5'-ACTGAGATAGTGGTTGAAGGCCTGG-3'
HPRT_fwd	5'-GCTATAAATTCTTTGCTGACCTGCTG-3'
HPRT_rev	5'-AATTACTTTTATGTCCCCTGTTGACTGG-3'

### Repeat-primed PCR

RP-PCR assay was performed as previously described [1], in a reaction volume of 15 µl, containing 100 ng genomic DNA, 8 µl Extensor mastermix (Thermo Scientific), 6 µl of 5 M Betaine, 20 µM of both forward and repeat specific primer and 2 µM of reverse primer. The reactions were subjected to touchdown PCR programme, then 1 µl of each PCR product was mixed with 8 µl of Formamide, and 0.5 µl GeneScan size standard (Applied Biosystem). Then samples were analysed on an ABI3730 DNA Analyzer and visualised using GeneMapper software. Primer sequences used:

RP1-F FAM-TGTA AACGACGGCCAGTCAAGGAGGGAAACAACCGCAGCC;

RP1-R CAGGAAACAGCTATGACC;

RP1-Repeat:

CAGGAAACAGCTATGACCGGGCCCGCCCCGACCACGCCCCGGCCCCGGCC  
CCGG

### **Electron microscopy**

Cells grown on polyester filters were immersion fixed with 2.5% glutaraldehyde in 0.1M phosphate buffer (pH7.2) for 10 minutes at room temperature and prepared for electron microscopy by standard methods. Briefly, filters were stained with osmium tetroxide (1% v/v in 0.1M phosphate buffer) and uranyl acetate (2% w/v in distilled water), dehydrated through increasing concentrations of ethanol (70-100%) and acetone and embedded in Spurr's resin (Agar scientific, Reading UK). Ultrathin sections (50-80nm) were prepared by use of a Reichart-Jung ultracut microtome and mounted on nickel grids (Agar Scientific, Stanstead, Essex, UK). Sections were counterstained with lead citrate and uranyl acetate and examined on a JOEL 1010 transmission electron microscope (JOEL USA Inc., Peabody, MA, USA).

### **Statistical analyses**

All quantitative data were analysed using GraphPad Prism. Groups of 3 or more samples were subject to One-Way ANOVA, with Bonferroni or Dunnett's *post hoc* comparisons. Alpha was set at 0.05 for all ANOVAs and *post hoc* tests.

### **Dataset deposit**

The Illumina HT12v4 transcriptome array results have been deposited in the Gene Expression Omnibus (GEO) under accession number GSE64583. SNP datasets have been deposited in GEO under the accession number GSE64582. These are both contained in the Superseries GSE64584.

## **Acknowledgments**

We thank Dr. Adrian Isaacs and Dr. Sarah Mizielinska for their support and advice on the RNA FISH technique. We thank Dr. Alex Clarke for his advice on calcium imaging experiments. We thank the High-Throughput Genomics Group at the Wellcome Trust Centre for Human Genetics (funded by Wellcome Trust grant reference 090532/Z/09/Z) for the generation of the genotyping and transcriptome data.

## **Conflict of interest statement**

The authors declare no conflict of interest.

## **Funding**

This work was made possible by a charitable donation from the Lights Up Asia Foundation. RM is funded by the MND Association (107/516). NA is supported by a scholarship from the Government of Jordan. JS is funded by a Medical Research Council/Motor Neurone Disease Association Lady Edith Wolfson Clinical Research Training Fellowship (MR/LOO2167/1). AGLD is funded by a Wellcome Trust Clinical PhD Fellowship (AVRVTB0). MRT is funded by a Medical Research Council/Motor Neurone Disease Association Lady Edith Wolfson Senior Clinical

Fellowship (MR/K01014X/1). The Oxford Martin School (LC0910-004) and the Wellcome Trust (WTISSF121302) provide core support to the James Martin Stem Cell Facility within the Sir William Dunn School of Pathology. Control samples were supplied by the Oxford Parkinson's Disease Centre (OPDC) study, funded by the Monument Trust Discovery Award from Parkinson's UK, a charity registered in England and Wales (2581970) and in Scotland (SC037554), with the support of the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre based at Oxford University Hospitals NHS Trust and University of Oxford, and the NIHR Comprehensive Local Research Network.

## References

1. DeJesus-Hernandez M, Mackenzie IR, Boeve BF et al. Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. **Neuron**. 2011;72:245-256.
2. Renton AE, Majounie E, Waite A et al. A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. **Neuron**. 2011;72:257-268.
3. Majounie E, Renton AE, Mok K et al. Frequency of the C9orf72 hexanucleotide repeat expansion in patients with amyotrophic lateral sclerosis and frontotemporal dementia: a cross-sectional study. **Lancet Neurol**. 2012;11:323-330.
4. van der Zee J, Gijssels I, Dillen L et al. A Pan-European Study of the C9orf72 Repeat Associated with FTL: Geographic Prevalence, Genomic Instability, and Intermediate Repeats. **Human mutation**. 2013;34:363-373.
5. Bieniek KF, Murray ME, Rutherford NJ et al. Tau pathology in frontotemporal lobar degeneration with C9ORF72 hexanucleotide repeat expansion. **Acta Neuropathol**. 2012.
6. Troakes C, Maekawa S, Wijesekera L et al. An MND/ALS phenotype associated with C9orf72 repeat expansion: abundant p62-positive, TDP-43-negative inclusions in cerebral cortex, hippocampus and cerebellum but without associated cognitive decline. **Neuropathology**. 2012;32:505-514.
7. Cooper-Knock J, Hewitt C, Highley J et al. Clinico-pathological features in amyotrophic lateral sclerosis with expansions in C9ORF72. **Brain**. 2012;135:751-764.

8. Haeusler AR, Donnelly CJ, Periz G et al. C9orf72 nucleotide repeat structures initiate molecular cascades of disease. **Nature**. 2014;507:195-200.
9. Gijssels I, Van Langenhove T, van der Zee J et al. A C9orf72 promoter repeat expansion in a Flanders-Belgian cohort with disorders of the frontotemporal lobar degeneration-amyotrophic lateral sclerosis spectrum: a gene identification study. **Lancet Neurol**. 2012;11:54-65.
10. Ash PE, Bieniek KF, Gendron TF et al. Unconventional translation of C9ORF72 GGGGCC expansion generates insoluble polypeptides specific to c9FTD/ALS. **Neuron**. 2013;77:639-646.
11. Mori K, Weng SM, Arzberger T et al. The C9orf72 GGGGCC Repeat Is Translated into Aggregating Dipeptide-Repeat Proteins in FTL/ALS. **Science**. 2013.
12. Almeida S, Gascon E, Tran H et al. Modeling key pathological features of frontotemporal dementia with C9ORF72 repeat expansion in iPSC-derived human neurons. **Acta Neuropathol**. 2013;126:385-399.
13. Sareen D, O'Rourke JG, Meera P et al. Targeting RNA foci in iPSC-derived motor neurons from ALS patients with a C9ORF72 repeat expansion. **Sci Transl Med**. 2013;5:208ra149.
14. Donnelly CJ, Zhang PW, Pham JT et al. RNA toxicity from the ALS/FTD C9ORF72 expansion is mitigated by antisense intervention. **Neuron**. 2013;80:415-428.
15. Grosskreutz J, Van Den Bosch L, Keller BU. Calcium dysregulation in amyotrophic lateral sclerosis. **Cell Calcium**. 2010;47:165-174.
16. Cleveland DW, Rothstein JD. From Charcot to Lou Gehrig: deciphering selective motor neuron death in ALS. **Nat Rev Neurosci**. 2001;2:806-819.
17. Yoshida H. ER stress and diseases. **FEBS J**. 2007;274:630-658.
18. Chen R, Valencia I, Zhong F et al. Bcl-2 functionally interacts with inositol 1,4,5-trisphosphate receptors to regulate calcium release from the ER in response to inositol 1,4,5-trisphosphate. **J Cell Biol**. 2004;166:193-203.
19. Rong YP, Aromolaran AS, Bultynck G et al. Targeting Bcl-2-IP3 receptor interaction to reverse Bcl-2's inhibition of apoptotic calcium signals. **Mol Cell**. 2008;31:255-265.
20. Hu BY, Zhang SC. Differentiation of spinal motor neurons from pluripotent human stem cells. **Nat Protoc**. 2009;4:1295-1304.
21. Qu Q, Li D, Louis KR et al. High-efficiency motor neuron differentiation from human pluripotent stem cells and the function of Islet-1. **Nat Commun**. 2014;5:3449.
22. Maury Y, Come J, Piskorowski RA et al. Combinatorial analysis of developmental cues efficiently converts human pluripotent stem cells into multiple neuronal subtypes. **Nat Biotechnol**. 2015;33:89-96.
23. Shi Y, Kirwan P, Livesey FJ. Directed differentiation of human pluripotent stem cells to cerebral cortex neurons and neural networks. **Nat Protoc**. 2012;7:1836-1846.
24. Giorgi C, Romagnoli A, Pinton P et al. Ca<sup>2+</sup> signaling, mitochondria and cell death. **Current molecular medicine**. 2008;8:119-130.
25. Buchan JR, Parker R. Eukaryotic stress granules: the ins and outs of translation. **Mol Cell**. 2009;36:932-941.
26. Mori K, Weng SM, Arzberger T et al. The C9orf72 GGGGCC Repeat Is Translated into Aggregating Dipeptide-Repeat Proteins in FTL/ALS. **Science**. 2013;339:1335-1338.

27. Renton AE, Majounie E, Waite A et al. A Hexanucleotide Repeat Expansion in C9ORF72 Is the Cause of Chromosome 9p21-Linked ALS-FTD. **Neuron**. 2011;72:257-268.
28. Tao Z, Wang H, Xia Q et al. Nucleolar stress and impaired stress granule formation contribute to C9orf72 RAN translation-induced cytotoxicity. **Hum Mol Genet**. 2015;24:2426-2441.
29. Buchan JR, Kolaitis RM, Taylor JP et al. Eukaryotic stress granules are cleared by autophagy and Cdc48/VCP function. **Cell**. 2013;153:1461-1474.
30. Kroemer G, Reed JC. Mitochondrial control of cell death. **Nature medicine**. 2000;6:513-519.
31. Thomenius MJ, Distelhorst CW. Bcl-2 on the endoplasmic reticulum: protecting the mitochondria from a distance. **J Cell Sci**. 2003;116:4493-4499.
32. Mutihac R, Alegre-Abarategui J, Gordon D et al. TARDBP pathogenic mutations increase cytoplasmic translocation of TDP-43 and cause reduction of endoplasmic reticulum Ca signaling in motor neurons. **Neurobiol Dis**. 2014.
33. van Loo G, Saelens X, van Gurp M et al. The role of mitochondrial factors in apoptosis: a Russian roulette with more than one bullet. **Cell death and differentiation**. 2002;9:1031-1042.
34. Groenendyk J, Michalak M. Endoplasmic reticulum quality control and apoptosis. **Acta biochimica Polonica**. 2005;52:381-395.
35. Kiskinis E, Sandoe J, Williams LA et al. Pathways disrupted in human ALS motor neurons identified through genetic correction of mutant SOD1. **Cell Stem Cell**. 2014;14:781-795.
36. Saxena S, Caroni P. Selective neuronal vulnerability in neurodegenerative diseases: from stressor thresholds to degeneration. **Neuron**. 2011;71:35-48.
37. Alexianu ME, Ho BK, Mohamed AH et al. The role of calcium-binding proteins in selective motoneuron vulnerability in amyotrophic lateral sclerosis. **Ann Neurol**. 1994;36:846-858.
38. Lips MB, Keller BU. Endogenous calcium buffering in motoneurons of the nucleus hypoglossus from mouse. **The Journal of physiology**. 1998;511 ( Pt 1):105-117.
39. Mattson MP, LaFerla FM, Chan SL et al. Calcium signaling in the ER: its role in neuronal plasticity and neurodegenerative disorders. **Trends in neurosciences**. 2000;23:222-229.
40. Ichimura Y, Kumanomidou T, Sou YS et al. Structural basis for sorting mechanism of p62 in selective autophagy. **J Biol Chem**. 2008;283:22847-22857.
41. Zatloukal K, Stumptner C, Fuchsbichler A et al. p62 Is a common component of cytoplasmic inclusions in protein aggregation diseases. **Am J Pathol**. 2002;160:255-263.
42. Kuusisto E, Salminen A, Alafuzoff I. Ubiquitin-binding protein p62 is present in neuronal and glial inclusions in human tauopathies and synucleinopathies. **Neuroreport**. 2001;12:2085-2090.
43. Nagaoka U, Kim K, Jana NR et al. Increased expression of p62 in expanded polyglutamine-expressing cells and its association with polyglutamine inclusions. **J Neurochem**. 2004;91:57-68.
44. Hartfield EM, Yamasaki-Mann M, Ribeiro Fernandes HJ et al. Physiological characterisation of human iPS-derived dopaminergic neurons. **PloS one**. 2014;9:e87388.

45. van Wilgenburg B, Browne C, Vowles J et al. Efficient, long term production of monocyte-derived macrophages from human pluripotent stem cells under partly-defined and fully-defined conditions. **PloS one**. 2013;8:e71098.
46. Fernandes HJ, Hartfield EM, Christian HC et al. ER Stress and Autophagic Perturbations Lead to Elevated Extracellular alpha-Synuclein in GBA-N370S Parkinson's iPSC-Derived Dopamine Neurons. **Stem cell reports**. 2016;6:342-356.
47. Nishimura K, Sano M, Ohtaka M et al. Development of Defective and Persistent Sendai Virus Vector. **Journal of Biological Chemistry**. 2011;286:4760-4771.
48. Nishimura T, Kaneko S, Kawana-Tachikawa A et al. Generation of rejuvenated antigen-specific T cells by reprogramming to pluripotency and redifferentiation. **Cell stem cell**. 2013;12:114-126.
49. Chia R, Achilli F, Festing M et al. The origins and uses of mouse outbred stocks. **Nature genetics**. 2005;37:1181-1186.
50. Gardner RL. Investigation of cell lineage and differentiation in the extraembryonic endoderm of the mouse embryo. **Journal of embryology and experimental morphology**. 1982;68:175-198.
51. Beers J, Gulbranson D, George N et al. Passaging and colony expansion of human pluripotent stem cells by enzyme-free dissociation in chemically defined culture conditions. **Nat Protocols**. 2012;7:2029-2040.
52. Livak K, Schmittgen T. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the  $2^{-\Delta\Delta CT}$  Method. **Methods**. 2001;25:402-408.
53. Koyanagi M, Takahashi J, Arakawa Y et al. Inhibition of the Rho/ROCK pathway reduces apoptosis during transplantation of embryonic stem cell-derived neural precursors. **J Neurosci Res**. 2008;86:270-280.
54. Zhang L, Valdez JM, Zhang B et al. ROCK inhibitor Y-27632 suppresses dissociation-induced apoptosis of murine prostate stem/progenitor cells and increases their cloning efficiency. **PloS one**. 2011;6:e18271.
55. Mizielinska S, Lashley T, Norona FE et al. C9orf72 frontotemporal lobar degeneration is characterised by frequent neuronal sense and antisense RNA foci. **Acta Neuropathol**. 2013;126:845-857.
56. Beck J, Poulter M, Hensman D et al. Large C9orf72 hexanucleotide repeat expansions are seen in multiple neurodegenerative syndromes and are more frequent than expected in the UK population. **Am J Hum Genet**. 2013;92:345-353.

## Figure Legends

### Figure 1. iPSc line characterisation.

Characterisation data for line iPSc C9 T2 7. A) Karyogram showing genome integrity of iPSc (Illumina SNP array data analysed using Karyostudio). Autosomal detected regions deviating from reference data are annotated with green (amplification) or

orange bands (deletion); the left hand panel shows the parental fibroblasts as reference. B) FACs analysis of iPSc for pluripotency markers Tra-1-60 and Nanog (black line); grey filled plot, isotype control. C) hiPSC show pluripotent stem cell-like morphology (cell-cell-contact-dependent clusters, with high nucleus to cytoplasm ratio) by phase microscopy; scale bar: 100  $\mu$ m. D) PluriTest analysis of Illumina HT12v4 transcriptome array data shows that all the iPSc cell lines used cluster with pluripotent stem cells (red cloud) and not with partly- or differentiated cells (blue clouds). Each circle represents one iPSc line, with previously published lines OX1-19 and NHDF-1 included as reference points [1 = AH017-13; 2 = OX3-9; 3 = OX1-61; 4 = C902-10; 5 = AH017-3; 6 = OX1-19; 7 = NHDF-1; 8 = 7245-1; 9 = 7245-3; 10 = T2-7; 11 = C902-02; 12 = T2-6] ; y axis Pluripotency score, x axis Novelty score. E) Assessment of Sendai clearance from iPSc lines reprogrammed with SeVdp(KOSM)302L , by qRT-PCR, relative to positive control.

**Figure 2: Functional characterization of *C9orf72* iPSC-derived MNs.**

A) Immunostaining for Islet1 and Tuj1 in control lines (OX1-19, OX3-9, AH017) and 6 patient lines (C9-T2-6, C9-T2-7, C9-7245-1, C9-7245-3, C902-2, C902-10). B) Quantification of Tuj1 positive cells shows 80-95% of cells are neuronal and C) quantification of Islet1 positive cells shows 23-40% of cells are motor neurons, with non-significant differences between the genotypes ( $*p>0.05$ , One-Way ANOVA). D) Whole cell recording reveals that trains of APs can be evoked reliably in both control MNs and E) *C9orf72* MNs. Representative voltage clamp data showing inward and outward currents recorded during step membrane depolarization of control OX3-9 (F) and C9-7245-3 (G) neurons. H) Spontaneous calcium oscillations were recorded from neurons after incubation with Fluo-4 AM and typical elevations in intracellular

calcium were detected on stimulation with D) KCl and H) kainate (control neurons are shown in this image). n=3 independent differentiations; m=10-15 neurons recorded per line; 5 neurons per line for ionic current measurements. Data are represented as average±SEM. Scale bar = 20 μm.

**Figure 3: RNA foci and RAN dipeptides are detected in *C9orf72* patient-derived MNs.**

A) [CCCCGG]<sub>4</sub> probe was used to detect sense RNA foci from the *C9orf72* hexanucleotide expansions; B) RNA foci were detected in up to 65% of neurons from *C9orf72* iPSC-derived MNs (121-157 neurons counted), while no foci were detected in the controls (\*\**p*<0.01, One-Way ANOVA). C) Dot blot analysis of GR, PR, GA and GP dipeptides in controls (OX3-9 and AH017) and patient lines (C902-2, C9-7245-3 and C902-3) at baseline and after treatment with MG-132 for 24 hrs. Scale bar = 5 μm. Data are represented as average±SEM.

**Figure 4: Increased ER Ca<sup>2+</sup> content is detected in *C9orf72* iPSC-derived MNs along with mitochondrial alterations and Bcl-2 protein imbalance.**

A) Images representing the iPSC-derived neurons before and after thapsigargin stimulation and B) their representative fluorescent traces. C) Measurement of the TG-evoked ER Ca<sup>2+</sup> release shows significantly higher ER Ca<sup>2+</sup> levels in C9-T2 and C9-7245 lines compared to OX3-9 (\*\*\**p*<0.001; \**p*<0.05; One-Way ANOVA with Dunnet's *post hoc* test); n=3 independent differentiations, m=150-200 neurons recorded per genotype. D) Immunoblotting shows elevated levels of GRP78/BiP in all patient lines and significantly upregulated in C9-7245-1 and C9-7245-3 (\**p*<0.05, One-Way ANOVA with Dunnet's *post hoc* test) (E). n=3 independent

differentiations, m=142-200 neurons recorded per genotype per differentiation. F) Immunoblotting for the anti-apoptotic Bcl-2 protein shows downregulation of levels in patient lines and G) significant reduction in C9-7245-1 and C9-7245-3 and C902-2 compared to control OX3-9, AH017-13 and OX1-61 ( $*p<0.05$ ,  $**p<0.01$ , One-Way ANOVA with Dunnett's *post hoc* test); n=3 independent differentiations (n=2 independent differentiations for AH017-13 and OX1-61), m=3-4 technical replicates. H) qPCR confirms downregulation of Bcl-X<sub>L</sub> and upregulation of D) BAK, specifically in the C9-7245 and C902-2 MN lines ( $*p<0.05$ ,  $**p<0.01$ ,  $***p<0.001$ , One-Way ANOVA with Dunnett's *post hoc* test); n=2 independent differentiations; m=3 technical replicates. J) Representative image of OX3-9 loaded with MitoTracker Red CMXRos and immunostained with Islet1 and Tuj1. K) Quantification of the mean fluorescence intensity of MitoTracker in Islet1<sup>+</sup>/Tuj1<sup>+</sup> neurons shows reduced mitochondrial membrane potential in all patient lines ( $***p<0.001$ , One-Way ANOVA with Dunnett's *post hoc* test); n=2 independent differentiations; m=17-31 neurons analyzed per genotype.

L) Electron micrographs show enlarged mitochondria and cristae in the patient lines C902-2 and C9-7245-3 compared to a healthy control (OX3-9). Arrows point to mitochondria. M) Quantification of EM images for mitochondrial alterations show 88% of C9orf72 neurons with swollen mitochondria ( $**p<0.01$ , ANOVA with Dunnett's *post hoc* test); N) Cytochrome *c* levels are elevated in all patient lines and significantly higher than healthy controls in C9-7245-1 ( $*p<0.05$ , One-Way ANOVA with Dunnett's *post hoc* test). n=2 independent differentiations, m=2 technical replicates each. O) Cortical neurons show elevated levels of the apoptotic marker cytochrome *c* in C902-10 ( $*p<0.05$ ,  $**p<0.01$ , One-Way ANOVA with Dunnett's

*post hoc* test). n=2 independent differentiations. Scale bars = 20  $\mu\text{m}$  (J), 1  $\mu\text{m}$  (L).  
Data are represented as average $\pm$ SEM.

**Figure 5: Cleaved caspase-3 is frequently detected in *C9orf72* MNs and CNs and reduced cell viability is found in *C9orf72* MNs and CNs.**

A,B) Immunostaining for cleaved caspase-3 shows increased frequency of apoptotic neurons in *C9orf72* MNs compared to control MNs ( $*p<0.05$ ,  $**p<0.01$ , One-Way ANOVA with Dunnett's *post hoc* test); n=3 independent differentiations; m=400 cells per genotype analyzed for each individual experiment; C) Reduced cell viability was detected in *C9orf72* MNs by propidium iodide (PI) staining ( $***p<0.001$ , One-Way ANOVA with Dunnett's *post hoc* test) n=2 independent differentiations; D,E) Immunostaining with cleaved caspase-3 shows significantly increased frequency of positive cortical neurons in C902-2 and C902-10 lines compared to 2 healthy control lines (OX3-9 and AH017-13) ( $*p<0.05$ ,  $***p<0.001$ , One-Way ANOVA with Dunnett's *post hoc* test). F) Reduced viability was detected by PI staining in the *C9orf72* patient ( $***p<0.001$ , One-Way ANOVA with Dunnett's *post hoc* test). n=3 independent differentiations. Data are represented as average $\pm$ SEM. Scale bar: 20  $\mu\text{m}$ .

**Figure 6: p62 is elevated in *C9orf72* neurons and PABP immunopositive stress granules form spontaneously in *C9orf72* neurons.**

A) Immunostaining for p62 in C9 MNs shows aggregates are formed under baseline conditions. B,C) Immunoblotting demonstrates high levels of p62 in C9-T2-6, C9-7245-3 and C902-2 MNs compared to controls under baseline conditions

(\* $p < 0.05$ , \*\* $p < 0.01$ , One-Way ANOVA with Dunnett's *post hoc* test);  $n=2$  independent differentiations,  $m=3$  technical replicates. D, E) p62 aggregates are detected more frequently in C9 CNs compared to controls (\*\* $p < 0.001$ , One-Way ANOVA with Dunnett's *post hoc* test). F,G) Immunoblotting shows elevated levels of LC3-II in both lines from the *C9orf72* patient C902 compared to controls (\*\* $p < 0.001$ , One-Way ANOVA with Dunnett's *post hoc* test); H) PABP<sup>+</sup> and TIA-1<sup>+</sup> stress granules are detected in all 5 C9 patient lines under baseline conditions and (I) PABP<sup>+</sup> MNs are significantly more frequent than in controls (\* $p < 0.05$ , \*\* $p < 0.01$ , One-Way ANOVA with Dunnett's *post hoc* test);  $n=3$  independent differentiations;  $m=387-918$  neurons analyzed per genotype. J, K) PABP<sup>+</sup> stress granules form in C9 CNs (\*\* $p < 0.001$ , One-Way ANOVA with Dunnett's *post hoc* test). Scale bars = 20  $\mu\text{m}$ ; scale bars in the bottom panels of H = 10  $\mu\text{m}$ . Data are represented as average  $\pm$  SEM.

### **Graphical abstract legend:**

This study presents a comprehensive characterization of cellular pathways that are contributing to cell death in iPSC-derived motor and cortical neurons from ALS/FTD patients carrying pathogenic hexanucleotide repeats in the *C9orf72* gene. The apoptotic pathway that we identified in the *C9orf72* neurons involves ER calcium elevation and stress, followed by mitochondrial alterations, the release of cytochrome *c* from the mitochondria and cleavage of caspase-3. Aggregates positive for p62 were detected in these neurons along with high frequency of stress granules, indicating altered proteostasis in the *C9orf72* motor and cortical neurons.