

BMJ Open

Understanding recruitment and retention in the NHS community pharmacy stop smoking service: perceptions of smoking cessation advisers

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2015-010921.R1
Article Type:	Research
Date Submitted by the Author:	20-Apr-2016
Complete List of Authors:	Sohanpal, Ratna; Queen Mary University of London, Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health Rivas, Carol; University of Southampton, Faculty of Health Sciences Steed, Liz; Queen Mary University of London, Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health MacNeill, Virginia; University of Oxford Kuan, Valerie; Queen Mary University of London, Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health Edwards, Elizabeth; Queen Mary University of London, Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health Griffiths, Chris; Queen Mary University of London, Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health Eldridge, Sandra; Queen Mary University of London, Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health Taylor, Stephanie; Queen Mary University of London, Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health Walton, Robert; Queen Mary University of London, Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health
Primary Subject Heading:	Health services research
Secondary Subject Heading:	Public health
Keywords:	Smoking cessation, Community Pharmacies, QUALITATIVE RESEARCH, Patient recruitment, Health Behaviour, Communication

SCHOLARONE™
Manuscripts

1
2
3 **Understanding recruitment and retention in the NHS community pharmacy stop smoking service:**
4 **perceptions of smoking cessation advisers**

5
6 Ratna Sohanpal¹

7
8 Carol Rivas²

9
10 Liz Steed¹

11
12 Virginia MacNeill³

13
14 Valerie Kuan¹

15
16 Elizabeth Edwards¹

17
18 Chris Griffiths¹

19
20 Sandra Eldridge¹

21
22 Stephanie Taylor¹

23
24 Robert Walton¹

25
26 ¹Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health, Barts and The
27 London School of Medicine and Dentistry, Queen Mary University of London, London, UK

28
29 ²Faculty of Health Sciences, University of Southampton, UK

30
31 ³University of Oxford, UK

32 **Corresponding author:**

33
34 Ratna Sohanpal

35
36 Research Fellow

37
38 Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health

39
40 Blizzard Institute

41
42 Queen Mary University of London

43
44 58 Turner Street, London

45
46 E1 2AB

47
48 Ph: +442078822492

49
50 Fax: +442078822552

51
52 Email: r.sohanpal@qmul.ac.uk

ABSTRACT

Objectives: To understand views of pharmacy advisers about smoker recruitment and retention in the National Health Service community pharmacy stop smoking programme.

Design: Thematic framework analysis of semi-structured, in-depth interviews applying the Theoretical Domains Framework and COM-B behaviour change model. We aimed to identify aspects of adviser behaviour that might be modified to increase numbers joining and completing the programme.

Participants: 25 Stop Smoking Advisers (13 pharmacists and 12 support staff).

Setting: 29 community pharmacies in three inner east London boroughs.

Results: Advisers had preconceived ideas about smokers likely to join or drop out and made judgements about smokers' readiness to quit. Actively recruiting smokers was accorded low priority due in part to perceived insufficient remuneration to the pharmacy and anticipated challenging interactions with smokers. Suggestions to improve smoker recruitment and retention included developing a more holistic and supportive approach using patient-centred communication. Training counter assistants was seen to be important as was flexibility to extend the programme duration to fit better with smokers needs.

Conclusions: Cessation advisers feel they lack the interpersonal skills necessary to engage well with smokers and help them to quit. Addressing advisers' behaviours about active engagement and follow-up of clients, together with regular skills training including staff not formally trained as cessation advisers could potentially boost numbers recruited and retained in the stop smoking programme. Adjustments to the pharmacy remuneration structure to incentivise recruitment and to allow personalisation of the programme for individual smokers should also be considered.

Keywords: Smoking cessation, Community Pharmacies, Qualitative research, Patient recruitment, Retention, Health Behaviour, Communication, Psychological Theory

Word count: 4,077

Strengths and weaknesses

- This is the first study to provide insight into smoker recruitment and retention in the NHS community pharmacy stop smoking programme from the adviser's perspective and therefore fills an important gap in knowledge.
- We used the Theoretical Domains Framework and COM-B behaviour change model as a framework to identify adviser attitudes and behaviours that could potentially be modified, thus providing a unique insight into smoker recruitment and retention in this clinical setting and facilitating development of interventions.
- In our analysis we achieved theoretical saturation, high inter-rater reliability and high agreement with a health psychologist about mapping thematic data onto constructs of the Theoretical Domains Framework.
- Whilst recruitment and retention are clearly two separate behaviours we analysed them together because we were interested in the combined result – namely increased throughput in the service. Future studies might profitably separate these behaviours to examine their unique motivations.
- Our findings will not be generally applicable, but could nevertheless be transferable to community pharmacies delivering the stop smoking programme in other socio-economically deprived communities across the UK. Certain findings such as low self-efficacy in consultation skills and the need to train all pharmacy staff to increase service throughput may well be transferrable to other healthcare systems, however this would need to be examined in the different settings.

INTRODUCTION

Improving public health through extending community pharmacy services has become a key part of UK National Health Service (NHS) operating strategy.[1-2] Smoking cessation is a public health priority[3] since smoking is the most important cause of premature morbidity and mortality worldwide.[4] In the UK, smoking cessation through pharmacy-led stop smoking services is now a key element of the Department of Health's tobacco control strategy.[2]

Community pharmacies are accessed by people both in health and illness, hence community pharmacy stop smoking advisers are well placed to reach large numbers of smokers.[5] Behavioural counselling/support, or a combination of nicotine replacement therapy and counselling by trained pharmacy staff are effective in helping smokers to quit.[2, 4, 6] However, the overall effectiveness of the pharmacy stop smoking programme does not only depend on the quit rates achieved but also on the number of smokers who participate and adhere to the programme.[7-8] In 2012-2013 of the estimated two million smokers in England,[9] only 149,034 smokers set a quit date in the pharmacy NHS stop smoking programme. Of these, 48% successfully quit at four weeks compared to the target of 70%.[2, 10] Quit rate was slightly lower than that achieved by the stop smoking services in primary care (50%) and lower than specialist stop smoking services (56%).[10] This apparent poor performance may result from differences in case-mix, however variations in staff training and environmental factors in pharmacies may make high cessation rates harder to achieve than in other settings. Nevertheless there are considerable disparities in quit rates between pharmacies suggesting potential for improvement.[2]

A recent review[6] suggested that pharmacists only target those they perceive to be ready to quit, thus reducing recruitment potential. Retention within the service is also poor, for example only 35% of the 4500 people who joined the stop smoking programme in one inner London borough, successfully quit (i.e. quit status biochemically verified by carbon monoxide testing at four weeks from the quit date)[11] and the remainder were lost to follow up.[12] Thus in addition to optimising quit rates, improving recruitment and retention to the stop smoking programme might raise overall numbers successfully giving up smoking.

In a systematic review of pharmacist views on delivery of public health services pharmacists recognised they should be more active in smoking cessation, however a number of barriers were suggested such as lack of time, staff and a designated space. Knowledge and skills were thought to be lacking and patients did not expect to receive health promotion advice from pharmacists. In addition there was reluctance to initiate conversations about health promotion because of fears of generating negative responses.[13]

1
2
3 No detailed investigation of stop smoking advisers' views on factors affecting recruitment
4 and retention in the NHS stop smoking service has been published to date. This is the first such UK
5 qualitative study aiming to: (1) understand stop smoking advisers' views relating to smoker
6 recruitment and retention in the NHS stop smoking programme in community pharmacies; (2)
7 identify factors that might be targeted in an intervention to increase smoker recruitment and
8 retention to maximise the effectiveness of the programme.
9
10

11
12 The results of this study will contribute to development of a complex intervention to
13 promote uptake and increase the effectiveness of the NHS pharmacy smoking cessation service. This
14 intervention will be evaluated in a cluster randomised trial in east London as part of the STOP
15 programme (Smoking Treatment Optimisation in Pharmacies).
16
17
18
19

20 21 **METHODS**

22 **Study design**

23
24 We conducted semi-structured, in-depth interviews with stop smoking advisers (comprising both
25 pharmacists and pharmacy support staff). For roles and responsibilities of staff and an outline of the
26 stop smoking programme see Box 1.
27
28

29 We used thematic framework analysis[14-15] applying the Theoretical Domains Framework
30 [16] within the COM-B model of behaviour change.[17] The Theoretical Domains Framework is
31 synthesised from multiple behaviour change theories and comprises 14 domains of theoretical
32 constructs that explain possible influences on behaviour.[16] Each domain fits within one of the
33 three components of the COM-B model - Capability Opportunity and Motivation.[17-18] For
34 example, 'Skills' in the Theoretical Domains Framework maps on to 'Capability' in the COM-B model,
35 'Professional Role and Identity' to 'Motivation', 'Environmental Context and Resources' to
36 'Opportunity'. [18] We used COM-B as a lens through which to view our data because it provides a
37 practical basis for designing interventions aimed at behaviour change, helping to identify the
38 behavioural target and the components of the behaviour system needing to be changed.[17]
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Box 1 Outline of the stop smoking training and the community pharmacy stop smoking programme

Training in smoking cessation[19-22]

The National Centre for Smoking Cessation and Training (NCSCT) in NHS England offers a range of training, assessment and certification programmes for both clinical and non-clinical health and social care workers to become more skilled in smoking cessation. The training programme is built around evidence-based behaviour change techniques that provide an understanding of the factors involved in smoking and smoking cessation.

The training includes:

- (1) Level 1 training or Very Brief Advice in Smoking Cessation training – this (online) training enables promotion of smoking cessation and can be undertaken as a minimum by health care professionals e.g. doctors, nurses, pharmacists including staff who advise people on how to quit smoking.
- (2) Level 2 training or Training and Assessment programme - this (online and face to face) training is for equipping health care professionals, who intend to become *stop smoking advisers*, with knowledge and skills to provide intensive one to one support in smoking cessation through delivery of the NHS Stop Smoking Programme (see below).

Delivery of the NHS Stop Smoking Programme in community pharmacy[23-24]

The pharmacy owners (contractors) are contracted by NHS England clinical commissioning groups and local authorities to deliver public health services including the NHS stop smoking services to meet the needs of the local population. The stop smoking services cover the range of activities from the proactive promotion of smoking cessation through to provision of the NHS Stop Smoking Programme.

In community pharmacy, the stop smoking programme can be delivered by stop smoking advisers who might be:

- Pharmacists: qualified experts in the use of medicines for treatment of disease. They offer a range of services such as Medicines Use Review and Prescription Intervention Service; New Medicine Service; Appliance Use Review Service; Public health services e.g. Stop smoking services, NHS Health Checks.
- Pharmacy support staff such as medicines counter assistant, dispenser, dispensing assistant, pharmacy technician and accredited checking technician who support the pharmacist in the selling of medicines and delivery of services.

Content of the NHS Stop Smoking Programme[2, 20]

The content includes delivery of behavioural support together with pharmaceutical treatments comprising Nicotine Replacement Therapy (NRT) e.g. patches, prescribed medication e.g. varenicline (Champix) or a combination of NRT and prescribed medication to help a smoker quit smoking. The duration of the programme, dependent upon commissioners, ranges between 6-12 weeks with quit status to be recorded at 4 weeks, verified by carbon monoxide (CO) monitoring and sent to NHS England for reporting of statistics, monitoring and commissioning purposes.

- Week 1
 - Introduction and set planned Quit Date (approx. ½ hour meeting)
 - Stop Smoking Adviser explains programme process
 - Gives service user/smoker information about 3 types of medication available (NRT, Champix (varenicline) tablets and Zyban (bupropion) tablets)
 - Discuss which is most suitable for service user
 - Adviser takes and records service user carbon monoxide (CO) reading
- Week 2-4
 - Brief meetings to check progress (each approx. ¼ hour and informal). Quit status CO monitoring is recorded at week 4 for NHS England statistics.[2]
- Week 5/6
 - Longer meeting to discuss motivations and techniques to avoid relapse.
- Week 6/7-12
 - Programme available to service user but the adviser is not obliged to follow up.

Setting and participants

The study was conducted in three inner east London boroughs: Tower Hamlets; Newham; City and Hackney. These boroughs include south Asian and African/Caribbean communities with high levels of tobacco use and persistent health problems linked to social and economic inequalities.[12, 25-26] Smoking prevalence in these deprived boroughs is close to the UK national average (20%), or higher (Newham 21%, 23% in Hackney, 37% in Tower Hamlets).[12, 25-26] Purposive sampling was used to obtain a diverse range of views.[27] We selected stop smoking advisers who differed by gender and duration of being an adviser. Independent community pharmacies were sent a letter and information sheet and contacted by telephone to arrange a face-to-face meeting in the pharmacy. Within pharmacies the pharmacist usually suggested the adviser (usually themselves or a member of other pharmacy support staff i.e. stop smoking adviser for interview). A member of the research team (CR) or a research assistant obtained written informed consent.

Data collection

Individual interviews were conducted from January to June 2014 by an experienced female qualitative researcher (RS, VM) using an interview schedule (Box 2). The interviewer was not known to the study participants and was not involved in recruiting participants to the qualitative study. Advisers were informed that study aim was to improve programme recruitment, retention and quit rates. Interviews took place in the consultation room of the community pharmacy and lasted 30-60 minutes. Recruitment and interviews continued until data saturation was achieved. All interviews were audio-recorded and fully transcribed. Nvivo10 was used for organisation of data and to facilitate analysis.

Box 2 Interview schedule

THE CONSULTATIONS

1. What helps/what do you do that makes smokers join the service (recruit)?
2. What helps/do you do that makes them keep coming back (retain)?

PROCESS

3. Can you describe the main issues involved in delivering the Stop Smoking Programme?
4. What prevents people from joining the stop smoking service?
5. What are the characteristics of people that do not join the service/ who join the service and quit/ who join the service and do not quit?
6. We know that many advisers select smokers most motivated to quit. What would make you take on the less motivated/interested?*

TRAINING

7. What would you like that is different to training you have had so far? (that can help more smokers to join the stop smoking service)

* This question was added after analysis had begun as an issue emerging from the data.

Data analysis and interpretation

A thematic framework analysis[14-15] was conducted. This method starts deductively with *a priori* codes from the study aims and objectives however, subsequent analysis is inductive and grounded in the accounts of the participants.[14] Researcher (RS) read and re-read the transcripts for familiarisation and data immersion. A thematic framework was formed following line-by-line coding and comprised useful memos or descriptive statements to develop categories. Comparisons were made within and between transcripts based on the thematic framework. The data were then lifted and charted across the thematic categories. This process was carried out initially with a few transcripts and discussed with the study team (CR, RW, LS) to assess reliability and agreement between two coders and to check validity of the analysis. Inter-rater reliability was 90.9% (kappa agreement) between the two coders (RS, CR) on 20% of the transcripts. Subsequently, the emergent themes were mapped (by RS) onto behavioural constructs of the TDF and the COM-B model, where applicable, to generate analytical themes.

The mapping was discussed and agreed with a health psychologist (LS) to ensure the mapped thematic data fitted with the domain definition and its content (Supplementary file 1). This process generated analytical themes or key domains influencing the engagement and the retention behaviour of advisers and helped us to identify domains that might be targeted to optimise adviser behaviour. Emergent themes are exemplified below with direct participant quotations.

RESULTS

Twenty-five interviews were conducted following approaches to 44 advisers in 29 community pharmacies. Reasons for non-participation were lack of interest (n=14) or unavailable/no answer (n=5). The participants were drawn from 15 pharmacies with six pharmacies contributing two or three interviewees. Over half of the advisers were of Asian ethnicity (56%) and a small proportion (20%) of pharmacies provided a multi-lingual service. Characteristics of the participants are shown in Table 1. All participants were smoking cessation advisers trained to level 2.

In the analysis the emergent themes mapped on to five of 14 TDF constructs with one independent theme. Figure 1 illustrates this using COM-B as headings and TDF as subheadings (*italics*). The independent theme and its potential relation to COM-B is highlighted with dashed lines. See supplementary file 1 for full definition of the COM-B and TDF constructs. In the text that follows the higher level heading is from the COM-B model and the subheadings represent relevant domains from the Theoretical Domains Framework that explain two separate adviser behaviours, recruitment and retention.

Table 1 Characteristics of study participants

Study ID*	Pharmacist or support staff	Gender	Duration being stop smoking adviser
S11A101	Pharmacist	Female	Not given
S19A182	Pharmacist	Female	3 years
S08A071	Pharmacist	Male	Not given
S13A121	Pharmacist	Male	10 years
S29A281	Pharmacist	Male	4 years
S04A031	Pharmacist	Male	Not given
S09A081	Pharmacist	Male	10 years
S23A221	Pharmacist	Male	Not given
S25A242	Pharmacist	Male	7 years
S26A253	Pharmacist	Male	Not given
S03A021	Pharmacist	Male	Not given
S05A041	Pharmacist	Male	5-6 years
S11A103	Support staff	Female	9 years
S02A012	Support staff	Female	Not given
S25A241	Support staff	Female	1 year
S26A251	Support staff	Female	7 years
S09A082	Support staff	Female	5 years
S25A243	Support staff	Female	Not given
S26A252	Support staff	Male	4 years
S01A001	Support staff	Male	Not given
S04A032	Support staff	Male	6 years
S11A102	Support staff	Male	4 years
S19A181	Support staff	Male	12 years
S24A231	Support staff	Male	Not given

* This code is used to identify the source of quotations in the text.

Motivation

(1) Professional role and identity

Many participants felt proud of their work and were satisfied that through their role as advisers they were able to help people quit smoking, which in turn helped with the health and well-being of the wider community.

I mean, you are always happy that you've helped someone do something. And there are many times that, this guy (client who has quit) will always pop in. ...and say "Oh, I'm just passing and I thought let me just say hi to you!" And there are many times ... This lady, it's nice. I mean, why would this lady bring her son over for me to advise? It's like they appreciate it and you have that feeling that, oh, you've also managed to help someone and it's great to do that. (Study ID S08A071, Pharmacist)

Identification and engagement of smokers into the programme

Advisers identified potential clients through: GP referrals; recommendations from people who had quit; opportunistically during medicine review or while conducting risk assessment; and when people bought products from the pharmacy, picked up regular prescriptions or bought over the counter medication. Despite this, many advisers selected and recruited only those smokers who mentioned their readiness to quit because they thought that these smokers were less likely to drop out. A few said they only recruited those who specifically asked for smoking cessation advice and were not interested in actively recruiting smokers to the NHS SSP.

She's like I'm not in the right frame of mind but I really want to give up smoking. So in that way we just tell them look, ... So we told her and with smoking you have to make sure that they're in the right frame of mind otherwise there's no point them joining the programme.

Otherwise they'll join and then quit after a week and there's no point. (S24A231, Pharmacy support staff)

I would say the recruitment. I mean it's a bit hard, most of it has to be walking and...yeah. Or maybe they see the poster. But we're not doing that much. (S19A182, Pharmacist)

Some participants said that recruitment into the programme was low priority. Advisers wanted to prioritise patients with long-term conditions or those that they thought had life-threatening conditions. One adviser felt giving up smoking was the responsibility of the smoker.

...so smoking is more or less a very very little part of our actual work here. We actually have people who have life threatening conditions, you can be dead tomorrow. So that's of a

1
2
3 *greater priority than smoking. Smoking is by choice, you want to give up, you don't want to*
4 *give up. (A01A001, Pharmacy support staff)*
5
6

7
8 Several participants also said they believed that all the advice, support and products they provided
9 could only help a smoker quit if the smoker was mentally prepared and had the willpower to follow
10 through the advice and make the necessary lifestyle/environmental changes conducive to stopping
11 smoking. Advisers felt they had most success with this type of client.
12

13
14 *... we're pretty straight with them (potential clients), ... if they're motivated to quit then*
15 *they've got a better chance of succeeding, things like ...NRT Champed (sic) only helps to a*
16 *certain extent ... and if they don't have the willpower it's not going to work... (S09A082,*
17 *Pharmacist)*
18
19
20

21 22 Encouragement and motivation to ensure attendance and adherence

23
24 Some participants stated that encouraging clients to continue attending the programme sessions
25 was important. One adviser mentioned that to prevent dropouts and indeed when engaging with
26 clients who had relapsed, identifying the reasons for stopping smoking and relapse were important.
27

28
29 *Oh I think you are always going to get that (people dropping out). ...I mean you are talking*
30 *about 1 in 2 not getting through so you just have to keep motivating and telling them you*
31 *know I mean you've been smoking for 20 or 30 years, you are not going to give up in 6 to 12*
32 *weeks, it's a long term process. And I think if you have like help, if you have re-started again,*
33 *again the important thing is to find the reasons why because I think if you can get to the*
34 *bottom of that and work on that rather than just giving them NRT and thinking, you know so*
35 *really that is most important. The NRT helps but you need to know why you want to stop*
36 *smoking. (S03A021, Pharmacist)*
37
38
39
40
41
42

43
44 A few participants did not see following up clients who missed sessions as part of their role. Some
45 said that they simply accepted it if smokers mentioned they were finding it difficult to not smoke
46 when they spoke to them by telephone or in person.
47

48
49 *Two persons I've seen, they can't manage (to quit), that's it. Sometimes if they come for their*
50 *medicines to pick up or anything, they can say, "Oh, sorry, I didn't come back, because I can't*
51 *manage at the moment, I'm not all right." So we say, "OK, whenever you feel OK."*
52 *(S02A012, Pharmacy support staff)*
53

54
55 *...when they don't come back it's like well, did you give them the wrong products? They didn't*
56 *suit them, they could've phoned us, we could've spoken to them, we could've swapped it. We*
57
58

1
2
3 *are pretty flexible. But if they don't come back to us there's no way of knowing, is there?*
4 (S25A241, Pharmacy support staff)
5
6

7
8 Some advisers acknowledged that more smoker engagement would result in higher quit rates. The
9 following were suggested to boost service recruitment and retention: identify potential smokers for
10 the service e.g. when they bought cough syrup; engage, inform, encourage and recommend the
11 programme to smokers e.g. through use of poster adverts; explain how the programme and adviser
12 involvement could help with quitting; place a holistic focus on the person.
13
14

15
16 *So if you're focused on the client rather than the process you're going to get a bigger*
17 *outcome rather than saying this is a tick form, I've got to tick this, I've got to fill this in, I've*
18 *got to fill that in and oh, I've missed out here, we've got to do a CO reading sort of thing.*
19 (S26A253, Pharmacist)
20
21
22

23 24 **(2) Reinforcement**

25 Remuneration

26
27 This process of remuneration was acknowledged as an issue by the adviser participants who were a
28 mix of owner pharmacists, employed pharmacists and other pharmacy support staff. Among the
29 owner pharmacists, a few said that the current remuneration for quitters was reasonable, while
30 some others felt that the remuneration did not take into account the time they spent with smokers
31 encouraging them to join and adhere to the service.
32
33

34
35 One owner pharmacist mentioned that the current payment system was a disincentive to spend
36 time with smokers, and needed revisiting by commissioners; payment based on quitters might
37 influence which smokers staff are willing to support.
38
39

40 *And things, we get penalised for non-quitters which are not really our fault.*

41
42 *Interviewer: ...How do you mean penalised?*

43 *So currently we may spend three or four hours actually going through these patients, and*
44 *because they don't quit we don't get the full £40 or £35. We end up with about £20. That's*
45 *the disincentive. There are certain health authorities now are saying that if a patient doesn't*
46 *quit there's no fees. So there is a lot of disincentives for a pharmacist to say look, I only get a*
47 *50 per cent success rate, I'm not going to spend three or four hours at a time and get nothing*
48 *for it. So they're (pharmacies are) not going to take on new clients and the way that the*
49 *health authorities and CCGs think about these programmes needs to change. (S26A253,*
50 *Pharmacist)*
51
52
53
54
55
56
57

1
2
3 Another owner pharmacist stated that perhaps smokers should be incentivised to attend their stop
4 smoking appointments as it could mean saving money for the NHS in the long run and some
5 pharmacy support staff suggested providing incentives/rewards to their clients that could be given
6 at programme completion.
7
8

9
10 *And I think if you can perhaps issue vouchers, but issue vouchers that are redeemable at the*
11 *end of the process, yeah? So it makes no sense for them to come to two or three*
12 *consultations, accumulate vouchers and then miss the fourth one, yeah, you know? So I*
13 *think there should be vouchers that can be redeemed for cash at the end of it, because most*
14 *people don't want to be restricted as to what they can do with a voucher or cash or anything.*
15 *That is one thing I think....at the end of the day, we need to put it in the balance; I mean, the*
16 *government feels that getting people to stop smoking leads to a greater saving on the NHS,*
17 *so it's something they should look at. (S29A281, Pharmacist)*
18
19
20
21
22

23
24 Several of the employed pharmacists and support staff acknowledged that delivery of the
25 programme was part of their job, although a few highlighted that if their pharmacy were to receive
26 an increase in the current remuneration for quitters then they might be allowed to spend more time
27 engaging with and following up clients to continue and complete the programme.
28
29

30
31 *I think so as well, because it (current remuneration) is a kind of barrier. You do encourage*
32 *them (smoker) to stop smoking, but then three weeks you spend with them, although they*
33 *don't stop smoking ... And for one engagement they give you (payment), but after that, if*
34 *they keep on giving us the money we (can) encourage more people, ... So if they encourage*
35 *me, I mean the pharmacy, by providing with more money or anything, no barriers regarding*
36 *money, then we can do more, much more. (S26A251, Pharmacy support staff)*
37
38
39
40
41

42 **(3) Beliefs about capabilities**

43 Interaction with smokers

44
45 Some advisers mentioned that engaging with smokers and motivating clients who enjoyed smoking
46 or who were not interested in quitting was a challenge to programme recruitment and retention. In
47 addition, one participant was unsure about the protocol to follow when a client smoked a cigarette
48 at the end of 12 week programme.
49
50
51

52 *It's more of a challenge when they say oh, but I love smoking, I love the taste of smoking.*
53 *Then it's a bit of a challenge because they've already got in their mind that...there's like a*
54 *mental block. Whatever you're going to say to me, I don't really care sort of thing. ..*
55 *(S24A231, Pharmacy support staff)*
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In contrast, a few participants felt confident to handle some of the above mentioned issues by talking about the harmful effects of smoking, showing the tar jar to smokers and suggesting to smokers they were available to support them when they are ready to quit.

Yeah, I mean there's no harm in recommending it (the service). You recommend it to everyone. I don't see anything bad about that. It's just whether they'll actually sign up or not. ... The only way we can deal with that is through awareness, like the tar outside and posters and our TV advert. That's the only way you're going to get those kind of people. (S26A252, Pharmacy support staff)

Capability

(4) Skills

Skills training

Several suggestions were given to improve the skill set of advisors (including counter assistants) to help with recruitment and retention in the stop smoking programme, such as reinforcement of motivational and communication skills, including how to approach people about smoking without causing offence, how to be supportive. Most suggestions were features of person-centred care which all health care professionals are expected to follow.[28] These were: how to personalise advice, how to elicit from smokers which things had worked and which did not work, how to work with smokers to identify from them what targets they would like to achieve and how they would achieve them.

Pharmacies are very prescriptive and I think when you're asking people to give up smoking you have to think about how they need to change their behaviour, and I think it's those ... skills that you need to apply, and that is very crucial and very important. Getting it out there. What is it they're able to do? You know what they need to do. It's only when they come up with solutions themselves they're more likely to follow it through. (S01A001, Pharmacy support staff)

In addition, a few participants stated there was need for regular training, some others suggested that it might be useful to include in any future training items such as visual scenarios, role plays or mock interviews to learn and practice their skills. Two participants also wanted to see experts in action with a client to help them learn and improve their skills.

I think sometimes you get a bit complacent. When you're doing it and you think everything's fine you give them the NRTs, tell them to come back next week if they have any problems.

1
2
3 *But they might actually have a bit of problems that they want to tell you but they can't for*
4 *whatever reason or they feel like if they tell you that they've failed then... So trying to get the*
5 *best out of their patients, so someone that could come in like a specialist to actually help,*
6 *that would be good. Because you always learn something when you're shadowing someone,*
7 *always. (S24A231, Pharmacy support staff)*
8
9
10

11 Opportunity

12 (5) Environmental context and resources

13
14
15
16 The study participants mentioned a number of structural/organisational challenges that affected
17 smoker recruitment and retention in the programme:

18 Lack of time because of delivery of other services

19
20 Participants mentioned that they forgot to remind clients about their appointment or they were
21 unable to keep tabs on clients' programme adherence. A few pharmacists, on account of being busy,
22 said they were reluctant to recruit smokers, if they were unable to provide details about the
23 programme and the procedure involved; this seemed to affect recruitment.
24
25
26

27 *Especially someone who says, "Can I join the scheme now?" and sometimes it's not possible;*
28 *when I'm so busy here, I can't leave the place and come in here. There's no way I can leave*
29 *it, so I have to give them an appointment. But sometimes people are not happy with that;*
30 *it's like they really want to join now! You get the point, they really want to join, so if you*
31 *can't do it for me now, then I can't come, so it's an issue. (S08A071, Pharmacist)*
32
33
34
35
36

37 In contrast, one pharmacy support staff participant, to avoid losing clients, started engaging people
38 on the same day of their pharmacy visit instead of giving them an appointment for a later date; this
39 strategy helped them to improve recruitment into the stop smoking programme. Another staff
40 member wanted a reminder system to enable client follow-up to help with retention.
41
42

43 *...if somebody comes in now, generally we do it straight away. That's how we found it works*
44 *better. If you give them a date they don't turn up because within this time they've got a*
45 *different commitment. (S11A102, Pharmacy support staff)*
46
47
48

49
50 To make time for smoker engagement, one pharmacist mentioned that having counter assistants to
51 help with the recruitment process or more staff to share with their dispensing workload could lead
52 them to engage and offer the stop smoking programme more frequently to smokers.
53
54
55
56
57

Inflexibility of the programme

Several participants felt that the duration of the programme was insufficient for some clients to quit and remain abstinent. One participant suggested that clients who relapsed going through the programme should be allowed to continue with the programme instead of asking them to re-join the programme, normally after a few weeks. In contrast, another participant mentioned that allowing dropouts to re-join the programme as soon as they were ready instead of using the “after six months” rule was not helpful in improving quit rates.

Other issues I've had have been people who kept coming back, like two or three times a year, and it's obvious that they are not serious, or I don't know. In the past, with the NRT, they used to say they can't do it again until after six months, but now as often as they want. But then with the Champix, I got people coming two or three times a year: stop/start, stop/start.
(S11A103, Pharmacist)

Working with budget cuts

The facility to offer two treatments concurrently to clients had been stopped due to budget cuts despite evidence of its effectiveness. In one of the participating boroughs the varenicline (Champix) license for pharmacists to prescribe had been taken away; one participant stated that this had affected recruitment into the programme.

Champix, the tablets. We haven't got a PGD (patient group direction) for it right now, so I had one person yesterday, she was a bit upset, she was like oh, I want the tablets, I want the tablets. And we don't have the licence for it. I think the whole of Newham, all the pharmacies...I think hopefully we'll be getting it back soon enough, ... So that's a problem, we've lost a few people... but we could've had those people as clients. So that's one thing, hopefully we'll try and get that back. If the providers can help us with that then that would be great. (S24A231, Pharmacy support staff)

Failing to take advantage of national campaigns

One participant felt that the ‘Stoptober’ campaign helped with smoker recruitment but not retention because they perceived smokers were not really committed to giving up smoking and only joined because of the advertisement. Some other participants felt that the No Smoking Day campaign went by too quickly to enable smoker recruitment.

Having it (No Smoking Day) spread over a week means you can maybe put up a lot more balloons up, you can put a lot more signs up, you can market it better, it gives you a longer time to market it and it will give you that whole week to engage with people. Because

1
2
3 *sometimes you might only see them once a day, but if you've missed that*
4 *opportunity...tomorrow's not a non-smoking day. Oh, I'm not doing it then. (S25A242,*
5 *Pharmacist)*
6
7

8
9
10 Other organisational factors such as inadequate stock of nicotine replacement products to meet
11 client demand, having only one consultation room that was often occupied were mentioned as
12 restrictions on smoker recruitment by two participants.
13

14 *...the room that I'm using is constantly being used up. I've got this one room. You know? And*
15 *things like that again – having resources [unclear] how to manage that. (S01A001, Pharmacy*
16 *support staff)*
17
18

19
20
21 **(6) Advisers' perceived characteristics of service 'joiners', 'non-joiners' and 'dropouts'**

22 Advisers' characterised smokers as those likely to join, not join or drop out of the stop smoking
23 programme. This categorisation of smokers then affected how likely the adviser would be to recruit
24 them into the stop smoking programme.
25

26
27 Eleven participants stated that 'Joiners' (including those who relapsed and wanted to quit again)
28 were those who were motivated, willing or determined to give up smoking to improve health/quality
29 of life, and were clear of their reasons for wanting to give up.
30
31

32 *Yeah, always ask them as well what's in your mind. Why do you want to do this (quit*
33 *smoking)?... So once they have that (reasons) in mind I will tell them okay, so have that in*
34 *mind, so even when the cravings come just hang onto that, because at least you have*
35 *something you want to achieve because of that. ... in fact, if you don't have it, for me you are*
36 *more likely to fail... (S19A182, Pharmacist)*
37
38
39
40

41
42 'Non-joiners' were characterised by eight participants as those who did not suffer from any health
43 related problems and hence were not ready or motivated to quit, they were either not interested or
44 in denial, thinking that they would not suffer any health-related consequences perhaps because
45 people they knew had been smoking for years and their health seemed unaffected. In contrast,
46 smokers living with a long-term condition such as cancer felt that it was too late for them to change
47 their behaviour as their illness progressed.
48
49
50

51 *Because like I said, in many years, I've seen so many patient ..., when they're smoking, end up*
52 *with a cancer. He was still coming to see me, went to the doctors, diagnosed cancer, came in*
53 *and said I'm not doing it (stopping smoking). Why? Because I've got cancer, what the hell,*
54 *I'm going to die. (S19A181, Pharmacy support staff)*
55
56
57
58

59
60
18

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

'Dropouts,' according to seven participants, were those who were fighting more than one addiction or going through some personal crisis and therefore were not ready to give up smoking. Such people, the participants suggested, might be easily tempted into smoking by family/friends who smoke, or might not perceive any benefits or might not be interested in quitting.

He was determined (to quit) because he got it from the GP, the letter, so I thought yeah, he's going to give up smoking. But he just...the first week I supply and then the second week I didn't see him there. Called him and he said no, I can't give up, it's not holding me now. So he wasn't ready really, I don't think he was ready. ... (S04A032, Pharmacy support staff)

DISCUSSION

Statement of principal findings

Recruitment and retention of smokers in the NHS stop smoking programme may be influenced by advisers' preconceived ideas about smoker types likely to join, not join or drop out of the programme. This early categorisation of smokers influenced perceived readiness to quit. Active smoker recruitment was often a low priority, partly because advisers considered the remuneration that the pharmacy received for each quitter was insufficient to justify use of time and because they anticipated challenging interactions with some smokers. There were also perceived structural/organisational challenges involving programme delivery.

To improve the programme, advisers suggested that they should adopt a more holistic and supportive approach. They further suggested that strengthening their belief in the importance of engagement with both new and relapsed clients would help to improve uptake of the programme and retention of smokers within the service. An increase in pharmacy remuneration for quitters was also thought to be beneficial. Advisers would welcome improvement in structural and organisational delivery of the programme and more regular training in person-centred communication including for counter assistants.

Comparison to other studies

The Theoretical Domains Framework and COM-B models have been used previously in studies of dental health professionals[29-30] and midwives.[31] A barrier to delivering smoking cessation services in these clinical settings was that healthcare workers did not consider smoking cessation part of their primary role. In contrast, all participants in our study were trained stop smoking advisers and wholeheartedly embraced the smoking cessation role. Nonetheless some advisers identified the need for more training, particularly to build capability in the recruitment

1
2
3 process, to maximise engagement with smokers and to bolster their communication skills. This need
4 for training was also felt by dental health professionals and midwives. The fear of negative patient
5 response, inadequate staffing, and lack of confidence in delivery of health behaviour advice which
6 we identified in our study has also been reported in other settings.[13, 32-33]
7
8

9
10 In line with the recent review on the effectiveness of pharmacy smoking cessation
11 interventions,[6] we found that many advisers were mostly accepting smokers who wanted to quit
12 because they did not want to risk dropouts. The National Centre for Smoking Cessation and Training
13 also recommends that if the smoker is not ready to make a serious quitting attempt, the adviser
14 should provide the programme's contact details and ask them to get in touch when they are
15 ready.[34] Another explanation for selection of those likely to quit may be pressure from service
16 targets.[35]
17
18
19

20
21 The perceived opportunity cost of engaging with smokers who did not express a desire to
22 quit or who fitted stereotypes that advisers associated with failure to quit, appeared to affect
23 adversely the recruitment of smokers into the service. Hoving[36] compared 'active' recruitment
24 (i.e. "asking each individual's smoking status and inviting smokers to participate in the service") and
25 passive (i.e. "leaving participation up to the smoker's initiative") in a community pharmacy setting.
26 The former approach was suggested as a way of increasing the total number of quitters. It has been
27 suggested elsewhere that passive recruitment strategies might reduce the likelihood of a quit
28 attempt.[37]
29
30
31
32
33

34 In addition to lack of time for health promotion which has been reported previously,[13, 32,
35 33, 38] problems with delivery of medicines and promotional material, linked to national stop
36 smoking campaigns, were perceived by advisers to affect smoker recruitment and retention.
37 Addressing these issues is crucial to facilitate adviser engagement behaviour,[39-40] and could add
38 to the success of national campaigns in increasing the number of people attempting to quit and
39 permanently quitting.[39, 41-42]
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Implications for clinical practice and policy makers

Advisers' perceptions of types of smokers who are likely not to join or to drop out of the programme and some advisers' experience of challenging interactions with smokers have been reported previously[33] including in socially deprived areas.[43-44] However, the community pharmacy stop smoking services in east London are effective in reaching socio-economically deprived communities[35] and the smokers in these communities are just as motivated to quit smoking as smokers in more affluent areas,[35, 45] and to want help with quitting.[46] Smokers' stage of change[47] and economic disadvantage should not preclude the use of active recruitment[8, 35, 45] and retention strategies.

Recent changes to the pharmacy smoking cessation services in one of the participating boroughs meant that pharmacists were no longer able to recommend co-therapy (nicotine patch plus gum/lozenge/inhalator) or varenicline to clients which advisers considered a problem. It seems likely that restricting access to effective treatment and thus compromising patient choice could adversely affect engagement of smokers and quit rates. In addition, several advisers felt the duration of the stop smoking programme was not long enough for some smokers to quit and to remain abstinent and suggested having flexibility in the programme to allow continuity of care, where appropriate to suit clients' needs, for example after relapse.

The remuneration for programme delivery is determined by commissioners on the basis of clear criteria e.g. the time and duration of intervention and treatment provided, carbon monoxide (CO) monitoring and data reporting. The payment is made to the pharmacy owner i.e. a contractor who may be a pharmacist sole trader, a partner with other pharmacists or who may appoint a superintendent pharmacist[2, 48] No payment is made to employed pharmacists or to pharmacy support staff. The current payment system for pharmacies appears to provide a disincentive to spend time with smokers - and this has been acknowledged in the community pharmacy stop smoking services guidance.[2]

The pharmacy cessation advisers derived professional satisfaction in helping individual smokers to quit and also saw their role in smoking cessation as benefitting the whole community. Advisers thought about wider aspects of the scheme and took a professional interest in how the scheme might develop – for example suggesting a voucher scheme and tailoring the duration of the smoking cessation programme to individual smokers. This source of intrinsic motivation may distinguish cessation advisers from other healthcare professionals and could be drawn upon further in pharmacy adviser training programmes.

Implications for future research

More research is needed with a broader range of participants including female advisers and advisers of other ethnicities delivering the programme in deprived and non-deprived areas. In addition, studies to understand in more detail the effects of the remuneration structure for pharmacists and support staff is warranted. We took recruitment and retention together in the analysis since we were interested in the joint outcome of these two behaviours, which leads to an increase in the total number of people completing the smoking cessation programme. Whilst this seems justified given the purpose of our work this is a potential limitation of our study since clearly the two behaviours may have different antecedents. Future work could usefully examine recruitment and retention separately leading to individual insights into these behaviours and how they might be modified to strengthen the NHS smoking cessation service.

Acknowledgments The authors thank participants and their community pharmacy. This paper presents independent research funded by the National Institute for Health Research (NIHR) under its Programme Grants of Applied Research Programme. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Contributors RW conceived the study. RS was involved in the data collection, data analysis and interpretation and writing the manuscript. VM was involved in the data collection. CR and LS were involved in the data analysis and interpretation. CR, VM, LS, EE, VK, CG, ST, SE and RW were involved in revising the article critically for important intellectual content. RW is the guarantor.

Competing interests None.

Funding: NIHR Programme grant RP-PG-0609-10181

Participant consent Written informed consent obtained.

Ethics approval Ethical approval for the study was obtained from the NRES Committee South Central - Berkshire B (reference number: 13/SC/0189).

Provenance and peer review Not commissioned; externally peer reviewed

1
2
3 **Data sharing statement** This study analyses qualitative data and the advisors did not consent to
4 have their full transcripts made publicly available. The advisors consented for their data to be stored
5 for five years, as part of the 5-year programme grant project and to be used in this or other studies
6 directly linked to the 5-year project. Supporting excerpts from the raw data (quotes from participant
7 interviews) are available with the text of the paper. Full transcripts of the interviews (with
8 identifying information removed) will be available, following completion of the project, on request
9 from the study guarantor, Robert Walton, r.walton@qmul.ac.uk
10
11
12
13

14
15
16 **Open Access** This is an Open Access article distributed in accordance with the terms of the Creative
17 Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build
18 upon this work, for commercial use, provided the original work is properly cited. See:
19 <http://creativecommons.org/licenses/by/4.0/>
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. NHS England. Improving health and patient care through community pharmacy - a call to action 2013. <http://www.england.nhs.uk/wp-content/uploads/2013/12/community-pharmacy-cta.pdf> (accessed 24 Feb 2015)
2. NHS Employers. *Pharmacy-based stop smoking services: optimising commissioning*. London: NHS Employers, 2009.
3. World Health Organization. *Tobacco Free Initiative*. 2015. http://www.who.int/tobacco/health_priority/en/ (accessed 6 Feb 2015)
4. Brose LS, West R, Michie S, et al. Changes in success rates of smoking cessation treatment associated with of a national evidence-based training programme. *Prev Med* 2014;69:1-4.
5. Prescribing and Primary care Health and Social Care Information Centre. *General Pharmaceutical Services England 2004-05 to 2013-14* Leeds 2014.
6. Mdege ND, Chindove S. Effectiveness of tobacco use cessation interventions delivered by pharmacy personnel : A systematic review. *Res Social Adm Pharm* 2014;10(1):21-44.
7. McDonald PW. Population-based recruitment for quit-smoking programs: an analytic review of communication variables. *Prev Med* 1999;28:545-57
8. Szklo AS. Review of strategies to recruit smokers for smoking cessation: a population impact perspective. *Cad Saude Publica* 2008;24:S621-S34.
9. Action on smoking and health. Facts at a glance. Smoking statistics. 2015. www.ash.org.uk/files/documents/ASH_93.pdf (accessed 24 Feb 2015)
10. Lifestyle statistics Health and Social Care Information Centre. *Statistics on NHS Stop Smoking Services: England, April 2012 – March 2013*, 2013.
11. Department of Health. *Stop Smoking Service Monitoring and Guidance Update 2012/13*. London, 2012.
12. Bevan P. Annual Report of the Director of Public Health – Health at the Heart of the Community. 2014. <http://democracy.cityoflondon.gov.uk/documents/s35538/DPH%20report%20to%20Health%20Social%20Care%20Scrutiny%20sub%20committee.pdf> (accessed 24 Feb 2015)
13. Eades C, Ferguson J, O'Carroll R. Public health in community pharmacy: A systematic review of pharmacist and consumer views. *BMC Public Health* 2011;11:582.
14. Pope C, Ziebland S, Mays N. Qualitative research in health care. Analysing qualitative data. *BMJ* 2000;320:114-6.
15. Rabiee F. Focus-group interview and data analysis. *Proc Nutr Soc* 2004;63:655-60.

- 1
2
3 16. Cane J, O'Connor D, Michie S. Validation of the theoretical domains framework for use in
4 behaviour change and implementation research. *Implement Sci* 2012;7:37.
5
6 17. Michie S, van Stralen MM, West R. The behaviour change wheel: A new method for
7 characterising and designing behaviour change interventions. *Implement Sci* 2011;6:42.
8
9 18. Michie S, Atkins L, West R. The Behaviour Change Wheel. A Guide to Designing Interventions.
10 London: Silverback Publishing, 2014.
11
12 19. National Centre for Smoking Cessation and Training. http://www.ncsct.co.uk/pub_training.php
13 (accessed 7 Aug 2015)
14
15 20. National Institute for Health and Care Excellence. *Training in smoking prevention and cessation*.
16 NICE, 2015.
17
18 21. National Pharmacy Association. <http://www.npa.co.uk/> (accessed 7 Aug 2015)
19
20 22. Robinson F. *Smoking cessation: Training providers*. 2014. [https://www.pcrs-uk.org/training-](https://www.pcrs-uk.org/training-providers)
21 providers (accessed 7 Aug 2015)
22
23 23. Pharmaceutical Services Negotiating Committee. *NHS Community Pharmacy Services - a*
24 *summary* 2013. [http://psnc.org.uk/wp-content/uploads/2013/08/CPCF-summary-July-](http://psnc.org.uk/wp-content/uploads/2013/08/CPCF-summary-July-2013.pdf)
25 2013.pdf (accessed 19 Jun 2015)
26
27 24. Pharmaceutical Services Negotiating Committee. *Public Health. A practical guide for community*
28 *pharmacists* 2013. [http://psnc.org.uk/wp-](http://psnc.org.uk/wp-content/uploads/2013/07/public_health_book_all.pdf)
29 content/uploads/2013/07/public_health_book_all.pdf (accessed 24 Feb 2015)
30
31 25. Stephenson-Glynn J. JSNA Factsheet: *Health and Wellbeing Tobacco Control Tower Hamlets Joint*
32 *Strategic Needs Assessment 2013-2015*.
33 [http://www.towerhamlets.gov.uk/idoc.ashx?docid=026c2fee-f37f-4479-8d6e-](http://www.towerhamlets.gov.uk/idoc.ashx?docid=026c2fee-f37f-4479-8d6e-4b9660a1e282&version=1)
34 [4b9660a1e282&version=1](http://www.towerhamlets.gov.uk/idoc.ashx?docid=026c2fee-f37f-4479-8d6e-4b9660a1e282&version=1) (accessed 24 Feb 2015)
35
36 26. Flowers R. *Newham joint strategic needs assessment 2011/12 September 2012 Update*.
37 [http://www.newham.gov.uk/Documents/Council%20and%20Democracy/JSNASept2012Upd](http://www.newham.gov.uk/Documents/Council%20and%20Democracy/JSNASept2012Update.pdf)
38 [ate.pdf](http://www.newham.gov.uk/Documents/Council%20and%20Democracy/JSNASept2012Update.pdf) (accessed 24 Feb 2015).
39
40 27. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a
41 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19:349-57.
42
43 28. Foot C, Gilbert H, Dunn P et al. *People in control of their own health and care. The state of*
44 *involvement*. King's Fund, London, 2014.
45
46 29. Amemori M, Michie S, Korhonen T, et al. Assessing implementation difficulties in tobacco use
47 prevention and cessation counselling among dental providers. *Implement Sci* 2011;6(1):50.
48
49 30. Amemori M, Korhonen T, Michie S, et al. Implementation of tobacco use cessation counseling
50 among oral health professionals in Finland. *J Public Health Dent* 2013;73:230-36.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 31. Beenstock J, Sniehotta F, White M, et al. What helps and hinders midwives in engaging with
4 pregnant women about stopping smoking? A cross-sectional survey of perceived
5 implementation difficulties among midwives in the North East of England. *Implement Sci*
6 2012;7(1):36.
7
8
9
10 32. Patwardhan PD, Chewning BA. Ask, advise and refer: Hypothesis generation to promote a brief
11 tobacco-cessation intervention in community pharmacies. *Int J Pharm Pract* 2009;17:221-29.
12
13 33. Morton K, Pattison H, Langley C, et al. A qualitative study of English community pharmacists'
14 experiences of providing lifestyle advice to patients with cardiovascular disease. *Res Social*
15 *Adm Pharm* 2015;11:e17-e29.
16
17 34. West R, McEwen A, Lorencatto F, et al. *NC SCT Training Standard. Learning Outcomes for Training*
18 *of Stop Smoking Practitioners* In: McEwen A, ed.: National Centre for Smoking Cessation and
19 Training, London, 2013.
20
21 35. Hiscock R, Bauld L. *Stop Smoking Services and Health Inequalities*. In: McEwen A, ed.: National
22 Centre for Smoking Cessation and Training (NCSCT), London 2013.
23
24 36. Hoving C, Mudde AN, de Vries H. Effect of recruitment method and setting on the composition of
25 samples consisting of adult smokers. *Patient Educ Couns* 2007;65(1):79-86.
26
27 37. Hammond D, McDonald PW, Fong GT, et al. Do smokers know how to quit? Knowledge and
28 perceived effectiveness of cessation assistance as predictors of cessation behaviour.
29 *Addiction* 2004;99:1042-48.
30
31 38. McDermott MS, Beard E, Brose LS, et al. Factors Associated With Differences in Quit Rates
32 Between "Specialist" and "Community" Stop-Smoking Practitioners in the English Stop-
33 Smoking Services. *Nicotine Tob Res* Published Online First: 13 December 2012. doi:
34 10.1093/ntr/nts262
35
36 39. Brown J, Kotz D, Michie S, et al. How effective and cost-effective was the national mass media
37 smoking cessation campaign 'Stoptober'? *Drug Alcohol Depend* 2014;135:52-58.
38
39 40. Kmietowicz Z. Action is needed to boost uptake of stop smoking services, say campaigners. *BMJ*
40 Published Online First: 23 August 2015. doi:10.1136/bmj.h4578
41
42 41. Kotz D, Stapleton JA, Owen L, et al. How cost-effective is 'No Smoking Day'? *Tob Control*
43 2011;20:302-04.
44
45 42. Owen L, Youdan B. 22 years on: the impact and relevance of the UK No Smoking Day. *Tob Control*
46 2006;15:19-25.
47
48 43. Oakes W, Chapman S, Borland R, et al. "Bulletproof skeptics in life's jungle": which self-
49 exempting beliefs about smoking most predict lack of progression towards quitting? *Prev*
50 *Med* 2004;39:776-82.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 44. Paul CL, Ross S, Bryant J, et al. The social context of smoking: A qualitative study of comparing
4 smokers of high versus low socioeconomic position. *BMC Public Health* 2010;10:211
5
6 45. Cunningham P. Patient engagement during medical visits and smoking cessation counseling.
7
8 *JAMA Intern Med* 2014;174:1291-98.
9
10 46. Hudmon KS, Prokhorov AV, Corelli RL. Tobacco cessation counseling: Pharmacists' opinions and
11 practices. *Patient Educ Couns* 2006;61:152-60.
12
13 47. Norcross JC, Krebs PM, Prochaska JO. Stages of Change. *J Clin Psychol* 2011;67:143-54.
14
15 48. Prescribing and Primary Care Team HSCIC. *General Pharmaceutical Services in England: 2004-05*
16 *to 2013-14*. Health and Social Care Information Centre, Leeds, 2014.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Figure legends**

4 Figure 1. Illustration of behavioural factors affecting adviser engagement and retention behaviour
5
6 (modified from the COM-B model)
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1 Illustration of behavioural factors affecting adviser engagement and retention behaviour (modified from the COM-B model)




Figure 1. Illustration of behavioural factors affecting adviser engagement and retention behaviour (modified from the COM-B model)
146x209mm (300 x 300 DPI)

Supplementary file 1 Theoretical Domains Framework (TDF) - Domain definitions

TDF (definition)	Theoretical Constructs	Possible Example
Knowledge <i>An awareness of the existence of something</i>	Knowledge (including knowledge of condition /scientific rationale); procedural knowledge; knowledge of task environment	Knowledge of why important to ask people if smoke; knowledge of nicotine patches so can deliver service well
Skills <i>An ability or proficiency acquired through practice</i>	Skills; Skills development; competence; ability; interpersonal skills; practice; skill assessment	Communication skills
Memory, attention and decision processes <i>The ability to retain information, focus selectively on aspects of the environment and choose between two or more alternatives</i>	Memory; attention; attention control; decision making; cognitive overload/tiredness	Remembering to deliver different parts of service
Behavioural regulation <i>Anything aimed at managing or changing objectively observed or measured actions</i>	Self-monitoring; breaking habit; action planning	Are there systems which help monitor whether something has been done or not
Environmental context and resources <i>Any circumstance of a person's situation or environment that discourages or encourages the development of skills and abilities, independence, social competence and adaptive behaviour</i>	Environmental stressors; resources/material resources; organisational culture/climate; salient events/critical incidents; person x environment interaction; barriers and facilitators	The lack of a consulting room impedes consultations
Social influences <i>Those interpersonal processes that can cause individuals to change their thoughts, feelings or behaviours</i>	Social pressure; social norms; group conformity; social comparisons; groups norms; social support; power; intergroup conflict; alienation; group identity; modelling	Senior pharmacist not prioritising smoking cessation
Professional/social role and identity <i>A coherent set of behaviours and displayed personal qualities of an individual in a social or work setting</i>	Professional identity; professional role; social identity; identity; professional boundaries; professional boundaries; professional confidence; group identity; leadership; organisational commitment	Identification that behaviour change is within a pharmacists remit
Beliefs about capabilities <i>Acceptance of the truth, reality or validity about an ability, talent or facility that a person can put to constructive use</i>	Self-confidence; perceived competence, self-efficacy; perceived behavioural control; beliefs; self-esteem; empowerment; professional confidence	Confidence that can raise question of smoking in non-threatening manner
Optimism <i>The confidence that things</i>	Optimism, pessimism; unrealistic optimism; identity	Belief that with support most people can give up smoking

1 2 3 4 5 6	<i>will happen for the best or that desired goals will be attained</i>		
7 8 9 10 11 12	Beliefs about consequences <i>Acceptance of truth, reality, or validity about outcomes of a behaviour in a given situation</i>	Beliefs; outcome expectancies; characteristics of outcome expectancies; anticipated regret; consequents	If don't stop smoking will die from cancer
13 14 15 16 17 18	Intentions <i>A conscious decision to perform a behaviour or a resolve to act in a certain way</i>	Stability of intentions; stages of change model; transtheoretical model and stages of change	Intention to give up smoking
19 20 21 22 23 24	Goals <i>Mental representations of outcomes or end states that an individual wants to achieve</i>	Goals (distal/proximal); goal priority; goal/target setting; goals (autonomous/controlled); action planning; implementation intention	Wanting to achieve increased uptake by x amount
25 26 27 28 29 30 31	Reinforcement <i>Increasing the probability of a response by arranging a dependent relationship, or contingency, between the response and a given stimulus</i>	Rewards (proximal/distal, valued/not valued, probable/improbable); incentives; punishment; consequents; reinforcement; contingencies; sanctions	Financial reward
32 33 34 35 36 37 38 39 40 41	Emotion <i>A complex reaction pattern, involving experiential, behavioural and physiological elements by which the individual attempts to deal with a personally significant matter or event</i>	Fear, anxiety; affect; stress; depression; positive/negative affect; burn-out	Discussing smoking raises distress due to family members recent smoking related death
42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60	Definitions of the COM-B model. Capability i) Physical Capability e.g. not having lost voice during smoking cessation consultation, ii) Psychological Capability e.g. having knowledge of NRT products to talk about them Opportunity i) Physical Opportunity afforded by environment involving time, resources, locations, cues e.g. pharmacist having sufficient time to speak with clients ii) Social Opportunity afforded by interpersonal influences, social cues, cultural norms influence way think about things e.g. pharmacist allows counter assistant time to become trained in stop smoking Motivation i) Reflective Motivation involving plans (intentions) and evaluations (beliefs about what is good and bad) e.g. intention to give up smoking because believe bad for health ii) Automatic Motivation includes automatic processes which involve emotional reactions, desires (wants and needs), impulses, inhibitions, drive states and reflex responses.		

1
2
3 Research Checklist: Consolidated criteria for reporting qualitative studies (COREQ): 32-item checklist
4

5 No Item Guide questions/description

6 **Domain 1: Research team and reflexivity**

7 Personal Characteristics

8 1. Interviewer/facilitator - Which author/s conducted the interview or focus group?

9 Author RS and VM conducted the interviews. Page 7.

10 2. Credentials - What were the researcher's credentials? E.g. PhD, MD

11 Author RS and VM have a PhD. Not mentioned in manuscript.

12 3. Occupation - What was their occupation at the time of the study?

13 Author VM was one of the qualitative researchers who conducted 11 interviews and left during the
14 course of the study. Author RS joined the study team as a qualitative researcher and conducted 14
15 interviews. Page 7 suggests 'qualitative researcher'.

16 4. Gender - Was the researcher male or female?

17 Both RS and VM are female. Page 7.

18 5. Experience and training - What experience or training did the researcher have?

19 Both RS and VM have expertise in qualitative research methods through education/training courses.
Page 7 mentions the term 'experienced'

20 Relationship with participants

21 6. Relationship established - Was a relationship established prior to study commencement?

22 No. The interviewers were not known to the study participants prior to study commencement and they
23 were not involved in participant recruitment. Page 7.

24 7. Participant knowledge of the interviewer - What did the participants know about the researcher?

25 e.g. personal goals, reasons for doing the research

26 The participants were made aware by the study team that a researcher would contact them to arrange
27 the interview. Prior to the start of the interview the purpose of the study and interview was explained
28 to each participant. Page 7.

29 8. Interviewer characteristics - What characteristics were reported about the interviewer/facilitator?

30 e.g. Bias, assumptions, reasons and interests in the research topic

31 The participants only knew that the interviewers were study researchers. Page 7.

32 **Domain 2: study design**

33 Theoretical framework

34 9. Methodological orientation and Theory - What methodological orientation was stated to underpin

35 the study? e.g. grounded theory, discourse analysis, ethnography, phenomenology, content analysis

36 We used thematic framework analysis applying the Theoretical Domains Framework (TDF) within the
37 COM-B (Capability, Opportunity, Motivation-Behaviour) model of behaviour change. Page 5.

38 Participant selection

39 10. Sampling - How were participants selected? e.g. purposive, convenience, consecutive, snowball

40 Purposive sampling was used to obtain a diverse range of views. We selected stop smoking advisers
41 who differed by gender and duration of being an adviser. Page 7.

42 11. Method of approach - How were participants approached? e.g. face-to-face, telephone, mail,
43 email.

44 Pharmacies were sent a letter and information sheet and contacted by telephone to arrange a face-
45 to-face meeting in the pharmacy. Page 7.

46 12. Sample size - How many participants were in the study?

47 Twenty-five participants were in the study. Page 9.

48 13. Non-participation - How many people refused to participate or dropped out? Reasons?

49 Nineteen advisers refused to participate. Reasons for non-participation were lack of interest (n=14) or
50 unavailable/no answer (n=5). Page 9.

51 Setting

52 14. Setting of data collection - Where was the data collected? e.g. home, clinic, workplace

53 The setting was the consultation room of the community pharmacy. Page 7.

54 15. Presence of non-participants - Was anyone else present besides the participants and
55 researchers?

56 No. Page 7.
57
58
59
60

16. Description of sample - What are the important characteristics of the sample? e.g. demographic data, date.

The following are characteristics of the study sample: whether interviewee/stop smoking adviser was a pharmacist or other pharmacy support staff, study ID, gender, duration of being a stop smoking adviser. Table 1 on Page 10. The proportion on ethnicity of advisers, proportion of multi-lingual service and training of advisers is on Page 9.

Data collection

17. Interview guide - Were questions, prompts, guides provided by the authors? Was it pilot tested? A semi-structured interview guide was used to conduct the interviews. A question was added after conduct of a few interviews as an issue emerging from the data. Page 8.

18. Repeat interviews - Were repeat interviews carried out? If yes, how many?
No repeat interviews were carried out.

19. Audio/visual recording - Did the research use audio or visual recording to collect the data?
Digital audio-recording was used to collect the data. Page 7.

20. Field notes - Were field notes made during and/or after the interview or focus group?
No.

21. Duration - What was the duration of the interviews or focus group?
Duration of the interviews lasted 30-60 minutes. Page 7.

22. Data saturation - Was data saturation discussed?
Yes. The study stopped recruitment once no new knowledge was being obtained from the interviews. Page 7.

23. Transcripts returned - Were transcripts returned to participants for comment and/or correction?
The transcripts (typed per verbatim) were not returned to participants for comment and/or correction as not planned for study.

Domain 3: analysis and findings

Data analysis

24. Number of data coders - How many data coders coded the data?
There were two data coders. RS coded all the data collected and author CR coded 20% of the data which was assessed using kappa agreement. In addition, the emergent themes from the data collected was mapped onto the behavioural theory to generate analytical themes. The mapping conducted by RS and was checked and discussed with author LS (health psychologist) to ensure the mapped thematic data fitted with the domain definition and its content. See supplementary file 1 for full definition of the COM-B and TDF constructs. Page 9.

25. Description of the coding tree - Did authors provide a description of the coding tree?
Yes, the description of the coding leading to the formation of categories and themes has been provided in the manuscript as per the thematic framework analysis. Page 9.

26. Derivation of themes - Were themes identified in advance or derived from the data?
As per the thematic framework analysis, this method starts deductively with a priori codes (not themes) from the study aims and objectives however, subsequent analysis is inductive and grounded in the accounts of the participants. So the themes were identified from the data and not identified in advance. Page 9.

27. Software - What software, if applicable, was used to manage the data?
Nvivo software was used to manage the data. Page 7.

28. Participant checking - Did participants provide feedback on the findings?
This was not planned in this study.

Reporting

29. Quotations presented - Were participant quotations presented to illustrate the themes / findings? Was each quotation identified? e.g. participant number
Yes. Participant quotations have been presented to illustrate the themes. Yes each quotation can be identified by the study ID given in Table 1 of the manuscript. Page 9 and 10.

30. Data and findings consistent - Was there consistency between the data presented and the findings?

Yes. The data analysis that included mapping onto theory to generate analytical themes led to the study findings, exemplified by participant quotations. The study findings include themes/high level finding from the COM-B model and the subheadings represent relevant domains from the Theoretical Domains Framework. Page 9.

31. Clarity of major themes - Were major themes clearly presented in the findings?

1
2
3 Yes. The emergent themes mapped on to five of 14 TDF constructs have been clearly presented in
4 the findings and has been illustrated in Figure 1 of the manuscript. Page 9.

5 32. Clarity of minor themes - Is there a description of diverse cases or discussion of minor themes?

6 Yes. An independent theme emerged following the analysis that was not mapped onto the TDF
7 constructs and yet was an important finding and has been clearly presented and illustrated in Figure 1
8 of the manuscript. Page 9.

9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1
2
3 **Understanding recruitment and retention in the NHS community pharmacy stop smoking service:**
4 **perceptions of smoking cessation advisers**

5
6 Ratna Sohanpal¹

7
8 Carol Rivas²

9
10 Liz Steed¹

11
12 Virginia MacNeill³

13
14 Valerie Kuan¹

15
16 Elizabeth Edwards¹

17
18 Chris Griffiths¹

19
20 Sandra Eldridge¹

21
22 Stephanie Taylor¹

23
24 Robert Walton¹

25
26 ¹Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health, Barts and The
27 London School of Medicine and Dentistry, Queen Mary University of London, London, UK

28
29 ²Faculty of Health Sciences, University of Southampton, UK

30
31 ³University of Oxford, UK

32 **Corresponding author:**

33
34 Ratna Sohanpal

35
36 Research Fellow

37
38 Asthma UK Centre for Applied Research, Centre for Primary Care and Public Health

39
40 Blizzard Institute

41
42 Queen Mary University of London

43
44 58 Turner Street, London

45
46 E1 2AB

47
48 Ph: +442078822492

49
50 Fax: +442078822552

51
52 Email: r.sohanpal@qmul.ac.uk

ABSTRACT

Objectives: To understand views of pharmacy advisers about smoker recruitment and retention in the National Health Service community pharmacy stop smoking programme.

Design: Thematic framework analysis of semi-structured, in-depth interviews applying the Theoretical Domains Framework and COM-B behaviour change model. We aimed to identify aspects of adviser behaviour that might be modified to increase numbers joining and completing the programme.

Participants: 25 Stop Smoking Advisers (13 pharmacists and 12 support staff).

Setting: 29 community pharmacies in three inner east London boroughs.

Results: Advisers had preconceived ideas about smokers likely to join or drop out and made judgements about smokers' readiness to quit. Actively recruiting smokers was accorded low priority due in part to perceived insufficient remuneration to the pharmacy and anticipated challenging interactions with smokers. Suggestions to improve smoker recruitment and retention included developing a more holistic and supportive approach using patient-centred communication. Training counter assistants was seen to be important as was flexibility to extend the programme duration to fit better with smokers needs.

Conclusions: Cessation advisers feel they lack the interpersonal skills necessary to engage well with smokers and help them to quit. [Addressing advisers' behaviours about active engagement and follow-up of clients, together with regular skills training including staff not formally trained as cessation advisers could potentially boost numbers recruited and retained in the stop smoking programme.](#) Adjustments to the [pharmacy remuneration](#) structure to incentivise recruitment and to allow personalisation of the programme for individual smokers should also be considered.

Keywords: Smoking cessation, Community Pharmacies, Qualitative research, Patient recruitment, Retention, Health Behaviour, Communication, Psychological Theory

Word count: 4,077

Strengths and weaknesses

- This is the first study to provide insight into smoker recruitment and retention in the NHS community pharmacy stop smoking programme from the adviser's perspective and therefore fills an important gap in knowledge.
- We used the Theoretical Domains Framework and COM-B behaviour change model as a framework to identify adviser attitudes and behaviours that could potentially be modified, thus providing a unique insight into smoker recruitment and retention in this clinical setting and facilitating development of interventions.
- In our analysis we achieved theoretical saturation, high inter-rater reliability and high agreement with a health psychologist about mapping thematic data onto constructs of the Theoretical Domains Framework.
- Whilst recruitment and retention are clearly two separate behaviours we analysed them together because we were interested in the combined result – namely increased throughput in the service. Future studies might profitably separate these behaviours to examine their unique motivations.
- Our findings will not be generally applicable, but could nevertheless be transferable to community pharmacies delivering the stop smoking programme in other socio-economically deprived communities across the UK. Certain findings such as low self-efficacy in consultation skills and the need to train all pharmacy staff to increase service throughput may well be transferrable to other healthcare systems, however this would need to be examined in the different settings.

INTRODUCTION

Improving public health through extending community pharmacy services has become a key part of UK National Health Service (NHS) operating strategy.[1-2] Smoking cessation is a public health priority[3] since smoking is the most important cause of premature morbidity and mortality worldwide.[4] In the UK, smoking cessation through pharmacy-led stop smoking services is now a key element of the Department of Health's tobacco control strategy.[2]

Community pharmacies are accessed by people both in health and illness, hence community pharmacy stop smoking advisers are well placed to reach large numbers of smokers.[5] Behavioural counselling/support, or a combination of nicotine replacement therapy and counselling by trained pharmacy staff are effective in helping smokers to quit.[2, 4, 6] However, the overall effectiveness of the pharmacy stop smoking programme does not only depend on the quit rates achieved but also on the number of smokers who participate and adhere to the programme.[7-8] In 2012-2013 of the estimated two million smokers in England,[9] only 149,034 smokers set a quit date in the pharmacy NHS stop smoking programme. Of these, 48% successfully quit at four weeks compared to the target of 70%.[2, 10] Quit rate was slightly lower than that achieved by the stop smoking services in primary care (50%) and lower than specialist stop smoking services (56%).[10] This apparent poor performance may result from differences in case-mix, however variations in staff training and environmental factors in pharmacies may make high cessation rates harder to achieve than in other settings. Nevertheless there are considerable disparities in quit rates between pharmacies suggesting potential for improvement.[2]

A recent review[6] suggested that pharmacists only target those they perceive to be ready to quit, thus reducing recruitment potential. Retention within the service is also poor, for example only 35% of the 4500 people who joined the stop smoking programme in one inner London borough, successfully quit (i.e. quit status biochemically verified by carbon monoxide testing at four weeks from the quit date)[11] and the remainder were lost to follow up.[12] Thus in addition to optimising quit rates, improving recruitment and retention to the stop smoking programme might raise overall numbers successfully giving up smoking.

In a systematic review of pharmacist views on delivery of public health services pharmacists recognised they should be more active in smoking cessation, however a number of barriers were suggested such as lack of time, staff and a designated space. Knowledge and skills were thought to be lacking and patients did not expect to receive health promotion advice from pharmacists. In addition there was reluctance to initiate conversations about health promotion because of fears of generating negative responses.[13]

1
2
3 No detailed investigation of stop smoking advisers' views on factors affecting recruitment
4 and retention in the NHS stop smoking service has been published to date. This is the first such UK
5 qualitative study aiming to: (1) understand stop smoking advisers' views relating to smoker
6 recruitment and retention in the NHS stop smoking programme in community pharmacies; (2)
7 identify factors that might be targeted in an intervention to increase smoker recruitment and
8 retention to maximise the effectiveness of the programme.
9

10
11
12 The results of this study will contribute to development of a complex intervention to
13 promote uptake and increase the effectiveness of the NHS pharmacy smoking cessation service. This
14 intervention will be evaluated in a cluster randomised trial in east London as part of the STOP
15 programme (Smoking Treatment Optimisation in Pharmacies).
16
17
18
19

20 21 **METHODS**

22 **Study design**

23
24 We conducted semi-structured, in-depth interviews with stop smoking advisers (comprising both
25 pharmacists and pharmacy support staff). For roles and responsibilities of staff and an outline of the
26 stop smoking programme see Box 1.
27
28

29 We used thematic framework analysis[14-15] applying the Theoretical Domains Framework
30 [16] within the COM-B model of behaviour change.[17] The Theoretical Domains Framework is
31 synthesised from multiple behaviour change theories and comprises 14 domains of theoretical
32 constructs that explain possible influences on behaviour.[16] Each domain fits within one of the
33 three components of the COM-B model - Capability Opportunity and Motivation.[17-18] For
34 example, 'Skills' in the Theoretical Domains Framework maps on to 'Capability' in the COM-B model,
35 'Professional Role and Identity' to 'Motivation', 'Environmental Context and Resources' to
36 'Opportunity'. [18] We used COM-B as a lens through which to view our data because it provides a
37 practical basis for designing interventions aimed at behaviour change, helping to identify the
38 behavioural target and the components of the behaviour system needing to be changed.[17]
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Setting and participants

The study was conducted in three inner east London boroughs: Tower Hamlets; Newham; City and Hackney. These boroughs include south Asian and African/Caribbean communities with high levels of tobacco use and persistent health problems linked to social and economic inequalities.[12, 25-26] Smoking prevalence in these deprived boroughs is close to the UK national average (20%), or higher (Newham 21%, 23% in Hackney, 37% in Tower Hamlets).[12, 25-26] Purposive sampling was used to obtain a diverse range of views.[27] We selected stop smoking advisers who differed by gender and duration of being an adviser. Independent community pharmacies were sent a letter and information sheet and contacted by telephone to arrange a face-to-face meeting in the pharmacy. Within pharmacies the pharmacist usually suggested the adviser (usually themselves or a member of other pharmacy support staff i.e. stop smoking adviser for interview). A member of the research team (CR) or a research assistant obtained written informed consent.

Data collection

Individual interviews were conducted from January to June 2014 by an experienced female qualitative researcher (RS, VM) using an interview schedule (Box 2). The interviewer was not known to the study participants and was not involved in recruiting participants to the qualitative study. Advisers were informed that study aim was to improve programme recruitment, retention and quit rates. Interviews took place in the consultation room of the community pharmacy and lasted 30-60 minutes. Recruitment and interviews continued until data saturation was achieved. All interviews were audio-recorded and fully transcribed. Nvivo10 was used for organisation of data and to facilitate analysis.

Data analysis and interpretation

A thematic framework analysis[14-15] was conducted. This method starts deductively with *a priori* codes from the study aims and objectives however, subsequent analysis is inductive and grounded in the accounts of the participants.[14] Researcher (RS) read and re-read the transcripts for familiarisation and data immersion. A thematic framework was formed following line-by-line coding and comprised useful memos or descriptive statements to develop categories. Comparisons were made within and between transcripts based on the thematic framework. The data were then lifted and charted across the thematic categories. This process was carried out initially with a few transcripts and discussed with the study team (CR, RW, LS) to assess reliability and agreement between two coders and to check validity of the analysis. Inter-rater reliability was 90.9% (kappa agreement) between the two coders (RS, CR) on 20% of the transcripts. Subsequently, the emergent themes were mapped (by RS) onto behavioural constructs of the TDF and the COM-B model, where applicable, to generate analytical themes.

The mapping was discussed and agreed with a health psychologist (LS) to ensure the mapped thematic data fitted with the domain definition and its content (Supplementary file 1). This process generated analytical themes or key domains influencing the engagement and the retention behaviour of advisers and helped us to identify domains that might be targeted to optimise adviser behaviour. Emergent themes are exemplified below with direct participant quotations.

RESULTS

Twenty-five interviews were conducted following approaches to 44 advisers in 29 community pharmacies. Reasons for non-participation were lack of interest (n=14) or unavailable/no answer (n=5). The participants were drawn from 15 pharmacies with six pharmacies contributing two or three interviewees. Over half of the advisers were of Asian ethnicity (56%) and a small proportion (20%) of pharmacies provided a multi-lingual service. Characteristics of the participants are shown in Table 1. All participants were smoking cessation advisors trained to level 2.

In the analysis the emergent themes mapped on to five of 14 TDF constructs with one independent theme. Figure 1 illustrates this using COM-B as headings and TDF as subheadings (*italics*). The independent theme and its potential relation to COM-B is highlighted with dashed lines. See supplementary file 1 for full definition of the COM-B and TDF constructs. In the text that follows the higher level heading is from the COM-B model and the subheadings represent relevant domains from the Theoretical Domains Framework that explain two separate adviser behaviours, recruitment and retention.

Table 1 Characteristics of study participants

Study ID*	Pharmacist or support staff	Gender	Duration being stop smoking adviser
S11A101	Pharmacist	Female	Not given
S19A182	Pharmacist	Female	3 years
S08A071	Pharmacist	Male	Not given
S13A121	Pharmacist	Male	10 years
S29A281	Pharmacist	Male	4 years
S04A031	Pharmacist	Male	Not given
S09A081	Pharmacist	Male	10 years
S23A221	Pharmacist	Male	Not given
S25A242	Pharmacist	Male	7 years
S26A253	Pharmacist	Male	Not given
S03A021	Pharmacist	Male	Not given
S05A041	Pharmacist	Male	5-6 years
S11A103	Support staff	Female	9 years
S02A012	Support staff	Female	Not given
S25A241	Support staff	Female	1 year
S26A251	Support staff	Female	7 years
S09A082	Support staff	Female	5 years
S25A243	Support staff	Female	Not given
S26A252	Support staff	Male	4 years
S01A001	Support staff	Male	Not given
S04A032	Support staff	Male	6 years
S11A102	Support staff	Male	4 years
S19A181	Support staff	Male	12 years
S24A231	Support staff	Male	Not given

* This code is used to identify the source of quotations in the text.

Motivation

(1) Professional role and identity

Many participants felt proud of their work and were satisfied that through their role as advisers they were able to help people quit smoking, which in turn helped with the health and well-being of the wider community.

I mean, you are always happy that you've helped someone do something. And there are many times that, this guy (client who has quit) will always pop in. ...and say "Oh, I'm just passing and I thought let me just say hi to you!" And there are many times ... This lady, it's nice. I mean, why would this lady bring her son over for me to advise? It's like they appreciate it and you have that feeling that, oh, you've also managed to help someone and it's great to do that. (Study ID S08A071, Pharmacist)

Identification and engagement of smokers into the programme

Advisers identified potential clients through: GP referrals; recommendations from people who had quit; opportunistically during medicine review or while conducting risk assessment; and when people bought products from the pharmacy, picked up regular prescriptions or bought over the counter medication. Despite this, many advisers selected and recruited only those smokers who mentioned their readiness to quit because they thought that these smokers were less likely to drop out. A few said they only recruited those who specifically asked for smoking cessation advice and were not interested in actively recruiting smokers to the NHS SSP.

She's like I'm not in the right frame of mind but I really want to give up smoking. So in that way we just tell them look, ... So we told her and with smoking you have to make sure that they're in the right frame of mind otherwise there's no point them joining the programme.

Otherwise they'll join and then quit after a week and there's no point. (S24A231, Pharmacy support staff)

I would say the recruitment. I mean it's a bit hard, most of it has to be walking and...yeah. Or maybe they see the poster. But we're not doing that much. (S19A182, Pharmacist)

Some participants said that recruitment into the programme was low priority. Advisers wanted to prioritise patients with long-term conditions or those that they thought had life-threatening conditions. One adviser felt giving up smoking was the responsibility of the smoker.

...so smoking is more or less a very very little part of our actual work here. We actually have people who have life threatening conditions, you can be dead tomorrow. So that's of a

1
2
3 *greater priority than smoking. Smoking is by choice, you want to give up, you don't want to*
4 *give up. (A01A001, Pharmacy support staff)*
5
6

7
8 Several participants also said they believed that all the advice, support and products they provided
9 could only help a smoker quit if the smoker was mentally prepared and had the willpower to follow
10 through the advice and make the necessary lifestyle/environmental changes conducive to stopping
11 smoking. Advisers felt they had most success with this type of client.
12

13
14 *... we're pretty straight with them (potential clients), ... if they're motivated to quit then*
15 *they've got a better chance of succeeding, things like ...NRT Champed (sic) only helps to a*
16 *certain extent ... and if they don't have the willpower it's not going to work... (S09A082,*
17 *Pharmacist)*
18
19
20

21 22 Encouragement and motivation to ensure attendance and adherence

23
24 Some participants stated that encouraging clients to continue attending the programme sessions
25 was important. One adviser mentioned that to prevent dropouts and indeed when engaging with
26 clients who had relapsed, identifying the reasons for stopping smoking and relapse were important.
27

28
29 *Oh I think you are always going to get that (people dropping out). ...I mean you are talking*
30 *about 1 in 2 not getting through so you just have to keep motivating and telling them you*
31 *know I mean you've been smoking for 20 or 30 years, you are not going to give up in 6 to 12*
32 *weeks, it's a long term process. And I think if you have like help, if you have re-started again,*
33 *again the important thing is to find the reasons why because I think if you can get to the*
34 *bottom of that and work on that rather than just giving them NRT and thinking, you know so*
35 *really that is most important. The NRT helps but you need to know why you want to stop*
36 *smoking. (S03A021, Pharmacist)*
37
38
39
40
41
42

43
44 A few participants did not see following up clients who missed sessions as part of their role. Some
45 said that they simply accepted it if smokers mentioned they were finding it difficult to not smoke
46 when they spoke to them by telephone or in person.
47

48
49 *Two persons I've seen, they can't manage (to quit), that's it. Sometimes if they come for their*
50 *medicines to pick up or anything, they can say, "Oh, sorry, I didn't come back, because I can't*
51 *manage at the moment, I'm not all right." So we say, "OK, whenever you feel OK."*
52 *(S02A012, Pharmacy support staff)*
53

54
55 *...when they don't come back it's like well, did you give them the wrong products? They didn't*
56 *suit them, they could've phoned us, we could've spoken to them, we could've swapped it. We*
57
58

59
60
10

1
2
3 *are pretty flexible. But if they don't come back to us there's no way of knowing, is there?*

4 (S25A241, Pharmacy support staff)
5
6
7

8 Some advisers acknowledged that more smoker engagement would result in higher quit rates. The
9 following were suggested to boost service recruitment and retention: identify potential smokers for
10 the service e.g. when they bought cough syrup; engage, inform, encourage and recommend the
11 programme to smokers e.g. through use of poster adverts; explain how the programme and adviser
12 involvement could help with quitting; place a holistic focus on the person.
13
14

15
16 *So if you're focused on the client rather than the process you're going to get a bigger*
17 *outcome rather than saying this is a tick form, I've got to tick this, I've got to fill this in, I've*
18 *got to fill that in and oh, I've missed out here, we've got to do a CO reading sort of thing.*

19 (S26A253, Pharmacist)
20
21
22

23 24 **(2) Reinforcement**

25 Remuneration

26
27 This process of remuneration was acknowledged as an issue by the adviser participants who were a
28 mix of owner pharmacists, employed pharmacists and other pharmacy support staff. Among the
29 owner pharmacists, a few said that the current remuneration for quitters was reasonable, while
30 some others felt that the remuneration did not take into account the time they spent with smokers
31 encouraging them to join and adhere to the service.
32
33

34
35 One owner pharmacist mentioned that the current payment system was a disincentive to spend
36 time with smokers, and needed revisiting by commissioners; payment based on quitters might
37 influence which smokers staff are willing to support.
38
39

40 *And things, we get penalised for non-quitters which are not really our fault.*

41
42 *Interviewer: ...How do you mean penalised?*

43 *So currently we may spend three or four hours actually going through these patients, and*
44 *because they don't quit we don't get the full £40 or £35. We end up with about £20. That's*
45 *the disincentive. There are certain health authorities now are saying that if a patient doesn't*
46 *quit there's no fees. So there is a lot of disincentives for a pharmacist to say look, I only get a*
47 *50 per cent success rate, I'm not going to spend three or four hours at a time and get nothing*
48 *for it. So they're (pharmacies are) not going to take on new clients and the way that the*
49 *health authorities and CCGs think about these programmes needs to change. (S26A253,*
50 *Pharmacist)*
51
52
53
54
55
56
57

1
2
3 Another owner pharmacist stated that perhaps smokers should be incentivised to attend their stop
4 smoking appointments as it could mean saving money for the NHS in the long run and some
5 pharmacy support staff suggested providing incentives/rewards to their clients that could be given
6 at programme completion.
7
8

9
10 *And I think if you can perhaps issue vouchers, but issue vouchers that are redeemable at the*
11 *end of the process, yeah? So it makes no sense for them to come to two or three*
12 *consultations, accumulate vouchers and then miss the fourth one, yeah, you know? So I*
13 *think there should be vouchers that can be redeemed for cash at the end of it, because most*
14 *people don't want to be restricted as to what they can do with a voucher or cash or anything.*
15 *That is one thing I think....at the end of the day, we need to put it in the balance; I mean, the*
16 *government feels that getting people to stop smoking leads to a greater saving on the NHS,*
17 *so it's something they should look at. (S29A281, Pharmacist)*
18
19
20
21
22

23
24 Several of the employed pharmacists and support staff acknowledged that delivery of the
25 programme was part of their job, although a few highlighted that if their pharmacy were to receive
26 an increase in the current remuneration for quitters then they might be allowed to spend more time
27 engaging with and following up clients to continue and complete the programme.
28
29

30
31 *I think so as well, because it (current remuneration) is a kind of barrier. You do encourage*
32 *them (smoker) to stop smoking, but then three weeks you spend with them, although they*
33 *don't stop smoking ... And for one engagement they give you (payment), but after that, if*
34 *they keep on giving us the money we (can) encourage more people, ... So if they encourage*
35 *me, I mean the pharmacy, by providing with more money or anything, no barriers regarding*
36 *money, then we can do more, much more. (S26A251, Pharmacy support staff)*
37
38
39
40
41

42 **(3) Beliefs about capabilities**

43 Interaction with smokers

44
45 Some advisers mentioned that engaging with smokers and motivating clients who enjoyed smoking
46 or who were not interested in quitting was a challenge to programme recruitment and retention. In
47 addition, one participant was unsure about the protocol to follow when a client smoked a cigarette
48 at the end of 12 week programme.
49
50
51

52 *It's more of a challenge when they say oh, but I love smoking, I love the taste of smoking.*
53 *Then it's a bit of a challenge because they've already got in their mind that...there's like a*
54 *mental block. Whatever you're going to say to me, I don't really care sort of thing. ..*
55 *(S24A231, Pharmacy support staff)*
56
57
58
59
60

12

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In contrast, a few participants felt confident to handle some of the above mentioned issues by talking about the harmful effects of smoking, showing the tar jar to smokers and suggesting to smokers they were available to support them when they are ready to quit.

Yeah, I mean there's no harm in recommending it (the service). You recommend it to everyone. I don't see anything bad about that. It's just whether they'll actually sign up or not. ... The only way we can deal with that is through awareness, like the tar outside and posters and our TV advert. That's the only way you're going to get those kind of people. (S26A252, Pharmacy support staff)

Capability

(4) Skills

Skills training

Several suggestions were given to improve the skill set of advisors (including counter assistants) to help with recruitment and retention in the stop smoking programme, such as reinforcement of motivational and communication skills, including how to approach people about smoking without causing offence, how to be supportive. **Most suggestions were features of person-centred care which all health care professionals are expected to follow.**[28] These were: how to personalise advice, how to elicit from smokers which things had worked and which did not work, how to work with smokers to identify from them what targets they would like to achieve and how they would achieve them.

Pharmacies are very prescriptive and I think when you're asking people to give up smoking you have to think about how they need to change their behaviour, and I think it's those ... skills that you need to apply, and that is very crucial and very important. Getting it out there. What is it they're able to do? You know what they need to do. It's only when they come up with solutions themselves they're more likely to follow it through. (S01A001, Pharmacy support staff)

In addition, a few participants stated there was need for regular training, some others suggested that it might be useful to include in any future training items such as visual scenarios, role plays or mock interviews to learn and practice their skills. Two participants also wanted to see experts in action with a client to help them learn and improve their skills.

I think sometimes you get a bit complacent. When you're doing it and you think everything's fine you give them the NRTs, tell them to come back next week if they have any problems.

1
2
3 *But they might actually have a bit of problems that they want to tell you but they can't for*
4 *whatever reason or they feel like if they tell you that they've failed then... So trying to get the*
5 *best out of their patients, so someone that could come in like a specialist to actually help,*
6 *that would be good. Because you always learn something when you're shadowing someone,*
7 *always. (S24A231, Pharmacy support staff)*
8
9
10

11 Opportunity

12 (5) Environmental context and resources

13
14
15
16 The study participants mentioned a number of structural/organisational challenges that affected
17 smoker recruitment and retention in the programme:

18 Lack of time because of delivery of other services

19
20 Participants mentioned that they forgot to remind clients about their appointment or they were
21 unable to keep tabs on clients' programme adherence. A few pharmacists, on account of being busy,
22 said they were reluctant to recruit smokers, if they were unable to provide details about the
23 programme and the procedure involved; this seemed to affect recruitment.
24
25
26

27 *Especially someone who says, "Can I join the scheme now?" and sometimes it's not possible;*
28 *when I'm so busy here, I can't leave the place and come in here. There's no way I can leave*
29 *it, so I have to give them an appointment. But sometimes people are not happy with that;*
30 *it's like they really want to join now! You get the point, they really want to join, so if you*
31 *can't do it for me now, then I can't come, so it's an issue. (S08A071, Pharmacist)*
32
33
34
35
36

37 In contrast, one pharmacy support staff participant, to avoid losing clients, started engaging people
38 on the same day of their pharmacy visit instead of giving them an appointment for a later date; this
39 strategy helped them to improve recruitment into the stop smoking programme. Another staff
40 member wanted a reminder system to enable client follow-up to help with retention.
41
42

43 *...if somebody comes in now, generally we do it straight away. That's how we found it works*
44 *better. If you give them a date they don't turn up because within this time they've got a*
45 *different commitment. (S11A102, Pharmacy support staff)*
46
47
48

49
50 To make time for smoker engagement, one pharmacist mentioned that having counter assistants to
51 help with the recruitment process or more staff to share with their dispensing workload could lead
52 them to engage and offer the stop smoking programme more frequently to smokers.
53
54
55
56
57

Inflexibility of the programme

Several participants felt that the duration of the programme was insufficient for some clients to quit and remain abstinent. One participant suggested that clients who relapsed going through the programme should be allowed to continue with the programme instead of asking them to re-join the programme, normally after a few weeks. In contrast, another participant mentioned that allowing dropouts to re-join the programme as soon as they were ready instead of using the “after six months” rule was not helpful in improving quit rates.

Other issues I've had have been people who kept coming back, like two or three times a year, and it's obvious that they are not serious, or I don't know. In the past, with the NRT, they used to say they can't do it again until after six months, but now as often as they want. But then with the Champix, I got people coming two or three times a year: stop/start, stop/start.
(S11A103, Pharmacist)

Working with budget cuts

The facility to offer two treatments concurrently to clients had been stopped due to budget cuts despite evidence of its effectiveness. In one of the participating boroughs the varenicline (Champix) license for pharmacists to prescribe had been taken away; one participant stated that this had affected recruitment into the programme.

Champix, the tablets. We haven't got a PGD (patient group direction) for it right now, so I had one person yesterday, she was a bit upset, she was like oh, I want the tablets, I want the tablets. And we don't have the licence for it. I think the whole of Newham, all the pharmacies...I think hopefully we'll be getting it back soon enough, ... So that's a problem, we've lost a few people... but we could've had those people as clients. So that's one thing, hopefully we'll try and get that back. If the providers can help us with that then that would be great. (S24A231, Pharmacy support staff)

Failing to take advantage of national campaigns

One participant felt that the ‘Stoptober’ campaign helped with smoker recruitment but not retention because they perceived smokers were not really committed to giving up smoking and only joined because of the advertisement. Some other participants felt that the No Smoking Day campaign went by too quickly to enable smoker recruitment.

Having it (No Smoking Day) spread over a week means you can maybe put up a lot more balloons up, you can put a lot more signs up, you can market it better, it gives you a longer time to market it and it will give you that whole week to engage with people. Because

15

1
2
3 *sometimes you might only see them once a day, but if you've missed that*
4 *opportunity...tomorrow's not a non-smoking day. Oh, I'm not doing it then. (S25A242,*
5 *Pharmacist)*
6
7

8
9
10 Other organisational factors such as inadequate stock of nicotine replacement products to meet
11 client demand, having only one consultation room that was often occupied were mentioned as
12 restrictions on smoker recruitment by two participants.
13

14 *...the room that I'm using is constantly being used up. I've got this one room. You know? And*
15 *things like that again – having resources [unclear] how to manage that. (S01A001, Pharmacy*
16 *support staff)*
17
18

19
20
21 **(6) Advisers' perceived characteristics of service 'joiners', 'non-joiners' and 'dropouts'**

22 Advisers' characterised smokers as those likely to join, not join or drop out of the stop smoking
23 programme. This categorisation of smokers then affected how likely the adviser would be to recruit
24 them into the stop smoking programme.
25

26
27 Eleven participants stated that 'Joiners' (including those who relapsed and wanted to quit again)
28 were those who were motivated, willing or determined to give up smoking to improve health/quality
29 of life, and were clear of their reasons for wanting to give up.
30
31

32 *Yeah, always ask them as well what's in your mind. Why do you want to do this (quit*
33 *smoking)?... So once they have that (reasons) in mind I will tell them okay, so have that in*
34 *mind, so even when the cravings come just hang onto that, because at least you have*
35 *something you want to achieve because of that. ... in fact, if you don't have it, for me you are*
36 *more likely to fail... (S19A182, Pharmacist)*
37
38
39
40

41
42 'Non-joiners' were characterised by eight participants as those who did not suffer from any health
43 related problems and hence were not ready or motivated to quit, they were either not interested or
44 in denial, thinking that they would not suffer any health-related consequences perhaps because
45 people they knew had been smoking for years and their health seemed unaffected. In contrast,
46 smokers living with a long-term condition such as cancer felt that it was too late for them to change
47 their behaviour as their illness progressed.
48
49
50

51 *Because like I said, in many years, I've seen so many patient ..., when they're smoking, end up*
52 *with a cancer. He was still coming to see me, went to the doctors, diagnosed cancer, came in*
53 *and said I'm not doing it (stopping smoking). Why? Because I've got cancer, what the hell,*
54 *I'm going to die. (S19A181, Pharmacy support staff)*
55
56
57
58

59
60
16

1
2
3
4
5 'Dropouts,' according to seven participants, were those who were fighting more than one addiction
6 or going through some personal crisis and therefore were not ready to give up smoking. Such
7 people, the participants suggested, might be easily tempted into smoking by family/friends who
8 smoke, or might not perceive any benefits or might not be interested in quitting.
9

10
11 *He was determined (to quit) because he got it from the GP, the letter, so I thought yeah, he's*
12 *going to give up smoking. But he just...the first week I supply and then the second week I*
13 *didn't see him there. Called him and he said no, I can't give up, it's not holding me now. So he*
14 *wasn't ready really, I don't think he was ready. ... (S04A032, Pharmacy support staff)*
15
16
17

18 19 **DISCUSSION**

20 **Statement of principal findings**

21
22 Recruitment and retention of smokers in the NHS stop smoking programme may be
23 influenced by advisers' preconceived ideas about smoker types likely to join, not join or drop out of
24 the programme. This early categorisation of smokers influenced perceived readiness to quit. Active
25 smoker recruitment was often a low priority, partly because advisers considered the remuneration
26 [that the pharmacy](#) received for each quitter was insufficient to justify use of time and because they
27 anticipated challenging interactions with some smokers. There were also perceived
28 structural/organisational challenges involving programme delivery.
29

30
31 To improve the programme, advisers suggested that they should adopt a more holistic and
32 supportive approach. They further suggested that strengthening their belief in the importance of
33 engagement with both new and relapsed clients would help to improve uptake of the programme
34 and retention of smokers within the service. An increase in [pharmacy remuneration](#) for quitters was
35 also thought to be beneficial. Advisers would welcome improvement in structural and organisational
36 delivery of the programme and more regular training in person-centred communication including for
37 counter assistants.
38
39

40 **Comparison to other studies**

41
42 The Theoretical Domains Framework and COM-B models have been used previously in
43 studies of dental health professionals[29-30] and midwives.[31] A barrier to delivering smoking
44 cessation services in these clinical settings was that healthcare workers did not consider smoking
45 cessation part of their primary role. In contrast, all participants in our study were trained stop
46 smoking advisers and wholeheartedly embraced the smoking cessation role. Nonetheless some
47 advisers identified the need for more training, particularly to build capability in the recruitment
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 process, to maximise engagement with smokers and to bolster their communication skills. This need
4 for training was also felt by dental health professionals and midwives. The fear of negative patient
5 response, inadequate staffing, and lack of confidence in delivery of health behaviour advice which
6 we identified in our study has also been reported in other settings.[13, 32-33]
7
8

9
10 In line with the recent review on the effectiveness of pharmacy smoking cessation
11 interventions,[6] we found that many advisers were mostly accepting smokers who wanted to quit
12 because they did not want to risk dropouts. The National Centre for Smoking Cessation and Training
13 also recommends that if the smoker is not ready to make a serious quitting attempt, the adviser
14 should provide the programme's contact details and ask them to get in touch when they are
15 ready.[34] Another explanation for selection of those likely to quit may be pressure from service
16 targets.[35]
17
18
19

20
21 The perceived opportunity cost of engaging with smokers who did not express a desire to
22 quit or who fitted stereotypes that advisers associated with failure to quit, appeared to affect
23 adversely the recruitment of smokers into the service. Hoving[36] compared 'active' recruitment
24 (i.e. "asking each individual's smoking status and inviting smokers to participate in the service") and
25 passive (i.e. "leaving participation up to the smoker's initiative") in a community pharmacy setting.
26 The former approach was suggested as a way of increasing the total number of quitters. It has been
27 suggested elsewhere that passive recruitment strategies might reduce the likelihood of a quit
28 attempt.[37]
29
30
31
32
33

34 In addition to lack of time for health promotion which has been reported previously,[13, 32,
35 33, 38] problems with delivery of medicines and promotional material, linked to national stop
36 smoking campaigns, were perceived by advisers to affect smoker recruitment and retention.
37 Addressing these issues is crucial to facilitate adviser engagement behaviour,[39-40] and could add
38 to the success of national campaigns in increasing the number of people attempting to quit and
39 permanently quitting.[39, 41-42]
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Implications for clinical practice and policy makers

Advisers' perceptions of types of smokers who are likely not to join or to drop out of the programme and some advisers' experience of challenging interactions with smokers have been reported previously[33] including in socially deprived areas.[43-44] However, the community pharmacy stop smoking services in east London are effective in reaching socio-economically deprived communities[35] and the smokers in these communities are just as motivated to quit smoking as smokers in more affluent areas,[35, 45] and to want help with quitting.[46] Smokers' stage of change[47] and economic disadvantage should not preclude the use of active recruitment[8, 35, 45] and retention strategies.

Recent changes to the pharmacy smoking cessation services in one of the participating boroughs meant that pharmacists were no longer able to recommend co-therapy (nicotine patch plus gum/lozenge/inhalator) or varenicline to clients which advisers considered a problem. It seems likely that restricting access to effective treatment and thus compromising patient choice could adversely affect engagement of smokers and quit rates. In addition, several advisers felt the duration of the stop smoking programme was not long enough for some smokers to quit and to remain abstinent and suggested having flexibility in the programme to allow continuity of care, where appropriate to suit clients' needs, for example after relapse.

The remuneration for programme delivery is determined by commissioners on the basis of clear criteria e.g. the time and duration of intervention and treatment provided, carbon monoxide (CO) monitoring and data reporting. The payment is made to the pharmacy owner i.e. a contractor who may be a pharmacist sole trader, a partner with other pharmacists or who may appoint a superintendent pharmacist[2, 48] No payment is made to employed pharmacists or to pharmacy support staff. The current payment system for pharmacies appears to provide a disincentive to spend time with smokers - and this has been acknowledged in the community pharmacy stop smoking services guidance.[2]

The pharmacy cessation advisers derived professional satisfaction in helping individual smokers to quit and also saw their role in smoking cessation as benefitting the whole community. Advisers thought about wider aspects of the scheme and took a professional interest in how the scheme might develop – for example suggesting a voucher scheme and tailoring the duration of the smoking cessation programme to individual smokers. This source of intrinsic motivation may distinguish cessation advisers from other healthcare professionals and could be drawn upon further in pharmacy adviser training programmes.

Implications for future research

More research is needed with a broader range of participants including female advisers and advisers of other ethnicities delivering the programme in deprived and non-deprived areas. In addition, studies to understand in more detail the effects of the remuneration structure for pharmacists and support staff is warranted. We took recruitment and retention together in the analysis since we were interested in the joint outcome of these two behaviours, which leads to an increase in the total number of people completing the smoking cessation programme. Whilst this seems justified given the purpose of our work this is a potential limitation of our study since clearly the two behaviours may have different antecedents. Future work could usefully examine recruitment and retention separately leading to individual insights into these behaviours and how they might be modified to strengthen the NHS smoking cessation service.

Acknowledgments The authors thank participants and their community pharmacy. This paper presents independent research funded by the National Institute for Health Research (NIHR) under its Programme Grants of Applied Research Programme. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Contributors RW conceived the study. RS was involved in the data collection, data analysis and interpretation and writing the manuscript. VM was involved in the data collection. CR and LS were involved in the data analysis and interpretation. CR, VM, LS, EE, VK, CG, ST, SE and RW were involved in revising the article critically for important intellectual content. RW is the guarantor.

Competing interests None.

Funding: NIHR Programme grant RP-PG-0609-10181

Participant consent Written informed consent obtained.

Ethics approval Ethical approval for the study was obtained from the NRES Committee South Central - Berkshire B (reference number: 13/SC/0189).

Provenance and peer review Not commissioned; externally peer reviewed

1
2
3 **Data sharing statement** This study analyses qualitative data and the advisors did not consent to
4 have their full transcripts made publicly available. The advisors consented for their data to be stored
5 for five years, as part of the 5-year programme grant project and to be used in this or other studies
6 directly linked to the 5-year project. Supporting excerpts from the raw data (quotes from participant
7 interviews) are available with the text of the paper. Full transcripts of the interviews (with
8 identifying information removed) will be available, following completion of the project, on request
9 from the study guarantor, Robert Walton, r.walton@qmul.ac.uk
10
11
12
13

14
15
16 **Open Access** This is an Open Access article distributed in accordance with the terms of the Creative
17 Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build
18 upon this work, for commercial use, provided the original work is properly cited. See:
19 <http://creativecommons.org/licenses/by/4.0/>
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. NHS England. Improving health and patient care through community pharmacy - a call to action 2013. <http://www.england.nhs.uk/wp-content/uploads/2013/12/community-pharmacy-cta.pdf> (accessed 24 Feb 2015)
2. NHS Employers. *Pharmacy-based stop smoking services: optimising commissioning*. London: NHS Employers, 2009.
3. World Health Organization. *Tobacco Free Initiative*. 2015. http://www.who.int/tobacco/health_priority/en/ (accessed 6 Feb 2015)
4. Brose LS, West R, Michie S, et al. Changes in success rates of smoking cessation treatment associated with of a national evidence-based training programme. *Prev Med* 2014;69:1-4.
5. Prescribing and Primary care Health and Social Care Information Centre. *General Pharmaceutical Services England 2004-05 to 2013-14* Leeds 2014.
6. Mdege ND, Chindove S. Effectiveness of tobacco use cessation interventions delivered by pharmacy personnel : A systematic review. *Res Social Adm Pharm* 2014;10(1):21-44.
7. McDonald PW. Population-based recruitment for quit-smoking programs: an analytic review of communication variables. *Prev Med* 1999;28:545-57
8. Szklo AS. Review of strategies to recruit smokers for smoking cessation: a population impact perspective. *Cad Saude Publica* 2008;24:S621-S34.
9. Action on smoking and health. Facts at a glance. Smoking statistics. 2015. www.ash.org.uk/files/documents/ASH_93.pdf (accessed 24 Feb 2015)
10. Lifestyle statistics Health and Social Care Information Centre. *Statistics on NHS Stop Smoking Services: England, April 2012 – March 2013*, 2013.
11. Department of Health. *Stop Smoking Service Monitoring and Guidance Update 2012/13*. London, 2012.
12. Bevan P. Annual Report of the Director of Public Health – Health at the Heart of the Community. 2014. <http://democracy.cityoflondon.gov.uk/documents/s35538/DPH%20report%20to%20Health%20Social%20Care%20Scrutiny%20sub%20committee.pdf> (accessed 24 Feb 2015)
13. Eades C, Ferguson J, O'Carroll R. Public health in community pharmacy: A systematic review of pharmacist and consumer views. *BMC Public Health* 2011;11:582.
14. Pope C, Ziebland S, Mays N. Qualitative research in health care. Analysing qualitative data. *BMJ* 2000;320:114-6.
15. Rabiee F. Focus-group interview and data analysis. *Proc Nutr Soc* 2004;63:655-60.

22

- 1
2
3 16. Cane J, O'Connor D, Michie S. Validation of the theoretical domains framework for use in
4 behaviour change and implementation research. *Implement Sci* 2012;7:37.
5
6 17. Michie S, van Stralen MM, West R. The behaviour change wheel: A new method for
7 characterising and designing behaviour change interventions. *Implement Sci* 2011;6:42.
8
9 18. Michie S, Atkins L, West R. The Behaviour Change Wheel. A Guide to Designing Interventions.
10 London: Silverback Publishing, 2014.
11
12 19. National Centre for Smoking Cessation and Training. http://www.ncsct.co.uk/pub_training.php
13 (accessed 7 Aug 2015)
14
15 20. National Institute for Health and Care Excellence. *Training in smoking prevention and cessation*.
16 NICE, 2015.
17
18 21. National Pharmacy Association. <http://www.npa.co.uk/> (accessed 7 Aug 2015)
19
20 22. Robinson F. *Smoking cessation: Training providers*. 2014. [https://www.pcrs-uk.org/training-](https://www.pcrs-uk.org/training-providers)
21 providers (accessed 7 Aug 2015)
22
23 23. Pharmaceutical Services Negotiating Committee. *NHS Community Pharmacy Services - a*
24 *summary* 2013. [http://psnc.org.uk/wp-content/uploads/2013/08/CPCF-summary-July-](http://psnc.org.uk/wp-content/uploads/2013/08/CPCF-summary-July-2013.pdf)
25 2013.pdf (accessed 19 Jun 2015)
26
27 24. Pharmaceutical Services Negotiating Committee. *Public Health. A practical guide for community*
28 *pharmacists* 2013. [http://psnc.org.uk/wp-](http://psnc.org.uk/wp-content/uploads/2013/07/public_health_book_all.pdf)
29 content/uploads/2013/07/public_health_book_all.pdf (accessed 24 Feb 2015)
30
31 25. Stephenson-Glynn J. JSNA Factsheet: *Health and Wellbeing Tobacco Control Tower Hamlets Joint*
32 *Strategic Needs Assessment 2013-2015*.
33 [http://www.towerhamlets.gov.uk/idoc.ashx?docid=026c2fee-f37f-4479-8d6e-](http://www.towerhamlets.gov.uk/idoc.ashx?docid=026c2fee-f37f-4479-8d6e-4b9660a1e282&version=1)
34 [4b9660a1e282&version=1](http://www.towerhamlets.gov.uk/idoc.ashx?docid=026c2fee-f37f-4479-8d6e-4b9660a1e282&version=1) (accessed 24 Feb 2015)
35
36 26. Flowers R. *Newham joint strategic needs assessment 2011/12 September 2012 Update*.
37 [http://www.newham.gov.uk/Documents/Council%20and%20Democracy/JSNASept2012Upd](http://www.newham.gov.uk/Documents/Council%20and%20Democracy/JSNASept2012Update.pdf)
38 [ate.pdf](http://www.newham.gov.uk/Documents/Council%20and%20Democracy/JSNASept2012Update.pdf) (accessed 24 Feb 2015).
39
40 27. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a
41 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19:349-57.
42
43 28. Foot C, Gilbert H, Dunn P et al. *People in control of their own health and care. The state of*
44 *involvement*. King's Fund, London, 2014.
45
46 29. Amemori M, Michie S, Korhonen T, et al. Assessing implementation difficulties in tobacco use
47 prevention and cessation counselling among dental providers. *Implement Sci* 2011;6(1):50.
48
49 30. Amemori M, Korhonen T, Michie S, et al. Implementation of tobacco use cessation counseling
50 among oral health professionals in Finland. *J Public Health Dent* 2013;73:230-36.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 31. Beenstock J, Sniehotta F, White M, et al. What helps and hinders midwives in engaging with
4 pregnant women about stopping smoking? A cross-sectional survey of perceived
5 implementation difficulties among midwives in the North East of England. *Implement Sci*
6 2012;7(1):36.
7
8
9
10 32. Patwardhan PD, Chewning BA. Ask, advise and refer: Hypothesis generation to promote a brief
11 tobacco-cessation intervention in community pharmacies. *Int J Pharm Pract* 2009;17:221-29.
12
13 33. Morton K, Pattison H, Langley C, et al. A qualitative study of English community pharmacists'
14 experiences of providing lifestyle advice to patients with cardiovascular disease. *Res Social*
15 *Adm Pharm* 2015;11:e17-e29.
16
17 34. West R, McEwen A, Lorencatto F, et al. *NC SCT Training Standard. Learning Outcomes for Training*
18 *of Stop Smoking Practitioners* In: McEwen A, ed.: National Centre for Smoking Cessation and
19 Training, London, 2013.
20
21 35. Hiscock R, Bauld L. *Stop Smoking Services and Health Inequalities*. In: McEwen A, ed.: National
22 Centre for Smoking Cessation and Training (NCSCT), London 2013.
23
24 36. Hoving C, Mudde AN, de Vries H. Effect of recruitment method and setting on the composition of
25 samples consisting of adult smokers. *Patient Educ Couns* 2007;65(1):79-86.
26
27 37. Hammond D, McDonald PW, Fong GT, et al. Do smokers know how to quit? Knowledge and
28 perceived effectiveness of cessation assistance as predictors of cessation behaviour.
29 *Addiction* 2004;99:1042-48.
30
31 38. McDermott MS, Beard E, Brose LS, et al. Factors Associated With Differences in Quit Rates
32 Between "Specialist" and "Community" Stop-Smoking Practitioners in the English Stop-
33 Smoking Services. *Nicotine Tob Res* Published Online First: 13 December 2012. doi:
34 10.1093/ntr/nts262
35
36 39. Brown J, Kotz D, Michie S, et al. How effective and cost-effective was the national mass media
37 smoking cessation campaign 'Stoptober'? *Drug Alcohol Depend* 2014;135:52-58.
38
39 40. Kmietowicz Z. Action is needed to boost uptake of stop smoking services, say campaigners. *BMJ*
40 Published Online First: 23 August 2015. doi:10.1136/bmj.h4578
41
42 41. Kotz D, Stapleton JA, Owen L, et al. How cost-effective is 'No Smoking Day'? *Tob Control*
43 2011;20:302-04.
44
45 42. Owen L, Youdan B. 22 years on: the impact and relevance of the UK No Smoking Day. *Tob Control*
46 2006;15:19-25.
47
48 43. Oakes W, Chapman S, Borland R, et al. "Bulletproof skeptics in life's jungle": which self-
49 exempting beliefs about smoking most predict lack of progression towards quitting? *Prev*
50 *Med* 2004;39:776-82.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 44. Paul CL, Ross S, Bryant J, et al. The social context of smoking: A qualitative study of comparing
4 smokers of high versus low socioeconomic position. *BMC Public Health* 2010;10:211
5
6 45. Cunningham P. Patient engagement during medical visits and smoking cessation counseling.
7
8 *JAMA Intern Med* 2014;174:1291-98.
9
10 46. Hudmon KS, Prokhorov AV, Corelli RL. Tobacco cessation counseling: Pharmacists' opinions and
11 practices. *Patient Educ Couns* 2006;61:152-60.
12
13 47. Norcross JC, Krebs PM, Prochaska JO. Stages of Change. *J Clin Psychol* 2011;67:143-54.
14
15 48. Prescribing and Primary Care Team HSCIC. *General Pharmaceutical Services in England: 2004-05*
16 *to 2013-14*. Health and Social Care Information Centre, Leeds, 2014.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Figure legends**

4 Box 1. Outline of the stop smoking training and the community pharmacy stop smoking programme

5
6 Box 2. Interview schedule

7
8 Figure 1. Illustration of behavioural factors affecting adviser engagement and retention behaviour

9
10 (modified from the COM-B model)

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only