

# PCCP

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: S. J. Elliott, L. J. Brown, J. Dumez and M. H. Levitt, *Phys. Chem. Chem. Phys.*, 2016, DOI: 10.1039/C6CP03619H.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

# Long-Lived Nuclear Spin States in Monodeuterated Methyl Groups

Stuart J. Elliott<sup>a</sup>, Lynda J. Brown<sup>a</sup>, Jean-Nicolas Dumez<sup>b</sup> and Malcolm H. Levitt<sup>a</sup>

<sup>a</sup>*School of Chemistry, University of Southampton, Southampton, SO17 1BJ, United Kingdom*

<sup>b</sup>*Institut de Chimie des Substances Naturelles, CNRS UPR2301, Univ. Paris-Sud, Université Paris-Saclay, 91190 Gif-sur-Yvette, France*

---

## Abstract

It is possible to access long-lived nuclear singlet order in monodeuterated methyl groups, in the case that a significant chemical shift difference exists between the CH<sub>2</sub>D protons. This occurs when the local environment is chiral, and the CH<sub>2</sub>D rotamers have different populations. An experimental demonstration is presented for the case of N-CH<sub>2</sub>D-2-methylpiperidine. The ratio of the singlet relaxation time constant  $T_S$  to the longitudinal relaxation time constant  $T_1$  is found to be equal to  $3.1 \pm 0.1$ , over a wide range of temperatures, solvents, and magnetic fields. The longest observed value of  $T_S$  approaches 1 minute. The relaxation mechanisms of the long-lived state are discussed, and a modified model of the CH<sub>2</sub>D geometry is proposed to explain the observed ratio of  $T_S$  to  $T_1$ .

**Keywords:** long-lived states, singlet state, methyl group, rotamer

---

## 1. Introduction

Long-lived states (LLS) are configurations of nuclear spins which are protected against relaxation in nuclear magnetic resonance (NMR) experiments [1–22]. Long-lived state lifetimes exceeding the conventional relaxation time  $T_1$  by a factor of 50 have been observed, with a lifetime exceeding 1 hour in room-temperature solution in one case [22]. The combination of LLS phenomena with hyperpolarization techniques is particularly promising. Techniques such as dissolution-DNP (dynamic nuclear polarization) [23] offer

large NMR signal enhancements, with wide-ranging implications, including the characterization of cancer in human patients [24]. However, the range of hyperpolarization applications is restricted by the finite lifetime of the enhanced magnetization, which is normally limited by  $T_1$ . The use of LLS offers a promising means to transcend this limitation [9, 25–28].

In systems of spin-1/2 pairs, the LLS is called *singlet order*. This consists of the population imbalance between the spin-0 singlet state and the spin-1 triplet states [1, 2]. The decay time constant of singlet order is denoted  $T_S$ . Access to nuclear singlet order is provided by a chemical shift difference between the participating spins, or by differences in spin-spin couplings to spins outside the pair [13].

LLS have also been observed in the 3-spin-1/2 systems of rapidly rotating methyl groups in solution [29–31]. In this case the LLS is given by the imbalance in populations between spin states spanning different irreducible representations of the  $C_3$  permutation group [32]. Some materials, such as  $\gamma$ -picoline, display quantum-rotor induced polarization (QRIP) effects, in which a large polarization of the methyl LLS is induced by dissolution of the material from cryogenic conditions [29, 33–35]. However, the hyperpolarized LLS only gives rise to observable NMR signals through an incoherent cross-relaxation mechanism involving a fourth nuclear spin, which greatly reduces the available signal enhancement [29–31].

Since methyl groups are ubiquitous in nature, the exploitation of methyl LLS is potentially attractive. In this communication we show that in certain cases, it is possible to achieve coherent access to a methyl LLS with a high conversion efficiency into observable NMR signals. The conditions are: (i) the methyl group is monodeuterated, and therefore contains a proton pair; (ii) the local environment is chiral, and (iii) the three methyl rotamers have sufficiently different populations, so that a small chemical shift difference is induced between the  $\text{CH}_2\text{D}$  protons after averaging over all populated states. We show that the small chemical shift difference allows coherent (and therefore efficient) access to the long-lived singlet order of the proton spin pair, using known radiofrequency pulse techniques [11, 12, 19, 20].

Chemical inequivalence of  $\text{CH}_2\text{D}$  protons has been observed in several molecular systems, most of which contain a  $\text{N-CH}_2\text{D}$  group in a chiral environment [36–38]. In this communication we report the observation of a proton LLS in the  $\text{N-CH}_2\text{D}$  group of  $\text{N-CH}_2\text{D-2-methylpiperidine}$  (Figure 1). We observe singlet lifetimes  $T_S$  with values between 20 and 55 seconds (dependent on temperature). The ratio of singlet relaxation time  $T_S$  to longi-

tudinal relaxation time  $T_1$  is found to be remarkably constant over a wide range of conditions, and equal to  $3.1 \pm 0.1$ . This result shows that is feasible to exploit methyl LLS without relying on weak cross-relaxation effects, in suitable cases. We provide evidence that singlet relaxation in this system is dominated by the dipolar interactions between the  $\text{CH}_2\text{D}$  protons and the  $\text{CH}_2\text{D}$  deuteron, and show that the singlet relaxation times cannot be explained by a model in which the three hydrogen nuclei are localised at the vertices of an equilateral triangle. We propose a modified geometrical model which is consistent with the experimental data.

## 2. Experiments

### 2.1. Proton Spectra

The methyl region of the proton NMR spectrum of (N- $\text{CH}_2\text{D}$ )-2-methylpiperidine is shown in Figure 1. This spectral region has a quartet-like appearance in the absence of deuteron decoupling (Figure 1a). The characteristic AB spectral pattern of an inequivalent proton pair appears when a deuteron decoupling field is used to remove the  $^2J_{\text{HD}}$  splittings (Figure 1b). This spectrum is consistent with a J-coupling of  $|^2J_{\text{HH}}| = 11.7 \pm 0.2$  Hz and a chemical shift difference of  $\Delta\delta = 13.5 \pm 0.4$  ppb between the  $\text{CH}_2\text{D}$  protons, as reported previously [36].

The existence of a small chemical shift difference has been attributed to (i) hyperconjugation between the nitrogen lone pair and the *anti*-methyl C-H(D)  $\sigma$ -bond, which allows the zero-point vibrational energies of the *anti* CH and CD bonds to influence the rotamer energies; as a result, the rotamer with the CD bond *anti* to the nitrogen lone pair is less populated than the other two rotamers in thermal equilibrium, and (ii) the chiral environment associated with the neighbouring methyl group, which causes a significant chemical shift difference between the two protons in each rotamer [36, 38]. In these circumstances, there remains a significant chemical shift difference between the  $\text{CH}_2\text{D}$  protons after averaging over all methyl rotamers. The observed chemical shift difference decreases as the temperature is increased, as shown in Figure 2a. This is consistent with the Boltzmann populations of the three rotamers becoming more similar at higher temperatures.

### 2.2. Singlet NMR

The small chemical shift difference allows access to the long-lived singlet order between the  $\text{CH}_2\text{D}$  protons by using the M2S [11, 12] or SLIC [19] pulse




Figure 1: Part of the experimental  $^1\text{H}$  spectrum of (N-CH<sub>2</sub>D)-2-methylpiperidine in CD<sub>2</sub>Cl<sub>2</sub> solution acquired at 11.7 T (500 MHz) with a single transient (for the full proton spectrum, see the Supporting Information). (a) Spectrum without deuteron decoupling; (b) Spectrum with deuteron decoupling (deuteron nutation frequency 500 Hz). The asterisk indicates a small signal from a non-deuterated N-methyl-2-methylpiperidine impurity, shifted in frequency by a secondary isotope effect. Inset: dominant di-equatorial conformation [36] of N-CH<sub>2</sub>D-2-methylpiperidine in solution, showing the nitrogen lone pair which is implicated in the inequivalence of the CH<sub>2</sub>D protons.

sequences, and their variants [20, 39]. In the current study, we used a variant of the SLIC (Spin-Lock-Induced Crossing) method, as shown in Figure 3.

This pulse sequence operates as follows: after the initial 90° pulse, a radiofrequency field is applied with a 90° phase shift. The amplitude of this field is selected so that the nutation frequency  $\omega_{\text{SLIC}}/2\pi$  matches the J-coupling  $^2J_{\text{HH}}$ . This establishes a resonance which causes the spin-locked magnetization to be converted into singlet order through the action of the chemical shift difference, with the conversion complete in a time  $\tau_{\text{SLIC}} \simeq 2^{-1/2}\Delta\nu^{-1}$ , where  $\Delta\nu$  is the chemical shift difference in Hertz, neglecting relaxation and other complications [19]. The singlet order is allowed to evolve for a variable interval  $\tau_{\text{EV}}$  under a larger-amplitude “spin-locking” rf field (nutation frequency  $\omega_{\text{LOCK}}/2\pi$ ), which suppresses singlet-triplet mixing [8].

A sequence of radiofrequency pulses and pulsed field gradients (known as a “T<sub>00</sub> filter”) suppresses NMR signals that do not pass through singlet order [40, 41]. A second SLIC sequence converts the singlet order into transverse proton magnetization, and the induced NMR signal is detected. The maximum amplitude of the singlet-filtered  $^1\text{H}$  NMR signal, relative to that induced by a single 90° pulse, was found to be 0.43, somewhat lower than the




Figure 2: (a) Chemical shift difference  $\Delta\delta$  between the  $\text{CH}_2\text{D}$  protons, and (b) singlet relaxation time  $T_S$  of 0.1 M N- $\text{CH}_2\text{D}$ -2-methylpiperidine in degassed  $\text{C}_6\text{D}_6$  solution, as a function of temperature. Experiments were performed at fields of (a) 11.7 T (500 MHz), and (b) 14.1 T (600 MHz).

theoretical maximum of  $2/3$  [42], the loss being attributed to radiofrequency field imperfections and relaxation. The decay of the long-lived singlet order is tracked by repeating the pulse sequence with different values of the singlet evolution interval  $\tau_{\text{EV}}$ . Experimental parameters and details of sample preparation are found in the Supporting Information (SI).




Figure 3: Pulse sequence used for accessing long-lived singlet order in monodeuterated methyl groups and measuring its decay. The experiments used the following parameters: nutation frequencies  $\omega_{\text{SLIC}}/2\pi = 11.7$  Hz and  $\omega_{\text{LOCK}}/2\pi = 300$  Hz. The duration of the SLIC pulse was 100 ms in the 500 MHz experiments and 73 ms in the 600 MHz experiments. The “ $T_{00}$  filter” sequence suppresses signals that do not pass through singlet order, as described in the Supporting Information. “MA” denotes the “magic angle” ( $54.7^\circ$ ). An interval of 90 s was used between successive transients.

A typical decay curve for singlet order is shown in Figure 4. This shows a single exponential decay with time constant  $T_S = 27.0 \pm 0.6$  s. This is approximately three times longer than the relaxation time for longitudinal



magnetization  $T_1 = 8.7 \pm 0.1$  s, as estimated from the inversion-recovery curve, also shown in Figure 4.




Figure 4: Experimental relaxation curves for 0.1 M N-CH<sub>2</sub>D-2-methylpiperidine in degassed CD<sub>2</sub>Cl<sub>2</sub> solvent (proton frequency 500 MHz, temperature 25°C). Open symbols, grey line, and right-hand axis: Decay of singlet order measured by the pulse sequence in Figure 3. Filled symbols, black line, and left-hand axis: Spin-lattice relaxation measured by inversion-recovery. All signal amplitudes were normalized to the first point. The fitted curves have a single-exponential form.

The singlet relaxation time constants  $T_S$  increase with increasing temperature, as shown in Figure 2b. Measured relaxation time constants  $T_1$  and  $T_S$  are presented for a variety of solvents, temperatures and magnetic fields in the Supporting Information. Figure 5 shows a plot of  $T_1^{-1}$  against  $T_S^{-1}$  for this disparate data set. The fit to a straight line with zero intercept and slope  $(3.1)^{-1} = 0.324$  is remarkably good. The ratio of  $T_S$  to  $T_1$  is remarkably consistent and given by  $3.1 \pm 0.1$  over a wide range of conditions.

The data shown in Figure 5 were all obtained for (N-CH<sub>2</sub>D)-2-methylpiperidine, except for a single point which was obtained for a compound with complete deuteration of the second methyl group, i.e. (N-CH<sub>2</sub>D)-2-CD<sub>3</sub>-piperidine (purple triangle). Clearly, deuteration of the second methyl group in N-CH<sub>2</sub>D-2-methylpiperidine does not have a strong effect on the relaxation behaviour.

### 2.3. Carbon-13 and Deuterium NMR

<sup>13</sup>C and <sup>2</sup>H spectra of N-CH<sub>2</sub>D-2-methylpiperidine, and  $T_1$  values for the ring <sup>13</sup>C sites, are shown in the Supporting Information, for the case of degassed CD<sub>2</sub>Cl<sub>2</sub> solution at 11.7 T (500 MHz) and 25°C. The ring CH<sub>2</sub> sites have similar <sup>13</sup>C relaxation time constants  $T_1$  of  $6.5 \pm 0.3$  s, with the ring CH site displaying a longer <sup>13</sup>C relaxation time constant of  $10.2 \pm 0.3$  s.

Under the same conditions, the  $^2\text{H}$  relaxation time constant for N-CH<sub>2</sub>D-2-methylpiperidine is given by  $T_1(^2\text{H}) = 0.75 \pm 0.01$  s.




Figure 5: Longitudinal relaxation rate constants ( $T_S^{-1}$ ) plotted against singlet relaxation rate constants ( $T_1^{-1}$ ) for 0.1 M N-CH<sub>2</sub>D-2-methylpiperidine over a wide range of solvents (degassed), temperatures and magnetic fields. The experimental conditions are as follows: (●) C<sub>6</sub>D<sub>6</sub> solution, at temperatures increasing from 25°C (rightmost point) to 60°C (leftmost point) in 5°C increments, at a field of 14.1 T (600 MHz); (○) C<sub>6</sub>D<sub>6</sub> solution, at magnetic fields increasing from 9.4 T (400 MHz, leftmost point) to 14.1 T (600 MHz, rightmost point) in 100 MHz increments, at a temperature of 25°C; (▲) CH<sub>2</sub>Cl<sub>2</sub> solution at 25°C and 11.7 T (500 MHz); (●) C<sub>6</sub>D<sub>6</sub> solution at 25°C and 11.7 T (500 MHz); (■) CD<sub>3</sub>CN solution at 25°C and 11.7 T (500 MHz); (▲) CD<sub>2</sub>Cl<sub>2</sub> solution at 25°C and 11.7 T (500 MHz); (◆) CD<sub>3</sub>OD solution at 25°C and 11.7 T (500 MHz); (▲) CD<sub>2</sub>Cl<sub>2</sub> solution of (N-CH<sub>2</sub>D)-2-(CD<sub>3</sub>)-piperidine at 25°C, 11.7 T (500 MHz). A constant ratio  $T_S/T_1 = 3.1 \pm 0.1$  is observed over a wide range of experimental conditions.

### 3. Relaxation Theory

The data show that long-lived singlet order may be accessed coherently, and with high efficiency, for the proton pair of the monodeuterated methyl group in N-CH<sub>2</sub>D-2-methylpiperidine. The singlet lifetime  $T_S$  is proportional to  $T_1$  of the same protons, with the proportionality constant given by  $3.1 \pm 0.1$ , over a wide range of experimental conditions. In this section we propose an explanation for the constant ratio of  $T_S$  to  $T_1$ , and investigate the value of the proportionality constant.

#### 3.1. Relaxation Mechanisms

A large variety of relaxation mechanisms may contribute to the proton  $T_S$  and  $T_1$  relaxation of the CH<sub>2</sub>D protons. As well as the magnetic dipole-dipole



interactions between the three magnetic nuclei of the CH<sub>2</sub>D group, there are also chemical shift anisotropy contributions, magnetic dipole-dipole interactions with other nuclei in the same molecule, and intermolecular dipole-dipole interactions. The quadrupolar relaxation of the deuterium nuclei might also contribute to proton relaxation through mechanisms such as scalar relaxation of the second kind [43, 44]. Furthermore, spin-rotation relaxation is known to be significant for many rotating methyl groups [45–48], and contributes to the LLS relaxation of  $\gamma$ -picoline in solution [29, 30]. Singlet relaxation may also be caused by state mixing due to a finite chemical shift difference between the participating protons, or by asymmetric J-couplings to other magnetic nuclei (“singlet-triplet leakage”) [14].

The excellent correlation between the  $T_S$  and  $T_1$  values shown in Figure 5 strongly supports the hypothesis that the longitudinal and singlet relaxation of the CH<sub>2</sub>D proton pair is driven by a common mechanism, with a common correlation function. This suggests that the mechanisms that dominate the  $T_1$  and  $T_S$  relaxation of the CH<sub>2</sub>D protons are internal to the CH<sub>2</sub>D group. This conclusion is supported by the following observations: (i) Deuteration of the 2-methyl group leads to only a small change in the relaxation times even though the protons of the 2-methyl group approach to within 229 pm of the CH<sub>2</sub>D protons; (ii) a large change in the deuteration level of the solvent only has a small influence on the relaxation times (see Figure 5 and the data in the Supporting Information).

In addition, one may discount major contributions from the spin-rotation, scalar relaxation, chemical shift anisotropy and singlet-triplet leakage mechanisms, for the following reasons:

1. *Spin-rotation.* The experimental finding that  $T_S$  increases with increasing temperature (Figure 2b) speaks strongly against a significant contribution from spin-rotation, since that mechanism usually increases in strength with increasing temperature [45–48]. Presumably, in the current case, the methyl rotation is too strongly hindered to permit a significant spin-rotation relaxation contribution.
2. *Scalar relaxation of the second kind.* The existence of a resolved deuteron splitting in the proton spectrum (Figure 1a), as well as direct measurements of deuteron relaxation (see section 2.3) indicate that deuteron relaxation is too slow to induce significant scalar relaxation of the coupled protons. Furthermore, the two  $^2J_{HD}$  couplings are identical to a good approximation, which precludes a scalar contribution to singlet

relaxation [44].

3. *Chemical shift anisotropy.* The weak dependence of  $T_1$  and  $T_S$  on magnetic field indicates a relatively small contribution from CSA.
4. *Singlet-triplet leakage.* Singlet-triplet leakage induced by the small chemical shift difference between the  $\text{CH}_2\text{D}$  protons can be a significant contribution to the singlet relaxation rate constant  $T_S^{-1}$  [14]. However, in the current experiments, this contribution is suppressed very effectively by the application of a resonant radiofrequency field during the singlet relaxation interval (see the Supporting Information).

We therefore postulate that both the singlet and the longitudinal relaxation of the proton pair in the monodeuterated methyl group of N- $\text{CH}_2\text{D}$ -2-methylpiperidine are dominated by the  $^1\text{H}$ - $^1\text{H}$  and  $^1\text{H}$ - $^2\text{D}$  dipole-dipole interactions within the methyl group itself, modulated by the internal rotation of the methyl group with respect to the rest of the molecule, and by the rotation of the molecule as a whole. A relaxation model is therefore constructed based on (i) a simplified description of the motion of the methyl group and the molecule as a whole, and (ii) a description of the vibrationally-averaged spin-spin interactions within the rotating methyl group.

### 3.2. Motional Model

A plausible motional model for the monodeuterated methyl group in N- $\text{CH}_2\text{D}$ -2-methylpiperidine consists of a hindered 3-fold rotor attached to a sphere undergoing isotropic rotational diffusion (Figure 6). Although the existence of a finite chemical shift difference between the  $\text{CH}_2\text{D}$  protons implies that the methyl rotamers have slightly unequal populations, we ignore this effect in the relaxation analysis, for the sake of simplicity. The thermally activated jumps between the methyl rotamers are assumed to be described by a rate constant  $\kappa$ , while the overall rotational diffusion of the molecule is described by a correlation time  $\tau_c$ . Both  $\kappa$  and  $\tau_c$  are in general temperature-dependent. We also assume that the correlation time  $\tau_c$  is short enough relative to the nuclear Larmor period to invoke the extreme narrowing approximation [43].

### 3.3. Dipole-Dipole Interactions

The three magnetic nuclei in the  $\text{CH}_2\text{D}$  group interact by the magnetic dipole-dipole interaction. In general these dipole-dipole interactions are described by traceless second-rank tensors, and are subjected to averaging over




Figure 6: The relaxation of the CH<sub>2</sub>D protons in N-CH<sub>2</sub>D-2-methylpiperidine is governed by the three-fold jump rate  $\kappa$  and overall rotational correlation time  $\tau_c$ .

local molecular vibrations and librations on a timescale fast compared to the methyl 3-fold jumps or the molecular tumbling. Spin relaxation is caused by motional modulation of the vibrationally-averaged dipole-dipole interaction tensors.

In general, the vibrationally-averaged tensors differ, both in magnitude and in orientation, from tensors derived from a naive geometrical model, for which point-like nuclei are located at the vertices of an equilateral triangle, with the magnitudes of the dipole-dipole interactions in exact proportion to the product of the gyromagnetic ratios, since the internuclear distances are all equal. As discussed below, this naive equilateral model of the CH<sub>2</sub>D group is inconsistent with the experimental results.

To maintain high generality, the relaxation theory is developed using three different interaction tensors for the vibrationally-averaged dipole-dipole interactions, leaving the magnitudes and orientations of the tensors as adjustable parameters. The two <sup>1</sup>H-<sup>2</sup>D interaction tensors are assumed to have the same principal values, by symmetry. For the sake of simplicity, the vibrationally-averaged interaction tensors are assumed to be axially symmetric, with the unique principal axes perpendicular to the N-C rotor axis. The angle between the unique principal axes of the two vibrationally-averaged <sup>1</sup>H-<sup>2</sup>D interaction tensors is denoted  $2\theta$ . This angle defines the cross-correlation of the two <sup>1</sup>H-<sup>2</sup>D interaction tensors, and is therefore important for the proton singlet relaxation. By symmetry, the unique principal axis of the vibrationally-averaged <sup>1</sup>H-<sup>1</sup>H interaction tensor is perpendicular to the bisector of the two <sup>1</sup>H-<sup>2</sup>D principal axes (see the Supporting Information). A point-nucleus equilateral geometry model would lead to the angle  $2\theta = 60^\circ$ , but this value is not

assumed in the following discussion.

### 3.4. Rate Expressions

As shown in the Supporting Information, the motional and intra-methyl interaction models described above lead to the following expressions for the proton longitudinal and singlet relaxation rate constants:

$$T_1^{-1}(\text{CH}_2\text{D}) = \frac{(4 + 3\kappa\tau_c)}{24(1 + 3\kappa\tau_c)} (16\omega_{\text{HD}}^2 + 9\omega_{\text{HH}}^2) \tau_c, \quad (1)$$

$$T_S^{-1}(\text{CH}_2\text{D}) = \frac{8 \sin^2 2\theta}{1 + 3\kappa\tau_c} \omega_{\text{HD}}^2 \tau_c. \quad (2)$$

The following expressions apply to the relaxation of the  $^{13}\text{C}$  nuclei in the  $\text{CH}_N$  groups on the six-membered ring (with  $N = 1$  or  $2$ ), and for the  $^2\text{H}$  relaxation of the deuteron in the monodeuterated methyl group:

$$T_1^{-1}(^{13}\text{CH}_N) = N\omega_{\text{CH}}^2 \tau_c, \quad (3)$$

$$T_1^{-1}(^2\text{H}) = \frac{3 \left( 32 + 33\kappa\tau_c + 9\kappa\tau_c (4 \cos(2\theta_Q) + 3 \cos(4\theta_Q)) \right)}{64(1 + 3\kappa\tau_c)} \omega_Q^2 \tau_c. \quad (4)$$

Here  $\omega_{\text{CH}}$  is the dipole-dipole coupling constant for the interaction between the proton and carbon nuclei,  $\omega_Q$  is the quadrupole coupling frequency of the deuteron, and  $\theta_Q$  is the angle between the 3-fold jump axis and the dominant principal axis of the  $^2\text{D}$  electric field gradient (EFG) tensor (assumed to be along the CD bond vector). For a tetragonal geometry of the  $\text{CH}_2\text{D}$  group:

$$T_1^{-1}(^2\text{H}) = \frac{(3 + \kappa\tau_c)}{2(1 + 3\kappa\tau_c)} \omega_Q^2 \tau_c. \quad (5)$$

These equations assume rigid-body rotational diffusion of the whole molecule (including the ring) and 3-fold jumps of the methyl group, with  $^{13}\text{C}$ - $^1\text{H}$  dipolar and quadrupolar relaxation dominating the  $^{13}\text{C}$  and  $^2\text{D}$  relaxation, respectively.

For the simple case of point nuclei (ignoring vibrational averaging), the coupling constants are defined as follows:

$$\omega_{\text{CH}} = -(\mu_0/4\pi) \gamma(^1\text{H})\gamma(^{13}\text{C})\hbar r_{\text{CH}}^{-3} \quad (6)$$

$$\omega_Q = \frac{e^2 q Q}{2\hbar} \quad (7)$$

where  $r_{\text{HC}}$  is the internuclear distance,  $Q$  is the electric quadrupolar moment of the deuterium nucleus, and  $eq$  is the electrical field gradient at the deuterium nucleus [49].

This relaxation model leads to the following expression for the ratio of  $T_S$  to  $T_1$ :

$$\frac{T_S}{T_1} = (4 + 3\kappa\tau_c) \frac{16\omega_{\text{HD}}^2 + 9\omega_{\text{HH}}^2}{192\omega_{\text{HD}}^2 \sin^2 2\theta}. \quad (8)$$

In general the ratio  $T_S/T_1$  depends on the jump rate  $\kappa$  and rotational correlation time  $\tau_c$ , and is expected to depend on temperature, solvent, and other factors. However, in the “slow-jump regime”  $\kappa\tau_c \ll 1$ , the ratio of  $T_S$  to  $T_1$  becomes independent of  $\kappa$  and  $\tau_c$ , and only depends on interaction parameters within the  $\text{CH}_2\text{D}$  group:

$$\frac{T_S}{T_1} \simeq \frac{16\omega_{\text{HD}}^2 + 9\omega_{\text{HH}}^2}{48\omega_{\text{HD}}^2 \sin^2 2\theta} \simeq \frac{3\omega_{\text{HH}}^2}{16\omega_{\text{HD}}^2 \sin^2 2\theta} \quad (9)$$

The fact that  $^1\text{H}$ - $^1\text{H}$  dipolar couplings are about 6 times stronger than  $^1\text{H}$ - $^2\text{D}$  dipolar couplings (at equal distances) has been invoked in the last approximation. The observed direct proportionality of  $T_S$  to  $T_1$  is consistent with the validity of the slow-jump regime over the explored range of experimental conditions.

### 3.5. Relaxation analysis

The  $^{13}\text{C}$  and  $^2\text{H}$  relaxation data allow an estimate of the rotational correlation time  $\tau_c$  and 3-fold jump rate  $\kappa$ . The following analysis refers to data obtained on 0.1 M N- $\text{CH}_2\text{D}$ -2-methylpiperidine in degassed  $\text{CD}_2\text{Cl}_2$  solution, at 500 MHz and 25°C.

The overall rotational correlation time  $\tau_c$  was estimated by analysing the experimental  $T_{1\text{C}}$  relaxation time constants for  $^{13}\text{C}$  nuclei in the  $\text{CH}_N$  groups on the six-membered ring, using Equation 3 which applies for extreme-narrowing isotropic rotational tumbling, dominated by the  $^{13}\text{C}$ - $^1\text{H}$  dipolar relaxation mechanism [50]. By assuming an internuclear  $^{13}\text{C}$ - $^1\text{H}$  distance of 108.9 pm, which corresponds to a dipole coupling constant of  $\omega_{\text{CH}}/2\pi = -23.4$  kHz, we obtain the following estimate of the rotational correlation time:  $\tau_c = 3.8 \pm 0.6$  ps.

The 3-fold jump rate constant  $\kappa$  may be estimated from the  $^2\text{D}$   $T_1$  relaxation time constant, by using Equation 5. The deuteron quadrupole coupling constant  $\omega_Q/2\pi = 83.5$  kHz has been estimated by solid-state NMR [51, 52].

The unique principal axis of the deuteron quadrupole coupling tensor is assumed to be along the C-D bond, at an angle of  $70.5^\circ$  with respect to the 3-fold jump axis. From comparing the experimental relaxation time  $T_1(^2\text{H}) = 0.75 \pm 0.01 \text{ s}$  with Equation 4, which was derived for the case of a hindered 3-fold rotor attached to a molecule undergoing isotropic rotational diffusion, and assuming that the quadrupolar mechanism dominates the deuteron relaxation, we estimate an upper limit on the 3-fold jump rate:  $\kappa \lesssim 3.2 \times 10^{10} \text{ s}^{-1}$ . The product of the rotational correlation time and the 3-fold jump rate constant is therefore given by  $\kappa\tau_c \lesssim 0.12$ , which supports the validity of the slow-jump regime and hence Equation 9.

Now consider the case where the three hydrogen nuclei of the  $\text{CH}_2\text{D}$  group are considered to be points, located at the vertices of an equilateral triangle. In this case  $2\theta = 60^\circ$ , the internuclear distances are all equal, and the dipolar couplings are in the ratio of the magnetogyric ratios  $\omega_{\text{HH}}/\omega_{\text{HD}} = \gamma(^1\text{H})/\gamma(^2\text{D}) = 6.51$ . This “equilateral triangle model” predicts the following relaxation time ratio:  $(T_S/T_1)_\Delta = 10.5$ . However, the observed value is quite different:  $(T_S/T_1)_{\text{obs}} = 3.1 \pm 0.1$ .

How can this discrepancy be explained? One approach would be to call into question the intra-methyl relaxation model: However, as discussed above, the evidence for the dominance of intra- $\text{CH}_2\text{D}$  interactions in the  $^1\text{H}$  and  $^2\text{D}$  relaxation is very strong. An alternative approach is to maintain the local intra-methyl relaxation model, but to modify the relative magnitudes and geometries of the dipole-dipole interactions within the rotating  $\text{CH}_2\text{D}$  group, to take into account differential vibrational averaging on a faster timescale than the molecular rotation or 3-fold jumps.

A detailed analysis of the effect of rapid vibrational motions on the dipolar interaction tensors in the  $\text{CH}_2\text{D}$  group would be a major project. For the sake of simplicity we employ a naive picture, in which the nuclei are still regarded as localized points, but with the  $^2\text{D}$  nucleus displaced from its original position, in order to account for differential vibrational averaging of the  $^2\text{D}$  and  $^1\text{H}$  interactions. In order to maintain symmetry, we consider a model in which the  $^2\text{D}$  nucleus is moved in the  $\text{CH}_2\text{D}$  plane along the line bisecting the  $^1\text{H}$ - $^1\text{H}$  vector (Figure 7a). This adjustment changes the internuclear distances as well as the angle  $\theta$ . The dependence of the theoretical relaxation rate ratio  $T_S/T_1$  on the deuterium displacement is shown in Figure 7b. There is a strong sensitivity to the deuterium displacement, with the observed ratio  $T_S/T_1 = 3.1$  being consistent with a displacement of 38.7 pm towards the






Figure 7: (a) Adjustment of the effective geometry of the  $\text{CH}_2\text{D}$  group to account for the observed relaxation rate ratio  $T_S/T_1 = 3.1$ . The three nuclei are initially at the vertices of an equilateral triangle of side length 179.6 pm (black circles). The deuterium nucleus is displaced towards the original centre of the triangle (open circle) by a distance  $\Delta r$ . (b) Dependence of the relaxation rate ratio  $T_S/T_1$  on the displacement  $\Delta r$ . The open circle shows the displacement needed for consistency with the observed rate ratio  $T_S/T_1 = 3.1$ . The black circle shows the predicted rate ratio from an equilateral triangle geometry.

original centre of the equilateral triangle. This represents a contraction in both  $^1\text{H}$ - $^2\text{D}$  distances by about 18%, and a change in the angle  $2\theta$  from  $60^\circ$  to  $75.1^\circ$ . A sketch of the adjusted geometry is shown in Figure 7a.

This degree of geometrical distortion is probably unrealistic. It is more likely that differential vibrational averaging of the  $^1\text{H}$ - $^1\text{H}$  and  $^1\text{H}$ - $^2\text{D}$  dipolar interactions is responsible for the observed  $T_S/T_1$  ratio. Vibrational motion out of the  $\text{CH}_2\text{D}$  plane is likely to be particularly effective. For example, the larger vibrational amplitudes of the  $^1\text{H}$  nuclei relative to the more massive  $^2\text{D}$  nucleus would reduce the  $^1\text{H}$ - $^1\text{H}$  dipolar interaction more than the  $^1\text{H}$ - $^2\text{D}$  dipolar interactions. This effect would lead to a correction in the right direction. We have not attempted a more sophisticated analysis of vibrational effects on the interaction parameters.

#### 4. Discussion

It is possible to populate long-lived nuclear singlet order in the proton pairs of monodeuterated methyl groups, under suitable conditions. This requires non-uniformity in the rotamer populations as well as a local chiral environment in order to induce a small isotropic shift difference between the  $\text{CH}_2\text{D}$  protons. Both conditions are fulfilled for the  $\text{CH}_2\text{D}$  group in  $\text{N-CH}_2\text{D-}$

2-methylpiperidine, where a hyperconjugation effect involving the nitrogen lone pair perturbs the vibrational energies and hence the rotamer populations, while the neighbouring methyl group provides a chiral environment.

In this system, the ratio of singlet relaxation time  $T_S$  to longitudinal relaxation time  $T_1$  is found to be  $3.1 \pm 0.1$  over a wide range of conditions, with the longest observed value of  $T_S$  approaching 1 minute at elevated temperature. The observation of a constant ratio of  $T_S$  to  $T_1$  supports a relaxation model in which dipolar interactions between the  $\text{CH}_2\text{D}$  protons and deuteron dominate the singlet relaxation. However, a naive model in which the  $^1\text{H}$  and  $^2\text{D}$  nuclei of the  $\text{CH}_2\text{D}$  group are viewed as point-like magnetic dipoles fixed at the vertices of an equilateral triangle predicts a much larger ratio of  $T_S$  to  $T_1$ . We explore the effects of a geometric distortion in which the  $^2\text{D}$  nucleus is displaced towards the centre of the triangle and find that a displacement of 38.7 pm is needed to explain the experimental data. This adjustment should not be viewed as a realistic structural proposal, but as a crude attempt to represent the differential effects of vibrational averaging and nuclear wavefunction delocalization within a simplistic point-nucleus geometric model.

The observed relaxation time ratio of  $T_S/T_1 \simeq 3.1$  is probably too small for most feasible applications. However, it should be noted that equation 8 permits a much larger ratio  $T_S/T_1$  in the case of fast 3-fold jumps, i.e.  $\kappa\tau_c \gtrsim 1$ . This does not occur in N- $\text{CH}_2\text{D}$ -2-methylpiperidine, where the rotational barrier appears to be relatively large. Nevertheless it may arise in other compounds displaying inequivalence of the  $\text{CH}_2\text{D}$  proton pair. It remains to be seen whether it is possible to maintain a sufficient differential between rotamer populations while still having a sufficiently low barrier to permit fast 3-fold jumps. We are currently exploring some other chemical candidates.

We have also attempted to observe long-lived singlet states in other chiral  $\text{CH}_2\text{D}$  systems, such as derivatives of 3- $^2\text{D}$ -lactic acid, and also (N- $\text{CH}_2\text{D}$ )-3-methylpiperidine (similar to the substance used above, but with a more remote  $\text{CH}_3$  group). We failed to access the  $\text{CH}_2\text{D}$  singlet state in both of these cases: the chemical inequivalence of the  $\text{CH}_2\text{D}$  protons is presumably too small to exploit. Apart from unusual circumstances [53], hyperconjugation between a lone pair on a neighbouring atom (such as N) and the methyl C-H/D bonds seems to be a requirement for obtaining a sufficient chemical shift difference of a few ppb or more. At the time of writing we have not attempted these experiments on other chiral compounds containing a X- $\text{CH}_2\text{D}$  moiety, where X is an atom other than N possessing a lone pair, such as P, O or S.

It should be noted that many interesting chiral molecules do contain a tetrahedral N-methyl group of suitable type. Examples include the psychoactive agents lysergic acid, codeine, morphine, cocaine, heroin, and methamphetamine. It is not yet known whether the methyl-monodeuterated versions of these systems possess an accessible long-lived singlet state.

## 5. Acknowledgements

This research was supported by the Engineering and Physical Sciences Research Council (UK), grant codes EP/N002482 and EP/L505067/1, and also the Wolfson Foundation, the Royal Society/CNRS exchange scheme and by Bruker Biospin (USA). The authors thank Daniel J. O'Leary (Pomona College) for discussions, Jim Kempf and Robert Krull (Bruker Biospin) for assistance with temperature dependent measurements, and Marcel Utz and Soumya Singha Roy for experimental help.

## 6. References

- [1] M. Carravetta, O. G. Johannessen and M. H. Levitt, *Phys. Rev. Lett.*, 2004, **92**, 153003.
- [2] M. Carravetta and M. H. Levitt, *J. Am. Chem. Soc.*, 2004, **126**, 6228–6229.
- [3] M. Carravetta and M. H. Levitt, *J. Chem. Phys.*, 2005, **122**, 214505.
- [4] G. Pileio and M. H. Levitt, *J. Magn. Reson.*, 2007, **187**, 141–145.
- [5] R. Sarkar, P. R. Vasos and G. Bodenhausen, *J. Am. Chem. Soc.*, 2007, **129**, 328–334.
- [6] G. Pileio, M. Carravetta, E. Hughes and M. H. Levitt, *J. Am. Chem. Soc.*, 2008, **130**, 12582–12583.
- [7] A. K. Grant and E. Vinogradov, *J. Magn. Reson.*, 2008, **193**, 177–190.
- [8] G. Pileio and M. H. Levitt, *J. Chem. Phys.*, 2009, **130**, 214501.
- [9] W. S. Warren, E. Jenista, R. T. Branca and X. Chen, *Science*, 2009, **323**, 1711–1714.

- [10] M. C. D. Tayler, S. Marie, A. Ganesan and M. H. Levitt, *J. Am. Chem. Soc.*, 2010, **132**, 8225–8227.
- [11] G. Pileio, M. Carravetta and M. H. Levitt, *Proc. Natl. Acad. Sci. U.S.A.*, 2010, **107**, 17135–17139.
- [12] M. C. D. Tayler and M. H. Levitt, *Phys. Chem. Chem. Phys.*, 2011, **13**, 5556–5560.
- [13] Y. Feng, R. M. Davis and W. S. Warren, *Nat. Phys.*, 2012, **8**, 831–837.
- [14] G. Pileio, J. T. Hill-Cousins, S. Mitchell, I. Kuprov, L. J. Brown, R. C. D. Brown and M. H. Levitt, *J. Am. Chem. Soc.*, 2012, **134**, 17494–17497.
- [15] M. H. Levitt, *Annu. Rev. Phys. Chem.*, 2012, **63**, 89–105.
- [16] M. C. D. Tayler, I. Marco-Rius, M. I. Kettunen, K. M. Brindle, M. H. Levitt and G. Pileio, *J. Am. Chem. Soc.*, 2012, **134**, 7668–7671.
- [17] G. Pileio, S. Bowen, C. Laustsen, M. C. D. Tayler, J. T. Hill-Cousins, L. J. Brown, R. C. D. Brown, J.-H. Ardenkjaer-Larsen and M. H. Levitt, *J. Am. Chem. Soc.*, 2013, **135**, 5084–5088.
- [18] Y. Feng, T. Theis, X. Liang, Q. Wang, P. Zhou and W. S. Warren, *J. Am. Chem. Soc.*, 2013, **135**, 9632–9635.
- [19] S. J. DeVience, R. L. Walsworth and M. S. Rosen, *Phys. Rev. Lett.*, 2013, **111**, 173002.
- [20] T. Theis, Y. Feng, T. Wu and W. S. Warren, *J. Chem. Phys.*, 2014, **140**, 014201.
- [21] G. Stevanato, S. Singha Roy, J. Hill-Cousins, I. Kuprov, L. J. Brown, R. C. D. Brown, G. Pileio and M. H. Levitt, *Phys. Chem. Chem. Phys.*, 2015, **17**, 5913–5922.
- [22] G. Stevanato, J. T. Hill-Cousins, P. Håkansson, S. S. Roy, L. J. Brown, R. C. D. Brown, G. Pileio and M. H. Levitt, *Angew. Chem. Int. Ed.*, 2015, **54**, 3740–3743.

- [23] J.-H. Ardenkjær-Larsen, B. Fridlund, A. Gram, G. Hansson, L. Hansson, M. H. Lerche, R. Servin, M. Thaning and K. Golman, *Proc. Natl. Acad. Sci. U.S.A.*, 2003, **100**, 10158–10163.
- [24] S. J. Nelson, J. Kurhanewicz, D. B. Vigneron, P. E. Z. Larson, A. L. Harzstark, M. Ferrone, M. van Criekinge, J. W. Chang, R. Bok, I. Park, G. Reed, L. Carvajal, E. J. Small, P. Munster, V. K. Weinberg, J. H. Ardenkjær-Larsen, A. P. Chen, R. E. Hurd, L.-I. Odegardstuen, F. J. Robb, J. Tropp and J. A. Murray, *Sci. Transl. Med.*, 2013, **5**, 198.
- [25] R. Buratto, D. Mammoli, E. Chiarparin, G. Williams and G. Bodenhausen, *Angew. Chem. Int. Ed.*, 2014, **53**, 11376–11380.
- [26] A. Bornet, X. Ji, D. Mammoli, B. Vuichoud, J. Milani, G. Bodenhausen and S. Jannin, *Chem. – A Eur. J.*, 2014, **20**, 17113–17118.
- [27] Y. Zhang, K. Basu, J. W. Canary and A. Jerschow, *Phys. Chem. Chem. Phys.*, 2015, **17**, 24370–24375.
- [28] T. Theis, G. X. Ortiz, A. W. J. Logan, K. E. Claytor, Y. Feng, W. P. Huhn, V. Blum, S. J. Malcolmson, E. Y. Chekmenev, Q. Wang and W. S. Warren, *Sci. Adv.*, 2016, **2**, 1501438.
- [29] B. Meier, J.-N. Dumez, G. Stevanato, J. T. Hill-Cousins, S. S. Roy, P. Håkansson, S. Mamone, R. C. D. Brown, G. Pileio and M. H. Levitt, *J. Am. Chem. Soc.*, 2013, **135**, 18746–18749.
- [30] J.-N. Dumez, P. Håkansson, S. Mamone, B. Meier, G. Stevanato, J. T. Hill-Cousins, S. S. Roy, R. C. D. Brown, G. Pileio and M. H. Levitt, *J. Chem. Phys.*, 2015, **142**, 044506.
- [31] S. S. Roy, J.-N. Dumez, G. Stevanato, B. Meier, J. T. Hill-Cousins, R. C. Brown, G. Pileio and M. H. Levitt, *J. Magn. Reson.*, 2015, **250**, 25–28.
- [32] P. R. Bunker and P. Jensen, *Molecular Symmetry and Spectroscopy*, 2nd Ed., NRC Research Press, Ottawa, 2006.
- [33] M. Icker and S. Berger, *J. Magn. Reson.*, 2012, **219**, 1–3.
- [34] M. Icker, P. Fricke and S. Berger, *J. Magn. Reson.*, 2012, **223**, 148–150.

- [35] M. Icker, P. Fricke, T. Grell, J. Hollenbach, H. Auer and S. Berger, *Magn. Reson. Chem.*, 2013, **51**, 815–820.
- [36] F. A. L. Anet and M. Kopelevich, *J. Am. Chem. Soc.*, 1989, **111**, 3429–3431.
- [37] B. D. Allen, and D. J. O’Leary, *J. Am. Chem. Soc.*, 2003, **125**, 9018–9019.
- [38] B. D. Allen, J.-C. Cintrat, N. Faucher, P. Berthault, B. Rousseau and D. J. O’Leary, *J. Am. Chem. Soc.*, 2005, **127**, 412–420.
- [39] A. S. Kiryutin, H. Zimmermann, A. V. Yurkovskaya, H.-M. Vieth and K. L. Ivanov, *J. Magn. Reson.*, 2015, **261**, 64–72.
- [40] M. C. D. Tayler and M. H. Levitt, *J. Am. Chem. Soc.*, 2013, **135**, 2120–2123.
- [41] D. Mammoli, B. Vuichoud, A. Bornet, J. Milani, J.-N. Dumez, S. Jannin and G. Bodenhausen, *J. Phys. Chem. B*, 2015, **119**, 4048–4052.
- [42] M. H. Levitt, *J. Magn. Reson.*, 2016, **262**, 91–99.
- [43] A. Abragam, *Principles of Nuclear Magnetism*, Clarendon Press, Oxford, 1961.
- [44] G. Pileio, *J. Chem. Phys.*, 2011, **135**, 174502.
- [45] A. S. Dubin and S. I. Chan, *J. Chem. Phys.*, 1967, **46**, 4533–4535.
- [46] T. Burke and S. I. Chan, *J. Magn. Reson. (1969)*, 1970, **2**, 120–140.
- [47] T. E. Bull, *J. Chem. Phys.*, 1976, **65**, 4802–4815.
- [48] G. B. Matson, *J. Chem. Phys.*, 1977, **67**, 5152–5161.
- [49] A. Jerschow, *Prog. Nucl. Magn. Reson. Spectrosc.*, 2005, **46**, 63–78.
- [50] J. Kowalewski and L. Mäler, *Nuclear Spin Relaxation in Liquids: Theory, Experiments, and Applications*, CRC Press, Boca Raton, 2006.
- [51] K. Beshah, E. T. Olejniczak and R. G. Griffin, *J. Chem. Phys.*, 1987, **86**, 4730–4736.



- [52] A. Mittermaier, and L. E. Kay, *J. Am. Chem. Soc.*, 1999, **121**, 10608–10613.
- [53] A. Restelli and J. S. Siegel, *J. Am. Chem. Soc.*, 1992, **114**, 1091–1092.