

This is the overview page

The ascending rank of CKD in the Global Burden of Disease Study

Journal:	<i>Nephrology Dialysis Transplantation</i>
Manuscript ID	NDT-01080-2016.R1
Manuscript Type:	NDT Perspectives - review (INVITED ONLY)
Date Submitted by the Author:	n/a
Complete List of Authors:	Jager, Kitty J.; Academic Medical Center, Amsterdam, Dept Medical Informatics Fraser, Simon; University of Southampton, Academic Unit of Primary Care and Population Sciences
Keyword list:	CKD, death rates, disability adjusted life years, global burden, prevalence, years of life lost

The ascending rank of CKD in the Global Burden of Disease Study

Kitty J. Jager MD PhD¹ and Simon D.S. Fraser BM DM²

¹ ERA-EDTA Registry, Dept. of Medical Informatics, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

² Academic Unit of Primary Care and Population Sciences. Faculty of Medicine, University of Southampton, Southampton General Hospital, Southampton, UK

Corresponding author

Kitty J. Jager MD PhD
Dept. of Medical Informatics
Academic Medical Center
Dept. of Medical Informatics, J1B-125
P.O. Box 22700
1100 DE Amsterdam
The Netherlands
Tel +31 20 566 7637
Email k.j.jager@amc.uva.nl

Running Title: Global Burden of CKD

Word Count: 3750

Abstract Word Count: 267

References: 35

Key words: CKD, death rates, disability adjusted life years, global burden, prevalence, years of life lost

Abstract:

General population based studies, the chronic kidney disease (CKD) prognosis consortium and renal registries worldwide have contributed to the description of the scale of CKD as a public health problem. Since 1990 CKD has been included in the list of non-communicable diseases investigated by the Global Burden of Disease (GBD) study. GBD represents a systematic, high quality, scientific effort to quantify the comparative magnitude of health loss from all major diseases, injuries and risk factors. This paper provides an outline of the place of CKD in the ranking of these diseases and the change over time.

Whereas age standardised death and disability adjusted life years (DALY) rates due to non-communicable diseases in general have been declining, such favourable trends do not exist for CKD. Altogether the GBD reports indicate increasing rates for death and DALYs due to CKD with huge variation across the globe. A substantial component of the observed increase in mortality attributable to CKD relates to that caused by diabetes mellitus and hypertension. For the increase in DALYs, CKD due to diabetes mellitus appears to be the main contributor.

It is possible that these trends are in part due to new data becoming available or different coding behaviour over time, including greater specificity of coding. Although some feel there is evidence of overdiagnosis, it seems clear that in many regions CKD and its risk factors are a growing public health problem and in some of them ranking very high as cause of YLL and DALYs. Therefore, public health policies to address this problem as well as secondary prevention in high risk groups remain highly needed.

Introduction

In 2002 chronic kidney disease (CKD) was classified into 5 stages [1] based on the presence of kidney damage (i.e. albuminuria) or decreased kidney function (i.e. glomerular filtration rate (GFR) <60 mL/min/1.73m²). Ten years later, as both kidney function and kidney damage had been shown to independently affect the prognosis of CKD, the Kidney Disease Improving Global Outcomes (KDIGO) modified this classification by proposing 6 categories of kidney function (eGFR, G1-G5) and 3 categories of kidney damage (albuminuria, A1-A3). This resulted in 18 categories that have been labelled as low, moderately increased, high or very high risk of complications like drug toxicity, endocrine and metabolic outcomes, cardiovascular disease and mortality [2]. These and other complications of CKD have been recognized to induce an important disease burden both for individual patients and for societies.

In almost every recent paper dealing with the implications of CKD the disease is referred to as an important public health problem. A number of general population based studies have been instrumental in describing the scale of this public health problem by assessing and monitoring the health of communities and populations at risk, while the CKD prognosis consortium has provided significant input to the understanding of the joint contribution of albuminuria and decreased kidney function to patient prognosis. Finally, for the past few decades renal registries have provided information on the access to appropriate, but costly, renal replacement therapy (RRT) for the treatment of end-stage kidney disease (ESKD).

European national surveys have shown CKD stage 3-5 prevalence to vary from 2.3% in Germany (18-79 year) [3]; 2.4% in Finland (20-74 years) [4], 4.0% in Spain (20-74 years) [4] to 5.2% in England in subjects of 16 years and older [5]. Outside Europe the range in prevalence of CKD stage 3-5 derived from nationally representative samples is slightly wider: from 1.7% in China (18 years and older) [6], 3.1% in adults from Canada [7], 5.8% in Australia (25 years and older) [8] to 6.7% in adults from the USA [9].

The CKD prognosis consortium has demonstrated that lower eGFR and higher albuminuria are risk factors for ESKD, acute kidney injury and progressive CKD in both general and high risk populations, independent of each other and of cardiovascular risk factors [10]. In addition, they have provided evidence that both are also risk factors for cardiovascular morbidity as well as cardiovascular and all-cause mortality, regardless of age [11].

Renal registries, on the other hand, have shown that in 2013 the access to RRT across different nations varied greatly - from a prevalence of 66 per million population (pmp) in Indonesia to 3138 pmp in Taiwan, a nearly 50-fold difference [12]. In 2010, the total number of RRT patients worldwide amounted to 2.6 million, whereas the number of patients needing this treatment was estimated to be between 4.9 and 9.7 million with the largest gap between need and supply of RRT in low-income countries, particularly in Asia and Africa [13].

Both the prevalence of CKD stage 3-5 and that of RRT are expected to increase further, although for the USA recent NHANES data have shown that the increase in CKD stage 3-5 may only be very modest [14]. Nevertheless, a worldwide increase can be expected due to population ageing and the increasing prevalence of obesity, hypertension and diabetes mellitus in the general population. At the same time worldwide use of RRT has been projected to more than double by 2030, with the largest growth in Asia [13]. This growing burden of RRT will likely be due to the continued improvement in survival on dialysis and transplantation [12,15] and, in developing countries, to an increasing number of patients taken on to RRT.

Since 1990 CKD has been included in the list of non-communicable diseases investigated by the Global Burden of Disease (GBD) study. GBD represents a systematic, high quality, scientific effort to quantify the comparative magnitude of health loss from all major diseases, injuries and risk factors. This paper aims to provide an outline of the place of CKD in the ranking of these diseases and the change over time.

GBD measures and analytical methods

The methods involved in the GBD studies are complex and this review will not attempt to cover them comprehensively [16-19]. The regions identified in the GBD study were the result of a collaboration of several hundred investigators reporting metrics such as adult and child mortality from across the world. Countries were divided into 21 regions according to similarity from an epidemiologic and geographical perspective [20]. Data availability and reliability varied widely between locations, therefore requiring several internal validity checks such as ensuring that the sum of individual cause of death numbers equalled the all-cause death totals [19].

The investigators have recognised important limitations of their data, such as insufficient information on certain causes of death from many countries. They therefore relied on complex statistical models to address gaps. For example, they used six different modelling strategies for causes of death depending on the quality and quantity of data available, and generated 95% confidence intervals to reflect the degree of uncertainty (95%UI) [19].

To summarize, they used the following steps to assess the quality of cause of death data:

1. Adjustment of cause of death data from vital registration systems for incompleteness
2. Development of maps to help translate causes of death in the collected data to the GBD 2013 cause list.
3. Identifying and redistributing 'garbage codes' (defined as 'codes for which deaths are assigned that cannot or should not be considered as the underlying cause of death').
4. Identifying data (such as verbal autopsy reports) where age or sex data were combined and using algorithms to split into specific age-sex groups.

- 1
- 2
- 3 5. Smoothing fluctuations in cause of death data for small countries (where number of
- 4 deaths for individual causes can fluctuate widely).
- 5
- 6 6. Exclusion of outliers on the basis of biological implausibility, where data from a country
- 7 was highly inconsistent with other studies for that country (or from countries with similar
- 8 sociodemographic profiles in the same region), and studies that led to abrupt changes in
- 9 trends that could not be explained by contextual factors [16].
- 10
- 11

12 The mortality estimation processes used in the GBD studies are complex due to the

13 widely varying degree of completeness of vital registration systems in different countries [21]. In

14 the 2013 GBD, the methods used to identify cause of death were the same as those used in the

15 2010 GBD with the exception of some new datasets that became available to give information

16 on 'deaths from CKD caused by glomerulonephritis' [16]. For those deaths that were medically

17 certified, this was used as part of the process of identifying cause of death. Even with the

18 introduction of standard ICD coding, however, there remains considerable variation in coding

19 practice and completeness globally. Where no death certification data were available, the

20 investigators used a variety of sources such as surveillance systems, demographic research

21 sites, surveys, censuses, disease registries, verbal autopsy data and police records. A summary

22 of the 'site-years' of data are provided that show the principle sources for each cause of death.

23 Where coding was available, CKD was identified by ICD10 codes (E10.2, E11.2, E12.2, E13.2,

24 I12.0, I12.9, I13.1, I13.2, I13.9, N02-N07, N15.0) and ICD9 codes (250.4, 403-404, 581-583,

25 589) (see Table a in Supplementary for ICD terms). A further limitation of GBD in relation to CKD

26 is that the latter was only defined as impaired GFR and did not take into account albuminuria.

27

28

29

30

31

32

33

34

35 To be able to make further meaningful comparison of the burden caused by different

36 diseases they calculated years of life lost due to premature death (YLL, calculated by multiplying

37 numbers of deaths by life expectancy at time of death in a reference population) and disability

38 adjusted life years (DALYs) in order to reflect the degree of disability such as pain, or functional

39 limitations caused by each condition. Assessing the impact of different conditions by these

40 metrics included the use of disability weights derived from general population surveys in five

41 countries and an open internet survey [22].

42

43

44

45

46 Finally, the GBD investigators calculated age-standardised death rates and DALY rates

47 using the world population age standard (WHO 2001) updated to a standard population structure

48 using population estimates for 2010-2035 (World Population Prospects, United Nations) [16].

49 Data for the charts in the next section were obtained from the Institute for Health Metrics and

50 Evaluation (IHME) 'GBD Compare' calculator' [23] except where referenced otherwise.

51 Throughout this paper death rate refers to deaths per 100,000 people and DALY rate to DALYs

52 per 100,000 people.

53

54

55

56

57

58

59

60

Global, national and regional variations in CKD burden over time

The change in CKD burden over time, globally and in different countries and regions as presented by GBD, can be considered in terms of both mortality and morbidity and by absolute and relative measures. The GBD considers absolute measures by describing changes in the crude and age-standardised death rate and giving absolute estimates for DALYs and YLLs. In addition, they provide relative ones by ranking deaths, DALYs and YLLs.

Mortality: deaths, death rates and YLL

Global deaths due to all cause CKD in 1990 were about 409,000 (363,844-433,380) compared with 956,000 (812,896-103,4491) in 2013 [16]. The upper panel of Figure 1 illustrates the change in death rate attributed to CKD over time, globally and by 21 regions. It shows that globally death due to CKD is increasing (rising from about 8 per 100,000 people to about 13 per 100,000 people between 1990 and 2013). However, these numbers hide considerable variation between the GBD regions. For some regions (such as Eastern Europe and Eastern and Central Sub-Saharan Africa) the death rate appears to have been static over this period. For others, particularly Latin America, high-income Asia Pacific and high income North America, the absolute increase is remarkable. The most notable is Central Latin America (which includes Columbia, Costa Rica, El Salvador, Guatemala, Honduras and Mexico) where deaths have tripled from 10.4 per 100,000 people in 1990 to 32.7 per 100,000 people in 2013. Within Mexico itself, the change is even more striking (from 12.2 in 1990 to 47.1 in 2013, not shown in the figure). Age-standardised death rates by region show a similar but somewhat diluted pattern compared to the crude death rates, although with greater prominence for some regions, such as Oceania and an even more noticeable difference between Central Latin America and other regions (Figure a in supplementary material) [23]. Table 1 shows the change in global age-standardised death rates for CKD according to cause. The GBD modelling suggests that the global increase in CKD mortality seen between 1990 and 2013 was 4.2 deaths per 100,000 people, of which 1.5 deaths (36%) were due to the increase in diabetes and hypertension-related CKD.

In relative terms, the GBD study shows the deaths attributable to CKD as a percentage of total deaths in a region (Figure 1, lower panel). Latin America still dominates the picture with an increase in Central Latin America to over 6% of all deaths attributable to CKD in 2013, from about 2% in 1990. Whereas globally CKD increased in the ranking of cause of death from 27th in

1
2
3 1990 to 13th in 2013, in Central Latin America over the same period it increased from 12th to 4th.
4 Similarly, CKD increased in the global ranking of YLL from 36th in 1990 to 19th in 2013 but from
5 17th to 5th in Central Latin America [16].
6
7
8
9

10 *Morbidity: DALYs*

11
12
13 In absolute terms, the total number of global DALYs associated with CKD has increased
14 from 19 million in 1990 to 33 million in 2013 [17]. The upper panel of Figure 2 illustrates the
15 change in DALY rate due to CKD over time, globally and in the 21 regions. The trend in DALY
16 rate shows a flatter pattern for many geographical areas compared with death rate, with an
17 apparent decrease in Eastern Sub-Saharan Africa. On the other hand, there is a striking
18 increase in DALY rate in Central Latin America since about 1995, resulting in having
19 approximately twice the global DALY burden in 2013. Table 1 shows the change in age
20 standardised global DALY rate, overall and by CKD cause. The GBD data suggest that morbidity
21 due to CKD is increasing globally and that CKD due to diabetes, hypertension and other causes
22 are all contributing to this increase. Nevertheless, the greatest percentage increase is in CKD
23 due to diabetes.
24
25
26
27
28
29

30 Similar to the pattern seen for mortality, the increasing percentage of DALYs due to CKD
31 is most marked in Central Latin America (from 1.34% in 1990 to 3.66% in 2013, Figure 2, lower
32 panel). There are also differences elsewhere. Western Europe for example recently shows a
33 greater proportion of DALYs attributable to CKD than most other regions (2.4% in 2013). The
34 steadily increasing pattern of CKD's relative position in the Caribbean was interrupted in 2010 by
35 the dominance of the effects of the Haiti earthquake, but appears to have resumed its previous
36 rising pattern following that humanitarian disaster. In the global ranking of the number of DALYs
37 CKD increased as a cause from 28th in 1990 to 20th in 2013 [17]. The change in Central Latin
38 America moved CKD from being the 18th cause of DALYs in 1990 to 7th in 2013. For Western
39 Europe, this rank change was modest, from 15th in 1990 to 12th in 2013 [23].
40
41
42
43
44
45
46
47
48

49 *Relative importance of CKD to YLLs and DALYs across nations*

50
51
52 The GBD report also highlights the top ten causes of YLLs and DALYs for each country
53 [16]. There are 51 countries for which CKD was among either the top ten causes of YLL (44
54 countries) or the top ten causes of DALYs (28 countries) or both (20 countries) in 2013. These
55 are shown in Table 2 with countries shown in blue that appear in both lists. These lists are again
56 dominated by Latin America, with the Caribbean, North Africa and the Middle East being strongly
57
58
59
60

1
2
3 represented in the top YLL list, and Europe in the DALY list. Latin America is particularly
4 prominent when considering the rank of CKD in each - it represents the second and third highest
5 causes of YLL and DALYs in Mexico and Nicaragua.
6
7
8
9

10 **Interpretation**

11
12
13 Whereas age standardised death and DALY rates due to non-communicable diseases in
14 general have been declining (median percent change: -18.6 during 1990-2013 and -6.3 during
15 2005-2013, respectively) [16-17], such favourable trends do not exist for CKD. Altogether the
16 different GBD reports indicate increasing rates for death and DALYs due to CKD, a substantial
17 part of which is resulting from CKD due to diabetes mellitus and hypertension-related CKD for
18 death and from CKD due to diabetes mellitus for DALYs. Death and DALYs from CKD due to
19 'other causes' also increased substantially, the reasons for which are unclear. The contribution
20 of acute kidney injury (AKI) to CKD incidence and progression is one potential cause, although
21 this cannot be inferred from these data [24]. As a result for 51 countries in the world CKD was
22 among the top ten causes of YLL, DALYs or both in 2013. On the other hand death and
23 disability as a result of CKD due to glomerulonephritis seem to be declining.
24
25
26
27
28
29

30 There is, however, substantial variation across the globe. The substantially worse
31 outcomes in Central and other parts of Latin America compared to many other countries are
32 striking, as are the other areas for which CKD appears in the top ten causes of YLL and DALYs.
33 Most notable among these are the Caribbean, Oceania, Mauritius, Thailand and parts of North
34 Africa and the Middle East. The increase in CKD in Latin America has been attributed to
35 demographic changes (aging population and rapid epidemiologic transition) as well as to an
36 epidemic of obesity and type 2 diabetes [25]. Others have suggested that a significant
37 contributor to the problem in Central Latin America is the high level of health care inequality, with
38 the greatest burden of CKD mortality and morbidity being born by those who are uninsured in
39 Mexico's health system [26]. Similarly, diabetes, hypertension and obesity are cited as key,
40 growing risk factors in the Arab world resulting in calls for public health action [27-28]. AKI has
41 also been highlighted as an important contributory factor in this population [29].
42
43
44
45
46
47
48

49 As far as DALY rates due to CKD are concerned, Western Europe gave up its leading
50 position around 2000 and now ranks second after Central Latin America. This may be due to
51 Western Europe's relatively old population compared to most other areas. However, United
52 Nations data projections estimate that for Latin America and the Caribbean, the population will
53 shift from just 11% aged 60 or over in 2015 to 26% by 2050, causing significant concern for the
54 prevalence and impact of CKD in these regions [30].
55
56
57
58
59
60

Methodological issues

The GBD study refers to a number of potential reasons as to why some causes of death – among them CKD – have shown such large changes in the number of global deaths. They point out that such changes may be because of new data, modifications of garbage coding algorithms and revised modelling strategies and acknowledge that the breakdown into specific causes is challenging in practice where patients have more than one associated condition, such as those with both diabetes and hypertension. [16]. This important limitation was also recognised by Rhee and Kovesdy in their review of these trends. They stressed the potential for increased recognition and coding of CKD (using ICD-9 and ICD-10 codes for cause of death) to mask true changes in absolute and relative mortality and morbidity [31]. While it is very difficult to establish this for certain, it is possible that some of the change in CKD mortality could be attributed to greater specificity of coding over time in certain areas. For example, some CKD-related death may have previously been recorded as cardiovascular death and use of death certification and renal registry data to attribute cause of death to CKD may underestimate its true impact. GBD data may at least lend some support to this. Whereas global CKD age standardised death rates have increased, the GBD study reports a substantial decrease in global age standardised death rate from cardiovascular disease (from 375.5 deaths per 100,000 people in 1990 to 293.2 deaths per 100,000 people in 2013; median percent change: -22 during 1990-2013) [16]. In Central Latin America, where the GBD study identified an approximately three-fold increase in crude CKD mortality, the crude CVD mortality increase has been considerably more modest (from 102 per 100,000 in 1990 to 131 per 100,000 in 2013). By contrast, the age-standardized death rates for CVD decreased from 239 per 100,000 to 204 per 100,000 in 2013, whereas the age-standardized death rate for CKD in Central Latin America more than doubled from 22 per 100,000 in 1990, to 48 per 100,000 in 2013. (Figure a in supplementary material). Moreover, the GBD calculator, which allows estimation of the percentage of cardiovascular deaths that are due to impaired GFR as risk factor, shows that approximately 8.5% of global ischaemic heart disease deaths and 8% of stroke-related deaths are attributable to low GFR [23].

CKD as a public health problem

To come back to some of the issues raised in the introduction. GBD data support the claims that CKD, as currently defined, is a global public health problem. As it is unlikely that the changes observed are solely due to different coding behaviour, we may assume that for many countries the burden of CKD is increasing in both absolute and relative terms resulting in greater

1
2
3 morbidity and mortality. Some countries and regions currently have a substantial problem, with
4 countries in Latin America, the Caribbean, North Africa and the Middle East, and Oceania
5 particularly at risk.
6

7 On the other hand dissenting opinions regarding labelling CKD as a disease have been
8 expressed. Some nephrologists and general practitioners feel that there is evidence of
9 overdiagnosis, given the large number of people with a CKD diagnosis who will never progress
10 to symptomatic forms of kidney disease. They feel clinicians should be sceptical about the
11 current definition of CKD and cautious about labelling, particularly older people [32].
12
13
14

15 What then should be the response of the global kidney and public health communities to
16 these data? Given the trends identified it seems unlikely that simply providing more RRT on
17 demand is going to be the solution, although improving equity of access to RRT is clearly an
18 important goal, particularly for countries in Central Latin America and other low income
19 countries. There is an urgent need for certain regions to develop their 'upstream' response to
20 these issues if their health systems are going to survive. Such response should be targeted both
21 at the population level and high risk groups [4,33]. For example – tackling the threats of obesity,
22 smoking and hypertension, addressing health inequalities and supporting efforts to prevent
23 avoidable AKI at population level, as well as targeting those at high risk of CKD progression [34].
24 The GBD data suggest that the highlighted countries in table 2 could be considered 'high risk
25 populations' and should become the focus of greater efforts to identify causal pathways and
26 support public health nephrology interventions. This is likely to require substantial investment,
27 strategic planning and political will, but will have co-benefits such as reducing the burden of
28 other long term conditions, which may be at least as important to population health gain as is the
29 prevention of CKD. Efforts to increase the availability and impact of low tech interventions
30 through public health campaigns (such as dietary modification (including lowering high salt
31 intake), smoking cessation and physical activity programmes) and primary care (such as
32 treatment of hypertension) may yield greater health gain than increasing the availability of
33 expensive high cost technologies. As far as high risk groups are concerned, it has been
34 suggested that the treatment of diabetes and hypertension has improved in high income
35 countries, as RRT incidence rates for these two causes are decreasing. Future studies need to
36 investigate which interventions are most effective in low and middle income countries [34].
37
38
39
40
41
42
43
44
45
46
47
48

49 In conclusion, GBD data suggest both death rates and DALY rates due to CKD are
50 increasing, be it with substantial variation across the globe. It is possible that this is in part due to
51 new data becoming available or different coding behaviour over time, including greater
52 specificity of coding. However, although some feel there is evidence of overdiagnosis, it seems
53 clear that in many countries and regions CKD is a growing public health problem and in some of
54 them ranking very high as cause of YLL and DALYs. Policies to address primary and secondary
55 prevention remain highly needed.
56
57
58
59
60

References

1. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification and stratification. *Am J Kidney Dis* 2002;39:S46-S75.
2. Kidney Disease Improving Global Outcomes (KDIGO) CKD Working Group. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int Suppl* 2013;3;S1-150.
3. Girndt M, Trocchi P, Scheidt-Nave C, Markau S, Stang A. The Prevalence of Renal Failure. *Dtsch Arztebl Int* 2016;113:85-91.
4. Brück K, Stel VS, Gambaro G et al. CKD Prevalence Varies across the European General Population. *J Am Soc Nephrol*. 2015 Dec 23. Epub ahead of print.
5. Fraser SDS, Aitken G, Taal MW et al. Exploration of Chronic Kidney Disease Prevalence Estimates Using New Measures of Kidney Function in the Health Survey for England. *PLoS ONE* 2015; 10(2): e0118676. doi:10.1371/journal.pone.0118676
6. Zhang L, Wang F, Wang L et al. Prevalence of chronic kidney disease in China: a cross-sectional survey. *Lancet*. 2012;379:815-22.
7. Arora P, Vasa P, Brenner D et al. Prevalence estimates of chronic kidney disease in Canada: results of a nationally representative survey. *CMAJ* 2013;185:E417-23.
8. White SL, Polkinghorne KR, Atkins RC, Chadban SJ. Comparison of the prevalence and mortality risk of CKD in Australia using the CKD Epidemiology Collaboration (CKD-EPI) and Modification of Diet in Renal Disease (MDRD) Study GFR estimating equations: the AusDiab (Australian Diabetes, Obesity and Lifestyle) Study. *Am J Kidney Dis* 2010;55:660-70.
9. Levey AS, Coresh J. Chronic kidney disease. *Lancet* 2012;379:165-80.
10. Gansevoort RT, Matsushita K, van der Velde M et al. Lower estimated GFR and higher albuminuria are associated with adverse kidney outcomes. A collaborative meta-analysis of general and high-risk population cohorts. *Kidney Int* 2011;80:93-104.

- 1
2
3 11. Hallan SI, Matsushita K, Sang Y et al. Age and association of kidney measures with mortality
4 and end-stage renal disease. *JAMA* 2012;308:2349-60.
5
6
- 7
8 12. United States Renal Data System. 2015 USRDS annual data report: Epidemiology of Kidney
9 Disease in the United States. National Institutes of Health, National Institute of Diabetes and
10 Digestive and Kidney Diseases, Bethesda, MD, 2015.
11
12
- 13
14 13. Liyanage T, Ninomiya T, Jha V et al. Worldwide access to treatment for end-stage kidney
15 disease: a systematic review. *Lancet* 2015;385:1975-82.
16
17
- 18
19 14. Chronic Kidney Disease Surveillance project at <http://www.cdc.gov/ckd>, accessed 28042016.
20
21
- 22
23 15. Pippias M, Jager KJ, Kramer A et al. The changing trends and outcomes in renal replacement
24 therapy: data from the ERA-EDTA Registry. *Nephrol Dial Transplant* 2016;31:831-41.
25
26
- 27
28 16. GBD 2013 Mortality and Causes of Death Collaborators. Global, regional, and national age-sex
29 specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic
30 analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;385:117-71.
31
32
- 33
34 17. GBD 2013 DALYs and HALE Collaborators. Global, regional, and national disability-adjusted
35 life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188
36 countries, 1990-2013: quantifying the epidemiological transition. *Lancet* 2015;386:2145-91.
37
38
- 39
40 18. Global Burden of Disease Study 2013 Collaborators. Global, regional, and national incidence,
41 prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188
42 countries, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*
43 2015;386:743-800.
44
45
- 46
47 19. Murray CJ, Lopez AD. Measuring the global burden of disease. *N Engl J Med* 2013;369:448-57.
48
49
- 50
51 20. Murray CJ, Vos T, Lozano R et al. Disability-adjusted life years (DALYs) for 291 diseases and
52 injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study
53 2010. *Lancet* 2012;380:2197-223.
54
55
- 56
57 21. Wang H, Dwyer-Lindgren L, Lofgren KT et al. Age-specific and sex-specific mortality in 187
58 countries, 1970-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*
59 2012;380:2071-94.
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
22. Salomon JA, Vos T, Hogan DR et al. Common values in assessing health outcomes from disease and injury: disability weights measurement study for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2129-43.
23. Institute for Health Metrics and Evaluation (IHME). GBD Compare. Seattle, WA: IHME, University of Washington, 2015 at <http://vizhub.healthdata.org/gbd-compare>, accessed 28042016.
24. Chawla LS, Eggers PW, Star RA, Kimmel PL. Acute kidney injury and chronic kidney disease as interconnected syndromes. *N Engl J Med* 2014; 371:58-66.
25. Cusumano AM, González Bedat MC. Chronic kidney disease in Latin America: time to improve screening and detection. *Clin J Am Soc Nephrol* 2008;3:594-600.
26. Kierans C, Padilla-Altamira C, Garcia-Garcia G, Ibarra-Hernandez M, Mercado FJ. When health systems are barriers to health care: challenges faced by uninsured Mexican kidney patients. *PLoS One* 2013;8:e54380. doi: 10.1371/journal.pone.0054380.
27. Farag YM, Kari JA, Singh AK. Chronic kidney disease in the Arab world: a call for action. *Nephron Clin Pract* 2012;121:c120-3.
28. Hassanien AA, Al-Shaikh F, Vamos EP, Yadegarfar G, Majeed A Epidemiology of end-stage renal disease in the countries of the Gulf Cooperation Council: a systematic review. *JRSM Short Rep* 2012;3:38.
29. Habibzadeh F. Kidney disease in the Middle East. www.lancet.com, Middle East edition, August 2013.
30. World population prospects; key findings and advance tables. United Nations 2015 Revision, United Nations, New York, 2015 at http://esa.un.org/unpd/wpp/Publications/Files/Key_Findings_WPP_2015.pdf, accessed 05052016
31. Rhee CM, Kovesdy CP. Epidemiology: Spotlight on CKD deaths—increasing mortality worldwide. *Nat Rev Nephrol*. 2015;11:199-200.

1
2
3 32. Moynihan R, Glassock R, Doust J. Chronic kidney disease controversy: how expanding
4 definitions are unnecessarily labelling many people as diseased. BMJ 2013 Jul 29;347:f4298. doi:
5 10.1136/bmj.f4298.
6
7

8
9 33. Brück K, Stel VS, Fraser S et al. Translational research in nephrology: chronic kidney disease
10 prevention and public health. Clin Kidney J. 2015;8:647-55.
11

12
13 34. Mehta RL, Cerdá J, Burdmann EA et al. International Society of Nephrology's Oby25 initiative
14 for acute kidney injury (zero preventable deaths by 2025): a human rights case for nephrology.
15 Lancet 2015;385:2616-43.
16
17

18
19 35. <http://ghdx.healthdata.org/global-burden-disease-study-2013-gbd-2013-data-downloads>.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Conflict of interest statement**
4

5
6 The results presented in this paper have not been published previously in whole or part.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 1. Global age-standardised death rate and age standardised DALY rate due to CKD over time, total and by cause (per 100,000 people) [35]

		1990	2013	% change and direction
Age-standardised death rate	CKD total	11.6 (10.4 - 12.3)	15.8 (13.5 - 17.1)	36% increase
	CKD due to diabetes	1.4 (1.0 - 1.6)	2.9 (2.3 - 3.5)	107% increase
	CKD due to hypertension	3.6 (2.8 - 4.4)	4.6 (3.3-5.6)	28% increase
	CKD due to glomerulonephritis	2.5 (2.1 - 2.9)	1.8 (1.5 - 2.3)	28% decrease
	CKD due to other causes	4.1 (3.3 - 4.8)	6.5 (4.9 - 7.5)	59% increase
Age standardised DALY rate	CKD total	443.4 (386.8 - 500.9)	497.3 (427.5 - 557.2)	12% increase
	CKD due to diabetes	59.3 (48.9 - 70.9)	90.9 (77.1 - 105.9)	53% increase
	CKD due to hypertension	116.7 (94.4-136.8)	121.1 (95.7 - 139.8)	4% increase
	CKD due to glomerulonephritis	114.4 (100.0 - 130.5)	88.2 (74.0 - 103.2)	23% decrease
	CKD due to other causes	153.0 (130.8 - 199.8)	197.2 (162.6 - 225.1)	29% increase

Table 2. Countries for which CKD was ranked in the top ten for YLL and DALYs in 2013

Years of Life Lost		Disability Adjusted Life Years	
Region / Country	CKD rank in country	Region / Country	CKD rank in country
Andean Latin America Ecuador	7	Andean Latin America Ecuador	8
Central Latin America Mexico Nicaragua Costa Rica El Salvador Panama Venezuela	2 3 4 4 7 7	Central Latin America Nicaragua Mexico Costa Rica El Salvador Panama Venezuela	2 3 7 5 10 10
Tropical Latin America Paraguay	9	Southern Latin America Chile	10
Caribbean Barbados Jamaica The Bahamas Dominica Grenada TTO (Trinidad and Tobago) Antigua and Barbuda Saint Lucia Cuba Dominican Republic VCT (St Vincent and the Grenadines) Suriname	5 6 7 8 8 8 9 9 10 10 10 10	Caribbean Barbados Antigua and Barbuda Dominica Jamaica The Bahamas TTO (Trinidad and Tobago)	7 9 9 9 9 8
North Africa and Middle East Iraq Saudi Arabia Kuwait Palestine Libya Algeria Bahrain Egypt Jordan Lebanon Oman Tunisia	6 6 7 7 8 9 9 9 9 10 10 10	North Africa and Middle East Iraq	9
Eastern sub-Saharan Africa Mauritius Seychelles	4 10	Eastern sub-Saharan Africa Mauritius	4
Oceania Fiji Samoa Marshall Islands FSM (Federated States of Micronesia)	7 7 8 9	Oceania Fiji Samoa Marshall Islands	8 9 10
High income Asia Pacific Singapore	7		
Southeast Asia Thailand Maldives Malaysia	7 9 10	Southeast Asia Thailand	8

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Western Europe Greece	10	Western Europe	
		Israel	6
		Malta	9
		Portugal	9
		Greece	9
		Cyprus	10
		Germany	10
		Central Europe	
		Macedonia	10
		Montenegro	10

For Peer Review

Supplementary table a. International Classification of Diseases (ICD) codes used by Global Burden of Disease (GBD)

GBD Name		ICD code		ICD term
Chronic kidney diseases	Chronic kidney disease due to diabetes mellitus	ICD10	E10.2	Type 1 diabetes mellitus with renal complications
			E11.2	Type 2 diabetes mellitus with renal complications
			E12.2	Malnutrition-related diabetes mellitus with renal complications
			E13.2	Other specified diabetes mellitus with renal complications
		ICD9	250.4	Diabetes with renal manifestations
	Chronic kidney disease due to hypertension	ICD10	I12.0	Hypertensive renal disease with renal failure
			I12.9	Hypertensive renal disease without renal failure
			I13.1	Hypertensive heart and renal disease with renal failure
			I13.2	Hypertensive heart and renal disease with both (congestive) heart failure and renal failure
			I13.9	Hypertensive heart and renal disease, unspecified
			ICD9	403
		404		Hypertensive heart and renal disease
	Chronic kidney disease unspecified	ICD10	N02	Recurrent and persistent haematuria
			N03	Chronic nephritic syndrome
			N04	Nephrotic syndrome
			N05	Unspecified nephritic syndrome
			N06	Isolated proteinuria with specified morphological lesion
			N07	Hereditary nephropathy, not elsewhere classified
			N15.0	Other renal tubulo-interstitial diseases
ICD9		581	Nephrotic syndrome	
		582	Chronic glomerulonephritis	
		583	Nephritis and nephropathy	
		589	Small kidney of unknown cause	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure legends

Figure 1. Crude death rate (upper panel) and percentage of total deaths attributed to CKD (lower panel) over time, by region. For clarity, the figure does not include 95%Uls.

Figure 2. Crude DALY rate (upper panel) and percentage of DALYs attributed to CKD (lower panel) over time, by region. For clarity, the figure does not include 95%Uls.

Supplementary material

Figure a. Age-standardised death rates by region. For clarity, the figure does not include 95%Uls.

For Peer Review




Figure 1. Crude death rate (upper panel) and percentage of total deaths attributed to CKD (lower panel) over time, by region. For clarity, the figure does not include 95%UIs.

120x116mm (600 x 600 DPI)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Crude DALY rate (upper panel) and percentage of DALYs attributed to CKD (lower panel) over time, by region. For clarity, the figure does not include 95%UIs.

109x98mm (600 x 600 DPI)






Figure a. Age-standardised death rates by region. For clarity, the figure does not include 95%UIs.

86x46mm (600 x 600 DPI)

Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60