

Challenging convention: the winter ecology of brown trout (Salmo trutta) in a productive and stable environment

Journal:	Freshwater Biology
Manuscript ID	FWB-P-Aug-16-0387.R1
Manuscript Type:	Standard Paper
Date Submitted by the Author:	17-Sep-2016
Complete List of Authors:	Kemp, Paul; University of Southampton, International Centre for Ecohydraulic Research, School of Civil Engineering and the Environment,; Vowles, Andrew; University of Southampton, Faculty of Engineering and the Environment Sotherton, Nick; Game and Wildlife Conservation Trust Roberts, Dylan; Game and Wildlife Conservation Trust Acreman, Mike; Centre for Ecology and Hydrology, ; KARAGEORGOPOULOS, Perikles; Environment Agency
Keywords:	Fresh waters < Habitat, Running water / rivers / streams < Habitat, Individual < Level of Organisation, Behaviour < Process / Approach / Methods, Fish < Taxonomic Group / Assemblage



Page 1 of 45

Freshwater Biology

2		
3	1	Challenging convention: the winter ecology of brown trout (Salmo trutta) in a
4 5	2	nuclustive and stable anniverment
6	2	productive and stable environment
7		
8	3	
9 10		
11	Л	PALL S KEMP [*] ANDREW S VOWLES [*] NICK SOTHERTON [†] DVI AN ROBERTS [†]
12	-	TROE 5. REAL TROPERTY 5. TO WEES, MOR SO THERION, DIENTRODERTS,
13	5	MICHAEL C. ACREMAN [‡] AND PERIKLES KARAGEORGOPOULOS ^{**}
14 15		
16	6	
17	6	International Centre for Econydraulics Research, University of Southampton, Southampton,
18	7	
19 20	,	
21	8	[†] Game and Wildlife Conservation Trust, Fordingbridge, UK;
22		
23	9	[‡] Centre for Ecology and Hydrology, Wallingford, UK;
24	4.0	
26	10	Environment Agency, worthing, UK.
27	11	
28		
30	12	Correspondence: Paul S. Kemp, International Centre for Ecohydraulics Research, Faculty of
31		
32	13	Engineering and the Environment, University of Southampton, Southampton, SO17 1BJ. UK.
33 34		
35	14	e-mail: p.kemp@soton.ac.uk
36	15	
37	15	
38 39		
40	16	Keywords: brown trout, chalk stream, habitat models, Habitat Template Concept, Salmonid,
41	47	
42	17	Site fidelity
43 44		
45	18	Running heading: The winter ecology of brown trout
46		
47	10	
48 49	19	
50		
51	20	
52		
53 54	21	
55	<u> </u>	
56		
57	22	
วช 59		
60		1

23 SUMMARY

25	1. Understanding of the winter ecology of stream salmonids is biased by research conducted
26	in northern temperate and boreal regions dominated by hard rock geology. Such systems are
27	driven by highly dynamic surface flow regimes and tend to be physically diverse, nutrient
28	poor, and influenced by ice. This study investigated how the behaviour of brown trout,
29	Salmo trutta, inhabiting a stable groundwater-fed, productive, and comparatively warm
30	southern English chalk stream differs from that described for other systems, and how this is
31	translated to performance, measured as growth.
32	2. Physical characteristics were mapped, and high-resolution temperature data collected using
33	a spatial array of data loggers installed throughout the study reach during the winter. A
34	combination of passive integrated transponder and radio telemetry was used to monitor
35	distribution, density, and movement of trout. Micro-archival data storage tags inserted in
36	some individuals provided information on temperature regimes experienced. Growth
37	performance was calculated for recaptured fish.
38	3. Trout density was positively related to depth and there was no evidence that temperature
20	
39	influenced microhabitat selection. Three patterns of movement were observed. Over three-
39 40	influenced microhabitat selection. Three patterns of movement were observed. Over three- quarters of tracked fish exhibited high site fidelity and tended to remain in a single focal
40 41	influenced microhabitat selection. Three patterns of movement were observed. Over three- quarters of tracked fish exhibited high site fidelity and tended to remain in a single focal position throughout the study. Fourteen percent of trout exploited more than one distinct
39404142	influenced microhabitat selection. Three patterns of movement were observed. Over three- quarters of tracked fish exhibited high site fidelity and tended to remain in a single focal position throughout the study. Fourteen percent of trout exploited more than one distinct location, while the remainder were detected at multiple locations and showed no preference
 39 40 41 42 43 	influenced microhabitat selection. Three patterns of movement were observed. Over three- quarters of tracked fish exhibited high site fidelity and tended to remain in a single focal position throughout the study. Fourteen percent of trout exploited more than one distinct location, while the remainder were detected at multiple locations and showed no preference for any one.
 39 40 41 42 43 44 	 influenced microhabitat selection. Three patterns of movement were observed. Over three-quarters of tracked fish exhibited high site fidelity and tended to remain in a single focal position throughout the study. Fourteen percent of trout exploited more than one distinct location, while the remainder were detected at multiple locations and showed no preference for any one. 4. Trout exhibited regular daily activity patterns and highly periodic local movements at dusk
 39 40 41 42 43 44 45 	 influenced microhabitat selection. Three patterns of movement were observed. Over three-quarters of tracked fish exhibited high site fidelity and tended to remain in a single focal position throughout the study. Fourteen percent of trout exploited more than one distinct location, while the remainder were detected at multiple locations and showed no preference for any one. 4. Trout exhibited regular daily activity patterns and highly periodic local movements at dusk and dawn and tended to experience positive growth performance during periods that included
 39 40 41 42 43 44 45 46 	 influenced microhabitat selection. Three patterns of movement were observed. Over three-quarters of tracked fish exhibited high site fidelity and tended to remain in a single focal position throughout the study. Fourteen percent of trout exploited more than one distinct location, while the remainder were detected at multiple locations and showed no preference for any one. 4. Trout exhibited regular daily activity patterns and highly periodic local movements at dusk and dawn and tended to experience positive growth performance during periods that included winter.

Freshwater Biology

5. This study challenges the conventional view of salmonid winter ecology, which is biased towards populations that inhabit hard rock surface-flow dominated rivers that experience the influence of ice. Despite inhabiting a distinctly different winter habitat template than more commonly studied populations, trout occupying a hydrologically stable and productive chalk stream exhibited behaviours similar to those described for elsewhere, yet performed considerably better.

55 Introduction

Physical and chemical environments vary with time and space, and the rate and extent by which they change characterise degree of habitat heterogeneity. The unequal challenges and opportunities imposed by heterogeneous habitats influence ecological organisation; the habitat acts as a template (Southwood 1977) that defines characteristics operating at the population and community (e.g. species diversity and abundance, Tews et al. 2004) levels (Poff and Ward 1990). Spatial and temporal heterogeneity drives the individual adaptive response adopted to enable exploitation of opportunities encountered to maximise fitness. Thus, the relationship between behavioural traits and habitat heterogeneity should be predictable.

66 Stream-dwelling salmonids provide a particularly apt model to compare behaviour 67 and performance among populations that experience differing degrees of habitat 68 heterogeneity. Throughout their range, salmonids inhabit diverse river types where they may 69 experience dynamic environments that impose severe ecological challenges, particularly

during the winter bottleneck (Armstrong et al. 2003). Understanding of the winter ecology of salmonids is primarily based on research conducted in highly heterogeneous habitats, with little comparison made with populations that inhabit relatively stable environments. The current study focused on a population of brown trout, *Salmo trutta*, inhabiting a southern English chalk stream during the winter, which when compared with systems that have received most attention, tend to be both spatially and temporally more physically stable, and substantially more productive.

Based on current understanding, the conditions imposed by the winter template are among the most critically challenging during the life history of stream dwelling salmonids. Stressful temporal environmental change may limit abundance (Cunjak et al. 1998) and productivity (Armstrong et al. 2003) through a variety of mechanisms. These include limiting the availability of suitable habitat (Cunjak 1996) and creating adverse conditions that isolate and damage fish when ice forms (Huusko et al. 2007), or displace them during scour and flooding events with the thaw (Cunjak et al. 1998). Furthermore, declines in temperature not only regulates physiological (e.g. maintenance of energy reserves; Finstad et al. 2004) and locomotory (e.g. Webb 1978) performance during the winter, but also a range of other biotic relationships. For example, shifts in predator prey dynamics in favour of homeothermic mammalian and avian piscivores has been described (Fraser et al. 1993), while production and drift of invertebrate food is typically reduced as temperature declines (Simpkins and Hubert 2000, but see Cunjak and Power 1987).

In response to the wintertime challenges imposed, salmonids display a multitude of adaptive behaviours. The benefits to juveniles of maintaining territories, as during the summer, are lost as costs of maintenance increase relative to energetic gains when food becomes limited (Dill et al. 1981). Aggressive interactions become less common (Heggenes et al. 1993) as fish become more gregarious and form aggregations in pools (e.g. Heggenes et

Page 5 of 45

1

Freshwater Biology

2	
3	
4	
5	
6	
7	
8	
a	
10	
10	
11	
12	
13	
14	
15	
16	
17	
18	
10	
20	
20	
21	
22	
23	
24	
25	
26	
27	
28	
20	
20	
24	
31	
32	
33	
34	
35	
36	
37	
38	
39	
10	
11	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
52	
23	
54	
55	
56	
57	
58	
59	
60	

95	al. 1999). Further, stream salmonids become increasingly nocturnal (Heggenes et al. 1993)
96	below some threshold temperature (Fraser et al. 1993), and commonly seek shelter during the
97	day (Heggenes et al. 1999) to avoid visual predators (Orpwood et al. 2006, 2010).
98	At the onset of winter, salmonids commonly seek alternative microhabitat that
99	provides refuge from both predators and unfavourable hydrodynamic and thermal conditions.
100	Compared to the summer and autumn, winter microhabitat is associated with slower water
101	velocities (Mäki-Petäys et al. 1997), greater depth (Heggenes et al. 1999), and access to
102	shelter provided by larger non-compacted substrate (Stickler et al. 2008), submerged woody
103	material (Swales et al. 1986), and overhead cover (Cunjak and Power 1986), including
104	surface ice (Linnansaari et al. 2009). The availability of suitable microhabitats can dictate
105	overwinter survival (e.g. Solazzi et al. 2000; Hedger et al. 2013), and thus influence
106	population abundance and productivity (Cunjak et al. 1998).
107	The generalised perspective of the winter ecology of stream salmonids so far
108	considered focuses predominantly on populations inhabiting systems that exhibit high
109	physical and chemical heterogeneity over both spatial and temporal scales (Table 1). Past

research has often focused on northern temperate and boreal rivers and streams based on a 110 111 hard rock geology and driven by surface flow, such as those that occur in Canada (e.g. Catamaran Brook, Cunjak et al. 1998; Linnansaari and Cunjak 2010) and Scandinavia (e.g. 112 113 Orkla River, Norway, Stickler et al. 2007; River Alta, Norway, Hedger et al. 2013). These 114 rivers experience dynamic flow regimes as they rapidly respond to precipitation and snow 115 melt events, have gravel beds with a wide range of substrate size classes, and well developed 116 frequent riffle-pool sequences. Salmonid populations that inhabit them often are adapted to 117 nutrient poor, low pH conditions, and, due to the wide fluctuations in temperatures experienced, may suffer extensive periods of freezing and the influence of ice dynamics. 118 119 Conversely, salmonids that inhabit lowland groundwater-fed systems experience quite

different environmental conditions. For example, in southern England, several lowland
salmon and trout rivers flow through chalk catchments, and do not share the same
heterogeneous physical, chemical, and ecological characteristics described for upland gravelbed rivers (Table 1). Referred to as chalk streams, these rivers are predominantly aquifer fed
and so maintain relatively constant annual flow and temperature regimes. Many chalk
streams exhibit a gravel bed matrix with limited geomorphological diversity owing to their
characteristically low stream power.

To improve understanding of how environmental variability influences behaviour, this study explored the winter ecology of a stream dwelling salmonid in a southern English chalk stream. We aimed to quantify: 1) trout behaviour as indicated by distribution relative to available habitat, patterns of movement, and levels of activity; and 2) trout performance, measured as growth. Compared with more dynamic systems, chalk streams during the winter are: a) spatially homogeneous, in terms of hydrodynamic and physical characteristics, b) warm and ice free, c) temporally constant in relation to flow and food abundance, and d) highly productive. Therefore, we predicted that under these less challenging conditions there would be: a) no need, and less opportunity, for fish to select refugia from adverse physical conditions (e.g. hydrodynamics and ice formation), and thus a lack of an association with deep, slow flowing areas; b) a positive relationship between fish density and areas of warm water, potentially associated with groundwater upwelling, to enhance growth; c) a high degree of site fidelity and limited movements of fish, including at night; and d) a high growth performance during the winter when compared with boreal systems.

- 142 Methods

Freshwater Biology

Site description, and physical and thermal characteristics

The study was conducted at a 500 m long reach of the River Lambourn (owned by the Centre for Ecology & Hydrology [CEH]) at Boxford (51.445542,-1.382947), Berkshire, UK (Figure 1). The site is situated approximately two-thirds (14 km) of the river length from the ephemeral source near the village of Lambourn. The river flows through a catchment (upstream area: 162 km^2) dominated by arable and mixed farmland prior to entering the River Kennet, a tributary of the River Thames. The catchment, dominated by cretaceous chalk geology with deposits of clays and flints, lacks impermeable strata and overland flow resulting in a low drainage density $(0.12 \text{ km km}^{-2})$ (Evans et al. 2003). The mean river discharge (\pm SD and range) at the site during the two study winters was 0.90 m³ s⁻¹ (winter 1: $\pm 0.23, 0.53-1.33$) and 0.52 m³ s⁻¹ (winter 2: $\pm 0.06, 0.36-0.98$) (Rameshwaran et al. 2015). Discharge exhibits low seasonal variation and is dominated by a stable groundwater-fed base-flow contribution, which typically exceeds 95% (Evans et al. 2003). The river water chemistry is driven by the calcium-bicarbonate component. The riverbed substrate comprises of gravel and coarse sub-angular flint and the channel gradient is 0.05° (Evans et al. 2003). A baseline survey was conducted between 23 and 26 August 2011 to quantify the spatial variability of key physical habitat characteristics throughout the study reach. Key parameters were recorded at approximately five equidistant points along transects that spanned the width of the channel at 10 m longitudinal intervals. Temporal variability was quantified by conducting a survey of a 270 m section of the study reach on 22 February 2011. Dominant substrate type was visually classified using the Wentworth grain size scale and ranged from silt (< 0.0063 cm) to pebble (1.6-6.4 cm), but was dominated by gravel (0.2 -1.6 cm (Table 2, Figure 2a). Water depth and mean (\pm SD) mid-column (60% depth) water

2
2
3
4
5
6
7
1
8
9
10
11
10
12
13
14
15
10
10
17
18
19
20
20
21
22
23
24
24
25
26
27
20
20
29
30
31
22
32
33
34
35
26
30
37
38
39
10
40
41
42
43
44
44
40
46
47
48
40
49
50
51
52
52
03
54
55
56
57
57
58
59
60

1

168	velocity was measured using a metre rule and electromagnetic flow meter (0.001 m s ⁻¹
169	resolution averaged over 10 seconds, Valeport Model 801, Totnes, UK), respectively.
170	Shallow, moderate velocity reaches dominated the upper and middle sections interspersed
171	with a few deep (≥ 1 m) low velocity areas, and a deep reach at the downstream end of the
172	site (Figure 2b, c). There was nearly continuous riparian cover along the true right bank, but
173	scarce vegetation on the left.
174	Data loggers accurate to 0.47°C (Hobo Temp/Light Pendant, Onset Computer
175	Corporation, MA, USA) recorded river temperature hourly at transects positioned
176	approximately every 20 m along the study reach. Three loggers were deployed at each
177	transect; one each at the left and right bank, and one in the centre of the channel. Loggers
178	recorded temperature at the substrate. Two additional loggers recorded ambient air
179	temperature in the shade at hourly intervals. Missing data points (e.g. due to loss of loggers,
180	delays in downloading, and equipment failure) were substituted with values (T_r) obtained by
181	regressing mean daily water against air temperature for each position. Winter temperature
182	regime obtained for the Svarttjønnbekken River (Norway) over the same period allowed
183	comparison between the River Lambourn and a northern boreal system.
184	During Winter 1 the UK experienced heavy snowfalls and record low temperatures,
185	with the coldest (mean air temperature: -1°C) December recorded since Met Office records
186	began in 1910 (Met Office data). On the 20 December 2010 a minimum ambient air
187	temperature of -13.72°C was recorded at the study site. Mean UK air temperatures for

188 December and January were 5.0°C and 0.3°C below average. The mean temperatures in the

189 UK were much milder during Winter 2, with December, January and February being 0.6°C,

190 1.3°C and 0.7°C above average (Met Office data). Despite this, a cold spell lasting 2 weeks in

191 England started at the end of January. This resulted in a minimum winter ambient air

Freshwater Biology

2
3
4
5
6
7
1
8
9
10
11
12
13
14
15
16
17
18
10
19
20
21
22
23
24
25
26
27
20
20
29
30
31
32
33
34
35
36
27
31
38
39
40
41
42
43
44
45
46
- 1 0 //7
+/ 40
48
49
50
51
52
53
54
55
56
50
э/ 50
58
59
60

192	temperature recorded at the study site of -12.81°C on 11 February 2012. River temperature
193	varied temporally during the two winter periods (Figure 3). Average mean river temperatures
194	during Winters 1 and 2 (recorded from start of December until end of February) were 7.55°C
195	and 7.84°C, respectively. This is consistently higher than the northern boreal system
196	(Svarttjønnbekken River, Norway), which remained close to zero (Figure 3). River
197	Lambourn water temperatures showed high variability (average SD Winter 1: 0.38; Winter 2:
198	0.52), being closely correlated with ambient air temperature throughout the two periods
199	(Winter 1: $\alpha = 6.5$, $\beta = 0.42$, $R^2 = 94.5\%$, $p < 0.001$; Winter 2: $\alpha = 6.3$, $\beta = 0.40$, $R^2 = 85.0\%$,
200	p < 0.001). River temperatures varied spatially (Figure 2d). For example, on the coldest day
201	in Winter 2 (11 February 2012, mean air temperature = -6.6°C) maximum and minimum river
202	temperatures were recorded as 8.1 and 2.7°C at locations 52.7 and 81.3 cm deep, respectively
203	
204	Fish distribution, movement, and activity
205	
206	The distribution, movement and activity of fish were recorded at coarse and meso-scale
207	resolution using a combination of electric fishing, mark recapture, and telemetry surveys.

208 Trout were captured through the entire study site by electric fishing upstream through

209 20 m long reaches (confined by stop nets) using a single pass on five occasions (Table 3).

210 Trout captured were anesthetised (0.3 ml L⁻¹ 2-phenoxyethanol), measured (fork length mm),

211 weighed (g), and scanned to identify recaptured individuals tagged in previous surveys. If

sufficiently large, trout were implanted with 12 mm full duplex Passive Integrated

213 Transponder (PIT) (Wyre Micro Design Ltd, Lancashire, UK; > 100 mm fork length), radio

214 (24 mm length, 1.9 g mass in air, estimated battery life = 8.7 months; Biotrack, Wareham,

2
2
3
4
5
6
7
8
9
10
10
11
12
13
14
15
16
17
18
10
19
20
21
22
23
24
25
26
20
21
28
29
30
31
32
33
24
34
35
36
37
38
39
10
11
41
42
43
44
45
46
47
48
10
73 50
50
51
52
53
54
55
56
50
57
58
59
60

1

215	UK; > greater than 150 g mass), and micro-archival data storage (DS) (25.4 mm length, 3.3 g
216	mass in air, estimated battery life = 18 months; DST micro-T, Star Oddi, Reykjavik, Iceland; >
217	300 g) tags (Table 3). The total tag burden did not exceed 2 % of the fish body mass. Fish
218	were not tagged during the final survey. Prior to surgery, the functionality and unique
219	frequency (between 173.199 and 173.994 MHz) of all radio tags was verified using a hand-
220	operated receiver, and DS tags were programmed to record fish body temperature every 15
221	minutes. Fish were allowed to recover (approx. 1 hour) in tanks containing aerated river
222	water before being returned to the electric fishing reach from which they were captured.
223	Standard tagging protocols were conducted in compliance with the UK Animals (Scientific
224	Procedures) Act 1986 under Home Office licence.
225	
226	Distribution and density: Trout density for each (20 m) electric-fishing reach was calculated
227	using data from the winter electric-fishing survey (February 2011) as the quotient of number
228	of fish captured and surface area (quantified using the GIS base map) of the reach. A linear
229	regression model was fitted to determine whether habitat variables (depth, velocity and water
230	temperature) were significant predictors of density during winter.
231	Coarse-scale site fidelity was described as the percentage of previously tagged fish
232	recaptured within ± 1 electric fishing reach during subsequent surveys.
233	To investigate whether trout selected and tracked thermal microhabitat to maximise
234	growth performance, i.e. areas exhibiting the highest temperatures (e.g. groundwater

235 upwelling or a tributary confluence), the difference between mean daily body and maximum

236 daily river temperatures (ΔT) was calculated for recaptured fish containing DS tags. To assess

- 237 how this changed during low temperature periods when fish may experience highest thermal
- 238 stress, ΔT was plotted against minimum daily river temperature. If trout exhibited

Freshwater Biology

behavioural thermoregulation they were expected to track patches that provided the greatest
opportunity for growth, i.e. those areas where maximum water temperatures were recorded.
In this case, no difference between maximum daily river and mean daily body temperature
was expected, and this would remain the case independent of minimum daily river
temperature. In the event that trout exhibit high site fidelity, a difference between maximum
daily river and mean daily body temperature should be observed, and this would unlikely
remain constant over time.

Movement: Meso-scale movements were assessed using telemetry during two study periods: November to April in 2010-2011 (Winter 1) and 2011–2012 (Winter 2). Radio tagged trout were located from the bank using a hand-held receiver (Sika, Biotrack, Wareham, UK) connected to a three-element Yagi antenna during systematic surveys of the entire site up to 3 times per week. Tagged fish were detected up to distances of approximately 250 m. Once located, the position of individual fish was estimated by reducing the gain on the receiver and moving longitudinally along the bank in the direction of increasing signal strength until the best fix was attained, with occasional validation through visual identification of some individuals. Fish positions were recorded relative to bankside landmarks of known location. Fish positions derived from mobile radio tracking were imported into ArcGIS, plotted on the base map of the river and underlying water depth data extracted. The relationship between fork length and average depth occupied during mobile radio tracking surveys was expressed using Pearson's correlation.

Using fish positions derived from mobile radio tracking, patterns of individual movement were quantified using two metrics: 1) distance moved (m day⁻¹), calculated as the distance between successive radio tracking fixes divided by number of days between fixes, and 2) longitudinal home range defined as the distance between the most up- and downstream locations recorded (Khan et al. 2004; Ovidio et al. 2002). Longitudinal home ranges

were calculated from 95% trimmed data. For each fish, the number of days between first and
last detection (detection period) and number of radio fixes was also recorded.

To identify factors influencing variability in patterns of individual movement, linear regression models were applied to determine whether the independent variables (fish length, mass, detection period and/or number of radio fixes) were significant predictors of either movement metric. Due to multicollinearity between fish length and mass (r = 0.92), these predictors were combined and replaced in the models by condition factor (*K*), calculated as:

$$K = 100 \cdot \frac{W}{L^3}$$

where *W* is fish mass (g) and *L* is fork length (cm). Number of radio fixes was also removed from the model due to multicollinearity with detection period (r = 0.92).

To investigate whether patterns of individual movement differed between the two winter periods Mann-Whitney U tests were used as data violated the assumption of normality, and could not be successfully corrected through transformation. Comparisons between the two winter periods were deemed valid as number of radio fixes and detection period (measures of sampling frequency) were not significant predictors of either movement metric. Activity: Four automated fixed radio receivers (SRX-DL Data loggers, Biotrack, Wareham, UK), installed during Winter 2 at the upper and lower ends of the site, and at two locations previously identified to contain a high number of tagged fish (Figure 2a), continually scanned for up to 20 frequencies. An autocorrelation function (ACF) was performed on mean signal strength per hour for individual fish using the 'correl' command in Microsoft Office Excel (Microsoft Corporation, Redmond, USA), with a one-hour time lag, for 336 hours (14 days) after detection for fish detected during $\geq 20\%$ of those hours. Autocorrelation was deemed to have occurred for fish with ACF peaks outside the limits of 95% confidence.

287

288

Fish performance

1

2	
3	
4	
5	
6	
7	
1	
8	
9	
10	
11	
12	
12	
13	
14	
15	
16	
17	
10	
10	
19	
20	
21	
22	
23	
20	
24	
25	
26	
27	
28	
20	
29	
30	
31	
32	
33	
34	
25	
30	
36	
37	
38	
39	
10	
-+U	
41	
42	
43	
44	
45	
ле 1	
40	
41	
48	
49	
50	
51	
ຮ່ວ	
52	
53	
54	
55	
56	
57	
50	
50	
59	
60	

Performance metrics were based on data collected during electric fishing surveys in July
2010 and February 2011 (which included Winter 1), and September 2011 and May 2012
(Winter 2).

292 Mean specific growth rate (% day⁻¹) for PIT tagged fish recaptured during electric 293 fishing surveys was calculated as:

 $G = 100 \cdot \left((\log_e W_2 - \log_e W_1) / t \right)$

where W_1 and W_2 are the initial and final fish mass (g), and *t* is the number of days between surveys (i.e. the growth period). For each fish, *G* was compared to an estimate of optimal growth (G_{op}) using the model developed by Elliott et al. (1995):

 $G_{\rm op} = c \cdot W_1^{-b} (T - T_{\rm lim}) / (T_{\rm M} - T_{\rm lim})$

where T is the mean water temperature during the growth period, and $T_{\rm M}$ and $T_{\rm lim}$ respectively 297 298 represent the temperatures at which growth is optimal (13.11°C) and ceases (limit). T_{lim} is 299 the lower or upper value at which growth rate is zero ($T_{\rm L}$ [3.56°C] or $T_{\rm U}$ [19.48°C]) depending on whether T is higher or lower than $T_{\rm M}$ (i.e. $T_{\rm lim} = T_{\rm L}$ if $T < T_{\rm M}$ or $T_{\rm lim} = T_{\rm U}$ if T >300 301 $T_{\rm M}$). The mass exponent b is the power transformation that produces linear growth with time 302 (0.308), and c is the growth rate of a 1 g trout at optimal temperature (2.803). All values were 303 obtained from Table 1 in Elliott et al. (1995). The growth model assumes fish fed to satiation 304 under laboratory conditions.

306	Results
307	
308	Fish distribution, movement, and activity
309	
310	Distribution and density: Trout density was positively related to depth; there was no
311	relationship with water temperature or velocity (Table 4; Figure 4).
312	Coarse resolution assessment of site fidelity indicated that 26% ($n = 5$) and 71% ($n =$
313	32) of trout that were recaptured were within ± 1 electric fishing reach during periods that
314	included Winter 1 (July 2010 – February 2011) and 2 (September 2011 – May 2012),
315	respectively.
316	For the five recaptured fish that contained DS tags the relationship between ΔT and
317	minimum daily river temperature was positive (Figure 5). These fish did not utilise and track
318	microhabitat based on selecting high temperature areas that may have maximised growth
319	during winter. Instead, they remained site attached at mean water depths that ranged from
320	41.7 cm to 79.8 cm with body temperature conforming to river temperatures at those
321	locations.
322	Movement: Of the 83 trout implanted with radio tags, 65 were detected during subsequent
323	tracking. Of these, eight were excluded from further analysis due to low levels of certainty of
324	their position or because they were detected once only. Of the remaining 57, 43 (75.4%) were
325	detected within the study site more than once during the winter periods.

Page 15 of 45

1

Freshwater Biology

2	
3	
4	
5	
6	
7	
0	
ð	
9	
10	
11	
12	
13	
11	
15	
10	
16	
17	
18	
19	
20	
21	
22	
22	
23	
24	
25	
26	
27	
28	
20	
29	
30	
31	
32	
33	
34	
35	
36	
27	
31	
38	
39	
40	
41	
42	
43	
⊿⊿	
74	
45	
46	
47	
48	
49	
50	
51	
51	
5Z	
53	
54	
55	
56	
57	
58	
50	
59	
$\mathbf{b}(\mathbf{l})$	

326	There was no relationship between fork length and the average water depth occupied
327	during mobile radio tracking surveys ($r = 0.11, p = 0.493$).

328	Differences in distance moved (median [IQR] = $2.92 [4.07] \text{ m day}^{-1}$) and longitudinal
329	home range (median [IQR] = 21.20 [45.20] m) among individual fish was not predicted by
330	either condition factor or detection period (distance moved: Multiple regression $R^2 = 0.06$,
331	$F_{2,40} = 1.30, p = 0.285$; home range: Multiple regression $R^2 = 0.05, F_{2,40} = 1.03, p = 0.365$).
332	Median distance moved and home range was low and did not differ between the two winter
333	periods (Mann-Whitney U: distance moved $U = 220$, $z = -0.24$, $p = 0.808$; home range $U =$
334	228, $z = -0.06$, $p = 0.951$). However, variability in behaviour among individual fish was
335	evident with patterns of movement categorised into three groups based on levels of site
336	attachment: 1) occupancy of a single position within the study reach (expressed by 76.7% of
337	tracked fish, $n = 33$; e.g. Figure 6a); 2) utilisation of more than one distinct location (14.0%)
338	of fish, $n = 6$; e.g. Figure 6b); or 3) multiple detections throughout the study site with no clean
339	single location for which fish exhibited high site fidelity (9.3% of fish, $n = 4$; e.g. Figure 6c).
340	Larger home ranges (> 65 m) generally indicated occupancy of more than one distinct
341	location within the study site (movement pattern 2 described above) rather than low site
342	fidelity.

Activity: All nine fish detected in the vicinity of the continually logging fixed radio receivers
during Winter 2 exhibited autocorrelated activity. Four fish displayed autocorrelation patterns
with lags roughly every 24 hours, suggesting that they displayed regular daily activity
patterns (e.g. Figure 7a). Two fish, both detected in the vicinity of deep low velocity areas,
displayed particularly high levels of autocorrelation with a consistent 24 hour lag and no
decay (Figure 7b). This is indicative of highly periodic behavioural rhythms. These fish

2	
3	
Δ	
-	
5	
6	
7	
8	
å	
3	
10	
11	
12	
13	
10	
14	
15	
16	
17	
18	
10	
19	
20	
21	
22	
23	
23	
24	
25	
26	
27	
20	
20	
29	
30	
31	
32	
22	
33	
34	
35	
36	
27	
31	
38	
39	
40	
41	
10	
42	
43	
44	
45	
46	
47	
47	
48	
49	
50	
51	
51	
52	
53	
54	
55	
55	
30	
57	
58	
59	
60	

349 moved in and out of the detection range of the receiver at dusk (17:00 h) and dawn (07:00 h),

350 respectively (Figure 7c).

351

1

352 Fish performance

353

On average trout growth during the periods that included winter was positive (Winter 1 mean 354 \pm SD = 0.08 \pm 0.09 % day⁻¹; Winter 2 = 0.22 \pm 0.10 % day⁻¹), but differed among years 355 356 (independent samples *t*-test: t = 5.97, d.f. = 90, p < 0.001), with higher rates in year 2. For 357 some fish, growth approached that predicted by the optimum growth model that assumes fish 358 are fed to satiation. Although rates were suboptimal during both years (Winter 1: U = 2.00, n = 38, z = -5.21, p < 0.001; Winter 2: U = 216.50, n = 145, z = -9.54, p < 0.001; Figure 8), 359 360 there was a positive relationship between observed and predicted growth (for Winter 1: r =0.66, p < 0.01; for Winter 2: r = 0.81, p < 0.001; Figure 8). Trout density and growth rates (% 361 increase in g day⁻¹) were unrelated (Pearson's correlation: r = -0.38, p = 0.055) during mid-362 363 winter (February 2011). 364 Discussion 365 366 367 Understanding of the winter ecology of stream salmonids is largely based on populations that 368 inhabit northern temperate and boreal surface flow dominated rivers with coarse-grained bed substrate situated on hard rock geology. Within this context, Cunjak (1996) provides a 369

Page 17 of 45

1 2

Freshwater Biology

3
4
5
6
7
י פ
0
9 10
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
20
21
20
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47 47
 /2
40
49 50
50
51
ວ∠
53
54
55
56
57
58
59
60

370	definition of winter that focuses on ecological and physical parameters, i.e. the period
371	immediately following egg deposition, which coincides with a decline in water temperature,
372	and that extends until the loss of all surface ice and the accompanying spate and snowmelt.
373	This definition is inappropriate, however, for different hydromorphological systems, such as
374	those fed by groundwater and that do not experience ice dynamics. In this study we
375	considered a groundwater fed chalk stream and defined winter simply as being the coldest
376	season of the year, which in the northern hemisphere extends from December to February.
377	Chalk streams in southern England generally experience a relatively mild climate and
378	consistently high mean annual groundwater temperatures (approximately 11°C), with little
379	variance, and no freeze-up and thaw events. Focusing on the relatively stable and physically
380	and chemically homogenous River Lambourn, this study investigated the behaviour
381	(distribution, movement and activity patterns) and performance (growth) of a local brown
382	trout population occupying an alternative habitat template (Southwood 1977) to those
383	commonly considered. The findings discussed relative to a suite of predictions that are based
384	on prevailing views provide further insight into the winter ecology of salmonids.

385 Salmonids in chalk streams do not face the adverse conditions imposed by freeze-up 386 and thaw and the associated hydrodynamic challenges experienced by fish inhabiting many 387 surface-flow dominated boreal rivers. For example, in the high energy dynamic systems that respond rapidly to precipitation and snowmelt, water velocities exceeding 5 m s⁻¹ may be 388 389 observed e.g. during surges of water and ice as ice dams rupture (Cunjak et al. 1998). Under 390 such conditions, salmonids must attempt to escape the negative effects of high shear stresses 391 and high-velocity scour. Perhaps as a consequence of adaptation to such pressures, stream-392 dwelling salmonids typically move to slower flowing (Mäki-Petäys et al. 1997), deeper areas 393 (Heggenes et al. 1999), with access to large non-compacted substrate (Stickler et al. 2008), and woody structure (Swales et al. 1986) during the winter. Partly due to a more benign 394

environment and the lack of structural complexity we did not expect brown trout to associate strongly with microhabitat based on either depth or velocity in this study. However, although trout distribution was not influenced by velocity, density was positively related to depth, an association that more likely reflected sheltering from predation than from adverse hydrodynamics. The value of deeper areas as refuge is apparent when the availability of alternative forms of shelter is considered. In many chalk streams sufficiently large particles and woody structure is limited, and the highly compacted and weathered nature of chalk bed materials means that the availability of interstitial space is low. Similarly, longitudinal pool and riffle frequency tends to be low in chalk streams, and thus where deeper areas do occur, they likely provide important and much sought after refugia in the absence of macrophyte cover during the winter. As temperature is the most pervasive environmental factor influencing the life history of fishes (Wootton 1998) we predicted that should the study site exhibit thermal heterogeneity, e.g. due to the presence of upwelling ground water or other inputs, then trout would select areas of high temperature in an attempt to maximise fitness. Further, trout were expected to enhance opportunities for growth by tracking favourable temperature habitat patches should they change over time. Although chalk streams are considered homeothermous in character, exhibiting relatively limited variability in water temperature over the annual cycle (Edwards 1979), during the winter much higher temporal variability is observed than for boreal surface flow rivers that experience freeze-up. For example, the Svarttjønnbekken River (Norway) remained close to 0°C throughout the two study winters; and even in the event of supercooling, water temperatures would have declined by only a few

417 hundredths to a tenth of a degree below zero Celsius (Brown et al. 2011). In comparison, the

- 418 mean daily temperatures recorded for the River Lambourn during the two study winters
- 419 (December to February) were much higher, ranging between 7.5°C and 8°C, and more

Page 19 of 45

1 2

Freshwater Biology

3
4
5
6
7
0
ð
9
10
11
12
13
14
15
16
17
10
10
19
20
21
22
23
24
25
20
20
21
28
29
30
31
32
33
34
35
26
30
31
38
39
40
41
42
43
44
45
40
40
41
48
49
50
51
52
53
54
55
55
00
5/
58
59
60

420	variable, being closely correlated with ambient air temperature. Even during unusually cold
421	periods for the UK for both winters, and especially December of winter 1, mean temperatures
422	rarely dropped below 5°C. Further, the study site exhibited spatial variability in local
423	temperatures, which was likely the result of warmer upwelling subsurface flow, the dominant
424	direction of hyporheic exchange at the study site (Pretty et al. 2006), and the contribution of a
425	tributary (Fig. 2d). Nevertheless, trout distribution was not influenced by either temporal or
426	spatial variation in water temperature, and there was no evidence, based on a small sample of
427	recaptured fish holding temperature sensitive data storage tags, that trout selected and then
428	tracked warmer areas of microhabitat that would have enhanced growth potential.

In a stable and productive chalk stream, in which energy intake (high) and shelter 429 430 opportunities (low) are predictable, we expected high site fidelity and low levels of patch 431 switching because the drive to explore new areas was likely to be weak. It is suggested that 432 high site fidelity represents an optimal foraging strategy in stable environments in which resource availability is high and predictable (Arthur et al. 2015). This is contrary to the 433 434 situation that occurs in heterogeneous "patchy" environments, such as that described for surface-flow dominated river systems in which the spatial distribution of drifting invertebrate 435 436 food is limited and temporally unpredictable. In such cases, stream-dwelling salmonids may 437 be predicted to switch patches as profitability of the foraging position declines and they 438 search for higher quality microhabitat. This assumption is supported conceptually (Marginal 439 Value Theorem; Charnov 1976) and through observation (for juvenile Atlantic salmon in 440 Scottish streams: Martin-Smith and Armstrong 2002). In line with our prediction, a large 441 majority of tracked trout exhibited high site fidelity and limited movement, with over three-442 quarters occupying a single location. Alternative strategies were demonstrated, but were rarer. 443 Less than 10% of trout tracked throughout the study area did not favour any distinct locations. 444 However, nearly 28% of the radio-tagged trout were recorded either only once or never again.

2	
3	
٨	
4	
5	
6	
0	
7	
8	
0	
9	
10	
11	
12	
12	
13	
14	
15	
16	
17	
17	
18	
19	
20	
21	
~ '	
22	
23	
∠4	
25	
26	
20	
27	
28	
20	
29	
30	
00	
31	
32	
20	
33	
34	
2.	
35	
36	
07	
37	
38	
20	
39	
40	
11	
42	
43	
11	
44	
45	
٨Ē	
40	
47	
۸۶	
+0	
49	
50	
50	
51	
52	
50	
53	
54	
6E	
55	
56	
57	
51	
58	
59	
60	
011	

1 2

> The fate of these fish remain unclear. They may have emigrated out of the study site and 445 446 travelled an undetermined distance, were predated, or suffered tag failure. Similarly high levels of site fidelity have been previously described for salmonids in other systems and at 447 448 other times of the year, including for a proportion of those populations in which patch 449 shifting was commonly observed (e.g. Martin-Smith and Armstrong, 2002), possibly 450 reflecting a strategy by some individuals to limit detection by predators. Thus, site fidelity is 451 not necessarily limited to spatially and temporally homogenous habitat templates during the 452 winter.

453 Where food is abundant and lower temperatures favour homeothermic mammalian and avian visual predators in attacks against ectothermic prey during the winter (Fraser et al. 454 455 1993; Orpwood et al. 2006, 2010), maximisation of energy intake during the night is likely to 456 represent an advantageous strategy. In other systems, stream-dwelling salmonids become 457 increasingly nocturnal with reductions in temperature during the winter and generally less active during the day when they commonly seek shelter (Cunjak 1988; Heggenes et al. 1993; 458 459 Heggenes et al. 1999). For juvenile Atlantic salmon the shift to increasingly nocturnal foraging appears to occur at some threshold temperature between 8-10°C (Fraser et al. 1993; 460 461 Rimmer et al. 1983). Although fish in this study tended to exhibit high site fidelity, there was 462 also evidence of high periodicity of movement exhibited by some fish, with changes in 463 position at dawn and dusk, which may suggest nocturnal foraging. This may not reflect a 464 strategy unique to the winter, however, as radio-tagged trout on the River Aisne were most 465 active at dusk in all seasons (Ovidio et al. 2002). Further research is needed to clarify how 466 salmonid foraging behaviour and anti-predator response varies with season and river type.

467 As chalk streams are highly productive aquatic environments it was predicted that 468 trout would exhibit high growth during the winter in this study. Indeed, the River Lambourn 469 supports elevated primary and secondary production, and exceptional invertebrate biomass

Freshwater Biology

3
4
5
6
7
0
ð
9
10
11
12
13
14
15
16
17
10
10
19
20
21
22
23
24
25
26
20
21
28
29
30
31
32
33
34
35
36
27
31
38
39
40
41
42
43
44
45
16
40 47
47
48
49
50
51
52
53
54
55
56
50
5/ 50
58
59
60

470	(Pretty et al. 2006). At Bagnor, approximately 3 km downstream of the study site,
471	invertebrate biomass tends to peak in December with total density found to range from
472	approximately 15,000 to 70,000 macroinvertebrates m ⁻¹ (data collected between 1971-1978,
473	Wright 1992). This high density of invertebrates during the winter reflects the energy
474	provided by abundant decaying macrophytes that act as a nucleus for the deposition of
475	allochthonous tree leaves in the autumn (Westlake et al. 1972). In contrast, boreal and
476	northern temperate surface-flow systems tend to be oligotrophic, and during the winter low
477	temperatures create a stressful period during which growth of salmonids is sufficiently low as
478	to be considered negligible (Cunjak and Power 1987), ceases altogether, or in some cases is
479	negative as indicated by a loss in mass (Egglishaw and Shackley 1977). The River Lambourn
480	trout exhibited positive growth during periods that included winters 1 and 2. It is
481	unsurprising that mean growth rates were below those predicted by the Elliott et al. (1995)
482	growth model that is more applicable to summer growth situations (Jensen et al. 2000) for
483	trout fed on maximum ration. Unfortunately, these results were influenced by growth periods
484	during autumn (year 1) and spring (year 2). Nevertheless, it is likely that the higher mean
485	ambient temperature regimes and high biomass of invertebrates resulted in elevated growth
486	rates when compared with other systems. This suggestion is supported by earlier studies that
487	quantified the annual instantaneous brown trout growth in which the River Lambourn yielded
488	the highest values of nine examples of UK rivers studied (Edwards et al. 1979).

489

490 **Conclusions**

491

492 Using passive integrated transponder and radio telemetry combined with information

493 obtained from data storage tags this study sheds light on the behaviour and performance of

2
2
3
4
5
6
7
1
8
9
10
11
10
12
13
14
15
16
10
17
18
19
20
21
21
22
23
24
25
20
26
27
28
29
20
30
31
32
33
34
34
35
36
37
38
20
39
40
41
42
43
44
44
45
46
47
10
40
49
50
51
52
52
53
54
55
56
57
57
58
59
60

1

494	brown trout inhabiting a comparatively stable, warm and productive habitat template
495	provided by a chalk stream during the winter. In this often ignored system, the
496	hydrodynamic, physical and thermal environment was mapped at a fine-resolution scale
497	appropriate to study the distribution and movement patterns of the target species.
498	Interestingly, several behaviours exhibited by trout were similar to those observed for other
499	systems and seasons. Distribution was strongly correlated with depth, but not velocity or
500	temperature, and the majority of radio-tracked fish exhibited high site fidelity and limited
501	movement. There was evidence for high periodicity of movement with changes in position at
502	dawn and dusk for some fish, potentially suggesting nocturnal foraging. Trout growth was
503	positive during the periods that included winter in both years, indicating higher performance
504	than for populations that suffer an intensely challenging winter bottleneck. Recognition of
505	regional variation in winter ecology of salmonids is important in developing appropriate
506	management strategies to preserve or rehabilitate winter habitat in the face of increasing
507	urbanisation, altered land use, and climate change.

508

509 Acknowledgments

510 We thank G. Old and P. Rameshwaran (CEH) for flow data for the site, K. Alfredsen

511 (Norwegian University of Science and Technology) for temperature data for the

512 Svarttjønnbekken River (Norway), W. Beaumont (GWCT) for advice regarding telemetry,

and local Environment Agency staff and volunteers who provided assistance during surveys.

514 We also thank the associate editor and two anonymous reviewers for comments on an earlier

515 draft of the manuscript. Data published in this paper are available from the University of

516 Southampton repository at DOI:10.5258/SOTON/392646.

517

1		
2		
3	518	
4		
5		
6	519	TABLES AND FIGURES
7	515	
8		
q	E 2 0	
10	520	
11		
12	F 2 4	Table 1. Comparison of concupliced physical and chemical habitat templates for barrel
12	521	radie 1. Comparison of generalised physical and chemical naditat templates for borear
17		
14	522	surface flow rivers on hard rock catchments and temperate ground water dominated
15		
16	523	chalk streams during the winter.
17		
18		
19		
20		
21		
22		
23		
24		
25		
26		
27		
28		
29		
30		
31		
32		
33		
34		
35		
36		
37		
38		
30		
40		
0 ⊿1		
 ⊿?		
42 /2		
43		
44		
40		
40		
4/		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		23

	Boreal surface flow hard rock rivers	Temperate ground water fed chalk streams
hysical		
- Hydrological regime	Moderate – high temporal variability. Dynamic and "flashy" – responsive to precipitation and snow-melt events.	Low temporal variability, relatively stable regime sustained for long periods by high baseflow component.
- Depth	Moderate to high spatial variability. Well defined riffle- pool sequences.	Low spatial variability. Lack of well-defined and low longitudina frequency of riffle-pool sequences. Frequently shallow cross sectional area in unmodified reaches.
 Hydrodynamics (velocity, turbulence etc.) 	Moderate to high spatial and temporal heterogeneity dictated by flow and substrate complexity.	Low meso-scale hydrodynamic heterogeneity. Fine scale complexity associated with aquatic plants.
- Substrate and structure	Potential for high spatial variability ranging from fine sediments to large boulders and large woody debris. Considerable potential for refuge sites including interstitial space.	Generally low variability and predominantly sands and flint gravels with a lack of large clasts. Fine sediment component is often high. The riverbed may be highly compacted and weathered with limited interstitial space. Infrequent gravel shoals and exposed river bed substrate.
 Channel planform and drainage 	High variability, ranging from natural channels with high sinuosity to those influenced by anthropogenic modification. Moderate to high drainage density and dendritic drainage patterns are common.	Often influenced by long history of anthropogenic modification and management. Channel frequently diverted along alternative paths running in parallel (e.g. mill lades). Low drainage density and limited tributary network. Headwaters may be ephemeral.
hemical		
 pH Nutrient concentrations and conductivity 	High acidity. pH ranges from 4.5 – 7.0. Often igneous or metamorphic geology in some instances dominated by Mg, Na, Ca, and K cations. Concentrations of ions and conductivity often considerably lower than in chalk rivers.	High alkalinity. pH typically 7.4 – 8.0. High concentration of dissolved calcium and bicarbonate ions and high conductivity. Expected P under pristine conditions may range from $0.02 - 0.06$ mg l ⁻¹ , but N and P levels are much highe in many chalk rivers due to agricultural practices.
- Temperature	High variability of river temperatures during the year, but low temperatures (0°C) for long periods during the winter	In southern England spring water temperature is typically 11°C. Temperatures remain relatively warm throughout the winter.

reach on the River Lambourn during two surveys (February 2011 and August 2011).

			Survey
Habitat variable	Measure	Values	date
		Gravel	
Substrata	Dominate	(35%)	Feb 2011
Substrate	size	Gravel	
		(57%)	Aug 2011
	Mean	31.9 cm	Feb 2011
Denth	Wear	38.3 cm	Aug 2011
Depth	Range	2-115 cm	Feb 2011
		13-98 cm	Aug 2011
	Mean	31.7 cm/s	Feb 2011
Velocity (cm s ⁻¹)	wicall	21.0 cm/s	Aug 2011
velocity (clifs)	Range	-4-89 cm/s	Feb 2011
	Nunge	1-54 cm/s	Aug 2011

528 Table 3. Telemetry survey data for brown trout, *Salmo trutta*, caught on the River

529 Lambourn by electric fishing.

		Number	Number	Number		
	Number	PIT	radio 🧹	DS	Fork length [mean	Weight [mean ± SD
Survey dates	caught	tagged	tagged	tagged	± SD (range), mm]	(range), g]
					229.2 ± 53.1 (148-	161.3 ± 95.9 (44-
15 to 16 July 2010	126	126	30	10	425)	418)
3 to 4 February					204.5 ± 72.7 (85-	126.5 ± 111.1 (7-
2011	318	203	10	0	387)	575)
					211.1 ± 63.7 (113-	124.9 ± 111.7 (12-
5 May 2011	220	209	19	9	390)	654)
9 September					197.1 ± 65.3 (83-	122.5 ± 112.2 (6-
2011	347	278	24	11	392)	631)
					216.6 ± 66.9 (67-	170.0 ± 165.7 (2-
31 May 2012	361	0	0	0	499)	1128)

531 Table 4. Regression statistics of physical habitat predictors of fish density at a coarse

532 scale in a 500 m reach of the River Lambourn (Berkshire, UK). 95% bias corrected and

533 accelerated confidence intervals are reported in parentheses. CIs, SEs and significance

values for all predictors are based on 1000 bootstrap samples.

Electric-fishing					
survey	Predictor	b	SE b	ß	р
	Depth	0.004 (0.002 - 0.005)	0.001	0.918	= 0.001
		0.002 (-0.001 -			
Feburary 2011	Velocity	0.004)	0.001	0.207	= 0.141
(Model: R ² = .71, F _{3,25}		-0.158 (-0.475 -			
= 20.10, <i>p</i> < 0.001)	Temperature	0.069)	0.136	-0.197	= 0.193

Figure 1. Site of a study conducted on the River Lambourn, a chalk stream in Southern
England (United Kingdom), to quantify habitat use by brown trout, *Salmo trutta*, during
the winters of 2010/11 and 2011/12. The 500 m reach of river is owned by the Centre
for Ecology and Hydrology.

Figure 2. Physical characteristics of a 500 m reach of the River Lambourn (Berkshire, UK) based on a topographic survey conducted in May 2012: a. dominant substrate, b. depth, c. velocity, and d. temperature (data for 11 February 2012, the coldest day during Winter 2 based on air temperature records [mean = -6.6° C]). Plots a – c represent data collected during a habitat survey conducted in August 2011. Stars represent the locations of the fixed radio receivers. Physical habitat and temperature contour plots were constructed using a spline interpolation within the boundaries of an ArcGIS base map.

Figure 3. River Lambourn temperatures recorded at Boxford during years 1 (a) and 2
(b). Solid green, red and blue lines represent mean, maximum and minimum
temperatures measured at the river bed, while the dashed line denotes the ambient air
temperature. For comparison, the dotted line represents mean instream temperatures
recorded at the Svarttjønnbekken River, a northern boreal system in Norway.

Freshwater Biology

2 3 4	555	
5 6 7	556	Figure 4. Relationship between density of fish (per m ²) and mean water depth for the
8 9	557	electric fishing reaches surveyed.
10 11 12 13	558	
14 15	559	Figure 5. Relationship between ΔT (the difference between mean daily fish and
16 17	560	maximum daily river temperatures) and minimum daily river temperature for brown
19 20	561	trout occupying a reach of the River Lambourn during winter.
21 22 23	562	
24 25 26	563	Figure 6. The majority of radio tagged brown trout were observed clustered around a
27 28	564	single position within the study site on the River Lambourn, UK (a). Occupancy of 2
29 30	565	distinct locations (b) or observation at multiple sites to which fidelity was low (c) were
31 32 33	566	less common.
34 35 36	567	
37 38 39	568	Figure 7. (a) Autocorrelation function (ACF) of mean signal strength with a one hour
40 41	569	time lag; the plots represent 2 fish exhibiting a daily activity pattern. (b) two different
42 43	570	fish exhibited extremely high levels of periodicity in activity as indicated by peaks in
44 45 46	571	correlation every 24 hours with no decay. (c) the mean signal strength per hour (date
40 47 48	572	plotted is 00:00 on 1st Nov 2011 – 23:00 on 7th Nov 2011, shaded area = night) for the
49 50 51	573	two fish represented in (b).
52 53 54	574	
55 56	575	Figure 8. The relationship between observed winter growth of brown trout and that
57 58	576	predicted by an optimal growth model that assumes fish are fed to satiation under

3
4
5
6
7
1
8
9
10
11
12
13
14
15
16
10
17
18
19
20
21
22
23
24
25
20
20
21
28
29
30
31
32
33
34
25
20
30
37
38
39
40
41
42
43
44
45
40 46
40
41
48
49
50
51
52
53
54
55
55
20
5/
58
59
60

577 laboratory conditions (Elliott et al. 1995). Solid and clear circles denote data for Winter

578 1 and 2, respectively. Points below the dashed line indicate suboptimal growth.

579

1 2

580 **References**

- Armstrong, J.D., Kemp, P.S., Kennedy, G.J.A, Ladle, M. and Milner, N.J. 2003. Habitat
 requirements of Atlantic salmon and brown trout in rivers and streams. *Fisheries Research*62, 143-170.
- 584 Arthur, B., Hindell, M., Bester, M., Trathan, P., Jonsen, I., Staniland, I., Oosthuizen, W.C.,

585 Wege, M. and Lea, M-A. 2015. Return customers: foraging site fidelity and the effect of

586 environmental variability in wide-ranging Antarctic fur seals. *PLoS ONE* **10**: e0120888.

- 587 doi:10.1371/journal.pone.0120888
- Brown, R.S., Power, G. and Beltaoa, S. 2001. Winter movements and habitat use of riverine
 brown trout, white sucker and common carp in relation to flooding and ice break-up. *Journal of Fish Biology* 59, 1126-1141.

591 Brown, R.S., Hubert, W.A., and Daly, S.F. 2011. A primer on winter, ice, and fish: What

592 fisheries biologists should know about winter ice processes and stream-dwelling fish.

593 *Fisheries* **36**, 8-36.

594 Charnov, E.L. 1976. Optimal foraging, the marginal value theorem. *Theoretical Population*595 *Biology* 9, 129–136.

596 Cunjak, R.A. 1996. Winter habitat of selected stream fishes and potential impacts from land-597 use activity. *Canadian Journal of Fisheries and Aquatic Sciences* **53** (S1), 267-282.

598	Cunjak, R.A. 1988. Behaviour and microhabitat of young Atlantic salmon (Salmo salar)
599	during winter. Canadian Journal of Fisheries and Aquatic Sciences 45, 2156-2160.
600	Cunjak, R.A. and Power, G. 1987. The feeding and energetics of stream-resident trout in
601	winter. Journal of Fish Biology 31, 493-511.
602	Cunjak, R.A. and Power, G. 1986. Winter habitat utilisation by stream resident brook trout
603	(Salvelinus fontinalis) and brown trout (Salmo trutta). Canadian Journal of Fisheries and
604	Aquatic Sciences 43 , 1970-1981.
605	Cunjak, R.A., Prowse, T.D. and Parrish, D.L. 1998. Atlantic salmon (Salmo salar) in winter:
606	"the season of parr discontent"? Canadian Journal of Fisheries and Aquatic Sciences 55 (S1),
607	161-180.
608	Dill, L.M., Ydenberg, R.C. and Fraser, A.H.G. 1981. Food abundance and territory size in
609	juvenile coho salmon (Oncorhynchus kisutch). Canadian Journal of Zoology 59, 1801-1809.
610	Edwards, R.W., Densem, J.W. and Russell, P.A. 1979. An assessment of the importance of
611	temperature as a factor controlling the growth rate of brown trout in streams. Journal of
612	Animal Ecology 48, 501–507.
613	Egglishaw, H.J. and Shackley, P.E. 1977. Growth, survival and production of juvenile salmon
614	and trout in a Scottish stream, 1966–75. Journal of Fish Biology 11, 647-672.
615	Elliott, J.M., Hurley, M.A. and Fryer, R.J. 1995. A new, improved growth model for brown
616	trout, Salmo trutta. Functional Ecology 9, 290-298.
617	Evans, D.J., Johnes, P.J. and Lawrence, D.S. 2003. Suspended and bed load sediment
618	transport dynamics in two lowland UK stream-storm integrated monitoring. Erosion and

2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
10	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
ר ב בי	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
22	
3Z 22	
33	
34	
35	
36	
37	
38	
39	
40	
41	
40 1	
+∠ 10	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
52	
55	
54 55	
55	
56	
57	
58	
59	
60	

1 2

619	Sediment Transport Measurements in Rivers: Technological and Methodological Advances
620	(Proceedings of the Oslo Workshop, June 2002) IAHS Publ. 283, 2003.

621 Fraser, N.H.C., Metcalfe, N.B. and Thorpe, J.E. 1993. Temperature-dependent switch

622 between diurnal and nocturnal foraging in salmon. *Proceedings of the Royal Society of*

623 London **B 252**, 135-139.

Hedger, R.D., Naesje, T.F., Fiske, P., Ugedal, O., Finstad, A.G. and Thorstad, E.B. 2013.

625 Ice-dependent winter survival of juvenile Atlantic salmon. *Ecol. Evol.* **3**, 523-535.

626 Heggenes, J., Baglinière, J.L. and Cunjak, R.A. 1999. Spatial niche variability for young

627 Atlantic salmon (*Salmo salar*) and brown trout (*S. trutta*) in heterogeneous streams. *Ecology*

628 *of Freshwater Fish* **8**, 1-21.

Heggenes, J., Krog, O.M.W., Lindås, O.R. and Dokk, J.G. 1993. Homeostatic behavioural
responses in a changing environment: brown trout (*Salmo trutta*) become nocturnal during

631 winter. Journal of Animal Ecology 62, 295-308.

632 Huusko, A., Greenberg, L., Stickler, M., Linnansaari, T., Nykanen, M., Vehanen, T.,

633 Koljonen, S., Louhi, P. and Alfredsen, K. 2007. Life in the ice lane: the winter ecology of

634 stream salmonids. *River Research and Applications* **23**, 469-491.

635 Jensen, A.J, Forseth, T. and Johnsen, B.O. 2000. Latitudinal variation in growth of young

636 brown trout Salmo trutta. *Journal of Animal Ecology* **69**, 1010–1020.

637 Khan, M.T., Khan, T.A. and Wilson, M.E. 2004. Habitat use and movement of river blackfish

638 (Gadopsis marmoratus R.) in a highly modified Victorian stream, Australia. Ecology of

639 *Freshwater Fish* **13**, 285-293.

1	
2	
3	
4	
5	
6	
0	
1	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
21 22	
22 22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
27	
20	
30	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
52	
53	
54	
22	
56	
57	
58	
59	
60	

640	Linnansaari, T. and Cunjak, R.A. 2010. Patterns in apparent survival of Atlantic salmon
641	(Salmo salar) parr in relation to variable ice conditions throughout winter. Canadian Journal
642	of Fisheries and Aquatic Sciences 67, 1744-1754.
643	Linnansaari, T., Alfredsen, K., Stickler, M., Arnekleiv, J.V., Harby, A. and Cunjak, R.A.
644	2009. Does ice matter? Site fidelity and movements by Atlantic salmon (Salmo salar L.) parr
645	during winter in a substrate enhanced river reach. River Research and Applications 25, 773-
646	787.
647	Mäki-Petäys, A., Muotka, T., Huusko, A., Tikkanen, P. and Kreivi, P. 1997. Seasonal
648	changes in habitat use and preference by juvenile brown trout, Salmo trutta, in a northern
649	boreal river. Canadian Journal of Fisheries and Aquatic Sciences 54, 520-530.
650	Martin-Smith, K.M. and Armstrong, J.D. 2002. Growth rates of wild stream-dwelling
651	Atlantic salmon correlate with activity and sex but not dominance. Journal of Animal
652	<i>Ecology</i> 71 , 413-423.
653	Orpwood, J.E., Armstrong, J.D. and Griffiths, S.W. 2010. Interactions between riparian
654	shading and food supply: a seasonal comparison of effects on time budgets, space use and
655	growth in Atlantic salmon Salmo salar. Journal of Fish Biology 77, 1835-1849.
656	Orpwood, J.E., Griffiths, S.W. and Armstrong, J.D. 2006. Effects of food availability on
657	temporal activity patterns and growth of Atlantic salmon. Journal of Animal Ecology 75, 677-
658	685.
659	Ovidio, M., Baras, E., Goffaux, D., Giroux, F. and Philippart, J.C. 2002. Seasonal variations

- of activity pattern of brown trout (*Salmo trutta*) in a small stream, as determined by radio-
- 661 telemetry. *Hydrobiologia* **470**, 195-202.

2	
3	
4	
5	
6	
7	
1	
8	
9	
10	
11	
12	
12	
13	
14	
15	
16	
17	
18	
19	
20	
20	
21	
22	
23	
24	
25	
26	
20	
21	
28	
29	
30	
31	
32	
22	
33	
34	
35	
36	
37	
38	
20	
39	
40	
41	
42	
43	
44	
45	
40	
40	
47	
48	
49	
50	
51	
51	
ວ <u>∠</u>	
53	
54	
55	
56	
57	
50	
00	
59	
60	

662 Poff, N.L. and Allan, J.D. 1995. Functional organisation of stream fish assemblages in

relation to hydrological variability. *Ecology* **76**, 606-627.

Poff, N.L and Ward, J.V. 1990. Physical habitat template of lotic systems: recovery in the
context of historical patterns of spatiotemporal heterogeneity. *Environmental Management*14, 629-645.

667 Pretty J., Hildrew A.G. and Trimmer M. 2006. Nutrient dynamics in relation to surface-

668 subsurface hydrological exchange in a groundwater fed chalk stream. *Journal of Hydrology*

669 330, 84-100.

670 Rameshwaran, P., Roberts, C., Old, G.H. and Naden, P. (2015). Hydraulics data for the River

671 Lambourn at Boxford, 2008 to 2014. NERC Environmental Information Data Centre. DOI:

672 10.5285/4ed6f721-b23b-454e-b185-02ba54d551f0

673 Simpkins, D.G. and Hubert, 2000. Drifting invertebrates, stomach contents, and body

674 conditions of juvenile rainbow trout from fall through winter in a Wyoming tailwater.

675 *Transactions of the American Fisheries Society* **129**, 1187-1195.

676 Solazzi, M.F., Nickelson, T.E., Johnson, S.L. and Rodgers, J.D. 2000. Effects of increasing

677 winter rearing habitat on abundance of salmonids in two coastal Oregon streams. *Canadian*

678 *Journal of Fisheries and Aquatic Sciences* 57, 906-914.

679 Southwood, T.R.E. 1977. Habitat, the templet for ecological strategies. *Journal of Animal*680 *Ecology* 46, 337-365.

581 Stickler, M., Enders, E.C., Pennell, C.J., Cote, D., Alfredsen, K.T. and Scruton, D.A. 2008.

682 Habitat use of Atlantic salmon *Salmo salar* part in a dynamic winter environment: the

683 influence of anchor-ice dams. *Journal of Fish Biology* **73**, 926-944.

Freshwater Biology

2	
3	
4	
5	
6	
0	
7	
8	
9	
10	
11	
11	
12	
13	
14	
15	
16	
17	
17	
18	
19	
20	
21	
22	
22	
23	
24	
25	
26	
27	
20	
20	
29	
30	
31	
32	
33	
24	
34	
35	
36	
37	
38	
20	
39	
40	
41	
42	
43	
11	
44	
45	
46	
47	
48	
40	
50	
50	
51	
52	
53	
54	
55	
55	
56	
57	
58	
59	

60

684 Stickler, M., Afredsen, K., Scruton, D.A., Pennell, C., Harby, A. and Okland, F. 2007. Mid-685 winter activity and movement of Atlantic salmon parr during ice formation events in a 686 Norwegian regulated river. Hydrobiologia 582, 81-89. 687 Swales, S., Lauzier, R.B. and Levings, C.D. 1986. Winter habitat preferences of juvenile 688 salmonids in two interior rivers in British Columbia. Canadian Journal of Zoology 64, 1506-689 1514. 690 Tews, J., Brose, U., Grimm, V., Tielborger, K., Wichmann, M.C., Schwager, M. and Jeltsch, 691 F. 2004. Animal species diversity driven by habitat heterogeneity/diversity: the importance 692 of keystone structures. Journal of Biogeography 31, 79-92. 693 Webb, P.W. 1978. Temperature effects on acceleration of rainbow trout, Salmo gairdneri. 694 Journal of the Fisheries Research Board of Canada 35, 1417-1422. 695 Westlake, D.F., Casey, H., Dawson, F.H., Ladle, M., Mann, R.H.K. and Marker, A.F.H. 1972. 696 The chalk stream ecosystem. In Z. Kajak and A. Hillbricht-Ilkowska (eds), Proceedings 697 I.B.P UNESCO Symposium on Productivity Problems of Freshwater, Kazimierz Dolnny. 698 Warszawa-Krakow: 615-635. 699 Wootton, R.J. 1998. Ecology of teleost fishes. Chapman and Hall, London, UK. 700 Wright, J.F. 1992. Spatial and temporal occurrence of invertebrates in a chalk stream,

701 Berkshire, England. *Hydrobiologia* **248**, 11-30.



Page 35 of 45

Freshwater Biology





Page 37 of 45





7 8







Charac	teristics	Boreal surface flow hard rock rivers	Temperate ground water fed chalk streams
Physica	al		
-	Hydrological regime	Moderate – high temporal variability. Dynamic and "flashy" – responsive to precipitation and snow-melt events.	Low temporal variability, relatively stable regime sustained for long periods by high baseflow component.
-	Depth	Moderate to high spatial variability. Well defined riffle- pool sequences.	Low spatial variability. Lack of well-defined and low longitudinal frequency of riffle-pool sequences. Frequently shallow cross sectional area in unmodified reaches.
-	Hydrodynamics (velocity, turbulence etc.)	Moderate to high spatial and temporal heterogeneity dictated by flow and substrate complexity.	Low meso-scale hydrodynamic heterogeneity. Fine scale complexity associated with aquatic plants.
-	Substrate and structure	Potential for high spatial variability ranging from fine sediments to large boulders and large woody debris. Considerable potential for refuge sites including interstitial space.	Generally low variability and predominantly sands and flint gravels with a lack of large clasts. Fine sediment component is often high. The riverbed may be highly compacted and weathered with limited interstitial space. Infrequent gravel shoals and exposed river bed substrate.
-	Channel planform and drainage	High variability, ranging from natural channels with high sinuosity to those influenced by anthropogenic modification. Moderate to high drainage density and dendritic drainage patterns are common.	Often influenced by long history of anthropogenic modification and management. Channel frequently diverted along alternative paths running in parallel (e.g. mill lades). Low drainage density and limited tributary network. Headwaters may be ephemeral.
Chemi	cal		
-	pH Nutrient concentrations and conductivity Temperature	 High acidity. pH ranges from 4.5 – 7.0. Often igneous or metamorphic geology in some instances dominated by Mg, Na, Ca, and K cations. Concentrations of ions and conductivity often considerably lower than in chalk rivers. High variability of river temperatures during the year, but low temperatures (0°C) for long periods during the winter months, especially in systems that experience freeze-up. 	High alkalinity. pH typically 7.4 – 8.0. High concentration of dissolved calcium and bicarbonate ions and high conductivity. Expected P under pristine conditions may range from $0.02 - 0.06$ mg l ⁻¹ , but N and P levels are much higher in many chalk rivers due to agricultural practices. In southern England spring water temperature is typically 11°C. Temperatures remain relatively warm throughout the winter.

Freshwater Biology

			Survey
labitat variable	Measure	Values	date
	Deminate	Gravel	5-k 2011
Substrate	Dominate	(35%) Craval	Feb 2011
	size	(57%)	Δυσ 2011
		(3776)	Aug 2011
	Mean	31.9 cm	Feb 2011
Depth		38.3 cm	Aug 2011
	Range	2-115 cm	Feb 2011
	-	13-98 cm	Aug 2011
	Mean	31.7 cm/s	Feb 2011
Velocity (cm s ⁻¹)	mean	21.0 cm/s	Aug 2011
velocity (ciri s)	Danga	-4-89 cm/s	Feb 2011
	Range	1-54 cm/s	Aug 2011

Survey dates	Number caught	Number PIT tagged	Number radio tagged	Number DS tagged	Fork length [mean ± SD (range), mm]	Weight [mean ± SD (range), g]
15 to 16 July 2010 3 to 4 February	126	126	30	10	425) 204.5 ± 72.7 (85-	418) 126.5 ± 111.1 (7-
2011	318	203	10	0	387) 211.1 ± 63.7 (113-	575) 124.9 ± 111.7 (12-
5 May 2011 9 September	220	209	19	9	390) 197.1 ± 65.3 (83-	654) 122.5 ± 112.2 (6-
2011	347	278	24	11	392) 216.6 ± 66.9 (67-	631) 170.0 ± 165.7 (2-
31 May 2012	361	0	0	0	499)	1128)

Electric-fishing						
survey	Predictor b		SE b	ß	р	
	Depth	0.004 (0.002 - 0.005)	0.001	0.918	= 0.001	
		0.002 (-0.001 -				
Feburary 2011	Velocity	0.004)	0.001	0.207	= 0.141	
(Model: $R^2 = .71$, $F_{3,25}$		-0.158 (-0.475 -				
= 20.10, <i>p</i> < 0.001)	Temperature	0.069)	0.136	-0.197	= 0.193	