


Accepted Manuscript

Postflight reconditioning for European astronauts – A case report of recovery after six months in space

Nora Petersen, Gunda Lambrecht, Jonathan Scott, Natalie Hirsch, Maria Stokes, Joachim Mester



PII: S2468-7812(16)30008-X

DOI: [10.1016/j.msksp.2016.12.010](https://doi.org/10.1016/j.msksp.2016.12.010)

Reference: MSKSP 24

To appear in: *Musculoskeletal Science and Practice*

Received Date: 15 August 2016

Revised Date: 17 October 2016

Accepted Date: 1 November 2016

Please cite this article as: Petersen, N., Lambrecht, G., Scott, J., Hirsch, N., Stokes, M., Mester, J., Postflight reconditioning for European astronauts – A case report of recovery after six months in space, *Musculoskeletal Science and Practice* (2017), doi: 10.1016/j.msksp.2016.12.010.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

For *Musculoskeletal Science and Practice* Supplement: “**Terrestrial neuro-musculoskeletal rehabilitation and astronaut reconditioning: reciprocal knowledge transfer**”

Guest editors: Maria Stokes, Simon Evetts, Julie Hides ”

Postflight reconditioning for European Astronauts – a case report of recovery after six months in space

Authors: Nora Petersen^{1,2,6}, Gunda Lambrecht¹, Jonathan Scott^{1,2}, Natalie Hirsch³, Maria Stokes^{4,5}; Joachim Mester⁶

Affiliations:

¹Wyle GmbH, Cologne, Germany;

²Space Medicine Office (HSO-AM), European Astronaut Centre Department, Directorate of Human Spaceflight and Operations (D/HSO), European Space Agency, Cologne, Germany;

³Canadian Space Agency, Canada;

⁴Faculty of Health Sciences, University of Southampton, UK;

⁵Arthritis Research UK Centre for Sport, Exercise and Osteoarthritis, UK;

⁶German Sport University, Cologne, Germany;

Corresponding Author:

Nora Petersen

Wyle GmbH, Cologne, Germany; Space Medicine Office, Department, European Space Agency, Cologne Germany.

Email: Nora.Petersen@esa.int

Tel. +49 2203 6001 406.

For *Manual Therapy Supplement*: “**Terrestrial neuro-musculoskeletal rehabilitation and astronaut reconditioning: reciprocal knowledge transfer**”

Guest editors: Maria Stokes, Simon Evetts, Julie Hides ”

Postflight reconditioning for European Astronauts – a case report of recovery after six months in space

Abstract

Background: Postflight reconditioning of astronauts is understudied. Despite a rigorous, daily inflight exercise countermeasures programme during six months in microgravity (μG) on-board the International Space Station (ISS), physiological impairments occur and postflight reconditioning is still required on return to Earth. Such postflight programmes are implemented by space agency reconditioning specialists.

Case Description and Assessments: A 38 year old male European Space Agency (ESA) crewmember’s pre- and postflight (at six and 21 days after landing) physical performance from a six-month mission to ISS are described. *Assessments:* muscle strength (squat and bench press 1 Repetition Maximum) and power (vertical jump), core muscle endurance and hip flexibility (Sit and Reach, Thomas Test).

Interventions: In-flight, the astronaut undertook a rigorous daily (2-hour) exercise programme. The 21 day postflight reconditioning exercise concept focused on motor control and functional training, and was delivered in close co-ordination by the ESA physiotherapist and exercise specialist to provide the crewmember with comprehensive reconditioning support.

Outcomes: Despite an intensive inflight exercise programme in this highly motivated crewmember, postflight performance showed impairments at R+6 for most parameters, all of which recovered by R+21 except muscular power (jump tests).

Conclusions: Regardless of intense inflight exercise countermeasures and excellent compliance to postflight reconditioning, postflight performance showed impairments at R+6 for most parameters. Complex powerful performance tasks took longer to return to preflight values. Research is needed to develop optimal inflight and postflight exercise programmes to overcome the negative effects of microgravity and return the astronaut to preflight status as rapidly as possible.

Keywords: spaceflight, astronaut reconditioning, rehabilitation, functional fitness assessment

ACCEPTED MANUSCRIPT

1. Introduction

The effects of prolonged exposure to microgravity (μG) on the human body have been well studied [1-4] and include reductions of muscle volume and strength, bone mass and aerobic capacity [1] as the human body adapts to its new environment. Reductions of muscle strength and stability are associated with lower physical performance capacity and have a number of health implications [1, 5]. There are alterations in muscle activity, proprioception and posture [5], disc hyperhydration associated with fluid shifts [6] and cardiovascular changes [7]. This is not a “sick” but an “adapted” status, which may negatively affect physical performance and capacity, especially on return into Earth's gravity.

To counteract the effects of μG exposure [1, 3, 4] and to adequately prepare for return to Earth during Long (approximately six months) Duration Missions to the International Space Station (ISS), a rigorous inflight exercise countermeasure programme is implemented [8, 9]. Although the exercise prescription is individualised, the on-board constraints of ISS limit exercise to resistance training, treadmill and cycle ergometer training [9-11]. Despite the inflight programme, which is performed on a daily basis, deconditioning cannot be completely prevented. Thus, astronauts returning from LDMs require a postflight reconditioning programme, which begins one day after return (R+1), to restore their physical condition to preflight levels.

The astronaut population is a heterogeneous, but healthy and active population, displaying both traits of patients and of professional athletes in their postflight physical condition.

The aims of this paper are:

1. To briefly describe the physical exercise component of the European Space Agency's (ESA) postflight reconditioning programme;
2. To present a case report describing the postflight performance of one ESA crewmember following a LDM to ISS.

2. Case Information

The case of an active 38 year old (stature: 1.85 m; body mass: 83 kg) ESA astronaut who spent six months on ISS is described. He provided informed consent regarding the publication of his data for this case study. Ethics approval was obtained from the North Rhine (Germany) Medical Association. Findings from assessments pre- and post-intervention are presented in the outcomes section below.

3. Medical and Astronaut Fitness Assessment (AFA)

All ESA crewmembers undergo standardised preflight and postflight medical evaluations for long duration ISS missions (Medical Evaluation Documents, [MED], Volume B; document not publically available). During the 21-day postflight reconditioning period, these assessments usually occur twice on fixed dates and include the ESA Astronaut Fitness Assessment (AFA), and evaluation of aerobic capacity (Periodic Fitness Evaluation – PFE) and isokinetic muscle strength.

The aim of the fitness assessments is twofold: they provide direct performance feedback to the crewmember and exercise specialist for reconditioning and training plan design, and contribute to the overall long-term documentation of the countermeasures programme and astronaut health after space missions [12, 13].

The ESA AFA is composed of 10 tests targeting muscle strength and power, hip flexibility, balance and cardiovascular performance. The AFA is conducted at launch minus 52 days (L-52 days), and postflight at R+6 days and R+21 days to capture changes in functional fitness capacities related to space flight. For this case report we present a selection of results including maximal (1 RM) strength assessment on squat and bench press [14], muscular power assessed by vertical jump, core endurance and hip flexibility (Appendix 1)[15] [13]. Although additional measurements were taken, for the scope of this article the focus is on the results most relevant to the neuro-musculoskeletal system.

4. Interventions

4.1 European Space Agency Reconditioning Programme - Exercise Component

The ESA Space Medicine Office (SMO) reconditioning team for each crewmember is composed of an experienced exercise specialist/sport scientist and a physiotherapist. This team supports crewmember health prior to, during and following the mission. The reconditioning programme integrates methods originating from physiotherapy and sports and exercise science, to provide a comprehensive, individualised 21-day programme. It includes a large spectrum of expertise, tools and methods to optimise the relatively short reconditioning time after flight. Exercise sessions focus on functionality, efficacy, safety and adequate intensity to promote a comprehensive neuro-musculoskeletal and cardiovascular response. Training locations are either at NASA's Johnson Space Center (JSC) Houston, USA, the Gagarin Cosmonaut Training Centre (GCTC) near Moscow, Russia, or at the European Astronaut Centre (EAC) in Cologne, Germany.

Exercise countermeasures to support the health and physical performance of crewmembers are composed of preflight preparation, inflight countermeasures and postflight reconditioning. The preflight phase is initiated on mission assignment, usually one to two years prior to launch, when individual-specific inflight protocols are developed. Preflight medical and fitness assessments are also implemented, one typically between one year to six months prior to launch, and another at L-30. Results are used as a baseline for comparison with inflight and postflight assessments to monitor the crewmember's condition and progress.

On-board ISS, daily 2-hour (2.5 hrs, including set-up time) intense countermeasure exercise is performed, guided by the exercise specialists ([8, 13, 16]). Since the inflight exercise countermeasures programme is not yet fully able to prevent deconditioning, a 21-day postflight reconditioning programme is undertaken. The objectives of the daily postflight intervention are to:

- 1) Prevent long-term health problems or injuries;
- 2) Return the crewmember to preflight physical condition, as assessed by postflight medical and fitness assessments.

4.2 Postflight physical exercise reconditioning concept

The present case report focuses on the physical activity and sport component of the ESA reconditioning programme, which is complementary to the physiotherapy concept reported by Lambrecht et al. in this special issue of *Manual Therapy* [17].

The reconditioning sessions are closely coordinated between the exercise specialist and the physiotherapist. Large differences occur in postflight condition between astronauts, so sessions are adapted for the individual in terms of their complexity and intensity, and include additional training techniques. During the daily 2-hour sessions, the first hour is dedicated to physiotherapy in a classical treatment room, and the second hour used for physical training with the sport scientist. During the first week, exercises are performed in a swimming pool in coordination with gym sessions. The feeling of neutral buoyancy in the water is perceived as comfortable by astronauts, reminding them of the absence of body weight in microgravity. As the crewmember progresses, gym time vs. pool time increases, and physiotherapy exercises are also performed in the gym. In the main phase of the reconditioning, more time is spent in the gym performing resistance exercises than in the pool or in the physiotherapy room.

The ultimate goal is for the astronaut to perform safe and effective weight lifting training, exercises and sports activities that he/she performed before the mission. At the end of the reconditioning period, sessions should be at or near preflight intensity. The concept then aims to support a continued neuro-musculoskeletal regeneration process over the following months. Since this process cannot be monitored as closely by the ESA specialists, an individual exercise programme for unsupervised training is provided to each crewmember to continue training safely, with the goal of maintaining and/or improve health and fitness after the supervised reconditioning. The training incorporates exercises (Appendix 2,4) that are individualised to each crewmember and therefore vary in timing and intensity. The order as listed is applicable for most crewmembers.

The scheduled interventions and physical fitness assessments in the postflight period are shown in Table 1.

Table 1. Sequence of interventions and assessments until R+21 for all ESA crewmembers

<i>Timeline</i>	R+0	R+1	R+2	R+3	R+4	R+5	R+6	R+7	R+8	R+9	R+10
<i>Intervention</i>	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab
<i>Fit test</i>					PFE	Isokin	AFA				

<i>Timeline</i>	R+11	R+12	R+13	R+14	R+15	R+16	R+17	R+18	R+19	R+20	R+21
<i>Intervention</i>	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab	Rehab
<i>Fit test</i>				PFE	Isokin						AFA

R+, Return date +number of days; *AFA*, Astronaut fitness assessment; *PFE*, Periodic Fitness Evaluation (Aerobic Capacity Evaluation), *Isokin*, Isokinetic measurement

4.3 Exercise Programmes Undertaken

4.3.1 Preflight training

During the preflight phase, ISS specific inflight training protocols (resistance training, cycle ergometer and treadmill) were prepared by the exercise specialist and in parallel, the ESA crewmember performed daily physical exercise for general fitness and Triathlon training. The preflight AFA was conducted 52 days prior to Launch (L-52 days). The crewmember was trained on flight-like ISS exercise hardware, with a focus on motor control, to be prepared for reduced proprioceptive feedback and altered muscle activation when exercising in μG [18] [19].

4.3.2 Inflight training

The astronaut performed daily intense and comprehensive training throughout the mission. He completed a total of 341 exercise sessions during his 6 months stay on ISS, of which 44% were dedicated to resistance training. Inflight physical condition, with the exception of cardiovascular performance on the cycle ergometer and training prescriptions, was not measured, since suitable methods are not yet available.

4.3.3 Postflight training

Reconditioning sessions were conducted along with medical and physical fitness assessments. Initially, ESA specialists implemented the reconditioning at NASA's

Johnson Space Center (JSC) astronaut gym. Training followed the ESA programme outlined above (section 4.2) and the crewmember's specific postflight reconditioning is shown in Appendix 3.

5. Outcomes

The astronaut's compliance with inflight countermeasure exercise was high. His at L-60 days preflight peak VO_2 of $45.2 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ ('excellent' compared to his age group according to ACSM guidelines [20]; internal confidential medical report) indicated that he was an active individual. Furthermore, during the postflight phase, he performed additional exercise to the supervised reconditioning programme and completed an 'Iron Man' event eleven months after return from space.

5.1 Astronaut Fitness Assessment results

In general, the crewmember's postflight physical performance measurements were reduced at R+6 and had partially or fully recovered at R+21 (Table 2). The most notable deficits were in jump performance and flexibility.

The height (cm) and force (N) of all three types of jump, countermovement (CMJ), squat (SJ) and drop jump (DJ), were reduced at R+6, with greatest deficits for both parameters seen in the DJ (height -31%; force -34%). None of the parameter for any jump had recovered by R+2, except CMJ force. The sit and reach flexibility results showed a 16% decrease from preflight distance at R+6, which had recovered to -3% a decrease by R+21.

Muscle strength (1RM and bench press) was less affected, reducing by approximately 8-9% at R+6 and recovering to a 1-3% deficit by R+21 compared with preflight measures. Core muscle endurance was preserved from preflight (260 sec) to R+6 (262 sec, +1%) and R+21 (251 sec, -3%). Balance test results for one leg standing with eyes closed for 15 seconds were reduced at R+6 and R+21 (9 secs and 8 secs respectively) and eyes closed on toes was 10 secs at R+6 and had recovered to 15 secs at R+21.

Table 2. Astronaut Fitness Assessment results comparing preflight (L-52) against postflight (R+6, R+21)

AFA	Parameter	L-52	R+6	R+21	% P#1	%P#2
Body	Body Mass [kg]	81	83	83	2%	2%
	BMI [bm/height]	24	24	24	0%	0%
1 RM	Squat [kg]	148	134	147	-9%	-1%
	Bench [kg]	108	99	105	-8%	-3%
Jumps height	CMJ [cm]	41	33	35	-20%	-15%
	SJ [cm]	35	32	32	-9%	-9%
	DJ [cm]	35	24	30	-31%	-14%
Jumps PF	CMJ [N]	2070	2141	1827	3%	-12%
	SJ [N]	2143	1782	1689	-17%	-21%
	DJ [N]	4727	3120	4381	-34%	-7%
Core endurance	Ventral [sec]	260	262	251	1%	-3%
Hip flexibility	Sit'n'Reach [cm]	37	31	36	-16%	-3%
	TT right [deg]	32	33	27	3%	-16%
	TT left [deg]	30	30	23	0%	-23%

AFA, Astronaut Fitness Assessment; L-52, 52 days prior launch; R+6/21; 6/21 days after return; 1 RM, repetition maximum; bm, body mass; BMI, body mass index; CMJ, countermovement jump; SJ, squat jump; DJ, drop jump; TT, Thomas Test; PF, peak force

6. Discussion

The pre- to postflight comparisons in the present data show the astronaut had neuromusculoskeletal impairments, particularly in the first week after return to Earth. Most, but not all aspects of performance recovered to preflight values by the end of the 21 day reconditioning phase, despite a high level of crew compliance to physical exercise before and during the mission, and despite good core endurance and peripheral strength. These observations support the reported findings in astronauts that, even in high performing and well-trained crewmembers who adhere to rigorous inflight countermeasures, it is not possible to preserve physical performance completely to prepare for everyday activities on return to Earth [21]. Intensive postflight reconditioning for 21 days was sufficient for most but not all aspects of function to recover fully.

6.1 Squat and Bench strength (1RM strength tests)

Loaded squatting is a complex movement requiring motor control and strength when lifting load against gravity. Squat 1RM performance was reduced by 9% at R+6, but recovered by the end of the reconditioning period. This deficit is larger to average values reported by NASA on similar, although not identical (leg press) postflight assessments (-2.8%), with day-to-day changes of 1 RM scores between 5-10% but they are not considered as physiologically relevant [12]. Squats showed slightly larger decreases than bench press for this crewmember, but also a very rapid and complete recovery. The crewmember's absolute pre-flight performance for bench press and sit and reach at R+6 were greater than average values reported by NASA for ISS crewmembers [12] but testing protocols between NASA and ESA vary slightly, so the results should be interpreted with caution.

In the postflight phase, while the body is re-adapting to gravity, crewmembers and exercise specialists are conservative when applying external loads to the body while postural stability is not yet fully restored [5]. Weight lifting training is usually inserted into the programme after R+5 or even later for most crewmembers. In the present case, controlled light load resistance exercise (also in preparation of the fitness assessment) was implemented earlier than usual at R+4, which is exceptional. The second postflight squat measurement (R+21 in Table 2) showed that the crewmember was able to lift similar weights as preflight, indicating successful reconditioning for that task.

Isokinetic testing on NASA astronauts has shown that reduction in knee extensors is greater than that of the flexors [21], which is consistent with extensors undergoing greater atrophy than flexors in MRI studies [22].

In addition to muscle strength, motor control may be an important factor for crew performance. Astronauts usually feel uncomfortable performing loaded squats shortly (R+6) after their mission, and therefore results may also be associated with a more careful and conservative approach to these exercises. As confidence, and postural stability and movement improve during reconditioning, this situation improves and thus may, in part, explain some of the apparent recovery in performance between R+6 and R+21.

6.2 Muscular power/Jump

Jump height was reduced at R+6 (between 9-31% for the three different jumps) and was not fully restored and even showed decreases in the second measurement (R+21) for CMJ and SJ peak force. This prolonged deficit may not be due entirely to loss of power but neuromuscular ability, as jump test performance is strongly dependant on inter-muscular coordination and technique proficiency [23].

Given that jumping is a complex task requiring stability, strength and movement control, the supervised postflight programme may be too short to achieve full recovery. Dynamic and powerful movements are introduced relatively late in the programme to allow spinal/trunk stabiliser muscles to recover prior to doing exercises requiring complex motor control. Astronauts do not perform fast or explosive movements/exercise such as jumping or sprinting whilst on ISS, nor do they need to perform any work tasks that require maximal or near maximal effort [24]. Instead of using their legs to move through the station, they use their hands. Movement patterns adapted for μ G need to be “unlearned” upon return to Earth, and previous, gravity-adapted movement patterns and reflexes must be re-acquired to successfully perform jumps.

Anecdotally we observe that many crewmembers are unable to perform a valid (*i.e.* movement controlled) jump at R+6. The ability to control the motion is critical. Prior to any loaded movement of the lower extremities, such as jumps, trunk muscles, especially deep abdominal/muscle corset muscles (transversus abdominis, then multifidus), are sequentially activated, preparing for the movement [23]. The sequential activation of muscle influences jump height [25]. Correctly coordinated contraction of leg and trunk muscles, for any powerful motion in daily life, needs to be retrained to ensure that crewmembers can respond to hazardous situations. This is a critical safety aspect of the programme, even with fit individuals.

These observations appear to complement findings from another case study of the same crewmember, which measured the intrinsic trunk muscles (multifidus, transversus abdominis, internal oblique) using ultrasound imaging and observed that muscle size was maintained at L2-L4, but not at L5 [19]. The phenomenon of muscle atrophy at the level of L5 after longer phases of inactivity occurs not only in bed rest participants and

astronauts [6, 26], but is also in highly trained professional athletes [27]. Incomplete re-training of intrinsic back muscles after inactivity may lead to compensation by superficial muscles without their inherent ability to effectively stabilise and protect spinal segments.

6.3 Flexibility

Hip flexibility was also impaired in the present astronaut, perhaps due to the lack of eccentric motion in μ G which is required for walking, running or jumping [28]. Stretching exercises are performed by some crewmembers on ISS in preparation for strength training. Specific stretching sessions are not prescribed by exercise specialists, but are encouraged. The sudden transition back to Earth's gravity may contribute to overall muscle stiffness compromising flexibility as a reaction to the new load/gravitational environment, which may persist during the 21 day postflight exercise period. Relaxation and flexibility training, involving fascial training [29] in parallel to individual physiotherapy treatments, was implemented with this astronaut as part of his postflight programme.

6.4 Core muscle endurance

Core muscle static endurance in a given position (measured in seconds) did not change significantly from pre to postflight, although changes in strength (squat, bench press, and jumping) were observed at R+6. The ventral core test requires the ability to maintain continuous muscle contraction in a bench-like position, but it is not a complex or dynamic motion such as a jump. Other tests of core endurance performed in different postures were found to be unreliable [12] so are no longer used as part of the AFA.

6.5 Balance

The Balance results of the AFA appear rather inconclusive. The reliability of this test showed high errors between measurements that are probably not attributable to performance changes [13]. Therefore the current test might not be suitable for assessing balance sufficiently for this occupational group and more appropriate tests are needed.

Regardless of the findings, the physical performance of this crewmember was never regarded as critical given his good overall condition. Complete recovery was expected

throughout the months following reconditioning due to his ambitious personal exercise schedule.

6.6 Future direction

The ESA postflight reconditioning programme is based on the best evidence available from terrestrial rehabilitation [17]. Future exploration missions of up to three years, such as to Mars, will be even more challenging to recover from and require research to develop optimal reconditioning programmes. Inflight 'Preconditioning' programmes may also be needed inflight, to prepare for planetary surface exploration during these missions [30]. Parallels with deconditioning in areas of terrestrial rehabilitation, such as sports injuries, neurological disorders and intensive care may be helpful to draw on in the absence of studies on astronauts [31]. Simple, non-invasive ways of monitoring neuro-musculoskeletal function are also needed. Conducting research in the relatively small astronaut population is challenging and possible solutions are discussed by Beard and Cook [32].

7. Conclusions

This case report has highlighted that, despite a high degree of compliance with arigorous daily inflight exercise countermeasure programme, postflight neuro-musculoskeletal impairments still occur. The recovery process was not complete at the end of the 21-day reconditioning programme for dynamic powerful movements, whereas muscle strength and core muscle endurance recovered more fully, suggesting that motor control of movement may require more specific/further retraining. Research to develop optimal reconditioning programmes for longer exploration-type missions is needed urgently.

References

1. Ploutz-Snyder L, Ryder J, English K, Haddad F, K B. NASA Evidence Report: Risk of Impaired Performance Due to Reduced Muscle Mass, Strength, and Endurance. HRP-47072. 2015.
2. Fitts R, Riley DA, Widrick J. Physiology of a Microgravity Environment. Invited Review: Micrograv Skelet Muscle. 2000.
3. Baker E.S. BMR, Wear M.L. Human Response to Space Flight. Principles of Clinical Medicine for Space Flight Herausgeber: Barratt, Michael R, Pool, Sam Lee (Eds).27 – 58.
4. Sibonga JD, Cavanagh PR, Lang TF, LeBlanc AD, Schneider VS, Shackelford L et al. Adaptation of the Skeletal System During Long-Duration Spaceflight. Clin Rev Bone Miner Metab. 2008;5(4):249-61.
5. Wood SJ, Loehr JA, Guilliams ME. Sensorimotor reconditioning during and after spaceflight. NeuroRehab. 2011;29(2):185-95. doi:10.3233/NRE-2011-0694.
6. Belavy DL, Adams M, Brisby H, Cagnie B, Danneels L, Fairbank J et al. Disc herniations in astronauts: What causes them, and what does it tell us about herniation on earth? Eur Spine J. 2015. doi:10.1007/s00586-015-3917-y.
7. Moore AD, Jr., Downs ME, Lee SM, Feiveson AH, Knudsen P, Ploutz-Snyder L. Peak exercise oxygen uptake during and following long-duration spaceflight. J Appl Physiol (1985). 2014;117(3):231-8. doi:10.1152/jappphysiol.01251.2013.
8. Loehr JA, Guilliams ME, Petersen N, Hirsch N, Kawashima S, Oshima H. Physical Training for Long-Duration Spaceflight. Aerosp Med Hum Perform. 2015;86(12 (Supplement)):14-23.
9. Hayes J. The first decade of ISS Exercise: Lessons Learned on Expeditions 1-25. Aerosp Med Hum Perform. 2015;86(12 Suppl):A1-A6.
10. Korth DW. Exercise Countermeasure Hardware Evolution on ISS: The First Decade. Aerosp Med Hum Perform. 2015;86 (12 Suppl). doi:doi: 10.3357/AMHP.EC02.2015.
11. Kozlovskaya IB, Yarmanova EN, Yegorov AD, Stepanstov VI, Fomina ES, Tomilovskaya ES. Russian Countermeasure Systems for Adverse Effects of Microgravity on Long-Duration ISS Flights. Aerosp Med Hum Perform. 2015;86(12, Suppl):A24-A31.
12. Laughlin MS, Guilliams ME, Nieschwitz BA, Hoellen D. Functional Fitness Testing Results Following Long-Duration ISS Missions. Aerosp Med Hum Perform. 2015;86(12, Suppl):A87-A91.
13. Petersen N, Thieschafer L, Ploutz-Snyder L, Damann V, Mester J. Reliability of a new test battery for fitness assessment of the European Astronaut corps. Extrem Physiol Med. 2015;4:12. doi:10.1186/s13728-015-0032-y.

14. Brzycki M. Strength testing: Predicting a one-rep max from reps to fatigue. *J Phys Educ Recreat Dance* e. 1993;46(1):88-90.
15. Tschopp M. Leistungsdiagnostik Kraft. Swiss Olympic Medical Centers. 2003.
16. Petersen N, Jaekel P, Rosenberger A, Weber T, Scott J, Castrucci F et al. Exercise in space: the European Space Agency approach to in-flight exercise countermeasures for long-duration missions on ISS. *Extreme Physiol Med*. 2016;5(9).
17. Lambrecht G, Petersen N, Weerts G, Pruett C, Evetts S, Stokes M. The role of physiotherapy in the European Space Agency Strategy for Preparation and Reconditioning of Astronauts before and after long term space flight. *Musculoskel Sci Pract Supplement on "Terrestrial neuro-musculoskeletal rehabilitation and astronaut reconditioning: reciprocal knowledge transfer"* 2017; xx: xx-xx
18. LeBlanc A, Lin C, Shackelford L, Sinitsyn V, Evans H, Belichenko O. Muscle volume, MRI relaxation times (T2), and body composition after spaceflight. *J Appl Physiol*. 2000(89):2158-64.
19. Hides J, Lambrecht G, Stanton W, Damann V. Changes in multifidus and abdominal muscle size in response to microgravity: possible implications for low back pain research. *Eur Spine J*. 2016(25 Suppl 1):175-82. doi:10.1007/s00586-015-4311-5.
20. ACSM. Guidelines for Exercise Testing and Prescription. American College of Sports Medicine (ACSM). 2013;9.
21. English K, Lee SMC, Loehr JA, Ploutz-Snyder RJ, Ploutz-Snyder LL. Isokinetic Strength Changes Following Long-Duration Spaceflight on ISS. *Aerosp Med Hum Perform*. 2015(86(12, Suppl)):A68-A77. doi:10.3357/AMHP.EC09.2015.
22. Danneels L VG, Cambier DC, Witvrouw EE, De Cuyper HJ, Danneels L. CT imaging of trunk muscles in chronic low back pain patients and healthy control subjects. *Eur Spine J* 2000; 9: 266-272. 2000.
23. Wilson JD, Dougherty CP, Ireland ML, Davis McClay I. Core stability and its relationship to lower extremity function and injury. *J Am Acad Orthop Surg*. 2005(13):316-25.
24. Hackney KJ, Scott JM, Hanson AM, English KL, Downs ME, Ploutz-Snyder LL. The Astronaut-Athlete: Optimizing Human Performance in Space. *J Strength Condition Res*. 2015;29(12)/3531–3545.
25. Pereira R, Machado M, Miragaya dos Santos M, Pereira LN, Sampaio-Jorge F. Muscle activation sequence compromises vertical jump performance. *Serb Journal Sports Sci*. 2008(2):85-90.

26. Johnston LS, Campbell MR, Scheuring RA, Feiveson AH, . Risk of Herniated Nucleus Pulposus Among US astronauts. *Aviat Space Environ Med.* 2010;81:566-74.
27. Hides J, Stanton W. Muscle imbalance among elite Australian rules football players: a longitudinal study of changes in trunk muscle size. *J Athl Train.* 2012;47(3):314-9. doi:10.4085/1062-6050-47.3.03.
28. McBride JM MG, Cormie P. Influence of preactivity and eccentric muscle activity on concentric performance during vertical jumping. *J Strength Condition Res* 2008(22(3)):750-7.
29. Schleip R, Muller DG. Training principles for fascial connective tissues: scientific foundation and suggested practical applications. *J Bodyw Mov Ther.* 2013;17(1):103-15. doi:10.1016/j.jbmt.2012.06.007.
30. Stokes M, Evetts S, Hides J. Guest Editorial on "Terrestrial neuro-musculoskeletal rehabilitation and astronaut reconditioning: reciprocal knowledge transfer" *Musculoskel Sci Pract (Supplement)* 2017; xx: xx-xx
31. Hides J RG, Cusack R, McKay C, Bloomberg J, Stokes M. Parallels between astronauts and terrestrial patients – taking physiotherapy rehabilitation “to infinity and beyond”. *Musculoskel Sci Pract Supplement on "Terrestrial neuro-musculoskeletal rehabilitation and astronaut reconditioning: reciprocal knowledge transfer"* 2017; xx: xx-xx
32. Beard D CJ. Methodology for astronaut reconditioning research. *Musculoskel Sci Pract Supplement on "Terrestrial neuro-musculoskeletal rehabilitation and astronaut reconditioning: reciprocal knowledge transfer"* 2017; xx: xx-xx

Abbreviations

AFA	Astronaut Fitness Assessment
ARED	Advanced Resistive Exercise Device
CSA	Canadian Space Agency
EAC	European Astronaut Centre
ESA	European Space Agency
GCTC	Gagarin Cosmonaut Training Centre
ISS	International Space Station
JSC	Johnson Space Center (NASA)
L-	Launch date minus (days to launch)
L	Vertebra level (spine)
LDM	Long Duration Mission
MCE	Motor Control Exercise
NASA	National Aeronautics and Space Administration
PFE	Periodic Fitness Evaluation
R+	Return date plus (days after return)
RM	Repetition maximum
μG	Microgravity

For *Musculoskeletal Science and Practice* Supplement: “**Terrestrial neuro-musculoskeletal rehabilitation and astronaut reconditioning: reciprocal knowledge transfer**”

Guest editors: Maria Stokes, Simon Evetts, Julie Hides ”

Postflight reconditioning for European Astronauts – a case report of recovery after six months in space

Acknowledgements

The authors thank: the exercise team members Patrick Jaekel and Andre Rosenberger (led by Nora Petersen) for conducting and reporting on astronaut fitness assessments; Dr Volker Damann, former Head of the ESA Space Medicine Office, and Dr Guillaume Weerts, the new Head for supporting the work of the exercise team; the US and Russian rehabilitation counterparts for their advice and support; and the ESA Astronauts for sharing their experience and ideas, and especially their trust in our work. Special thanks goes to the ESA Astronaut for sharing his personal performance data and his support in producing this case report.

For *Musculoskeletal Science and Practice* Supplement: “**Terrestrial neuro-musculoskeletal rehabilitation and astronaut reconditioning: reciprocal knowledge transfer**”

Guest editors: Maria Stokes, Simon Evetts, Julie Hides

Highlights

- Despite intensive training , most performance measures were impaired postflight
- Most functional abilities recovered 21 days postflight after daily exercise
- Powerful dynamic tasks (e.g. jumps) did not recover during the 21 day programme
- Reconditioning focussed on motor control and functional training for holistic recovery

12 October 2016

For *Manual Therapy Supplement*: “**Terrestrial neuro-musculoskeletal rehabilitation and astronaut reconditioning: reciprocal knowledge transfer**”

Guest editors: Maria Stokes, Simon Evetts, Julie Hides ”

**Postflight reconditioning for European Astronauts – a case report of recovery
after six months in space**

Appendices

ACCEPTED MANUSCRIPT

Appendix 1. Selected AFA tasks, parameters and measurement protocols

AFA element	Parameter	Device	Protocol	Reference
Body anthropometrics	Height	Seca scale	Metric measurement	
1 RM [kg]	<i>Maximal load</i>	Smith machine	Calculation from 2 sets of max 10 reps	[14]
Jump height	Best of 3 trials in [cm]	Templo software on Kistler force platform		[13]
Jumps peak force	Best of 3 trials in [N]	Templo software on Kistler force platform		[13]
Core endurance	Holding standardized position as long as possible			[15],[13]
Hip flexibility	Sit and reach test: reaching forward on SNR box Thomas Test, hip angle measurement			[13]

1 RM, repetition maximum (squat and bench press)

Appendix 2. Overview of ESA reconditioning exercises

Activity	Exercises	Application	Timing (NET)
2 <i>Pool workout</i>	Balancing on pool noodles, moving boards against water resistance, aqua jogging, water ball mini games. Hopping and jumping exercises, slow and fast movements with water training equipment	Enabling dynamic but safe exercise using water resistance for proprioceptive, movement control, local muscular endurance training and fast movements against water resistance	R+1
3 <i>Cardiovascular training</i>	Cycling, rowing, (suspended) running, swimming, exercise circuits	Training of cardio-vascular capacity in parallel, continuously adapting to crew level	R+1
4 <i>Body weight exercises</i>	Selection of gymnastics, Pilates, yoga, martial arts, flexibility (especially lower legs), gym ball exercises and ball games (eye-hand coordination, balance, rotation), concentric resistive exercises. Additional exercise tools (balls, bars, bags, ...) are used to challenge movement control	Functional training requiring postural and movement control, preparing for life and sports application Focus on functional training concept preparing for and successively integrating strength, power, local muscular endurance and cardio-vascular training (all throughout entire programme)	R+2
5 <i>Resistive training</i>	Weight lifting, squats, deadlift, seated-row, rotational cable pulls, lunges, dumbbells/Kettlebells	Loaded training to trigger muscle hypertrophy and bone strengthening, supporting a comprehensive re-adaptation to athletic training	R+5
6 <i>Plyometric and reactive training</i>	Hopping, jumping, eccentric movements, ball throws, ball catches during motion, responding to signals, dynamic lifting, Kettlebell training, martial arts elements	Recovery of athletic components of sports and activities of daily life. Challenging of postural and movement stability, based on preceding progress made, integration of dynamic/fast motions	R+15
1 <i>Physiotherapy</i>	Physiotherapy treatments are implemented in parallel to complement exercise sessions.	Musculoskeletal stress symptoms (e.g. on the AC joint or rotator cuff, plantar fascia, lower lumbar and/or	R+0

lower cervical spine) effects of the intense daily programme can be treated and thus immediately neutralized to enable the crewmember to continue the exercise training as foreseen.

NET, not earlier than

ACCEPTED MANUSCRIPT

Appendix 3. Case report reconditioning outline

Return date +	Device/Facility	Duration (min)	Exercises/Protocol	Activity	Intensity
R+1	Physio	120	Physiotherapy only	1	
	Physio	60	Physiotherapy	1	
R+2	Gym	20	Cardiovascular exercise	3	low-medium
	Gym	40	(Gym) Ball exercises, stretching	4	medium
R+3	Gym	10	Physiotherapy	1	
	Gym	110	(Gym) Ball exercises, stretching, CINDY	4	medium
R+4	Gym	120	Resistance exercise	5	medium
R+5	Physio	120	Mat exercises, ball exercises	4	medium
R+6	Gym	30	Alter G® treadmill running	3	medium-hard
	Gym/lab	120	AFA		medium-hard
R+7	Gym	120	Functional Fitness, stretching	4	low-medium
R+8	Gym	120	Resistance exercise	5	medium
R+9	Gym	20	Mat exercises	4	low-medium
	Gym	75	Cardiovascular exercise: Parcours	3,6	medium-hard
	Pool	15	Swimming	2	low-medium
R+10	Gym	45	Functional Fitness, agility (jumps, vision)	6	medium
	Gym	175	Resistance exercise	5	medium
R+14	Gym	120	Resistance exercise	5	medium-hard
R+15	Gym	120	Circuit training	3	medium
			<i>continued unsupervised training programme</i>		
R+21	At EAC Gym/lab		AFA		
			<i>continued unsupervised training programme</i>		

R+, Return date +; AFA, Astronaut fitness assessment; EAC, European Astronaut Centre; Alter G®, suspended treadmill system; lab, laboratory; CINDY; cross-fit protocol

The activity numbers refer to the table in Appendix 2, which presents exercises more specifically. For more details see Appendix 4.

Appendix 4. Objectives, methods and exercise examples of ESA reconditioning programme

Time period (Phase/R+)	Objective(s)	Exercise Type/Method	Example	Intensity	Volume
Phase 1					
R+1	Physiotherapy: <ul style="list-style-type: none"> - status quo Training of multifidus and m transversus <ul style="list-style-type: none"> - Check posture, tonus, tension, mobility - Diagnostics (US) - Reorganization of stabilizing chain - Posture correction - Tonus normalization - Muscle balance - Active postural control 	MCE (Motor control exercise) Isolated exercises for m multifidus and m transversus Exercises for all stabilizer muscles Closed chain exercises Axial training Massage	Prone position: breath in and out with isolated/segmental activation of multifidus (hand feedback), Supine position: draw-in belly button Trunk exercises	Low	90 min
R+2	<ul style="list-style-type: none"> - Progression on R+1 exercise - Isolated training of stabilizers - Integration of stabilizing muscle system into complete muscle corset - Muscle balance - Active postural control 	MCE with rotational elements: Axial training Upright position (sitting, standing) Individual physiotherapy treatments Closed chain exercises	Kneeling: rotate basket ball (upper body rotation) Sitting: lift med ball towards ceiling Trunk and arm exercises Rotational exercises Physical exercise session	Low	90 min
R+3	<ul style="list-style-type: none"> - Progression on R+1 exercise - Isolated training of stabilizers - Integration of stabilizing muscle system into complete muscle corset 	MCE with rotational elements: Axial training Balance training Coordination training Increasing movement velocity (w increased stability of CM) Training of shoulder axis	Standing: rotate ball side to side Trunk, arm and leg exercises Walking on unstable underground Decrease of base of support Physical exercise session	Low	90 min

	<ul style="list-style-type: none"> - Muscle balance - Active postural control 				
R+5	<ul style="list-style-type: none"> - Integration of postural control into more complex exercises - Increased demand and complexity of movement - Approximation to functional exercise movements 	Manual therapy Fascial training Low intensity functional exercises Balance/Coordination training Local muscular endurance Whole body loading Stretching Closed eyes exercises (MCE)	Flexibar exercise on balance board Physical exercise session Endurance training with reduced body weight (pool, Alter-G), Resistance training (low loading, controlled motion – continued supervision of posture by coach) Trunk rotation movements (e.g. such as in Tai Chi)	Low	90 min
R+7	<ul style="list-style-type: none"> - Integration of postural control into more complex exercises - Increased demand and complexity of movement - Approximation to functional exercise movements - Increase of complexity and intensity 	Manual therapy Fascial training Reaction training Core stabilization Balance/Coordination training Local muscular endurance Whole body loading Stretching	Complex coordinative ball handling exercises Initiation of combination moves Squats with low loads and (supervised) focus on form	Low-medium	90 min
R+10	<ul style="list-style-type: none"> - Increased complexity and intensity of preceding exercises 	Same as before <ul style="list-style-type: none"> - Continued integration of physiotherapy exercise preparations into exercise programme - Motion control in exercise movements 	Physical exercise session Complex combination of forward steps, balance movements and sudden stops	Low-medium	
R+11	<ul style="list-style-type: none"> - Based previous results, 	Same as stated above/before and	Physical exercise session Complex combinations of	Medium	90 min

	<p>increased integration of functional exercises/movements</p> <ul style="list-style-type: none"> - Increased complexity - Increased intensity 	<p>additionally</p> <ul style="list-style-type: none"> - Integration of reactive/eccentric exercises 	<p>forward steps, balance movements and sudden stops (movement control throughout entire motion)</p> <p>Plank exercises Push-ups on gym ball (core strength). Hopping exercises w coordination ladder, steps Rowing (Endurance training) Stretching</p>		
R+13	<ul style="list-style-type: none"> - Based previous results, increased integration of functional exercises/movements - Increased complexity - Increased intensity - Construction of "target movements" 	<p>Same as before and</p> <ul style="list-style-type: none"> - Agility - Extended 3D-spatial orientation (+management of additional tools: ball, ...) 	<p>Physical exercise session Combination of jumping and throwing Moving in larger radius Parcours (combine moving and tasks) Circle training (endurance, coordination)</p>	Medium	90 min
R+15	<ul style="list-style-type: none"> - Based previous results, increased integration of functional exercises/movements - Increased complexity - Increased intensity - Construction of "target movements" 	<ul style="list-style-type: none"> - Coordination and endurance - Construct target final movement (isolated elements) - Training of muscular strength (incl. stability) - Endurance - Agility - Plyometric basics 	<p>Physical exercise session Coordination and endurance Ball dribbling (varied situations with partner) Resistance exercises: Squat, Back extensions/deadlift, Hopping exercises (plyometric training prep/controlled) Parcours</p>	Medium	90 min
R+18	<ul style="list-style-type: none"> - Integration of optimized posture and movement strategies into 	<ul style="list-style-type: none"> - Training of elements of more complex movements 	<ul style="list-style-type: none"> - Step-ups with passing/catching / throwing a ball, - Work on constructing 	Medium	90 min

	<p>more complex functional (athletic and daily life) motion</p> <ul style="list-style-type: none"> - Posture protection & - Movement safety - Continued body loading 	<p>with many repetitions (eye-hand coordination, distracted attention (2 simultaneous tasks))</p> <ul style="list-style-type: none"> - Training of muscular strength (incl. stability) – progressing intensity - Endurance - Agility 	<p>elements of more complex movements into one</p> <ul style="list-style-type: none"> - Loaded squats/deadlifts/seated row 		
R+21	<ul style="list-style-type: none"> - Achieve target exercises/final target movement performance (complex combination with basketball) - Successful and complete integration of postural optimization info functional movement - Bone loading - Achieving complex balance/movement stability - Enable crew to return into complex and intense physical exercise and - “Crew physical functional empowerment” - Pick-up a “normal” as preflight exercise programme 	<p>Athletic movements from ball/game sports, resistance & strength training, Pilates, Martial arts fitness, Endurance (Running, Cycling, Swimming, ...), Plyometric exercises (controlled)</p>	<p>Target exercise movements/positions:</p> <ul style="list-style-type: none"> - Basketball dribble and shot - Balance complex position (Tai Chi/Yoga) (Standwaage) - Work/Life: lift & carry (heavy) load - Sports : MMA/“Crossfit”? /functional fitness movement variations - Sports: Pilates/Yoga - Sports: Climbing (if possible) 	<p>Medium-high (in complexity, not yet extensive loading) Significant for endurance and strength training</p> <p>Control of movement over intensity (priority)</p>	90 min