

Genomics of Myeloproliferative Neoplasms

Katerina Zoi¹ and Nicholas C.P. Cross^{2,3}

1. Haematology Research Laboratory, Biomedical Research Foundation, Academy of Athens, Athens, Greece
2. Wessex Regional Genetics Laboratory, Salisbury District Hospital, Salisbury, UK
3. Faculty of Medicine, University of Southampton, Southampton, UK

Corresponding author:

Professor Nicholas C.P. Cross
Wessex Regional Genetics Laboratory
Salisbury District Hospital
Salisbury SP2 8BJ, UK

Tel +44 1722 429080

Fax +44 1722 331531

Email: ncpc@soton.ac.uk

Abstract

Myeloproliferative neoplasms (MPN) are a group of related clonal hematological disorders characterized by excess accumulation of one or more myeloid cell lineages and a tendency to transform to acute myeloid leukemia. Deregulated JAK2 signaling has emerged as the central phenotypic driver of *BCR-ABL1*-negative MPN and a unifying therapeutic target. In addition, MPN show unexpected layers of genetic complexity, with multiple abnormalities associated with disease progression, interactions between inherited factors and phenotype driver mutations, and effects related to the order in which mutations are acquired. Although morphology and clinical laboratory analysis continues to play an important role in defining these conditions, genomic analysis is providing a platform for better disease definition, more accurate diagnosis, direction of therapy and refined prognostication. There is an emerging consensus with regard to many prognostic factors, but a clear need to synthesize genomic findings into robust, clinically actionable and widely accepted scoring systems, and well as the need to standardize the laboratory methodologies that are employed.

Introduction

The World Health Organization classification¹ recognizes seven subtypes of myeloproliferative neoplasms (MPN). Chronic myeloid leukemia is unique in that is defined by the presence of a specific somatic abnormality, the *BCR-ABL1* fusion gene. Of the *BCR-ABL1*-negative MPN (the focus of this review), polycythemia vera (PV), essential thrombocythemia (ET) and primary myelofibrosis (PMF) are often referred to collectively as classic MPN. Chronic neutrophilic leukemia (CNL), chronic eosinophilic leukemia (CEL) and MPN unclassifiable (MPN-U) are much less common and referred to as atypical or non-classic MPN. During their diagnostic work up, MPN need to be carefully distinguished from other myeloid entities by a combination of morphologic, clinical laboratory, cytogenetic, molecular genetic and, increasingly, genomic analysis.

Classic *BCR-ABL1* negative MPN

Deregulation of JAK2/STAT signalling

Direct or indirect dysregulation of JAK2 signalling by somatically acquired mutations has emerged as a central phenotypic driver of classic MPN. JAK2 is a non-receptor tyrosine kinase which plays an essential role in transducing signals from class 1 cytokine receptors critical for normal myelopoiesis, notably the erythropoietin receptor, the thrombopoietin receptor (TPOR; encoded by the *MPL* gene) and the granulocyte colony-stimulating factor receptor.^{2,3}

The principal driver mutations in MPN are indicated on Figure 1. JAK2 is directly activated by acquisition of the *JAK2* V617F mutation, seen in the majority of MPN.⁴⁻⁷ *JAK2* V617F mutation burdens are usually higher in PV and PMF compared to ET, with many cases demonstrating a homozygous mutant clone.⁸ Mutations of *JAK2* exon 12, typically complex insertion/deletion events, are seen in roughly one third of cases of *JAK2* V617F-negative PV.⁹

Indirect dysregulation of JAK2 signalling occurs principally by activating mutations in *MPL* or *CALR*. *MPL* mutations are located in exon 10 and most commonly result in amino acid substitutions at W515 of the thrombopoietin receptor (TPOR).^{10,11} W515 is located within a small amphipathic motif of the human receptor at the junction between the transmembrane and cytoplasmic domains that is required for maintaining TPOR in its inactive conformation and prevents its autonomous activation; mutation of this residue abrogates this function thereby constitutively activating JAK2 and downstream signalling.^{10,12}

CALR exon 9 mutations are found in about 25-30% of patients with ET and PMF.^{13,14} All pathogenic *CALR* result in a common frameshift encoding a novel C-terminus with a high positive charge, with 80% of mutants being type-1 (52bp deletion) or type-2 (5bp insertion).^{13,14} Calreticulin is a highly conserved chaperone protein that directs the correct conformation and intracellular trafficking of glycoproteins, as well as in homeostatic control of calcium levels. Remarkably, calreticulin mutants specifically activate TPOR, and thereby JAK2, after binding to its extracellular N-glycosylation residues.¹⁵⁻¹⁷

Triple negative MPN

ET and PMF negative for standard phenotype driver mutations (*JAK2*, *CALR* and *MPL*) are referred to as 'triple negative'. High throughput sequencing identified non-canonical activating *JAK2* or *MPL* mutations in a minority of triple negative MPN, some of which were somatic but others were inherited, thus identifying cases with hereditary thrombocythemia misdiagnosed as MPN. Similarly, some cases had no evidence for a clonal disorder suggesting that they had also been misdiagnosed. Other cases, particularly those with a diagnosis of PMF, had clonal disease associated with a range of mutations characteristic of MDS/MPN and other myeloid disorders e.g. in *CBL*, *SH2B3*, *TET2*, *ASXL1* and *SRSF2*.^{14,18,19}

MPN genomes: cytogenetics, arrays and exome sequencing

Karyotypic abnormalities detected by conventional cytogenetic analysis are seen in a minority of PV and ET (<10%), but about a third of PMF.²⁰ Gain of chromosome 9 is seen recurrently in PV and associated with *JAK2* V617F (*JAK2* is located at 9p24). Interstitial deletions of 13q and 20q are seen in all MPN subtypes suggesting the presence of one or more tumour suppressor genes in these regions. Of these, 20q deletion is the most frequent and may target the imprinted genes *L3MBTL1* and *SGK2*.²¹ Other recurrent abnormalities in MPN for which the molecular basis is unknown include +8 and der(6)t(1;6).²²

Single nucleotide polymorphism (SNP) array analysis has confirmed that PV and ET genomes are relatively simple with very few somatic copy number changes.^{23,24} By contrast, copy number changes are more common in PMF and MF evolving from PV or ET.^{23,25} SNP arrays detect frequent regions of acquired uniparental isodisomy (aUPD) in MPN. Such regions arise from mitotic recombination and enable genetic variants to progress from a heterozygous to a homozygous state without change in DNA copy number. Chromosome 9p is most commonly affected and is almost always associated with homozygosity for *JAK2* V617F.⁶ Other chromosomes are affected less frequently, e.g. aUPD at 4q, 7q and 11q are associated with mutations in *TET2*, *EZH2* and *CBL*, respectively, whereas aUPD14q is associated with an imprinting defect.²⁶⁻²⁹

Exome sequencing identified a median of 6.5 total somatic mutations in ET and PV, increasing to 13 in MF, consistent with the latter being a more advanced stage of disease.¹⁴ Most of these mutations are passengers that almost certainly play no role in the disease process. RNAseq has not thus far identified any new in-frame oncogenic fusion genes, although this technique may detect loss of function events.³⁰ Whole genome sequencing for MPN has not been reported, but may, potentially reveal new somatic changes in non-coding regions of the genome. Overall, however, it is anticipated that the MPN genome is largely understood, at least in terms of the spectrum and frequency of somatic mutations. By contrast, the epigenome remains poorly understood.

Initiating mutations and phenotype modifying mutations

JAK2 V617F can be detected in the general population at a prevalence of 0.1-0.2%, roughly 10x the prevalence of diagnosed MPN.³¹⁻³⁵ Although some of these mutated individuals turn out to have a previously undiagnosed MPN, many have blood counts within the normal range and thus represent examples of what is now termed age related clonal hematopoiesis (ARCH) or clonal hematopoiesis of indeterminate potential (CHIP). Some of these individuals subsequently develop an MPN but many maintain normal blood counts. A *JAK2* V617F mutation burden of >2% has been proposed as suggestive of an MPN,³⁶ but this is not a clean cut off and some individuals with ARCH /CHIP have higher *JAK2* mutation burdens.

Broad population-based mutations screens indicate that ARCH/CHIP is associated with mutations in several different genes with most affected individuals having only one discernible mutation.³⁷⁻³⁹ Up to 50% of *JAK2* V617F mutated PV and ET cases also have no additional pathogenic mutations^{40,41} and thus whilst it appears that *JAK2* V617F is sufficient to drive clonal hematopoiesis and an MPN phenotype in many cases, additional factors may play a role in promoting the likelihood of an MPN phenotype developing.

About 50% of *JAK2* V617F mutated ET and PV cases have pathogenic mutations in additional genes, most commonly *DNMT3A* and *TET2*. The great majority of *JAK2* V617F positive PMF cases have additional mutations, particularly those encoding chromatin modifiers and splicing factors, suggesting that these abnormalities promote an MF phenotype.^{40,41}

Remarkably, the order in which mutations are acquired is not fixed. In some patients *JAK2* V617F is acquired early in the disease and then additional mutations appear as subsequent subclonal events. In other patients, clinically covert clonal hematopoiesis is believed to be established by mutations in *DNMT3A*, *TET2* or other abnormalities such as del(20q) or aUPD14q, and *JAK2* V617F is acquired as a later event, triggering the onset of an MPN. The order in which mutations are acquired influences the disease phenotype: patients who acquired *JAK2* V617F before *TET2*, *DNMT3A* or aUPD14q were more likely to develop PV than ET compared to cases where *JAK2*

V617F was acquired late.^{26,42,43} In addition, detailed analysis of *TET2/JAK2* co-mutated patients indicated that the order in which mutations are acquired influences tumor biology as well as the behaviour of stem and progenitor cells.⁴³

In contrast to *JAK2 V617F*, *CALR* mutations show a high degree of clonal dominance throughout hematopoietic development and are believed to be the initiating event in most cases. Additional mutations are thus secondary events associated with disease evolution.^{13,14,40,44} The principal genetic abnormalities and evolution of MPN are shown on Figure 2.

Factors determining disease phenotype

Although MPN phenotype driver mutations all induce the activation of JAK2, they do not lead to the same disorder. *JAK2 V617F* is associated with PV, ET and PMF while *MPL* and *CALR* mutations are associated with ET and PMF. *JAK2 V617F* mutant ET patients tend to present many of the phenotypic features of PV and an increased risk of thrombosis compared to ET negative for *JAK2 V617F*.^{45,46} Cases with *MPL* mutations are on average older and have higher platelet counts, lower hemoglobin levels and reduced bone marrow cellularity at diagnosis than patients with *JAK2 V617F*, although, no significant differences in clinical outcome are apparent.⁴⁷

CALR mutations in ET are associated with higher platelet counts, lower haemoglobin levels, lower leukocyte counts and younger age at presentation, while the incidence of thrombotic events is lower than *JAK2 V617F* positive ET.⁴⁸⁻⁵⁰ Although no differences in survival have been demonstrated for the different driver mutation subgroups,⁵⁰ ET patients with type-1 *CALR* mutations progress to MF much more frequently than those with type-2 mutations.⁵¹

In PMF the nature of the MPN driver mutation has a stronger prognostic significance. Compared to *JAK2 V617F* mutated cases, patients with *CALR* mutations (most commonly type-1 in PMF) present at a younger age, with higher platelet counts and lower leukocyte counts, but with reduced anemia and transfusion dependency, while they have a better prognosis regarding overall and leukemia-free survival.

Importantly, triple-negative patients for both PMF and post-MPN MF have the worst prognosis and a more severe anemia.^{48,52,53} Compared to PV and ET, the *JAK2* V617F and *CALR* mutant allele burden is significantly higher in MF evolving from either ET or PV, indicating a role for the accumulation of mutated alleles in the process of transformation.⁴⁸

The reason why some *JAK2* V617F positive patients present with PV whereas others present with ET or other myeloid disorders is likely to be multifactorial. As discussed above, the *JAK2* mutant allele burden is higher in PV compared to ET and the order in which mutations are acquired can influence the disease phenotype. Constitutional genetics also influences the disease phenotype, with inherited variation in the *HBS1L-MYB* region influencing whether *JAK2* V617F mutant cases develop ET or PV.²⁴ Phenotypic determinants in MPN may also include factors other than acquired mutations and constitutional genetic modifiers. PV and PMF are more common in males and ET is predominant in females,⁵⁴ whilst *CALR*-mutated ET is more prevalent in men and *JAK2* V617F is more common in women,⁵⁵ suggesting that gender-specific factors may influence MPN phenotype. Hormones could be an explanation for these differences, as oestrogens can inhibit the *JAK2* V617F mutated stem cells,⁵⁶ an observation that may explain the observation that *JAK2* V617F allele burdens are lower in women compared to men.⁵⁷ Iron status, especially in premenopausal women, is another factor relevant to MPN phenotype.⁵⁴

Disease progression and prognostic significance of disease modifying mutations

Considerable effort has been put into the identification of genetic predictors of transformation from MPN to AML. Robust molecular prognostic markers are still lacking for patients with PV and ET, and risk stratification for these patients is still based mainly on clinical criteria and the presence of phenotypic driver mutations. The mutational burden is certainly higher on transformation of MPN to AML with the appearance of variants in a range of genes that are not unique to MPN but are also seen in other myeloid malignancies (Table 1). Gene expression profiling with adjustment for gender-specific effects has been suggested to provide prognostic information in PV,⁵⁸ but this approach remains unvalidated. Overall, whilst no strong

predictors of transformation have been identified, some clinically useful prognostic factors have emerged, particularly for the management of patients with MF.

Mutations in genes involved in transcription or DNA damage response are clearly associated with leukemic transformation, particularly *RUNX1* and *TP53* but also *IKZF1* and *CUX1*.^{25,41,59,60} Mutations in *TP53* may persist in a heterozygous state for an extended period of time during chronic phase MPN without clonal expansion. However, after loss of the wild-type allele by either chromosomal deletion or aUPD, the hemizygous or homozygous *TP53* clone rapidly expands, leading to leukemic transformation.⁴¹ Mutations in *RUNX1* and other transcription factors are usually acquired at the time transformation, thus limiting their use as predictive factors.⁶⁰

Impairment of polycomb repressive complex 2 (PRC2) function also appears to be an important pathogenic mechanism that promotes disease progression. Both *EZH2* and *ASXL1* mutations are associated with an increased risk of leukemic transformation and inferior survival.^{41,61-63} In PMF, *ASXL1* is an important prognostic indicator, with *CALR* wild type, *ASXL1* mutated cases having the worst prognosis.⁶⁴

Mutations in genes encoding chromatin modifiers are not consistently associated with prognosis. Whereas *DNMT3A* mutations are not considered to be prognostically significant, there are conflicting data regarding the association between *TET2* mutations and the risk of leukemic transformation.^{41,65} Rather than promoting disease progression, mutations in these genes may be considered as landscaping events that create a 'fertile ground', i.e. an environment more conducive to the development of a myeloid neoplasm.⁶⁶ Early studies indicated, unexpectedly, that *JAK2* V617F mutated MPN may evolve to *JAK2* unmutated AML.^{67,68} Most likely this is due to the presence of a pre-existing, clonal phase driven by *DNMT3A* or *TET2* mutations with this clone independently acquiring *JAK2* V617F (leading to an MPN) or other mutations (leading to AML). Alternatively, it is possible that some individuals have an inherited tendency to develop myeloid malignancies and thus develop MPN and AML as truly independent clones.

In PMF the presence of mutations in any one of *ASXL1*, *SRSF2*, *IDH1/2* or *EZH2* has been shown to be associated with reduced overall survival and increased risk for leukemic transformation.⁶⁹ Specifically, the number of these “high-molecular-risk” (HMR) mutations was inversely correlated with median survival in PMF independently of standard risk scores.⁶³ Notably, HMR mutations are enriched in triple negative PMF, while spliceosome mutations (*SRSF2*, *SF3B1*, *U2AF1*) are infrequent in *CALR*-mutant patients compared with the *JAK2/MPL*-mutant and triple-negative patients.⁷⁰ Mutational analysis in PMF patients who received ruxolitinib revealed that patients with ≥ 3 mutations of any type had significantly lower odds of a spleen response and a shorter time to treatment discontinuation. Patients with ≥ 1 mutations in *ASXL1*, *EZH2*, *IDH1* or *IDH2* had shorter survival and time to treatment discontinuation, and were significantly less likely to have a spleen response.⁷¹ Additional mutations may correlate with other relevant clinical factors, for example in PMF the presence of *U2AF1* mutations is strongly associated with severe anemia and thrombocytopenia.⁷² Overall, these data suggest that broad mutational analysis, for PMF patients at least, should be integrated into routine management.

Genetic predisposition

Familial predisposition to MPN may occur in up to 5-8% of apparently sporadic cases.⁷³ MPN phenotype driver mutations are never inherited but are frequently acquired somatically in familial cases, as they are in sporadic cases.⁷³⁻⁷⁵ As noted above, inherited, relatively weakly activating *JAK2* and *MPL* mutations are associated with hereditary erythrocytosis or thrombocytosis, disorders that are non-clonal but which may occasionally be mistaken for true MPN.⁷⁶⁻⁷⁸

The causes of familial predisposition to MPN are largely unknown. The inheritance patterns are heterogeneous and attempts to aggregate families for linkage analysis have largely failed, suggesting there are probably a variety of different germline mutations of variable penetrance driving the effect.⁷⁹ Thus far, only one clear explanation for high penetrance germline predisposition has emerged: analysis of four large families identified a 14q duplication including *ATG2P* and *GSKIP* that segregated with the MPN phenotype (ET/AML) and provided a ‘fertile ground’ for

MPN development.⁸⁰ Other abnormalities have also been associated with familial MPN, e.g. germline mutations in the *RBBP6* gene in about 5% of familial cases.⁸¹

Large epidemiological studies have suggested the presence of weakly penetrant, common risk factors in the general population.⁸² Several of these factors have been identified, but just two (*JAK2* 46/1 and *TERT*) account for the major part of the population attributable risk for development of an MPN, although the underlying reasons for this effect remain obscure. *JAK2* 46/1 (also known as GGCC) strongly predisposes to *JAK2*-mutated MPN but also, to a much lesser degree, to MPN with other mutations. *TERT* variants strongly predispose to all MPN.^{24,31,83-87} Although these effects are strong in genetic epidemiological terms, their penetrance is very low and consequently they cannot be used to predict whether any individual is likely to develop an MPN. Nevertheless, aggregation of these factors may contribute to apparent familial clustering.⁷⁹

Non-classic or atypical MPN

Chronic neutrophilic leukemia

Frequent oncogenic mutations of *CSF3R* are seen in CNL, a rare MPN subtype sharing overlapping features with atypical CML.⁸⁸⁻⁹⁰ The initial study found that 50-60% of patients with CNL or atypical CML harboured *CSF3R* mutations⁸⁸ with subsequent reports indicating that such abnormalities were restricted to CNL and essentially absent in aCML and MDS/MPN-unclassified.^{89,90} Co-operative mutations in *SETBP1* and *ASXL1* appear to be of prognostic significance and correlate with disease progression.⁹¹

CSF3R encodes the receptor for granulocyte-colony stimulating factor 3, which requires *JAK2* to function. Mutations fall into 2 types: nonsense or frameshift mutations leading to premature truncation of the cytoplasmic tail of the receptor and point mutations in the extracellular domain, most commonly T618I, an abnormality that strongly activates the *JAK/STAT* signalling. *CSF3R* mutations are

almost always acquired, although a recent report described a child with CNL and an inherited T618I mutation.⁹² Ongoing studies are evaluating whether CNL patients are amenable to treatment with JAK2 or other inhibitors, as suggested by initial reports.^{88,93}

Chronic eosinophilic leukemia and MPN-unclassified

CEL (not otherwise specified) is distinguished from idiopathic hypereosinophilic syndrome largely by the finding of a marker of clonality. Most commonly this has been achieved by cytogenetics, although only a very small proportion of suspected cases turn out to have a karyotypic abnormality. Cytogenetics and/or fluorescent *in situ* hybridization (FISH) is also critical to help identify cases with *FIP1L1-PDGFR*A or other tyrosine kinase fusions such as those involving *PDGFR*A, *PDGFR*B, *FGFR*1 or *JAK*2, abnormalities that are amenable to treatment with specific targeted therapies.

A recent study has demonstrated that targeted next-generation sequencing helps to establish clonality in a subset of patients with cytogenetically normal hypereosinophilia that would otherwise have been classified as idiopathic hypereosinophilic syndrome. Importantly, the survival of these cases was significantly shorter than cases without mutations and was indistinguishable from cases with cytogenetically defined CEL, clearly indicating the value of genomic analysis in this disorder.⁹⁴ There are no molecular markers for MPN-U but it is possible that wide mutation screens may help to better define this heterogeneous condition.

Concluding remarks

The 2016 revision of WHO classification has incorporated testing for MPN phenotype driver mutations as essential components in the diagnostic workup of MPN. The presence of *BCR-ABL*1 defines CML but this fusion needs to be excluded in the diagnosis of classic MPN and other tyrosine kinase fusions need to be excluded in

cases with eosinophilia. It is also suggested that testing for the most frequent disease modifying mutations may assist in the determination of the clonal nature of the disease.¹ and, as discussed above, many of these abnormalities have prognostic value. Currently, many centers perform a combination of cytogenetics, FISH, reverse transcriptase polymerase chain reaction (RT-PCR), Sanger sequencing and NGS panels in the diagnostic and prognostic work up of MPN. Clearly the time is ripe for genomics-based approaches, for example RNAseq, to supplant these diverse techniques and provide a platform for improved management and understanding of MPN.

References

1. Arber DA, Orazi A, Hasserjian R, et al: The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood* 127:2391-405, 2016
2. Cross NC: Genetic and epigenetic complexity in myeloproliferative neoplasms. *Hematology Am Soc Hematol Educ Program* 2011:208-14, 2011
3. Vainchenker W, Constantinescu SN: JAK/STAT signaling in hematological malignancies. *Oncogene* 32:2601-13, 2013
4. Baxter EJ, Scott LM, Campbell PJ, et al: Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet* 365:1054-61, 2005
5. James C, Ugo V, Le Couedic JP, et al: A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. *Nature* 434:1144-8, 2005
6. Kralovics R, Passamonti F, Buser AS, et al: A gain-of-function mutation of JAK2 in myeloproliferative disorders. *N Engl J Med* 352:1779-90, 2005
7. Levine RL, Loriaux M, Huntly BJ, et al: The JAK2V617F activating mutation occurs in chronic myelomonocytic leukemia and acute myeloid leukemia, but not in acute lymphoblastic leukemia or chronic lymphocytic leukemia. *Blood* 106:3377-9, 2005
8. Godfrey AL, Nangalia J, Baxter EJ, et al: Nongenetic stochastic expansion of JAK2V617F-homozygous subclones in polycythemia vera? *Blood* 124:3332-4, 2014
9. Scott LM, Tong W, Levine RL, et al: JAK2 exon 12 mutations in polycythemia vera and idiopathic erythrocytosis. *N Engl J Med* 356:459-68, 2007
10. Defour JP, Chachoua I, Pecquet C, et al: Oncogenic activation of MPL/thrombopoietin receptor by 17 mutations at W515: implications for myeloproliferative neoplasms. *Leukemia* 30:1214-6, 2016
11. Pikman Y, Lee BH, Mercher T, et al: MPLW515L is a novel somatic activating mutation in myelofibrosis with myeloid metaplasia. *PLoS Med* 3:e270, 2006
12. Defour JP, Itaya M, Gryshkova V, et al: Tryptophan at the transmembrane-cytosolic junction modulates thrombopoietin receptor dimerization and activation. *Proc Natl Acad Sci U S A* 110:2540-5, 2013
13. Klampfl T, Gisslinger H, Harutyunyan AS, et al: Somatic mutations of calreticulin in myeloproliferative neoplasms. *N Engl J Med* 369:2379-90, 2013
14. Nangalia J, Massie CE, Baxter EJ, et al: Somatic CALR mutations in myeloproliferative neoplasms with nonmutated JAK2. *N Engl J Med* 369:2391-405, 2013
15. Chachoua I, Pecquet C, El-Khoury M, et al: Thrombopoietin receptor activation by myeloproliferative neoplasm associated calreticulin mutants. *Blood* 127:1325-35, 2016
16. Elf S, Abdelfattah NS, Chen E, et al: Mutant Calreticulin Requires Both Its Mutant C-terminus and the Thrombopoietin Receptor for Oncogenic Transformation. *Cancer Discov* 6:368-81, 2016

17. Marty C, Pecquet C, Nivarthi H, et al: Calreticulin mutants in mice induce an MPL-dependent thrombocytosis with frequent progression to myelofibrosis. *Blood* 127:1317-24, 2016
18. Cabagnols X, Favale F, Pasquier F, et al: Presence of atypical thrombopoietin receptor (MPL) mutations in triple-negative essential thrombocythemia patients. *Blood* 127:333-42, 2016
19. Milosevic Feenstra JD, Nivarthi H, Gisslinger H, et al: Whole-exome sequencing identifies novel MPL and JAK2 mutations in triple-negative myeloproliferative neoplasms. *Blood* 127:325-32, 2016
20. Bench AJ, White HE, Foroni L, et al: Molecular diagnosis of the myeloproliferative neoplasms: UK guidelines for the detection of JAK2 V617F and other relevant mutations. *Br J Haematol* 160:25-34, 2013
21. Aziz A, Baxter EJ, Edwards C, et al: Cooperativity of imprinted genes inactivated by acquired chromosome 20q deletions. *J Clin Invest* 123:2169-82, 2013
22. Dingli D, Grand FH, Mahaffey V, et al: Der(6)t(1;6)(q21-23;p21.3): a specific cytogenetic abnormality in myelofibrosis with myeloid metaplasia. *Br J Haematol* 130:229-32, 2005
23. Stegelmann F, Bullinger L, Griesshammer M, et al: High-resolution single-nucleotide polymorphism array-profiling in myeloproliferative neoplasms identifies novel genomic aberrations. *Haematologica* 95:666-9, 2010
24. Tapper W, Jones AV, Kralovics R, et al: Genetic variation at MECOM, TERT, JAK2 and HBS1L-MYB predisposes to myeloproliferative neoplasms. *Nat Commun* 6:6691, 2015
25. Klampfl T, Harutyunyan A, Berg T, et al: Genome integrity of myeloproliferative neoplasms in chronic phase and during disease progression. *Blood* 118:167-76, 2011
26. Chase A, Leung W, Tapper W, et al: Profound parental bias associated with chromosome 14 acquired uniparental disomy indicates targeting of an imprinted locus. *Leukemia* 29:2069-74, 2015
27. Delhommeau F, Dupont S, Della Valle V, et al: Mutation in TET2 in myeloid cancers. *N Engl J Med* 360:2289-301, 2009
28. Ernst T, Chase AJ, Score J, et al: Inactivating mutations of the histone methyltransferase gene EZH2 in myeloid disorders. *Nat Genet* 42:722-6, 2010
29. Grand FH, Hidalgo-Curtis CE, Ernst T, et al: Frequent CBL mutations associated with 11q acquired uniparental disomy in myeloproliferative neoplasms. *Blood* 113:6182-92, 2009
30. Schischlik F, Milosevic Feenstra, J.D., Rumi, E., Pietra, D., Gisslinger, B., Schalling, M., Bogner, E., Gisslinger, H., Cazzola, M., Kralovics, R.: Fusion Gene Detection Using Whole Transcriptome Analysis in Patients with Chronic Myeloproliferative Neoplasms and Secondary Acute Myeloid Leukemia. *Blood* 126:4093, 2015
31. Hinds DA, Barnholt KE, Mesa RA, et al: Germ line variants predispose to both JAK2 V617F clonal hematopoiesis and myeloproliferative neoplasms. *Blood* 128:1121-8, 2016
32. McKerrell T, Park N, Moreno T, et al: Leukemia-associated somatic mutations drive distinct patterns of age-related clonal hemopoiesis. *Cell Rep* 10:1239-45, 2015

33. Nielsen C, Birgens HS, Nordestgaard BG, et al: The JAK2 V617F somatic mutation, mortality and cancer risk in the general population. *Haematologica* 96:450-3, 2011
34. Roaldsnes C, Holst R, Frederiksen H, et al: Myeloproliferative neoplasms: trends in incidence, prevalence and survival in Norway. *Eur J Haematol*, 2016
35. Xu X, Zhang Q, Luo J, et al: JAK2(V617F): Prevalence in a large Chinese hospital population. *Blood* 109:339-42, 2007
36. Nielsen C, Bojesen SE, Nordestgaard BG, et al: JAK2V617F somatic mutation in the general population: myeloproliferative neoplasm development and progression rate. *Haematologica* 99:1448-55, 2014
37. Busque L, Patel JP, Figueroa ME, et al: Recurrent somatic TET2 mutations in normal elderly individuals with clonal hematopoiesis. *Nat Genet* 44:1179-81, 2012
38. Genovese G, Kahler AK, Handsaker RE, et al: Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *N Engl J Med* 371:2477-87, 2014
39. Jaiswal S, Fontanillas P, Flannick J, et al: Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med* 371:2488-98, 2014
40. Delic S, Rose D, Kern W, et al: Application of an NGS-based 28-gene panel in myeloproliferative neoplasms reveals distinct mutation patterns in essential thrombocythaemia, primary myelofibrosis and polycythaemia vera. *Br J Haematol*, 2016
41. Lundberg P, Karow A, Nienhold R, et al: Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. *Blood* 123:2220-8, 2014
42. Nangalia J, Nice FL, Wedge DC, et al: DNMT3A mutations occur early or late in patients with myeloproliferative neoplasms and mutation order influences phenotype. *Haematologica* 100:e438-42, 2015
43. Ortmann CA, Kent DG, Nangalia J, et al: Effect of mutation order on myeloproliferative neoplasms. *N Engl J Med* 372:601-12, 2015
44. Lundberg P, Nienhold R, Ambrosetti A, et al: Somatic mutations in calreticulin can be found in pedigrees with familial predisposition to myeloproliferative neoplasms. *Blood* 123:2744-5, 2014
45. Antonioli E, Guglielmelli P, Pancrazzi A, et al: Clinical implications of the JAK2 V617F mutation in essential thrombocythemia. *Leukemia* 19:1847-9, 2005
46. Campbell PJ, Scott LM, Buck G, et al: Definition of subtypes of essential thrombocythaemia and relation to polycythaemia vera based on JAK2 V617F mutation status: a prospective study. *Lancet* 366:1945-53, 2005
47. Vannucchi AM, Antonioli E, Guglielmelli P, et al: Characteristics and clinical correlates of MPL 515W>L/K mutation in essential thrombocythemia. *Blood* 112:844-7, 2008
48. Rotunno G, Pacilli A, Artusi V, et al: Epidemiology and clinical relevance of mutations in postpolycythemia vera and postessential thrombocythemia myelofibrosis: A study on 359 patients of the AGIMM group. *Am J Hematol* 91:681-6, 2016

49. Rumi E, Pietra D, Ferretti V, et al: JAK2 or CALR mutation status defines subtypes of essential thrombocythemia with substantially different clinical course and outcomes. *Blood* 123:1544-51, 2014
50. Tefferi A, Guglielmelli P, Larson DR, et al: Long-term survival and blast transformation in molecularly annotated essential thrombocythemia, polycythemia vera, and myelofibrosis. *Blood* 124:2507-13; quiz 2615, 2014
51. Pietra D, Rumi E, Ferretti VV, et al: Differential clinical effects of different mutation subtypes in CALR-mutant myeloproliferative neoplasms. *Leukemia* 30:431-8, 2016
52. Rumi E, Pietra D, Pascutto C, et al: Clinical effect of driver mutations of JAK2, CALR, or MPL in primary myelofibrosis. *Blood* 124:1062-9, 2014
53. Tefferi A, Lasho TL, Finke CM, et al: CALR vs JAK2 vs MPL-mutated or triple-negative myelofibrosis: clinical, cytogenetic and molecular comparisons. *Leukemia* 28:1472-7, 2014
54. Godfrey AL, Chen E, Pagano F, et al: Clonal analyses reveal associations of JAK2V617F homozygosity with hematologic features, age and gender in polycythemia vera and essential thrombocythemia. *Haematologica* 98:718-21, 2013
55. Cabagnols X, Defour JP, Ugo V, et al: Differential association of calreticulin type 1 and type 2 mutations with myelofibrosis and essential thrombocythemia: relevance for disease evolution. *Leukemia* 29:249-52, 2015
56. Sanchez-Aguilera A, Arranz L, Martin-Perez D, et al: Estrogen signaling selectively induces apoptosis of hematopoietic progenitors and myeloid neoplasms without harming steady-state hematopoiesis. *Cell Stem Cell* 15:791-804, 2014
57. Stein BL, Williams DM, Wang NY, et al: Sex differences in the JAK2 V617F allele burden in chronic myeloproliferative disorders. *Haematologica* 95:1090-7, 2010
58. Spivak JL, Considine M, Williams DM, et al: Two clinical phenotypes in polycythemia vera. *N Engl J Med* 371:808-17, 2014
59. Cerquozzi S, Tefferi A: Blast transformation and fibrotic progression in polycythemia vera and essential thrombocythemia: a literature review of incidence and risk factors. *Blood Cancer J* 5:e366, 2015
60. Milosevic JD, Puda A, Malcovati L, et al: Clinical significance of genetic aberrations in secondary acute myeloid leukemia. *Am J Hematol* 87:1010-6, 2012
61. Brecqueville M, Rey J, Bertucci F, et al: Mutation analysis of ASXL1, CBL, DNMT3A, IDH1, IDH2, JAK2, MPL, NF1, SF3B1, SUZ12, and TET2 in myeloproliferative neoplasms. *Genes Chromosomes Cancer* 51:743-55, 2012
62. Guglielmelli P, Biamonte F, Score J, et al: EZH2 mutational status predicts poor survival in myelofibrosis. *Blood* 118:5227-34, 2011
63. Guglielmelli P, Lasho TL, Rotunno G, et al: The number of prognostically detrimental mutations and prognosis in primary myelofibrosis: an international study of 797 patients. *Leukemia*, 2014
64. Tefferi A, Guglielmelli P, Lasho TL, et al: CALR and ASXL1 mutations-based molecular prognostication in primary myelofibrosis: an international study of 570 patients. *Leukemia* 28:1494-500, 2014

65. Tefferi A, Pardanani A, Lim KH, et al: TET2 mutations and their clinical correlates in polycythemia vera, essential thrombocythemia and myelofibrosis. *Leukemia* 23:905-11, 2009
66. Corces-Zimmerman MR, Hong WJ, Weissman IL, et al: Preleukemic mutations in human acute myeloid leukemia affect epigenetic regulators and persist in remission. *Proc Natl Acad Sci U S A* 111:2548-53, 2014
67. Campbell PJ, Baxter EJ, Beer PA, et al: Mutation of JAK2 in the myeloproliferative disorders: timing, clonality studies, cytogenetic associations, and role in leukemic transformation. *Blood* 108:3548-55, 2006
68. Theoharides A, Boissinot M, Girodon F, et al: Leukemic blasts in transformed JAK2-V617F-positive myeloproliferative disorders are frequently negative for the JAK2-V617F mutation. *Blood* 110:375-9, 2007
69. Vannucchi AM, Lasho TL, Guglielmelli P, et al: Mutations and prognosis in primary myelofibrosis. *Leukemia* 27:1861-9, 2013
70. Guglielmelli P, Rotunno G, Fanelli T, Pacilli A, Corbizi-Fattori G, Calabresi L, Pancrazzi A, Vannucchi A.M.: A greater underlying mutational complexity may contribute to the differential prognostic impact of phenotypic-driver mutation in primary myelofibrosis. *EHA21 Congress. Copenhagen, Haematologica*, 2016, pp 8-9
71. Patel KP, Newberry KJ, Luthra R, et al: Correlation of mutation profile and response in patients with myelofibrosis treated with ruxolitinib. *Blood* 126:790-7, 2015
72. Tefferi A, Finke CM, Lasho TL, et al: U2AF1 mutations in primary myelofibrosis are strongly associated with anemia and thrombocytopenia despite clustering with JAK2V617F and normal karyotype. *Leukemia* 28:431-3, 2014
73. Rumi E, Passamonti F, Della Porta MG, et al: Familial chronic myeloproliferative disorders: clinical phenotype and evidence of disease anticipation. *J Clin Oncol* 25:5630-5, 2007
74. Bellanne-Chantelot C, Chaumarel I, Labopin M, et al: Genetic and clinical implications of the Val617Phe JAK2 mutation in 72 families with myeloproliferative disorders. *Blood* 108:346-52, 2006
75. Rumi E, Harutyunyan AS, Pietra D, et al: CALR exon 9 mutations are somatically acquired events in familial cases of essential thrombocythemia or primary myelofibrosis. *Blood* 123:2416-9, 2014
76. Ding J, Komatsu H, Wakita A, et al: Familial essential thrombocythemia associated with a dominant-positive activating mutation of the c-MPL gene, which encodes for the receptor for thrombopoietin. *Blood* 103:4198-200, 2004
77. Kapralova K, Horvathova M, Pecquet C, et al: Cooperation of germ line JAK2 mutations E846D and R1063H in hereditary erythrocytosis with megakaryocytic atypia. *Blood* 128:1418-23, 2016
78. Mead AJ, Chowdhury O, Pecquet C, et al: Impact of isolated germline JAK2V617I mutation on human hematopoiesis. *Blood* 121:4156-65, 2013
79. Jager R, Harutyunyan AS, Rumi E, et al: Common germline variation at the TERT locus contributes to familial clustering of myeloproliferative neoplasms. *Am J Hematol* 89:1107-10, 2014

80. Saliba J, Saint-Martin C, Di Stefano A, et al: Germline duplication of ATG2B and GSKIP predisposes to familial myeloid malignancies. *Nat Genet* 47:1131-40, 2015
81. Harutyunyan AS, Giambruno R, Krendl C, et al: Germline RBBP6 mutations in familial myeloproliferative neoplasms. *Blood* 127:362-5, 2016
82. Landgren O, Goldin LR, Kristinsson SY, et al: Increased risks of polycythemia vera, essential thrombocythemia, and myelofibrosis among 24,577 first-degree relatives of 11,039 patients with myeloproliferative neoplasms in Sweden. *Blood* 112:2199-204, 2008
83. Jones AV, Campbell PJ, Beer PA, et al: The JAK2 46/1 haplotype predisposes to MPL-mutated myeloproliferative neoplasms. *Blood* 115:4517-23, 2010
84. Jones AV, Chase A, Silver RT, et al: JAK2 haplotype is a major risk factor for the development of myeloproliferative neoplasms. *Nat Genet* 41:446-9, 2009
85. Kilpivaara O, Mukherjee S, Schram AM, et al: A germline JAK2 SNP is associated with predisposition to the development of JAK2(V617F)-positive myeloproliferative neoplasms. *Nat Genet* 41:455-9, 2009
86. Oddsson A, Kristinsson SY, Helgason H, et al: The germline sequence variant rs2736100_C in TERT associates with myeloproliferative neoplasms. *Leukemia* 28:1371-4, 2014
87. Olcaydu D, Harutyunyan A, Jager R, et al: A common JAK2 haplotype confers susceptibility to myeloproliferative neoplasms. *Nat Genet* 41:450-4, 2009
88. Maxson JE, Gotlib J, Pollyea DA, et al: Oncogenic CSF3R mutations in chronic neutrophilic leukemia and atypical CML. *N Engl J Med* 368:1781-90, 2013
89. Pardanani A, Lasho TL, Laborde RR, et al: CSF3R T618I is a highly prevalent and specific mutation in chronic neutrophilic leukemia. *Leukemia* 27:1870-3, 2013
90. Wang SA, Hasserjian RP, Fox PS, et al: Atypical chronic myeloid leukemia is clinically distinct from unclassifiable myelodysplastic/myeloproliferative neoplasms. *Blood* 123:2645-51, 2014
91. Elliott MA, Tefferi A: Chronic neutrophilic leukemia 2016: Update on diagnosis, molecular genetics, prognosis, and management. *Am J Hematol* 91:341-9, 2016
92. Druhan LJ, McMahon DP, Steuerwald N, et al: Chronic neutrophilic leukemia in a child with a CSF3R T618I germline mutation. *Blood*, 2016
93. Gotlib J, Maxson JE, George TI, et al: The new genetics of chronic neutrophilic leukemia and atypical CML: implications for diagnosis and treatment. *Blood* 122:1707-11, 2013
94. Wang SA, Tam W, Tsai AG, et al: Targeted next-generation sequencing identifies a subset of idiopathic hypereosinophilic syndrome with features similar to chronic eosinophilic leukemia, not otherwise specified. *Mod Pathol* 29:854-64, 2016
95. Lasho TL, Pardanani A, Tefferi A: LNK mutations in JAK2 mutation-negative erythrocytosis. *N Engl J Med* 363:1189-90, 2010
96. Tefferi A, Lasho TL, Abdel-Wahab O, et al: IDH1 and IDH2 mutation studies in 1473 patients with chronic-, fibrotic- or blast-phase essential thrombocythemia, polycythemia vera or myelofibrosis. *Leukemia* 24:1302-9, 2010

97. Oh ST, Simonds EF, Jones C, et al: Novel mutations in the inhibitory adaptor protein LNK drive JAK-STAT signaling in patients with myeloproliferative neoplasms. *Blood* 116:988-92, 2010

98. Score J, Hidalgo-Curtis C, Jones AV, et al: Inactivation of polycomb repressive complex 2 components in myeloproliferative and myelodysplastic/myeloproliferative neoplasms. *Blood* 119:1208-13, 2012

Table 1. Approximate frequencies of acquired mutations in MPNs

Gene	Genomic location	Mutation location	Protein	Mutation effect	Mutation frequency in MPN, % of cases			
					PV	ET	PMF	Blast phase
<i>Phenotypic/Driver Mutations</i>								
<i>JAK2</i>	9q24	exon 14	JAK2	JAK/STAT signalling	95-97	60	60	50
<i>JAK2</i>	9q24	exon 12	JAK2	JAK/STAT signalling	1-2	0	Rare	nk
<i>MPL</i>	1q34	exon 10	TPOR	JAK/STAT signalling	0	3-5	5-10	nk
<i>CALR</i>	19q13	exon 9	CALR	JAK/STAT signalling	0	25	30	nk
<i>Initiation/Landscaping mutations</i>								
<i>TET2</i>	4q24	all coding regions	TET2	Epigenetic regulation	10-20	5	10-20	20
<i>DNMT3A</i>	2p23	exons 7-23	DNMT3A	Epigenetic regulation	5-10	1-5	8-12	10-20
<i>Disease Modifying/Progression Mutations</i>								
<i>IDH1/2</i>	2q33/15q26	exon 4 exon 4 all coding regions	IDH1/2	Epigenetic regulation	1-2	1-2	5-6	10-30
<i>EZH2</i>	7q35-q36	all coding regions	EZH2	Epigenetic regulation	1-2	1-2	7-10	5
<i>ASXL1</i>	20q11	exon 12	ASXL1	Epigenetic regulation	2	2-5	10-35	20-25
<i>SF3B1</i>	2q33	exons 12-16	SF3B1	mRNA splicing	2	2	5	5
<i>SRSF2</i>	17q25	exon 1	SRSF2	mRNA splicing	Rare	Rare	5-17	15-25
<i>U2AF1</i>	21q22	exons 2-7	U2AF1	mRNA splicing	Rare	Rare	2-8	nk
<i>ZRSR2</i>	Xp22	all coding regions	ZRSR2	mRNA splicing	Rare	Rare	1	nk
<i>RUNX1</i>	21q22	all coding regions	RUNX1	Transcription	nk	nk	nk	10-15

IKZF1	7p12	all coding regions	Ikaros	Transcription	Rare	Rare	Rare	20
TP53	17p13	all coding regions	TP53	DNA repair	Rare	Rare	Rare	25
CUX1	7q22	all coding regions	CUX1	Cell cycle	nk	nk	nk	5
CBL	11q23	exons 8-9	CBL	JAK/STAT signalling	Rare	Rare	5-10	6-9
SH2B3	12q24	exon 2	LNK	JAK/STAT signalling	1-2	3-6	3-6	10

Klampfl et al.²⁵, Delhommeau et al.²⁷, Grand et al.²⁹, Delic et al.⁴⁰, Lundberg et al.⁴¹, Milosevic et al.⁶⁰, Brecqueville et al.⁶¹, Guglielmelli et al.⁶³, Tefferi et al.⁶⁵, Vannucchi et al.⁶⁹, Lasho et al.⁹⁵, Tefferi et al.⁹⁶, Oh et al.⁹⁷, Score et al.⁹⁸.

Figure legends

Figure 1. Phenotypic driver mutations in classic, *BCR-ABL1* negative MPN

Figure 2. Pathways and principal genetic abnormalities in the evolution of *BCR-ABL1* negative MPN




Figure 1




Figure 2