

Phenotypic and genetic aspects of epithelial barrier function in asthmatic patients



Matthew Loxham, PhD, and Donna E. Davies, PhD *Southampton, United Kingdom*

The bronchial epithelium is continuously exposed to a multitude of noxious challenges in inhaled air. Cellular contact with most damaging agents is reduced by the action of the mucociliary apparatus and by formation of a physical barrier that controls passage of ions and macromolecules. In conjunction with these defensive barrier functions, immunomodulatory cross-talk between the bronchial epithelium and tissue-resident immune cells controls the tissue microenvironment and barrier homeostasis. This is achieved by expression of an array of sensors that detect a wide variety of viral, bacterial, and nonmicrobial (toxins and irritants) agents, resulting in production of many different soluble and cell-surface molecules that signal to cells of the immune system. The ability of the bronchial epithelium to control the balance of inhibitory and activating signals is essential for orchestrating appropriate inflammatory and immune responses and for temporally modulating these responses to limit tissue injury and control the resolution of inflammation during tissue repair. In asthmatic patients abnormalities in many aspects of epithelial barrier function have been identified. We postulate that such abnormalities play a causal role in immune dysregulation in the airways by translating gene-environment interactions that underpin disease pathogenesis and exacerbation. (*J Allergy Clin Immunol* 2017;139:1736-51.)

Key words: Asthma, tight junction, innate immunity, cytokine, homeostasis

ASTHMA HETEROGENEITY

Asthma is a common chronic inflammatory disorder of the conducting airways, which undergo distinct structural and functional changes leading to nonspecific bronchial hyperresponsiveness (BHR) and variable airflow obstruction. Recruitment and careful clinical characterization of large cohorts of asthmatic

Abbreviations used

AhR:	Aryl hydrocarbon receptor
AJ:	Adherens junction
BHR:	Bronchial hyperresponsiveness
CDHR3:	Cadherin-related family member 3
DC:	Dendritic cell
DUOX1:	Dual oxidase 1
EGFR:	Epidermal growth factor receptor
eQTL:	Expression quantitative trait locus
GC:	Gene cluster
GST:	Glutathione-S-transferase
GWAS:	Genome-wide association study
ILC:	Innate lymphoid cell
ILC2:	Type 2 innate lymphoid cell
NK:	Natural killer
ORMDL3:	Orosomucoid-like 3
PCDH1:	Protocadherin 1
SC:	Subject cluster
SNP:	Single nucleotide polymorphism
TJ:	Tight junction
TLR:	Toll-like receptor
TSLP:	Thymic stromal lymphopoietin

patients has established beyond doubt that asthma is a heterogeneous disease in terms of phenotype, endotype (ie, underlying pathogenic mechanism), response to treatment, and/or long-term clinical outcomes.¹ Cluster analysis has enabled identification of 4 to 5 phenotypic clusters that have differences in sex, asthma onset, lung function, atopic status, asthma control, health care use, and exacerbation frequency.²⁻⁵ Molecular phenotyping of blood, induced sputum, and epithelial brushings has identified additional heterogeneity, especially in patients with severe asthma,⁶⁻⁹ who are a major economic burden on the health care system because of poor responses to traditional asthma

From Clinical and Experimental Sciences and the Southampton NIHR Respiratory Biomedical Research Unit, University of Southampton Faculty of Medicine, Sir Henry Wellcome Laboratories, University Hospital Southampton.

Supported by the Medical Research Council (MRC [UK]), the National Centre for Reduction Refinement and Replacement of Animals in Research (NC3Rs [UK]), Asthma UK, the Asthma Allergy and Inflammation Research Charity (AAIR), the National Institute for Health Research (NIHR [UK]), and the Biotechnology and Biological Sciences Research Council (BBSRC).

Disclosure of potential conflict of interest: M. Loxham received a fellowship from the Biotechnology and Biological Sciences Research Council (BBSRC) UK. D. E. Davies has received grants from the Medical Research Council UK, National Centre for Reduction, Refinement, and Replacement of Animals in Research (NC3Rs), Asthma Allergy and Inflammation Research Charity, Asthma UK, and the NIHR; has received consulting fees from Synairgen Research; has received travel support from the American Academy of Allergy, Asthma & Immunology and the World Immune Regulation Meeting; has a patent through the University of Southampton; received

royalties from the University of Southampton; is cofounder and shareholder of Synairgen; and is chair of the Trustees of the Asthma Allergy and Inflammation Research Charity (AAIR).

Received for publication February 20, 2017; revised April 13, 2017; accepted for publication April 14, 2017.

Corresponding author: Donna E. Davies, PhD, Clinical and Experimental Sciences, Faculty of Medicine, University of Southampton, Southampton General Hospital, Tremona Road, Southampton SO16 6YD, United Kingdom. E-mail: donnad@soton.ac.uk.

The CrossMark symbol notifies online readers when updates have been made to the article such as errata or minor corrections
0091-6749

© 2017 The Authors. Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

<http://dx.doi.org/10.1016/j.jaci.2017.04.005>

Terms in boldface and italics are defined in the glossary on page 1737.

medications. Some of the differences in asthma clusters might reflect underlying genetic differences; for example, there are differences in genetic risk in early-onset compared with *late-onset asthma*,¹⁰ whereas others might reflect differences in environment and lifestyle or, perhaps most likely, a combination of both gene and environment effects.¹¹ Many, but not all, asthmatic patients have T_H2 inflammation in their airways, and clinical trials with mAbs to IL-5, IL-13, or IL-4 receptor (α chain) have identified a type 2 endotype.¹² Thus patient stratification with type 2 relevant biomarkers has enabled effective targeting of these treatments to subsets of patients with moderate and severe asthma.¹³⁻¹⁷ However, although clinical trials have shown that type 2 inflammation is an important disease modifier in some patients, they have also highlighted that non-type 2 inflammatory pathways must contribute to certain forms of asthma.¹⁸ These can include pathways associated with obesity or neutrophilia or with susceptibility to environmental factors, such as infection and air pollution, but disease mechanisms/endotypes are not well understood. We postulate that a dynamic interaction between a genetically susceptible epithelium and environmental risk factors for asthma is important for the development of asthma and its subphenotypes.¹⁹

BRONCHIAL EPITHELIAL BARRIER STRUCTURE AND FUNCTION

Given the multitude of challenges imposed on the airway epithelium, it is not surprising that it combines structural and

functional protective mechanisms with innate immunologic mechanisms to maintain healthy barrier homeostasis and to minimize inflammation and cellular dysregulation. Structurally, the bronchial epithelium is *pseudostratified*, comprising mainly columnar multiciliated, secretory (goblet), and undifferentiated cells that overlie smaller basal cells with the capacity for self-renewal.²⁰ Rare cell types include pulmonary neuroendocrine cells^{21,22} and brush (tuft) cells²³ that can have neurosensory or chemosensory functions, but information on these cells is limited.

On the epithelial surface, the mucociliary apparatus is a crucial primary innate defense mechanism that protects the lungs from deleterious effects of inhaled pollutants such as noxious gases and *particulate matter*, allergens, and pathogens. Surface epithelial cells and submucosal glands produce secretions comprising a superficial gel or mucus layer and a layer of periciliary fluid that contacts the epithelial surface. Mucus contains hydrated gel-forming mucins and a range of host defense and cytoprotective molecules, including *defensins*, IgA, lactoperoxidase, catalase, superoxide dismutase, and low-molecular-weight antioxidants.²⁴ The viscoelastic properties of the mucus are dictated in large part by the oligomeric secreted mucins MUC5AC and MUC5B,²⁵ multifunctional glycoproteins that provide the structural framework of the mucus barrier. These bronchial secretions shield the epithelial surface, detoxify noxious agents, and trap many inhaled particles, allowing clearance through the action of the mucociliary escalator. MUC5B might also contribute to immune homeostasis by means of direct regulation of leukocyte functions.^{26,27}

GLOSSARY

CIGARETTE SMOKE: Gases, hydrocarbon vapors, and particulate matter generated by burning tobacco. Cigarette smoke contains around 4000 substances, more than 60 of which have been identified as carcinogens. Cigarette smoke promotes increased local elastase production, which contributes to lung tissue injury likely caused by increased epithelial permeability through loss of tight junction integrity.

DEFENSINS: A distinct family of antimicrobial peptides produced by epithelial cells of mucosal surfaces, as well as by neutrophils, natural killer cells, and cytotoxic T lymphocytes. Defensins have direct antimicrobial activity, as well as the ability to activate inflammatory responses.

EPISTASIS: The expression of one gene is influenced by the expression of 1 or more independently inherited (nonallelic) genes.

GLUTATHIONE: An intracellular antioxidant whose principal site of synthesis is the liver. Glutathione serves as a cofactor for glutathione peroxidase, which is responsible for detoxifying lipid peroxides.

INNATE LYMPHOID CELLS: Cells possessing lymphoid morphology but without antigen receptors. Their subpopulations are divided into groups that resemble T helper subsets. Group 2 innate lymphoid cells produce type 2 cytokines, such as IL-4, IL-5, IL-9, and IL-13, on stimulation with epithelium-derived cytokines, such as IL-33, IL-25, and TSLP.

LATE-ONSET ASTHMA: Often defined as asthma that develops in adolescence or adulthood. Most cases of asthma are diagnosed in childhood. Late-onset asthma tends to be more common in women than in men. Occupational asthma and aspirin-exacerbated respiratory disease are subtypes of late-onset asthma. Smoking and passive smoke exposure appear to be risk factors for late-onset asthma.

NOD-LIKE RECEPTORS (NLRs): A family of cytoplasmic receptors that recognize bacterial products, such as bacterial peptidoglycan. NLRs include an N-terminal effector region and a central nucleotide oligomerization domain (NOD), and most have C-terminal leucine-rich

repeats for ligand (PAMP) binding. Once activated, NODs assemble signaling proteins, resulting in nuclear factor κ B and mitogen-activated protein kinase activation, and control the activation of inflammatory caspases.

PARTICULATE MATTER (PM): PM is a term describing airborne dust particles originating from a range of sources and processes, such as fossil fuel combustion, waste incineration, cigarette smoking, and erosion, which can contain black carbon (soot), metals, polyaromatic hydrocarbons, and anions, such as sulfate and nitrate, among others. If sufficiently small to be inhaled (aerodynamic diameter <10 μ m), PM can settle in the airways and exert a range of effects.

PATHOGEN-ASSOCIATED MOLECULAR PATTERN (PAMP): Molecules associated with groups of pathogens that are recognized as “danger signals” by pattern recognition receptors, such as Toll-like receptors or NOD-like receptors. Many types of molecules can serve as PAMPs, including LPSs, peptidoglycans, lipoteichoic acid, nucleic acid variants, and flagellin.

POLYAROMATIC HYDROCARBONS (PAHs): Also known as polycyclic aromatic hydrocarbons, PAHs are volatile substances produced by cooking oils and coal-burning or petroleum products. Indoor biomass burning generates PAHs and is associated with chronic obstructive pulmonary disease development in women.

PSEUDOSTRATIFIED: Although showing features of layering in an epithelium, all cells are still attached to the basement membrane.

TOLL-LIKE RECEPTORS (TLRs): TLRs are a family of innate pattern recognition receptors that respond to a variety of structurally conserved molecules derived from pathogens (ie, PAMPs). They are single, membrane-spanning, noncatalytic receptors whose signaling pathways are finely regulated by Toll/IL-1 receptor homologous region (TIR) domain-containing adaptors. Differential use of these adaptor proteins provides specificity of individual TLR-mediated signaling pathways.




FIG 1. **A**, Schematic representation of a pseudostratified bronchial epithelial cell layer (comprising a goblet cell, 2 ciliated cells, and 2 basal cells) showing the junctional complexes and their interactions with the cytoskeleton or basement membrane to form a robust sheet-like structure. **B**, Illustration of the TJ and AJ complexes showing how they mediate cell-cell contact and interact with the actin cytoskeleton. *ECM*, Extracellular matrix; *JAM*, junctional adhesion molecule; *ZO*, zonula occludens.

In addition to secreting mucus, the bronchial epithelium forms a sheet-like structure that acts as a physical barrier to protect the internal milieu of the tissue. Individual epithelial cells contact each other through a range of cell-cell adhesion complexes (tight junctions [TJs], adherens junctions [AJs], and desmosomes) that control the permeability of the epithelial sheet and link with the cytoskeleton to resist mechanical stress (Fig 1); in addition, gap junctions directly connect the cytoplasm of adjacent cells,

allowing cell-cell communication.²⁸⁻³⁰ The apical-most adhesive complexes are the TJs, which are formed by transmembrane and intracellular proteins that link to the actin cytoskeleton (Fig 1, B).³¹ TJs seal the epithelium, regulating paracellular passage of ions, water, and various macromolecules. They also maintain cell polarity by preventing lateral diffusion and intermixing of molecules in the apical membrane with those in the lateral membrane. Proteins of the TJs include tricellulin and

occludin, which regulate the passage of macromolecules through the TJs,³² and claudins, which are responsible for the size- and charge-selective conductance properties of the TJ paracellular pathway.³³

Expression of barrier or sealing claudins that selectively decrease paracellular cation permeability has been reported in normal human adult lung (claudins 1, 3, 4, 5, 7, and 18),³⁴ and the expression profile varies with anatomic location and function.^{35,36} Claudin-2, a pore-forming claudin, is also detected in the lung, and its presence is thought to increase ionic permeability by acting as a cation-selective pore.³⁶

Located below the TJs are the AJs, which link to the actin cytoskeleton^{37,38}; desmosomes, which link to the intermediate filaments³⁹; and hemidesmosomes,⁴⁰ containing $\alpha_6\beta_4$ integrins that facilitate attachment to the basement membrane (Fig 1, A). AJs and desmosomes are critical for providing the adhesive force to ensure the integrity of the cell layer. Cadherin-catenin complexes comprise the core of the AJs, bridging neighboring cells and the actin-myosin cytoskeleton and contributing to mechanical coupling between cells. In addition to its adhesive function, E-cadherin physically interacts with several receptor tyrosine kinases and affects their signaling abilities. Similarly, β -catenin, which is an integral structural component of the AJs, is also the key nuclear effector of canonical Wnt signaling in the nucleus.⁴¹ This coupling of cell-cell adhesion with signaling functions ensures that AJs can be extremely plastic, allowing the cell to adapt rapidly to its changing environment. Like AJs, the TJ plaque also contains many signaling molecules,^{42,43} allowing proteins involved in cell-cell and cell-matrix adhesion to integrate and coordinate epithelial responses.⁴⁴ Therefore perturbation in the turnover and concentration of junctional proteins is likely to have important implications for the maintenance and stability of the epithelium and the permeability barrier.

Junctional adhesion molecules also serve as sites for interaction of the epithelium with cells involved in immune surveillance. For example, TJ proteins interact directly with dendritic cells (DCs) to allow them to sample the airway lumen without disruption of the epithelial barrier,^{45,46} whereas E-cadherin is a ligand for $\alpha_E\beta_7$ integrin (CD103)-expressed T cells^{47,48} and DCs.⁴⁹ In addition to structural adhesion molecules, the bronchial epithelium expresses inducible adhesion molecules, such as intercellular adhesion molecules 1 and 2, which have essential functions in the clearance of T cells from the lung during resolution of inflammation.⁵⁰

Airway epithelial cells express an array of pattern recognition receptors, including *Toll-like receptors (TLRs)*, *NOD-like receptors*, retinoic acid-inducible gene I (RIG-I)-like receptors, and a variety of natural killer (NK) cell receptor ligands. These enable detection of a wide variety of microbial and nonmicrobial agents, resulting in production of many different soluble and cell-surface molecules, collectively termed the epimune (cytokines, chemokines, damage-associated molecular pattern molecules), and major histocompatibility complex [MHC] gene products,⁵¹ that recruit and activate cells, such as macrophages and neutrophils, involved in inflammation and induction of adaptive immunity. Together, these responses enable many infections to be controlled by the immune system with limited damage to host tissues; however, it is important to note that both innate and adaptive immune-signaling events are involved in mediating tissue damage.⁵² For example, macrophages, neutrophils, and eosinophils release a range of molecules, including

cytotoxic cytokines, cationic proteins, lipid mediators, metalloproteinases, and reactive oxygen species, that induce tissue damage or malfunction. Therefore the ability of the epithelium to control the balance of inhibitory and activating signals is essential not only for initiating an appropriate immune response to environmental challenges, if required (Fig 2), but also for temporally orchestrating these responses to limit tissue injury and control the resolution of inflammatory reactions through cell-surface molecules and release of inhibitory cytokines and lipids during tissue repair.

In vitro and *in vivo* studies have shown that epithelial cells can modulate a variety of immune cells. For example, epithelium-derived TGF- β is chemoactive for *innate lymphoid cells* (ILCs),⁵³ which might provide early defense against pathogens and intervene in repair of damaged tissues. TGF- β secreted by bronchial epithelial cells has a direct inhibitory effect on T-lymphocyte proliferation, and epithelial cell-conditioned T lymphocytes show increased differentiation toward IL-10-producing T_H1 cells.⁵⁴ Epithelial cell secretions also inhibit proinflammatory responses of monocytes, macrophages, and DCs; increase DC expression of the negative regulatory programmed death ligand 1 (CD274); decrease the ability of DCs to induce T-lymphocyte proliferation⁵⁴; and suppress human lung mast cell histamine secretion.⁵⁵ Epithelial cells express CD200, which binds to the inhibitory immune receptor CD200R, which is expressed at high levels on lung macrophages. This not only maintains a strong threshold for response in the context of inhaled nonpathogenic antigens⁵⁶ but also dampens macrophage responses in the context of infection. Thus in CD200 knockout mice there is increased macrophage activity and severe immune-mediated lung damage after influenza infection.⁵⁷

The activation status of NK cells is also controlled by the balance of various inhibitory and activation receptors.^{58,59} For example, the NK cell-activating receptor NKG2D is ligated by molecules, such as MHC class I polypeptide-related sequences A and B or UL16-binding proteins, which are only expressed on stressed airway epithelial cells,^{60,61} resulting in killing of the target cells and ultimately leading to protection from infection. The importance of NK cells and NKG2D in allergic airways responses has been suggested by the findings that mice lacking NKG2D are resistant to induction of allergic inflammation. Although adoptive transfer of wild-type NK cells was able to restore the response, granzyme B-deficient NK cells could not.⁶²

One common link between both infectious and noninfectious triggers of type 2 immunity is that many induce some level of physical trauma that breaches the protective barrier of the body. Tissue damage, at least in the absence of strong type 1-promoting *pathogen-associated molecular pattern* signaling, appears to be a potent mechanism driving type 2 immunity. This involves rapid release of several epithelium-derived cytokine alarmins, such as IL-1, IL-33, thymic stromal lymphopoietin (TSLP), and IL-25, all of which can drive downstream type 2 immunity.⁶³ These cytokines invoke an immune response, involving mast cells, basophils, eosinophils, type 2 innate lymphoid cells (ILC2s), and alternatively activated macrophages, which has evolved to respond to a parasitic infection by generating proinflammatory mediators, toxin-neutralizing enzymes, and helminth-killing toxins, which also have endogenous tissue-damaging properties. A number of studies have identified many environmental agents linked to asthma that have the potential to cause epithelial barrier




FIG 2. Schematic representation of epithelial barrier function illustrating protective and immunoregulatory functions. Under basal conditions, the epithelium maintains homeostasis by limiting exposure of the airway tissue to components of the inhaled environment and by balancing immunoregulatory signals. However, when compromised, the epithelium responds by releasing innate cytokines that help to orchestrate appropriate innate and adaptive immune responses. CS, Cigarette smoke; NO_x, nitrogen oxides; O[·], oxygen radicals; PM, particulate matter; Treg, regulatory T cell.

disruption and tissue injury in the airways, including the house dust mite allergen Der p 1,⁶⁴ fungal allergens,⁶⁵ rhinovirus,⁶⁶ cigarette smoke,^{67,68} and air pollutants.^{69,70}

Nonetheless, a key question arising from these observations is the following: Why are the airways of asthmatic subjects more susceptible than normal to these relatively ubiquitous agents? As detailed below, it is likely that the explanation lies in a combination of (1) decreased epithelial barrier defenses reducing the threshold for epithelial damage, (2) dysregulated innate immune or immunoregulatory responses that contribute to ongoing barrier dysfunction, and (3) impaired epithelial barrier repair, leading to failure to resolve inflammatory responses.

DYSREGULATION OF THE EPITHELIAL BARRIER IN ASTHMATIC PATIENTS

Targeted studies of the bronchial epithelium have demonstrated a range of abnormalities at many levels of barrier function and innate immunity (Fig 3). However, unbiased transcriptomic approaches are now enabling in-depth analysis of epithelial gene expression profiles^{8,9} to provide evidence of molecular mechanisms that might eventually define specific epithelial endotypes of asthma. We will first summarize key abnormalities identified in the epithelial barrier in asthmatic patients and then put these into the context of newer clusters that have been identified and how these relate to genetic susceptibilities.




FIG 3. Schematic representation of the epithelial barrier in asthmatic patients highlighting abnormalities in protective and immunoregulatory functions (gray boxes). Persistent airway inflammation most likely arises as a consequence of impaired barrier defenses (altered cytoprotective secretions and reduced cell-cell adhesion), leading to epithelial susceptibility to injury and dysregulated immune responses. In parallel, impaired repair might contribute to maintenance of epithelial activation and chronicity of responses. The relative contribution of each aspect of barrier dysfunction is likely to influence the overall phenotype of the epithelium and might manifest as distinct subgroups of asthma. CS, Cigarette smoke; NO_x, nitrogen oxides; O₃, oxygen radicals; PM, particulate matter.

The mucociliary apparatus is modified in asthmatic patients, as evidenced by an increase in the number of goblet cells with increased mucin gene expression, an increase in MUC5AC protein relative to MUC5B, and a reduction in ciliated cell numbers.⁷¹⁻⁷³ In addition, decreased ciliary beat frequency, dyskinesia, and ciliary disorientation have been reported in patients with severe asthma.⁷⁴ Together, mucus hypersecretion and ciliary dysfunction in asthmatic patients can result in

stimulation of neural receptors that result in cough⁷⁵ and mucous plugging, which, over time, can lead to severe airflow obstruction.

The increase in expression of MUC5AC relative to MUC5B seen in asthmatic patients has been postulated to affect mucus clearance, reduce eosinophil apoptosis,⁷⁶ and/or contribute to abnormal innate immune responses.⁵⁷ Reprogramming of epithelial differentiation toward a hypersecretory phenotype has been linked to increased expression of the epidermal growth

factor receptor (EGFR)⁷² and to the activity of T_H2 cytokines, including IL-13 and IL-9.^{77,78} Consistent with this, patients with T_H2-high asthma have significantly increased airway mucin gene expression.⁷⁹ T_H2 cytokines also significantly decrease epithelial expression of the antimicrobial peptide human β -defensin 2 *in vitro*, and mice with allergic airway inflammation have significantly more viable bacteria in their lungs after infection.⁸⁰ In contrast, atopic asthmatic patients with type 2–high asthma have been reported to harbor significantly lower bronchial bacterial burden,⁸¹ and in patients with severe asthma, no taxa were associated with a T_H2-related epithelial gene expression signature.⁸² These differences might reflect long-term changes and treatment effects and contrast with the acute responses seen after infection of mice with allergic airways inflammation.⁸⁰

There is considerable evidence for an association between levels of particulate pollutants and asthma exacerbations,^{83–85} asthma pathogenesis, and poorer lung function outcomes.^{86–88} Exposure to air pollutants can lead to oxidative stress in the airways, and there is compelling evidence that asthmatic airways are deficient in antioxidant defenses.⁸⁹ Furthermore, the antioxidant capacity of the lungs is inversely related to asthma severity.⁹⁰ In addition to lower levels of superoxide dismutase and catalase,⁸⁹ it has been shown that goblet cells express the high-affinity sodium ascorbate cotransporter, which is involved in vitamin C uptake into cells, and that expression of sodium ascorbate cotransporters is inversely related to lung lining fluid vitamin C levels.⁹¹ There is also considerable evidence that polymorphisms in *glutathione*-cycling enzymes can result in increased susceptibility to air pollution.^{92–94} Glutathione-S-transferase (GST)-pi is predominantly expressed in airway epithelial cells, and expression is decreased in the airways of children with asthma.⁹⁵ In view of the increased susceptibility of the asthmatic bronchial epithelium to oxidant-induced apoptosis *in vitro*⁹⁶ and the observation that increased levels of oxidants can reduce the anti-inflammatory effects of budesonide, an inability to control oxidative stress might not only drive epithelial damage but also confound treatment responses.⁹⁷

Polyaromatic hydrocarbons are a key toxic component of air pollution. Polyaromatic hydrocarbon levels are increased in the plasma of asthmatic children and linked to a number of asthma markers.⁹⁸ The aryl hydrocarbon receptor (AhR), which plays a key role in detoxification of environmental pollutants, also regulates multiciliogenesis.⁹⁹ Importantly, although air exposure triggers AhR targeting of genes important for multiciliogenesis, toxic AhR ligands induce detoxifying cytochromes, with no overlap in target gene induction. These mutually exclusive responses suggest a potential pathophysiologic mechanism whereby AhR ligands in air pollutants disrupt AhR-mediated ciliogenesis to contribute to disruption of barrier defenses in asthmatic patients.⁹⁹

Epithelial fragility¹⁰⁰ and epithelial shedding¹⁰¹ in asthmatic patients have been recognized for many years, but this remains a controversial area.¹⁰² Nonetheless, through use of specific markers of response to injury, such as increased expression of EGFR, epithelial damage has been confirmed in bronchial biopsy specimens from asthmatic adults¹⁰³ and children.¹⁰⁴ Many studies have reported disruption of adhesive mechanisms in asthmatic patients, including loss of TJ proteins,^{67,105,106} reduction in AJ proteins,¹⁰⁵ and reduction in desmosome length.¹⁰⁷ Membrane

expression of caveolin-1, a stabilizer of AJs, is significantly lower in airway epithelia of asthmatic patients, and *in vitro* loss of caveolin-1 causes loss of junctional E-cadherin and β -catenin expression and disrupted epithelial barrier function.¹⁰⁸ Consistent with reduced adhesion, functional studies comparing epithelial cultures from asthmatic or healthy donors indicate that there is increased permeability and sensitivity to environmental stressors in asthmatic patients⁶⁷ and increased susceptibility to oxidant stress.⁹⁶ Increased barrier permeability might not only promote allergic sensitization but also reduce the threshold for epithelial damage and activation of a type 2 response, which itself might affect barrier function. Thus, in addition to their effects on goblet cell differentiation, T_H2 cytokines have a disruptive effect on epithelial barrier function¹⁰⁹ and lead to a distinct profile of epithelial gene expression, both *in vitro* and in T_H2-high asthmatic patients *in vivo*.⁷⁹

Claudin-18, a lung-specific barrier claudin, has been shown to be expressed in bronchial epithelium, and its levels are reduced in asthmatic patients, being lowest in patients with T_H2-high asthma.¹⁰⁶ In the same studies IL-13 downregulated claudin-18 *in vitro*, and targeted knockdown of claudin-18 increased epithelial permeability. Furthermore, claudin-18–null mice had significantly higher serum IgE levels and increased airway responsiveness after intranasal *Aspergillus* species sensitization, suggesting loss of claudin-18 can promote sensitization and airway hyperresponsiveness.¹⁰⁶

Because mast cells are important sources of IL-13 and are in close proximity to the bronchial epithelium in asthma,¹¹⁰ it is noteworthy that IL-33–activated mast cells, as well as ILC2s, are able to drive a predominantly IL-13–regulated pattern of gene expression in normal human bronchial epithelial cells *in vitro*.¹¹¹ Furthermore, ILC2s have been shown to directly impair epithelial barrier integrity through IL-13,¹¹² whereas T_H2 cells cause barrier leakiness through IL-4 and IL-13, an effect that can be prevented by inhibition of histone deacetylases.¹¹³

Consistent with the evidence of epithelial disruption in asthmatic patients, levels of epithelium-derived cytokine alarmins, such as IL-33, TSLP, and IL-25, are increased in asthmatic patients.^{114,115} IL-33, a member of the IL-1 cytokine family, has gained prominence in type 2 immunity by virtue of the genetic association of both *IL33* and its receptor, *IL1RL1* (ST2), with asthma^{10,116} and by its functional effects on ILC2 cells, T_H2 cells, mast cells, basophils, and alternatively activated macrophages.¹¹⁷ IL-33 is normally localized in the nucleus, where it is a transcriptional regulator¹¹⁸ and can act as an extracellular cytokine by binding to its receptor, ST2.¹¹⁹ Full-length IL-33 binds ST2 and is biologically active, although activity can be increased after cleavage by inflammatory proteases,¹²⁰ whereas caspase cleavage leads to inactivation.¹²¹ IL-33 can be released by nonprogrammed cell death, or it can be actively secreted through vesicular transport from the Golgi complex.¹²² Stimulation of bronchial epithelial cells with allergen or ATP results in active release of IL-33, which depends on the NADPH oxidase dual oxidase 1 (DUOX1)–mediated activation of Src and EGFR signaling through cysteine oxidation.¹²³ Nasal epithelial cells from asthmatic patients display enhanced DUOX1 expression, as well as allergen-induced IL-33 secretion, compared with healthy control subjects, suggesting that increased expression and activation of DUOX1 might be an important feature of enhanced IL-33 secretion in asthmatic patients.¹²³ In addition to full-length IL-33,

alternative splicing of the IL-33 transcript can result in deletion of exons 3 and 4 (Δ exon 3,4) to confer cytoplasmic localization and facilitate extracellular secretion without cell death, while retaining signaling capacity. Analyses of epithelial brush RNA suggest that Δ exon 3,4 is strongly associated with airway type 2 inflammation, whereas full-length IL-33 is not.¹²⁴ These results suggest that therapeutic IL-33 inhibitors will need to block all biologically active isoforms.

TSLP is an IL-7–like cytokine that can trigger DC-mediated T_H2 inflammatory responses¹²⁵ and T_H2 cytokine production by mast cells.¹²⁶ A variety of stimuli, including double-stranded RNA and allergens, stimulate TSLP expression in bronchial epithelial cells, and this is enhanced by inflammatory cytokines.¹²⁷ Challenge of cultured epithelial cells from asthmatic donors with double-stranded RNA results in a skewed response favoring more TSLP and less type 1 interferon compared with healthy cells.¹²⁸ Allergen-specific T cells also enhance TSLP production by epithelial cells from asthmatic donors, suggesting that T cell–airway epithelium interactions can lead to maintenance and amplification of allergic inflammation.¹²⁹ In a double-blind, placebo-controlled study, treatment with a human mAb to TSLP resolved airway inflammation and attenuated allergen-induced bronchoconstriction, findings consistent with TSLP as a therapeutic target in patients with allergic asthma.¹³⁰ However, in addition to its effects on immune cells, it is noteworthy that TSLP drives an IL-13–dependent increase in bronchial epithelial cell proliferation¹³¹ and increases TJ expression to enhance nasal epithelial barrier function, suggesting a role for TSLP in restoration of epithelial barrier integrity.¹³² In contrast, TSLP has been reported to disrupt TJs in 16HBE bronchial epithelial cells.¹³³ Furthermore, a short and constitutively expressed form of TSLP has been detected in the skin and gut; this variant cannot activate signal transducer and activator of transcription (STAT) 5 but has potent antimicrobial activity.¹³⁴ Recent studies suggest that the short and constitutively expressed form of TSLP can protect against bronchial epithelial barrier disruption *in vitro* and house dust mite– or toluene diisocyanate–induced airway inflammation *in vivo*.^{133,135} Consequently, optimal therapeutic antibody targeting might need to be directed specifically to the long form of TSLP.

IL-25 belongs to the IL-17 cytokine family and is secreted by T_H2 cells, mast cells, basophils, and eosinophils, as well as epithelial cells.¹³⁶ It can drive airway remodeling in allergic models of airway inflammation,¹³⁷ and, in combination with IL-33, can promote the development of ILC2s, which appear critical in early initiation of the T_H2 response.¹³⁸ Expression of IL-25 has been reported to be increased in epithelial cells from patients with asthma and can be induced further by rhinovirus infections.¹³⁹ Others have found increased systemic levels of IL-25 in subgroups of patients with T_H2 -high asthma.¹⁴⁰ Furthermore, the IL-25 receptor (IL-17RB) is upregulated on myeloid and plasmacytoid DCs in blood and sputum 24 hours after allergen challenge.¹⁴¹ IL-25 upregulated TLR9 expression by plasmacytoid DCs and orchestrated the responses to TLR9 ligation, suggesting that IL-25 can act as a link between the adaptive and innate immune responses.¹⁴¹

Viral respiratory tract infections, especially rhinovirus infection, are the main triggers of asthma exacerbations.^{142,143} Several,^{144–146} but not all,^{147,148} studies have shown that bronchial epithelial cells from asthmatic donors respond abnormally to rhinovirus infection involving an insufficiency of IFN- β and

IFN- λ . This has been linked to increased TGF- β 2 production by epithelial cells from asthmatic patients¹⁴⁹ and suppression of cytokine signaling expression¹⁵⁰; however, it is also of interest that rhinovirus-induced EGFR activation can suppress IFN- λ production and increase viral infection.¹⁵¹ The importance of decreased antiviral immunity in asthmatic patients has been tested in a clinical trial with inhaled IFN- β : the drug was found to improve asthma control and reduce exacerbations in patients with difficult-to-treat asthma.¹⁵²

It is well known that mechanical forces are critical to lung development and that abnormal mechanical stresses can lead to pathologic lung injury.¹⁵³ In asthmatic patients constriction of bronchial smooth muscle during an acute asthma attack causes the airway wall to buckle, resulting in folding and compression of the bronchial epithelium.¹⁵³ *In vitro* studies have shown that airway epithelial cells respond rapidly and robustly to compressive stress with changes in goblet cell numbers and production of profibrogenic growth factors.^{154,155} The relevance of these findings has been demonstrated *in vivo*, where induction of bronchoconstriction with methacholine caused airway remodeling involving goblet cell metaplasia and subepithelial fibrosis without evidence of inflammation.¹⁵⁶ Although these changes might simply be due to the hyperresponsive properties of bronchial smooth muscle in asthmatic patients, there is evidence that bronchial epithelial cells from asthmatic donors respond abnormally to compression with increased release of TGF- β and GM-CSF,¹⁵⁷ suggesting that bronchoconstriction can skew epithelial innate immune responses in asthmatic patients. Because the asthma susceptibility gene a disintegrin and metalloprotease 33 (*ADAM33*) has been linked to BHR¹⁵⁸ and has been shown to cause bronchial smooth muscle contraction,¹⁵⁹ there is the potential for multifactorial indirect genetic effects on epithelial barrier function.

Increased expression of the EGFR in bronchial biopsy specimens from asthmatic adults¹⁰³ and children¹⁰⁴ is consistent with an ongoing response to injury, and this is highly correlated with epithelial IL-8 expression.¹⁶⁰ However, expression of the cyclin-dependent kinase inhibitor p21^{waf} might be indicative of impaired proliferation or ongoing epithelial stress in asthmatic patients.^{104,161} During epithelial repair, neighboring epithelial cells become migratory in response to growth factors, such as TGF- β or epidermal growth factor. This repair phenotype is characterized by downregulation of TJs and increased expression of matrix metalloproteases and extracellular matrix components, as observed in asthmatic patients. Studies with cultures of epithelial cells from asthmatic children suggest that the airway epithelium displays a dysregulated repair response, taking longer to repair mechanically induced wounds¹⁶² and undergoing a more extensive epithelial-mesenchymal transition in response to TGF- β than cultures from nonasthmatic donors.¹⁶³ Recently, it has been reported that IL-22 can promote a repair phenotype in the presence of TGF- β 1, causing a marked reduction in E-cadherin but only in cells obtained from donors with severe asthma.¹⁶⁴

EPITHELIAL CLUSTERS AND ASTHMA HETEROGENEITY

Use of large-scale transcriptomic approaches in large cohorts of well-characterized asthmatic and healthy control volunteers has enabled unbiased in-depth analysis of gene expression profiles in epithelial brushings and allowed clustering into distinct

Gene cluster	Subject Cluster	SC1	SC2	SC3	SC4	SC5
% mod to severe asthma		16	73	74	50	43
GC1 (Innate immunity/antibacterial function; Cell proliferation/apoptosis; Lymphocyte activation/migration)		Red	Blue	Red	Red	Red
GC2 Cilia structure/function, other		Red	Blue	Red	Red	Red
GC3TNF- α ; Muscle		Blue	Red	Red	Red	Blue
GC4 Notch signalling; Neuronal function; Dystrophin family; WNT family, Ion channels; Other		Red	Blue	Blue	Red	Red
GC5 'no obvious function'		Red	Red	Blue	Red	Red
GC6 Microtubules; Mitochondrial; Actin related; Neuronal; Other		Blue	Red	Red	Red	Red
GC7 Interferons; Apoptosis; P38 related; Keratins; Sialyl Lewis antigen; Cell matrix interactions; Other		Blue	Red	Red	Red	Blue
GC8 Cysteine metabolism; Mucins; Mast cells; Vasoconstrictors (possibly MC); Glycolipid antigen presentation; Other		Blue	Red	Blue	Blue	Red
GC9; mitochondria; Intracellular trafficking; O-linked glycosylation; N-linked glycosylation; "Type 2 genes"; Other		Blue	Red	Red	Red	Red




FIG 4. Pictorial representation of the SCs and GCs found in a transcriptomic analysis of epithelial brushings from 155 donors. Red indicates high, pink indicates medium, and blue indicates low expression of genes within the cluster. The bar chart indicates the percentage of healthy control subjects and patients with mild, moderate, or severe asthma in each SC, and the width of the bar is proportional to the number of subjects in the cluster. Findings are summarized from Modena et al.⁸

phenotypes. Analysis of transcriptomic data from 155 donors in combination with exhaled nitric oxide has identified 5 molecularly defined and clinically distinct subject clusters (SCs) with distinct expression of gene clusters (GCs),⁸ as summarized in Fig 4. The majority (73%) of all healthy control subjects were located in SC1, which was distinguished by high expression of GCs involved in processes including “innate immunity/antibacterial function” and “Notch signaling” and low expression of GCs, including “interferons/stress” and “type 2 immunity.” In contrast, the largest group of patients with severe asthma (SC2) showed a diametrically opposite pattern with low expression of both “innate immunity/antibacterial function” and “Notch signaling” GCs and high expression of “interferons/stress” and “type 2 immunity” GCs. In addition, “cilia structure and function” was low in SC2 with severe asthma. It is interesting to note an apparent paradox that gene signatures for both cilia-related genes and Notch signaling are reduced in SC2. Because Notch signaling inhibits ciliated cell differentiation *in vitro* by repressing multicilin and forkhead box J1,¹⁶⁵ low Notch levels might suggest increased ciliogenesis, but this was not the case.

However, it has been shown that IL-13 inhibits ciliated cell differentiation independent of Notch signaling,¹⁶⁶ suggesting 2 distinct signaling pathways can affect ciliated cell differentiation, which might be of relevance in the different SCs of severe asthma. The other SCs showed some overlap with SC2, but each exhibited distinct profiles illustrating the heterogeneity of the epithelial gene signature across the spectrum of asthma severity. Further analysis of the same data using weighted gene coexpression network analysis (WGCNA) highlighted that genes in modules linked to epithelial growth and repair and neuronal function were markedly decreased in patients with severe asthma.⁹ Of particular note, low expression of epithelial growth and repair and neuronal function genes was more strongly associated with severe asthma than type 2 inflammation, suggesting that epithelial integrity and related processes are of primary importance to the development of asthma and severe asthma.

Assuming that these phenotypes are stable rather than fluctuations because of disease activity, these data illustrate the complexity of the epithelial phenotype. Reinforcement of these findings with longitudinal studies should provide a basis for

hypothesis-driven research that allows precise definition of epithelial endotypes in asthmatic patients. Nonetheless, based on the evidence to date, further consideration of strategies that promote epithelial repair and restore epithelial homeostasis might provide novel therapeutic approaches for the treatment of asthma.²⁴ For example, the protective effects of growth factors, such as epidermal growth factor, have been recognized for many years (reviewed by Swindle et al²⁴). However, novel strategies include potential use of the macrolide antibiotic azithromycin, which has been shown to decrease ionic permeability of human airway epithelia by changing the processing of TJ proteins,¹⁶⁷ or histone deacetylase inhibition with JNJ-26481585, which has been shown to ameliorate the effects of T_H2 cells on barrier function.¹¹³

FROM ASTHMA GENES TO FUNCTION

Genome-wide association studies (GWASs) of asthma have identified novel risk alleles and loci, with many of the asthma susceptibility genes being expressed in the airway epithelium.¹⁶⁸ Among susceptibility factors for asthma, the genes *IL1RL1*/*IL18R1*, *IL33*, and *TSLP* have emerged as some of the most important associated with development of the disease,¹⁰ linking epithelium-derived cytokines to type 2 inflammation. Furthermore, a number of genes associated with epithelial homeostasis, differentiation, or barrier immunity have been identified, including protocadherin 1 (*PCDH1*),¹⁶⁹ cadherin-related family member 3 (*CDHR3*),¹⁷⁰ *HLA-DQ*,¹⁰ *SPINK5*,¹⁷¹ *GPRA*,¹⁷² and orosomucoid-like 3 (*ORMDL3*)/*GSDMB*¹⁰ at the 17q12-21 locus. However, it should be noted that asthma-associated alleles have small effect sizes and account for little of the prevalence of asthma, and it is likely that a significant portion of the genetic risk for asthma and its exacerbations results from genotype-specific responses to environmental exposures, including allergens, pollution, and viral infections, especially at particular stages of life.¹⁷³⁻¹⁷⁷ Here we have attempted to place some of the asthma susceptibility genes in the context of epithelial barrier dysregulation, with a view to highlighting potential epithelial endotypes of disease linked to reduced barrier defenses, dysregulated immune responses, and/or abnormal repair responses (Fig 5).

Epidemiologic and genetic evidence have implicated epithelial susceptibility to environmental insults in asthma pathogenesis. However, clear functional relationships are not always easy to identify, perhaps reflecting the need for assessment in the context of an appropriate environmental trigger. For example, although 2 common deletion polymorphisms of the GST genes *GSTM1* and *GSTT1* and the *GSTP1* Ile105Val polymorphism have been associated with asthma in children and adults, a meta-analysis has revealed extreme between-study heterogeneity,¹⁷⁸ suggesting more focused study in the context of environmental oxidative exposures would be more informative.

Genes, such as the cadherin family members *CDHR3*¹⁷⁰ and *PCDH1*,¹⁶⁹ appear to play roles in adhesion. Several single nucleotide polymorphisms (SNPs) in *PCDH1* have been linked to asthma and BHR. These include Ala750Ala and IVS3_116, which are localized in the 3' untranslated region of exon 3 and might affect mRNA stability or splicing, whereas Ala514Thr is localized in the fifth cadherin repeat of the extracellular domain and can affect cell-cell adhesion¹⁶⁹; however, the functional consequences of this mutation have not been explored. Protocadherin 1 (*PCDH1*) colocalizes with E-cadherin in airway epithelial cells,

and it has been implicated in the barrier-enhancing properties of glucocorticoids¹⁷⁹ and suppression of TGF- β signaling.¹⁸⁰ Because gene-passive smoking interactions have been found to be relevant for the association of *PCDH1* with asthma,^{169,181} the contribution of *PCDH1* gene variants to asthma might only become evident in the context of smoke exposure.¹⁸² *CDHR3* was originally identified as an asthma susceptibility gene linked to childhood exacerbation.¹⁷⁰ The asthma-associated SNP (rs6967330) causes a nonsynonymous mutation (G>A; C529Y) in the fifth cadherin repeat of *CDHR3*, which affects cellular localization.¹⁷⁰ Subsequent studies showed that *CDHR3* is a receptor for rhinovirus C, suggesting that the increased localization of Y529 *CDHR3* on the bronchial epithelial cell surface increases susceptibility for rhinovirus C infection and replication.¹⁸³ However, the normal cellular function of *CDHR3* is still unknown.

ORMDL3 has been shown to be associated with early-onset asthma susceptibility in multiple independent genome-wide and candidate-gene association studies.¹⁷³ It is regulated by STAT6 and can be induced by IL-13 or IL-4,¹⁸⁴ and SNPs in *ORMDL3* correlate with changes in T_H2 cytokine levels.¹⁸⁵ *ORMDL3* is found in the endoplasmic reticulum and is involved in maintaining sphingolipid homeostasis and in the unfolded protein response,¹⁸⁶ but *in vitro* studies involving underexpression or overexpression of *ORMDL3* did not show a significant role in modulating innate immune responses and the unfolded protein response.¹⁸⁷ However, in mice overexpression of *ORMDL3* decreases serum sphingolipid levels and increases inflammatory markers, airway remodeling, and BHR in response to allergic stimuli.¹⁸⁸ Furthermore, pulmonary epithelial expression of *ORMDL3* is sufficient for induction of *Alternaria* species-induced allergic airways disease.¹⁸⁹

As already described, polymorphisms in genes, including *IL33*, *IL1RL1*, and *TSLP*, have been linked to epithelial activation/damage and type 2 immunity, although detailed studies are still revealing new levels of complexity involving alternative splicing.¹²⁴ In the case of *TSLP*, multiple SNPs are correlated with the expression levels of *TSLP*, and some alleles are protective.¹⁹⁰ Of note, in subjects with 1 or more *SPINK5* risk alleles, the absence of the *TSLP* protective minor alleles has been associated with a significant increase in asthma.¹⁹¹ Thus, in addition to gene-environment effects, *epistasis* adds another level of complexity to asthma pathogenesis. Other immune regulators might be relevant to exacerbation-prone asthma: these include suppressor of cytokine signaling 1 (*SOC1*)¹⁹² and IL-1 receptor-associated kinase M (*IRAK-M*),¹⁹³ both of which suppress IFN- β signaling and antiviral responses.^{150,194}

The focus on epithelial repair genes in asthmatic patients has been limited to date, but promoter variants in *TGFB1* and *TGFB2*, which increase TGF- β expression, are associated with asthma^{195,196} and airflow obstruction.¹⁹⁷ It is also interesting to note that genes, such as hedgehog interacting protein (*HHIP*) and patched homolog 1 (*PTCH1*), which might play a role in epithelial repair have been identified through genetic association with reduced lung function,¹⁹⁸ suggesting that impaired repair might drive extracellular matrix deposition and tissue remodeling.

Most of the asthma-associated SNPs identified by using GWASs are not coding-change variants. Therefore expression quantitative trait loci (eQTLs) analysis has been adopted to identify functional SNPs regulating expression levels of disease-associated genes in a cell type-specific fashion. Applying this




FIG 5. Potential mechanisms of asthma defined by epithelial barrier dysfunction. Identification of potential links with asthma susceptibility genes and their interaction with environmental stimuli are shown.

analysis to bronchial epithelial cells has revealed SNPs in *TSLP*, *GSDMB*, *IL33*, *HLA-DQB1*, *C11orf30*, *DEXI*, *CDHR3*, and *ZBTB10* that affect asthma risk by allowing *cis*-regulation of its gene expression in an epithelial specific manner.¹⁹⁰ In the case of *IL33*, all asthma-associated SNPs in this region of the genome are located in the 5' or first intron of *IL33*, and eQTL analysis has revealed that SNPs in the promoter region of *IL33* are correlated with *IL-33* expression in bronchial epithelial cells. The same study identified an eQTL SNP for *CDHR3* (rs17152490) in bronchial epithelial cells, which is in linkage

disequilibrium with the GWAS SNP (rs6967330, G>A; C529Y), suggesting *cis*-regulation of *CDHR3* expression can also contribute to the asthma risk. SNPs in pituitary tumor-transforming 1 interacting protein (*PTTG1IP*) and mastermind-like 3 (*MAML3*) have been reported to be associated with BHR severity in adult asthma,¹⁹⁹ and eQTL analyses indicate higher tissue expression with less severe BHR. These gene products might be particularly relevant to epithelial repair because *PTTG1IP* is coexpressed with vimentin and E-cadherin 1, whereas *MAML3* is coexpressed with *MAML2*, both of which

are involved in Notch signaling, a repair pathway that was deficient in the transcriptomic studies of severe asthma.

CONCLUDING COMMENTS

Taken together, the evidence for epithelial dysregulation in asthmatic patients is compelling. Genomic studies have revealed the extent of epithelial heterogeneity in asthmatic patients and have provided considerable insight into expression profiles, pathways, and processes that can drive epithelial dysfunction. Further understanding of asthma endotypes will come from integration of findings from these large data sets with the function and regulation of asthma genes and how these are modified by interaction with environmental factors, including the airway microbiome. However, the stability of the asthma phenotypes identified in molecular studies still needs to be addressed in longitudinal studies. In addition, the appreciation that changes in gene expression are also evident in epithelial cells harvested from peripheral airways of patients with severe asthma raises new questions about gene dysregulation in the smaller airways, which comprise the majority of the airway surface area, and the need for better-targeted therapies for the peripheral airways.²⁰⁰ Furthermore, there is a lack of critical information about epithelial heterogeneity and its role in childhood asthma. Crucially, we still lack detailed information about the functions of many asthma genes and how genetic polymorphism of these genes drives asthma susceptibility. The high costs of transgenic and gene-deletion mouse models has restricted progress in this area. Thus it would be timely to investigate the potential of nonmammalian models, such as *Drosophila* species or zebrafish, as tools to investigate gene function because the genetic tractability and low cost of rearing these organisms are major advantages.^{201,202} Better understanding of epithelial dysfunction and its interrelationship with airways inflammation and structural remodeling should help to define specific epithelial endotypes in asthmatic patients. Through development and use of therapeutic approaches that restore epithelial barrier homeostasis, it might be possible to prevent or modify the disease course by intervening close to the origin of the disease.

REFERENCES

- Muraro A, Lemanske RF Jr, Hellings PW, Akdis CA, Bieber T, Casale TB, et al. Precision medicine in patients with allergic diseases: airway diseases and atopic dermatitis—PRACTALL document of the European Academy of Allergy and Clinical Immunology and the American Academy of Allergy, Asthma & Immunology. *J Allergy Clin Immunol* 2016;137:1347-58.
- Moore WC, Meyers DA, Wenzel SE, Teague WG, Li H, Li X, et al. Identification of asthma phenotypes using cluster analysis in the Severe Asthma Research Program. *Am J Respir Crit Care Med* 2010;181:315-23.
- Loza MJ, Adcock I, Auffray C, Chung KF, Djukanovic R, Sterk PJ, et al. Longitudinally stable, clinically defined clusters of patients with asthma independently identified in the ADEPT and U-BIOPRED asthma studies. *Ann Am Thorac Soc* 2016;13(suppl 1):S102-3.
- Chang TS, Lemanske RF Jr, Mauger DT, Fitzpatrick AM, Sorkness CA, Szefer SJ, et al. Childhood asthma clusters and response to therapy in clinical trials. *J Allergy Clin Immunol* 2014;133:363-9.
- Denlinger LC, Phillips BR, Ramratnam S, Ross K, Bhakta NR, Cardet JC, et al. Inflammatory and co-morbid features of patients with severe asthma and frequent exacerbations. *Am J Respir Crit Care Med* 2017;195:302-13.
- Bigler J, Boedigheimer M, Schofield JP, Skipp PJ, Corfield J, Rowe A, et al. A severe asthma disease signature from gene expression profiling of peripheral blood from U-BIOPRED cohorts. *Am J Respir Crit Care Med* 2016 [Epub ahead of print].
- Lefauieux D, De MB, Loza MJ, Peffer N, Rowe A, Baribaud F, et al. U-BIOPRED clinical adult asthma clusters linked to a subset of sputum omics. *J Allergy Clin Immunol* 2016 [Epub ahead of print].
- Modena BD, Tedrow JR, Milosevic J, Bleecker ER, Meyers DA, Wu W, et al. Gene expression in relation to exhaled nitric oxide identifies novel asthma phenotypes with unique biomolecular pathways. *Am J Respir Crit Care Med* 2014;190:1363-72.
- Modena BD, Bleecker ER, Busse WW, Erzurum SC, Gaston BM, Jarjour NN, et al. Gene expression correlated to severe asthma characteristics reveals heterogeneous mechanisms of severe disease. *Am J Respir Crit Care Med* 2016 [Epub ahead of print].
- Moffatt MF, Gut IG, Demenais F, Strachan DP, Bouzigon E, Heath S, et al. A large-scale, consortium-based genomewide association study of asthma. *N Engl J Med* 2010;363:1211-21.
- Vawda S, Mansour R, Takeda A, Funnell P, Kerry S, Mudway I, et al. Associations between inflammatory and immune response genes and adverse respiratory outcomes following exposure to outdoor air pollution: a HuGE systematic review. *Am J Epidemiol* 2014;179:432-42.
- Fajt ML, Wenzel SE. Asthma phenotypes and the use of biologic medications in asthma and allergic disease: the next steps toward personalized care. *J Allergy Clin Immunol* 2015;135:299-310.
- Corren J, Busse W, Meltzer EO, Mansfield L, Bensch G, Fahrenholz J, et al. A randomized, controlled, phase 2 study of AMG 317, an IL-4/RA1pha antagonist, in patients with asthma. *Am J Respir Crit Care Med* 2010;181:788-96.
- Corren J, Lemanske RF, Hanania NA, Korenblat PE, Parsey MV, Arron JR, et al. Lebrikizumab treatment in adults with asthma. *N Engl J Med* 2011;365:1088-98.
- Piper E, Brightling C, Niven R, Oh C, Faggioni R, Poon K, et al. A phase II placebo-controlled study of tralokinumab in moderate-to-severe asthma. *Eur Respir J* 2013;41:330-8.
- Pavord ID, Korn S, Howarth P, Bleecker ER, Buhl R, Keene ON, et al. Mepolizumab for severe eosinophilic asthma (DREAM): a multicentre, double-blind, placebo-controlled trial. *Lancet* 2012;380:651-9.
- Haldar P, Brightling CE, Hargadon B, Gupta S, Monteiro W, Sousa A, et al. Mepolizumab and exacerbations of refractory eosinophilic asthma. *N Engl J Med* 2009;360:973-84.
- Fahy JV. Type 2 inflammation in asthma—present in most, absent in many. *Nat Rev Immunol* 2015;15:57-65.
- Davies DE, Wicks J, Powell RM, Puddicombe SM, Holgate ST. Airway remodelling in asthma—new insights. *J Allergy Clin Immunol* 2003;111:215-25.
- Rock JR, Randell SH, Hogan BL. Airway basal stem cells: a perspective on their roles in epithelial homeostasis and remodeling. *Dis Model Mech* 2010;3:545-56.
- Reynolds SD, Giangreco A, Power JH, Stripp BR. Neuroepithelial bodies of pulmonary airways serve as a reservoir of progenitor cells capable of epithelial regeneration. *Am J Pathol* 2000;156:269-78.
- Gosney JR. Pulmonary neuroendocrine cell system in pediatric and adult lung disease. *Microsc Res Tech* 1997;37:107-13.
- Reid L, Meyrick B, Antony VB, Chang LY, Crapo JD, Reynolds HY. The mysterious pulmonary brush cell: a cell in search of a function. *Am J Respir Crit Care Med* 2005;172:136-9.
- Swindle EJ, Collins JE, Davies DE. Breakdown in epithelial barrier function in patients with asthma: identification of novel therapeutic approaches. *J Allergy Clin Immunol* 2009;124:23-34.
- Thornton DJ, Rousseau K, McGuckin MA. Structure and function of the polymeric mucins in airways mucus. *Annu Rev Physiol* 2008;70:459-86.
- Roy MG, Livraghi-Butrico A, Fletcher AA, McElwee MM, Evans SE, Boerner RM, et al. Muc5b is required for airway defence. *Nature* 2014;505:412-6.
- Janssen WJ, Stefanski AL, Bochner BS, Evans CM. Control of lung defence by mucins and macrophages: ancient defence mechanisms with modern functions. *Eur Respir J* 2016;48:1201-14.
- Kast JJ, Wanke K, Soyka MB, Wawrzyniak P, Akdis D, Kingo K, et al. The broad spectrum of interepithelial junctions in skin and lung. *J Allergy Clin Immunol* 2012;130:544-7.
- Rezaee F, Georas SN. Breaking barriers. New insights into airway epithelial barrier function in health and disease. *Am J Respir Cell Mol Biol* 2014;50:857-69.
- Georas SN, Rezaee F. Epithelial barrier function: at the front line of asthma immunology and allergic airway inflammation. *J Allergy Clin Immunol* 2014;134:509-20.
- Tsukita S, Furuse M, Itoh M. Multifunctional strands in tight junctions. *Nat Rev Mol Cell Biol* 2001;2:285-93.
- Krug SM, Schulzke JD, Fromm M. Tight junction, selective permeability, and related diseases. *Semin Cell Dev Biol* 2014;36:166-76.
- Krause G, Winkler L, Mueller SL, Haseloff RF, Piontek J, Blasig IE. Structure and function of claudins. *Biochim Biophys Acta* 2008;1778:631-45.

34. Schlingmann B, Molina SA, Koval M. Claudins: gatekeepers of lung epithelial function. *Semin Cell Dev Biol* 2015;42:47-57.
35. Niimi T, Nagashima K, Ward JM, Minoo P, Zimonjic DB, Popescu NC, et al. claudin-18, a novel downstream target gene for the T/EBP/NKX2.1 homeodomain transcription factor, encodes lung- and stomach-specific isoforms through alternative splicing. *Mol Cell Biol* 2001;21:7380-90.
36. Kaarteenaho-Wiik R, Soini Y. Claudin-1, -2, -3, -4, -5, and -7 in usual interstitial pneumonia and sarcoidosis. *J Histochem Cytochem* 2009;57:187-95.
37. Ivanov AI, Naydenov NG. Dynamics and regulation of epithelial adherens junctions: recent discoveries and controversies. *Int Rev Cell Mol Biol* 2013;303:27-99.
38. Nelson WJ. Regulation of cell-cell adhesion by the cadherin-catenin complex. *Biochem Soc Trans* 2008;36:149-55.
39. Garrod D, Chidgey M. Desmosome structure, composition and function. *Biochim Biophys Acta* 2008;1778:572-87.
40. Nievers MG, Schaapveld RQ, Sonnenberg A. Biology and function of hemidesmosomes. *Matrix Biol* 1999;18:5-17.
41. Gangl K, Reininger R, Bernhard D, Campana R, Pree I, Reisinger J, et al. Cigarette smoke facilitates allergen penetration across respiratory epithelium. *Allergy* 2009;64:398-405.
42. Gonzalez-Mariscal L, Tapia R, Chamorro D. Crosstalk of tight junction components with signaling pathways. *Biochim Biophys Acta* 2008;1778:729-56.
43. Balda MS, Matter K. Tight junctions and the regulation of gene expression. *Biochim Biophys Acta* 2009;1788:761-7.
44. Tsukita S, Yamazaki Y, Katsuno T, Tamura A, Tsukita S. Tight junction-based epithelial microenvironment and cell proliferation. *Oncogene* 2008;27:6930-8.
45. Veres TZ, Voedisch S, Spies E, Tschernig T, Braun A. Spatiotemporal and functional behavior of airway dendritic cells visualized by two-photon microscopy. *Am J Pathol* 2011;179:603-9.
46. Blank F, Wehrli M, Lehmann A, Baum O, Gehr P, Von GC, et al. Macrophages and dendritic cells express tight junction proteins and exchange particles in an in vitro model of the human airway wall. *Immunobiology* 2011;216:86-95.
47. Pauls K, Schon M, Kubitz RC, Homey B, Wiesenborn A, Lehmann P, et al. Role of integrin alphaE(CD103)beta7 for tissue-specific epidermal localization of CD8+ T lymphocytes. *J Invest Dermatol* 2001;117:569-75.
48. Cepek KL, Shaw SK, Parker CM, Russell GJ, Morrow JS, Rimm DL, et al. Adhesion between epithelial cells and T lymphocytes mediated by E-cadherin and the alpha E beta 7 integrin. *Nature* 1994;372:190-3.
49. Kim TH, Lee HK. Differential roles of lung dendritic cell subsets against respiratory virus infection. *Immune Netw* 2014;14:128-37.
50. Porter JC, Hall A. Epithelial ICAM-1 and ICAM-2 regulate the egression of human T cells across the bronchial epithelium. *FASEB J* 2009;23:492-502.
51. Swamy M, Jamora C, Havran W, Hayday A. Epithelial decision makers: in search of the "epimune." *Nat Immunol* 2010;11:656-65.
52. Rouse BT, Sehrawat S. Immunity and immunopathology to viruses: what decides the outcome? *Nat Rev Immunol* 2010;10:514-26.
53. Denney L, Byrne AJ, Shea TJ, Buckley JS, Pease JE, Herledan GM, et al. Pulmonary epithelial cell-derived cytokine TGF-beta1 is a critical cofactor for enhanced innate lymphoid cell function. *Immunity* 2015;43:945-58.
54. Mayer AK, Bartz H, Fey F, Schmidt LM, Dalpke AH. Airway epithelial cells modify immune responses by inducing an anti-inflammatory microenvironment. *Eur J Immunol* 2008;38:1689-99.
55. Martin N, Ruddick A, Arthur GK, Wan H, Woodman L, Brightling CE, et al. Primary human airway epithelial cell-dependent inhibition of human lung mast cell degranulation. *PLoS One* 2012;7:e43545.
56. Gwyer FE, Hussell T. Macrophage-mediated inflammation and disease: a focus on the lung. *Mediators Inflamm* 2012;2012:140937.
57. Snelgrove RJ, Goulding J, Didierlaurent AM, Lyonga D, Vekaria S, Edwards L, et al. A critical function for CD200 in lung immune homeostasis and the severity of influenza infection. *Nat Immunol* 2008;9:1074-83.
58. Vivier E, Nunes JA, Vely F. Natural killer cell signaling pathways. *Science* 2004;306:1517-9.
59. Bryceson YT, March ME, Ljunggren HG, Long EO. Activation, coactivation, and costimulation of resting human natural killer cells. *Immunol Rev* 2006;214:73-91.
60. Obeidy P, Sharland AF. NKG2D and its ligands. *Int J Biochem Cell Biol* 2009;41:2364-7.
61. Borchers MT, Harris NL, Wesselkamper SC, Vitucci M, Cosman D. NKG2D ligands are expressed on stressed human airway epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 2006;291:L222-31.
62. Farhadi N, Lambert L, Triulzi C, Openshaw PJ, Guerra N, Culley FJ. Natural killer cell NKG2D and granzyme B are critical for allergic pulmonary inflammation. *J Allergy Clin Immunol* 2014;133:827-35.
63. Hammad H, Lambrecht BN. Barrier epithelial cells and the control of type 2 immunity. *Immunity* 2015;43:29-40.
64. Wan H, Winton HL, Soeller C, Tovey ER, Gruenert DC, Thompson PJ, et al. Derp 1 facilitates transepithelial allergen delivery by disruption of tight junctions. *J Clin Invest* 1999;104:123-33.
65. Tai HY, Tam MF, Chou H, Peng HJ, Su SN, Perng DW, et al. Pen h 13 allergen induces secretion of mediators and degradation of occludin protein of human lung epithelial cells. *Allergy* 2006;61:382-8.
66. Sajjan U, Wang Q, Zhao Y, Gruenert DC, Hershenson MB. Rhinovirus disrupts the barrier function of polarized airway epithelial cells. *Am J Respir Crit Care Med* 2008;178:1271-81.
67. Xiao C, Puddicombe SM, Field S, Haywood J, Broughton-Head V, Puxeddu I, et al. Defective epithelial barrier function in asthma. *J Allergy Clin Immunol* 2011;128:549-56.
68. Olivera DS, Boggs SE, Beenhouwer C, Aden J, Knall C. Cellular mechanisms of mainstream cigarette smoke-induced lung epithelial tight junction permeability changes in vitro. *Inhal Toxicol* 2007;19:13-22.
69. London NR Jr, Tharakan A, Rule AM, Lane AP, Biswal S, Ramanathan M Jr. Air pollutant-mediated disruption of sinonasal epithelial cell barrier function is reversed by activation of the Nrf2 pathway. *J Allergy Clin Immunol* 2016;138:1736-8.
70. Ghio AJ, Devlin RB. Inflammatory lung injury after bronchial instillation of air pollution particles. *Am J Respir Crit Care Med* 2001;164:704-8.
71. Fahy JV. Goblet cell and mucin gene abnormalities in asthma. *Chest* 2002;122(suppl):320S-6S.
72. Takeyama K, Fahy JV, Nadel JA. Relationship of epidermal growth factor receptors to goblet cell production in human bronchi. *Am J Respir Crit Care Med* 2001;163:511-6.
73. Lachowicz-Scroggins ME, Yuan S, Kerr SC, Dunican EM, Yu M, Carrington SD, et al. Abnormalities in MUC5AC and MUC5B Protein in Airway Mucus in Asthma. *Am J Respir Crit Care Med* 2016;194:1296-9.
74. Thomas B, Rutman A, Hirst RA, Haldar P, Wardlaw AJ, Bankart J, et al. Ciliary dysfunction and ultrastructural abnormalities are features of severe asthma. *J Allergy Clin Immunol* 2010;126:722-9.
75. Nadel JA. Mucous hypersecretion and relationship to cough. *Pulm Pharmacol Ther* 2013;26:510-3.
76. Kiwamoto T, Katoh T, Evans CM, Janssen WJ, Brummet ME, Hudson SA, et al. Endogenous airway mucins carry glycans that bind Siglec-F and induce eosinophil apoptosis. *J Allergy Clin Immunol* 2015;135:1329-40.
77. Kondo M, Tamaoki J, Takeyama K, Nakata J, Nagai A. Interleukin-13 induces goblet cell differentiation in primary cell culture from Guinea pig tracheal epithelium. *Am J Respir Cell Mol Biol* 2002;27:536-41.
78. Vermeer PD, Harson R, Einwalter LA, Moninger T, Zabner J. Interleukin-9 induces goblet cell hyperplasia during repair of human airway epithelia. *Am J Respir Cell Mol Biol* 2003;28:286-95.
79. Woodruff PG, Modrek B, Choy DF, Jia G, Abbas AR, Ellwanger A, et al. T-helper type 2-driven inflammation defines major subphenotypes of asthma. *Am J Respir Crit Care Med* 2009;180:388-95.
80. Beisswenger C, Kandler K, Hess C, Garn H, Felgentreff K, Wegmann M, et al. Allergic airway inflammation inhibits pulmonary antibacterial host defense. *J Immunol* 2006;177:1833-7.
81. Durack J, Lynch SV, Nariya S, Bhakta NR, Beigelman A, Castro M, et al. Features of the bronchial bacterial microbiome associated with atopy, asthma, and responsiveness to inhaled corticosteroid treatment. *J Allergy Clin Immunol* 2016 [Epub ahead of print].
82. Huang YJ, Nariya S, Harris JM, Lynch SV, Choy DF, Arron JR, et al. The airway microbiome in patients with severe asthma: associations with disease features and severity. *J Allergy Clin Immunol* 2015;136:874-84.
83. Schwartz J, Slater D, Larson TV, Pierson WE, Koenig JQ. Particulate air pollution and hospital emergency room visits for asthma in Seattle. *Am Rev Respir Dis* 1993;147:826-31.
84. Norris G, YoungPong SN, Koenig JQ, Larson TV, Sheppard L, Stout JW. An association between fine particles and asthma emergency department visits for children in Seattle. *Environ Health Perspect* 1999;107:489-93.
85. Malig BJ, Green S, Basu R, Broadwin R. Coarse particles and respiratory emergency department visits in California. *Am J Epidemiol* 2013;178:58-69.
86. Clark NA, Demers PA, Karr CJ, Koehoorn M, Lencar C, Tamburic L, et al. Effect of early life exposure to air pollution on development of childhood asthma. *Environ Health Perspect* 2010;118:284-90.
87. McConnell R, Islam T, Shankardass K, Jerrett M, Lurmann F, Gilliland F, et al. Childhood incident asthma and traffic-related air pollution at home and school. *Environ Health Perspect* 2010;118:1021-6.
88. Young MT, Sandler DP, DeRoo LA, Vedral S, Kaufman JD, London SJ. Ambient air pollution exposure and incident adult asthma in a nationwide cohort of U.S. women. *Am J Respir Crit Care Med* 2014;190:914-21.

89. Erzurum SC. New insights in oxidant biology in asthma. *Ann Am Thorac Soc* 2016;13(suppl 1):S35-9.
90. Ahmad A, Shameem M, Husain Q. Relation of oxidant-antioxidant imbalance with disease progression in patients with asthma. *Ann Thorac Med* 2012;7:226-32.
91. Larsson N, Rankin GD, Bicer EM, Roos-Engstrand E, Pourazar J, Blomberg A, et al. Identification of vitamin C transporters in the human airways: a cross-sectional in vivo study. *BMJ Open* 2015;5:e006979.
92. Amaral AF, Ramasamy A, Castro-Giner F, Minelli C, Accordini S, Sorheim IC, et al. Interaction between gas cooking and GSTM1 null genotype in bronchial responsiveness: results from the European Community Respiratory Health Survey. *Thorax* 2014;69:558-64.
93. Bowatte G, Lodge CJ, Perret JL, Matheson MC, Dharmage SC. Interactions of GST polymorphisms in air pollution exposure and respiratory diseases and allergies. *Curr Allergy Asthma Rep* 2016;16:85.
94. Bowatte G, Lodge CJ, Knibbs LD, Lowe AJ, Erbas B, Dennekamp M, et al. Traffic-related air pollution exposure is associated with allergic sensitization, asthma, and poor lung function in middle age. *J Allergy Clin Immunol* 2017;139:122-9.
95. Schroer KT, Gibson AM, Sivaprasad U, Bass SA, Ericksen MB, Wills-Karp M, et al. Downregulation of glutathione S-transferase pi in asthma contributes to enhanced oxidative stress. *J Allergy Clin Immunol* 2011;128:539-48.
96. Buccheri F, Puddicombe SM, Lordan JL, Richter A, Buchanan D, Wilson SJ, et al. Asthmatic bronchial epithelium is more susceptible to oxidant-induced apoptosis. *Am J Respir Cell Mol Biol* 2002;27:179-85.
97. Heijink I, van OA, Kliphuis N, Jonker M, Hoffmann R, Telenga E, et al. Oxidant-induced corticosteroid unresponsiveness in human bronchial epithelial cells. *Thorax* 2014;69:5-13.
98. Al-Daghri NM, Alokail MS, Abd-Alrahman SH, Draz HM, Yakout SM, Clerici M. Polycyclic aromatic hydrocarbon exposure and pediatric asthma in children: a case-control study. *Environ Health* 2013;12:1.
99. Villa M, Crotta S, Dingwell KS, Hirst EM, Gialitakis M, Ahlfors H, et al. The aryl hydrocarbon receptor controls cyclin O to promote epithelial multiciliogenesis. *Nat Commun* 2016;7:12652.
100. Naylor B. The shedding of the mucosa of the bronchial tree in asthma. *Thorax* 1962;17:69-72.
101. Laitinen LA, Heino M, Laitinen A, Kava T, Haahtela T. Damage of the airway epithelium and bronchial reactivity in patients with asthma. *Am Rev Respir Dis* 1985;131:599-606.
102. Ordonez C, Ferrando R, Hyde DM, Wong HH, Fahy JV. Epithelial desquamation in asthma: artifact or pathology? *Am J Respir Crit Care Med* 2000;162:2324-9.
103. Puddicombe SM, Polosa R, Richter A, Krishnam MT, Howarth PH, Holgate ST, et al. Involvement of the epidermal growth factor receptor in epithelial repair in asthma. *FASEB J* 2000;14:1362-74.
104. Fedorov IA, Wilson SJ, Davies DE, Holgate ST. Epithelial stress and structural remodelling in childhood asthma. *Thorax* 2005;60:389-94.
105. de Boer WI, Sharma HS, Baelemans SM, Hoogsteden HC, Lambrecht BN, Braunstahl GJ. Altered expression of epithelial junctional proteins in atopic asthma: possible role in inflammation. *Can J Physiol Pharmacol* 2008;86:105-12.
106. Sweerus K, Lachowicz-Scroggins M, Gordon E, LaFemina M, Huang X, Parikh M, et al. Claudin-18 deficiency is associated with airway epithelial barrier dysfunction and asthma. *J Allergy Clin Immunol* 2017;139:72-81.
107. Shahana S, Bjornsson E, Ludviksdottir D, Janson C, Nettelbladt O, Venge P, et al. Ultrastructure of bronchial biopsies from patients with allergic and non-allergic asthma. *Respir Med* 2005;99:429-43.
108. Hackett TL, de Bruin HG, Shaheen F, Van Den Berge M, van Oosterhout AJ, Postma DS, et al. Caveolin-1 controls airway epithelial barrier function. Implications for asthma. *Am J Respir Cell Mol Biol* 2013;49:662-71.
109. Saatian B, Rezaee F, Desando S, Emo J, Chapman T, Knowlden S, et al. Interleukin-4 and interleukin-13 cause barrier dysfunction in human airway epithelial cells. *Tissue Barriers* 2013;1:e24333.
110. Bradding P, Walls AF, Holgate ST. The role of the mast cell in the pathophysiology of asthma. *J Allergy Clin Immunol* 2006;117:1277-84.
111. Nagarkar DR, Ramirez-Carrozzi V, Choy DF, Lee K, Soriano R, Jia G, et al. IL-13 mediates IL-33-dependent mast cell and type 2 innate lymphoid cell effects on bronchial epithelial cells. *J Allergy Clin Immunol* 2015;136:202-5.
112. Sugita K, Steer CA, Martinez-Gonzalez I, Altunbulakli C, Morita H, Castro-Giner F, et al. Type 2 innate lymphoid cells disrupt bronchial epithelial barrier integrity by targeting tight junctions via IL-13 in asthma. *J Allergy Clin Immunol* 2017 [Epub ahead of print].
113. Wawrzyniak P, Wawrzyniak M, Wanke K, Sokolowska M, Bendelja K, Ruckert B, et al. Regulation of bronchial epithelial barrier integrity by type 2 cytokines and histone deacetylases in asthmatic patients. *J Allergy Clin Immunol* 2017;139:93-103.
114. Prefontaine D, Nadigel J, Chouiali F, Audusseau S, Semlali A, Chakir J, et al. Increased IL-33 expression by epithelial cells in bronchial asthma. *J Allergy Clin Immunol* 2010;125:752-4.
115. Shikotra A, Choy DF, Ohri CM, Doran E, Butler C, Hargadon B, et al. Increased expression of immunoreactive thymic stromal lymphopoietin in patients with severe asthma. *J Allergy Clin Immunol* 2012;129:104-11.
116. Torgerson DG, Ampleford EJ, Chiu GY, Gauderman WJ, Gignoux CR, Graves PE, et al. Meta-analysis of genome-wide association studies of asthma in ethnically diverse North American populations. *Nat Genet* 2011;43:887-92.
117. Garlanda C, Dinarello CA, Mantovani A. The interleukin-1 family: back to the future. *Immunity* 2013;39:1003-18.
118. Carriere V, Roussel L, Ortega N, Lacorre DA, Americh L, Aguilar L, et al. IL-33, the IL-1-like cytokine ligand for ST2 receptor, is a chromatin-associated nuclear factor in vivo. *Proc Natl Acad Sci U S A* 2007;104:282-7.
119. Smith DE. The biological paths of IL-1 family members IL-18 and IL-33. *J Leukoc Biol* 2011;89:383-92.
120. Lefrancais E, Roga S, Gautier V, Gonzalez-de-Peredo A, Monsarrat B, Girard JP, et al. IL-33 is processed into mature bioactive forms by neutrophil elastase and cathepsin G. *Proc Natl Acad Sci U S A* 2012;109:1673-8.
121. Cayrol C, Girard JP. The IL-1-like cytokine IL-33 is inactivated after maturation by caspase-1. *Proc Natl Acad Sci U S A* 2009;106:9021-6.
122. Daniels MJ, Brough D. Unconventional pathways of secretion contribute to inflammation. *Int J Mol Sci* 2017;18.
123. Hristova M, Habibovic A, Veith C, Janssen-Heininger YM, Dixon AE, Geiszt M, et al. Airway epithelial dual oxidase 1 mediates allergen-induced IL-33 secretion and activation of type 2 immune responses. *J Allergy Clin Immunol* 2016;137:1545-56.
124. Gordon ED, Simpson LJ, Rios CL, Ringel L, Lachowicz-Scroggins ME, Peters MC, et al. Alternative splicing of interleukin-33 and type 2 inflammation in asthma. *Proc Natl Acad Sci U S A* 2016;113:8765-70.
125. Wang WL, Li HY, Zhang MS, Gao PS, He SH, Zheng T, et al. Thymic stromal lymphopoietin: a promising therapeutic target for allergic diseases. *Int Arch Allergy Immunol* 2013;160:18-26.
126. Nagarkar DR, Poposki JA, Comeau MR, Biyasheva A, Avila PC, Schleimer RP, et al. Airway epithelial cells activate TH2 cytokine production in mast cells through IL-1 and thymic stromal lymphopoietin. *J Allergy Clin Immunol* 2012;130:225-32.
127. Hallstrand TS, Hackett TL, Altemeier WA, Matute-Bello G, Hansbro PM, Knight DA. Airway epithelial regulation of pulmonary immune homeostasis and inflammation. *Clin Immunol* 2014;151:1-15.
128. Uller L, Bedke N, Sammut D, Green NG, Lau L, Howarth PH, et al. Disproportionate thymic stromal lymphopoietin versus interferon-beta production by asthmatic bronchial epithelial cells challenged with double-stranded RNA. *Thorax* 2010;65:626-32.
129. Hui CC, Murphy DM, Neighbour H, Al-Sayegh M, O'Byrne S, Thong B, et al. T cell-mediated induction of thymic stromal lymphopoietin in differentiated human primary bronchial epithelial cells. *Clin Exp Allergy* 2014;44:953-64.
130. Gauvreau GM, O'Byrne PM, Boulet LP, Wang Y, Cockcroft D, Bigler J, et al. Effects of an anti-TSLP antibody on allergen-induced asthmatic responses. *N Engl J Med* 2014;370:2102-10.
131. Semlali A, Jacques E, Koussih L, Gounni AS, Chakir J. Thymic stromal lymphopoietin-induced human asthmatic airway epithelial cell proliferation through an IL-13-dependent pathway. *J Allergy Clin Immunol* 2010;125:844-50.
132. Kojima T, Go M, Takano K, Kurose M, Ohkuni T, Koizumi J, et al. Regulation of tight junctions in upper airway epithelium. *Biomed Res Int* 2013;2013:947072.
133. Dong H, Hu Y, Liu L, Zou M, Huang C, Luo L, et al. Distinct roles of short and long thymic stromal lymphopoietin isoforms in house dust mite-induced asthmatic airway epithelial barrier disruption. *Sci Rep* 2016;6:39559.
134. Bjerkas L, Sonesson A, Schenck K. Multiple functions of the new cytokine-based antimicrobial peptide thymic stromal lymphopoietin (TSLP). *Basel: Pharmaceuticals*; 2016:9.
135. Wang Y, Le Y, Zhao W, Lin Y, Wu Y, Yu C, et al. Short TSLP attenuates toluene diisocyanate (TDI)-induced airway inflammation and inhibits HMGB1-RAGE and long TSLP expression. *Toxicol Sci* 2017 [Epub ahead of print].
136. Angkasekwinai P, Park H, Wang YH, Wang YH, Chang SH, Corry DB, et al. Interleukin 25 promotes the initiation of proallergic type 2 responses. *J Exp Med* 2007;204:1509-17.
137. Gregory LG, Jones CP, Walker SA, Sawant D, Gowers KH, Campbell GA, et al. IL-25 drives remodelling in allergic airways disease induced by house dust mite. *Thorax* 2013;68:82-90.

138. Saenz SA, Noti M, Artis D. Innate immune cell populations function as initiators and effectors in Th2 cytokine responses. *Trends Immunol* 2010;31:407-13.
139. Beale J, Jayaraman A, Jackson DJ, Macintyre JD, Edwards MR, Walton RP, et al. Rhinovirus-induced IL-25 in asthma exacerbation drives type 2 immunity and allergic pulmonary inflammation. *Sci Transl Med* 2014;6:256ra134.
140. Cheng D, Xue Z, Yi L, Shi H, Zhang K, Huo X, et al. Epithelial interleukin-25 is a key mediator in Th2-high, corticosteroid-responsive asthma. *Am J Respir Crit Care Med* 2014;190:639-48.
141. Tworek D, Smith SG, Salter BM, Baatjes AJ, Scime T, Watson R, et al. IL-25 receptor expression on airway dendritic cells after allergen challenge in subjects with asthma. *Am J Respir Crit Care Med* 2016;193:957-64.
142. Johnston SL, Pattemore PK, Sanderson G, Smith S, Lampe F, Josephs L, et al. Community study of role of viral infections in exacerbations of asthma in 9-11 year old children. *BMJ* 1995;310:1225-9.
143. Gern JE. Rhinovirus respiratory infections and asthma. *Am J Med* 2002;112(suppl 6A):19S-27S.
144. Wark PA, Johnston SL, Bucchieri F, Powell R, Puddicombe S, Laza-Stanca V, et al. Asthmatic bronchial epithelial cells have a deficient innate immune response to infection with rhinovirus. *J Exp Med* 2005;201:937-47.
145. Contoli M, Message SD, Laza-Stanca V, Edwards MR, Wark PA, Bartlett NW, et al. Role of deficient type III interferon-lambda production in asthma exacerbations. *Nat Med* 2006;12:1023-6.
146. Kicic A, Stevens PT, Sutanto EN, Kicic-Starcevic E, Ling KM, Looi K, et al. Impaired airway epithelial cell responses from children with asthma to rhinoviral infection. *Clin Exp Allergy* 2016;46:1441-55.
147. Lopez-Souza N, Favoreto S, Wong H, Ward T, Yagi S, Schnurr D, et al. In vitro susceptibility to rhinovirus infection is greater for bronchial than for nasal airway epithelial cells in human subjects. *J Allergy Clin Immunol* 2009;123:1384-90.
148. Bochkov YA, Hanson KM, Keles S, Brockman-Schneider RA, Jarjour NN, Gern JE. Rhinovirus-induced modulation of gene expression in bronchial epithelial cells from subjects with asthma. *Mucosal Immunol* 2010;3:69-80.
149. Bedke N, Sammut D, Green B, Kehagia V, Dennison P, Jenkins G, et al. Transforming growth factor-beta promotes rhinovirus replication in bronchial epithelial cells by suppressing the innate immune response. *PLoS One* 2012;7:e44580.
150. Gielen V, Sykes A, Zhu J, Chan B, Macintyre J, Regamey N, et al. Increased nuclear suppressor of cytokine signaling 1 in asthmatic bronchial epithelium suppresses rhinovirus induction of innate interferons. *J Allergy Clin Immunol* 2015;136:177-88.
151. Ueki IF, Min-Oo G, Kalinowski A, Ballon-Landa E, Lanier LL, Nadel JA, et al. Respiratory virus-induced EGFR activation suppresses IRF1-dependent interferon lambda and antiviral defense in airway epithelium. *J Exp Med* 2013;210:1929-36.
152. Djukanovic R, Harrison T, Johnston SL, Gabbay F, Wark P, Thomson NC, et al. The effect of inhaled interferon-beta on worsening of asthma symptoms caused by viral infections: a randomised trial. *Am J Respir Crit Care Med* 2014;190:145-54.
153. Park JA, Fredberg JJ, Drazen JM. Putting the squeeze on airway epithelia. *Physiology* (Bethesda) 2015;30:293-303.
154. Tschumperlin DJ, Shively JD, Kikuchi T, Drazen JM. Mechanical stress triggers selective release of fibrotic mediators from bronchial epithelium. *Am J Respir Cell Mol Biol* 2003;28:142-9.
155. Park JA, Tschumperlin DJ. Chronic Intermittent Mechanical Stress Increases MU-C5AC Protein Expression. *Am J Respir Cell Mol Biol* 2009;41:459-66.
156. Grainge CL, Lau LC, Ward JA, Dulay V, Lahiff G, Wilson S, et al. Effect of bronchoconstriction on airway remodeling in asthma. *N Engl J Med* 2011;364:2006-15.
157. Grainge C, Dennison P, Lau L, Davies D, Howarth P. Asthmatic and normal respiratory epithelial cells respond differently to mechanical apical stress. *Am J Respir Crit Care Med* 2014;190:477-80.
158. Van Eerdewegh P, Little RD, Dupuis J, Del Mastro RG, Falls K, Simon J, et al. Association of the ADAM-33 gene with asthma and bronchial hyper-responsiveness. *Nature* 2002;418:426-30.
159. Duan Y, Long J, Chen J, Jiang X, Zhu J, Jin Y, et al. Overexpression of soluble ADAM33 promotes a hypercontractile phenotype of the airway smooth muscle cell in rat. *Exp Cell Res* 2016;349:109-18.
160. Hamilton LM, Torres-Lozano C, Puddicombe SM, Richter A, Kimber I, Dearman RJ, et al. The role of the epidermal growth factor receptor in sustaining neutrophil inflammation in severe asthma. *Clin Exp Allergy* 2003;33:233-40.
161. Puddicombe SM, Torres-Lozano C, Richter A, Bucchieri F, Lordan JL, Howarth PH, et al. Increased expression of p21(waf) cyclin dependent kinase inhibitor in asthmatic bronchial epithelium. *Am J Respir Cell Mol Biol* 2003;28:61-8.
162. Stevens PT, Kicic A, Sutanto EN, Knight DA, Stick SM. Dysregulated repair in asthmatic paediatric airway epithelial cells: the role of plasminogen activator inhibitor-1. *Clin Exp Allergy* 2008;38:1901-10.
163. Hackett TL, Warner SM, Stefanowicz D, Shaheen F, Pechkovsky DV, Murray LA, et al. Induction of epithelial-mesenchymal transition in primary airway epithelial cells from patients with asthma by transforming growth factor-beta1. *Am J Respir Crit Care Med* 2009;180:122-33.
164. Johnson JR, Nishioka M, Chakir J, Risse PA, Almaghlouth I, Bazarbashi AN, et al. IL-22 contributes to TGF-beta1-mediated epithelial-mesenchymal transition in asthmatic bronchial epithelial cells. *Respir Res* 2013;14:118.
165. Gervac BJ, Valencia M, Baumlin N, Salathe M, Conner GE, Fregien NL. Submersion and hypoxia inhibit ciliated cell differentiation in a notch-dependent manner. *Am J Respir Cell Mol Biol* 2014;51:516-25.
166. Gervac BJ, Fregien NL. IL-13 inhibits multicilin expression and ciliogenesis via janus kinase/signal transducer and activator of transcription independently of Notch cleavage. *Am J Respir Cell Mol Biol* 2016;54:554-61.
167. Asgrimsson V, Gudjonsson T, Gudmundsson GH, Baldursson O. Novel effects of azithromycin on tight junction proteins in human airway epithelia. *Antimicrob Agents Chemother* 2006;50:1805-12.
168. Zhang Y, Moffatt MF, Cookson WO. Genetic and genomic approaches to asthma: new insights for the origins. *Curr Opin Pulm Med* 2012;18:6-13.
169. Koppelman GH, Meyers DA, Howard TD, Zheng SL, Hawkins GA, Ampleford EJ, et al. Identification of PCDH1 as a novel susceptibility gene for bronchial hyperresponsiveness. *Am J Respir Crit Care Med* 2009;180:929-35.
170. Bonnelykke K, Sleiman P, Nielsen K, Kreiner-Moller E, Mercader JM, Belgrave D, et al. A genome-wide association study identifies CDHR3 as a susceptibility locus for early childhood asthma with severe exacerbations. *Nat Genet* 2014;46:51-5.
171. Meng JF, Rosenwasser LJ. Unraveling the genetic basis of asthma and allergic diseases. *Allergy Asthma Immunol Res* 2010;2:215-27.
172. Laitinen T, Polvi A, Rydman P, Vendelin J, Pulkkinen V, Salmikangas P, et al. Characterization of a common susceptibility locus for asthma-related traits. *Science* 2004;304:300-4.
173. Ober C. Asthma Genetics in the Post-GWAS Era. *Ann Am Thorac Soc* 2016;13(suppl 1):S85-90.
174. Moheimani F, Hsu AC, Reid AT, Williams T, Kicic A, Stick SM, et al. The genetic and epigenetic landscapes of the epithelium in asthma. *Respir Res* 2016;17:119.
175. Gaval ML, Bertics PJ, Gern JE. Rhinoviruses, allergic inflammation, and asthma. *Immunol Rev* 2011;242:69-90.
176. Kelly FJ, Fussell JC. Air pollution and airway disease. *Clin Exp Allergy* 2011;41:1059-71.
177. Jackson DJ, Evans MD, Gangnon RE, Tisler CJ, Pappas TE, Lee WM, et al. Evidence for a causal relationship between allergic sensitization and rhinovirus wheezing in early life. *Am J Respir Crit Care Med* 2012;185:281-5.
178. Minelli C, Granell R, Newson R, Rose-Zerilli MJ, Torrent M, Ring SM, et al. Glutathione-S-transferase genes and asthma phenotypes: a Human Genome Epidemiology (HuGE) systematic review and meta-analysis including unpublished data. *Int J Epidemiol* 2010;39:539-62.
179. Kozu Y, Gon Y, Maruoka S, Kazumichi K, Sekiyama A, Kishi H, et al. Protocadherin-1 is a glucocorticoid-responsive critical regulator of airway epithelial barrier function. *BMC Pulm Med* 2015;15:80.
180. Faura TG, Vandepoele K, Brouwer U, Koning H, Elderman RM, Hackett TL, et al. Protocadherin-1 binds to SMAD3 and suppresses TGF-beta1-induced gene transcription. *Am J Physiol Lung Cell Mol Physiol* 2015;309:L725-35.
181. Mortensen LJ, Kreiner-Moller E, Hakonarson H, Bonnelykke K, Bisgaard H. The PCDH1 gene and asthma in early childhood. *Eur Respir J* 2014;43:792-800.
182. Faura TG, Nawijn MC, Koppelman GH. Protocadherin-1: epithelial barrier dysfunction in asthma and eczema. *Eur Respir J* 2014;43:671-4.
183. Bochkov YA, Watters K, Ashraf S, Griggs TF, Devries MK, Jackson DJ, et al. Cadherin-related family member 3, a childhood asthma susceptibility gene product, mediates rhinovirus C binding and replication. *Proc Natl Acad Sci U S A* 2015;112:5485-90.
184. Qiu R, Yang Y, Zhao H, Li J, Xin Q, Shan S, et al. Signal transducer and activator of transcription 6 directly regulates human ORM DL3 expression. *FEBS J* 2013;280:2014-26.
185. Schedel M, Michel S, Gaertner VD, Toncheva AA, Depner M, Binia A, et al. Polymorphisms related to ORM DL3 are associated with asthma susceptibility, alterations in transcriptional regulation of ORM DL3, and changes in TH2 cytokine levels. *J Allergy Clin Immunol* 2015;136:893-903.
186. Worgall TS. Sphingolipids, ORM DL3 and asthma: what is the evidence? *Curr Opin Clin Nutr Metab Care* 2016;20:99-103.
187. Hsu KJ, Turvey SE. Functional analysis of the impact of ORM DL3 expression on inflammation and activation of the unfolded protein response in human airway epithelial cells. *Allergy Asthma Clin Immunol* 2013;9:4.
188. Miller M, Rosenthal P, Beppu A, Mueller JL, Hoffman HM, Tam AB, et al. ORM DL3 transgenic mice have increased airway remodeling and airway responsiveness characteristic of asthma. *J Immunol* 2014;192:3475-87.

189. Loser S, Gregory LG, Zhang Y, Schaefer K, Walker SA, Buckley J, et al. Pulmonary ORMDL3 is critical for induction of *Alternaria*-induced allergic airways disease. *J Allergy Clin Immunol* 2016 [Epub ahead of print].
190. Li X, Hastie AT, Hawkins GA, Moore WC, Ampleford EJ, Milosevic J, et al. eQTL of bronchial epithelial cells and bronchial alveolar lavage deciphers GWAS-identified asthma genes. *Allergy* 2015;70:1309-18.
191. Biagini Myers JM, Martin LJ, Kovacic MB, Mersha TB, He H, Pilipenko V, et al. Epistasis between serine protease inhibitor Kazal-type 5 (SPINK5) and thymic stromal lymphopoietin (TSLP) genes contributes to childhood asthma. *J Allergy Clin Immunol* 2014;134:891-9.
192. Harada M, Nakashima K, Hirota T, Shimizu M, Doi S, Fujita K, et al. Functional polymorphism in the suppressor of cytokine signaling 1 gene associated with adult asthma. *Am J Respir Cell Mol Biol* 2007;36:491-6.
193. Balaci L, Spada MC, Olla N, Sole G, Loddo L, Anedda F, et al. IRAK-M is involved in the pathogenesis of early-onset persistent asthma. *Am J Hum Genet* 2007;80:1103-14.
194. Wu Q, van Dyk LF, Jiang D, Dakhama A, Li L, White SR, et al. Interleukin-1 receptor-associated kinase M (IRAK-M) promotes human rhinovirus infection in lung epithelial cells via the autophagic pathway. *Virology* 2013;446:199-206.
195. Yao YS, Chang WW, He LP, Jin YL, Li CP. An updated meta-analysis of transforming growth factor-beta1 gene: three well-characterized polymorphisms with asthma. *Hum Immunol* 2016;77:1291-9.
196. Hatsushika K, Hirota T, Harada M, Sakashita M, Kanzaki M, Takano S, et al. Transforming growth factor-beta(2) polymorphisms are associated with childhood atopic asthma. *Clin Exp Allergy* 2007;37:1165-74.
197. Ueda T, Niimi A, Matsumoto H, Takemura M, Yamaguchi M, Matsuoka H, et al. TGFB1 promoter polymorphism C-509T and pathophysiology of asthma. *J Allergy Clin Immunol* 2008;121:659-64.
198. Li X, Howard TD, Moore WC, Ampleford EJ, Li H, Busse WW, et al. Importance of hedgehog interacting protein and other lung function genes in asthma. *J Allergy Clin Immunol* 2011;127:1457-65.
199. Nieuwenhuis MA, Vonk JM, Himes BE, Sarnowski C, Minelli C, Jarvis D, et al. PTTG1IP and MAML3, novel genomewide association study genes for severity of hyperresponsiveness in adult asthma. *Allergy* 2017;72:792-801.
200. Singhanian A, Rupani H, Jayasekera N, Lumb S, Hales P, Gozzard N, et al. Altered epithelial gene expression in peripheral airways of severe asthma. *PLoS One* 2017;12:e0168680.
201. Roeder T, Isermann K, Kallsen K, Uliczka K, Wagner C. A *Drosophila* asthma model—what the fly tells us about inflammatory diseases of the lung. *Adv Exp Med Biol* 2012;710:37-47.
202. Progatzyk F, Cook HT, Lamb JR, Bugeon L, Dallman MJ. Mucosal inflammation at the respiratory interface: a zebrafish model. *Am J Physiol Lung Cell Mol Physiol* 2016;310:L551-61.