

DNA methylation profiles between airway epithelium and proxy tissues in children

Journal:	<i>Acta Paediatrica</i>
Manuscript ID	SPAE-2017-0361
Manuscript Type:	Regular Article
Date Submitted by the Author:	08-May-2017
Complete List of Authors:	Brugha, Rossa; Imperial College London, National Heart and Lung Institute Lowe, Robert; Barts and The London School of Medicine and Dentistry Blizard Institute Henderson, John; University of Bristol, Holloway, John; Human Development and Health, Faculty of Medicine, University of Southampton Rakyan, Vardhman; Barts and The London School of Medicine and Dentistry Blizard Institute Wozniak, Eva; Barts and The London School of Medicine and Dentistry Blizard Institute Mahmud, Nadiya; Barts and The London School of Medicine and Dentistry Blizard Institute Seymour, Kay; Barts Health NHS Trust Grigg, Jonathan; Barts and The London School of Medicine and Dentistry Blizard Institute Shaheen, Seif; Barts and The London School of Medicine and Dentistry Blizard Institute
Keywords:	Epigenetics, asthma, wheeze

DNA methylation profiles between airway epithelium and proxy tissues in children

Rossa Brugha¹, Robert Lowe², A. John Henderson³, John W. Holloway⁴, Vardhman Rakyan²,
Eva Wozniak⁵, Nadiya Mahmud⁵, Kay Seymour⁶, Jonathan Grigg² Seif O. Shaheen².

¹National Heart and Lung Institute, Imperial College London, London, UK.

²Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary,
University of London, London, UK

³School of Social and Community Medicine, University of Bristol, Bristol, UK.

⁴Human Development and Health, Faculty of Medicine, University of Southampton,
Southampton, UK.

⁵Genome Centre, Barts and The London School of Medicine and Dentistry, London, UK.

⁶Ear, Nose and Throat Surgery, Barts Health NHS Trust, London, UK.

Correspondence and reprint requests to: Prof Seif Shaheen, Department of Primary Care and
Public Health, Blizard Institute, Barts and The London School of Medicine and Dentistry, 58
Turner Street, London E1 2AB.

Email: s.shaheen@qmul.ac.uk

Tel: +44 (0)20 7882 2480; Fax: +44 (0)20 7882 2552

Key words: Epigenetics, asthma and early wheeze

Short title: Airway DNA methylation Word count: 2396

Abstract

Background

Epidemiological studies of DNA methylation in airway disease have largely been conducted using blood or buccal samples. However, given tissue specificity of DNA methylation, these surrogate tissues may not allow reliable inferences about methylation in the lung.

Objective

To compare the pattern of DNA methylation in blood, buccal and nasal epithelial cells to that in airway epithelial cells of children with and without asthma.

Methods

Samples of blood, and buccal, nasal and airway epithelium were obtained from six children undergoing elective anaesthesia for adenotonsillectomy. DNA methylation was assessed at 450,000 sites using the Illumina HumanMethylation450 array.

Results

Eighteen samples from all sites were suitable for analysis. Hierarchical clustering demonstrated that the methylation profile in nasal epithelium was most representative of that in airway epithelium; the profile in buccal cells was moderately similar; and that in blood was least similar.

Conclusion

DNA methylation in blood poorly reflects methylation in airway epithelium. Future epidemiological studies of DNA methylation and airway diseases should measure methylation either in buccal cells or, preferably, in nasal epithelial cells.

Article summary

DNA methylation is tissue specific, and primary tissue may not be available for large scale studies of epigenetic markers in airways disease. We sought to compare nasal, buccal, airway and blood DNA methylation in order to establish a best proxy for airway cells. We demonstrate that DNA methylation in nasal epithelium is the most representative of airway DNA methylation.

For Peer Review Only

Introduction

There is growing interest in the role of epigenetic control of gene expression in the lung [1], and in the likely importance of epigenetic mechanisms in the aetiology of asthma [2-4].

Consequently there have been calls to include epigenetics in epidemiological studies of asthma [5] and evidence has been gradually accumulating to implicate altered DNA methylation, the most readily studied epigenetic process in population-based studies, in childhood asthma. For example, differential methylation has been reported in relation to childhood wheezing and asthma [6, 7], and to risk factors such as maternal smoking in pregnancy [8, 9], although epidemiological data confirming whether the associations between early life risk factors and asthma are mediated through altered DNA methylation are still sparse. Experimental evidence in mouse models has provided further evidence for a potential role of altered DNA methylation in asthma [10-12].

Epidemiological studies of epigenetics and asthma to date have tended to measure DNA methylation predominantly using DNA extracted from peripheral blood leukocytes [6, 7, 9], and sometimes buccal samples [8], as these sources of DNA are readily accessible and direct lower airway sampling is generally not feasible in population-based studies. A potential difficulty with this approach is that DNA methylation tends to be highly tissue-specific [13] and even cell-type specific [14], and hence these tissues may not allow reliable inferences about methylation in the lung. It has been argued that methylation measured in buccal DNA samples may better reflect methylation in the lung than methylation measured in peripheral blood leukocyte DNA [15], and we recently showed that buccal samples are likely to be a more informative surrogate tissue than

1
2
3 blood for epigenome-wide association studies (EWAS) of non-blood based diseases/phenotypes,
4
5 especially those linked to epithelial disorders such as paediatric eosinophilic oesophagitis [16].
6
7
8 However, the most informative proxy tissue may be nasal epithelium, given that the nose and
9
10 airways form a continuous tract, and that nasal epithelial cells have been used as surrogates for
11
12 bronchial epithelial cells in studies of airway inflammation [17]. To date, no comparison has
13
14 been made between the DNA methylation profiles of airway and nasal epithelial cells, buccal
15
16 cells and blood to confirm the best proxy tissue for epidemiological studies of airway diseases.
17
18
19 We have therefore sought to establish which proxy tissue exhibits the closest pattern of
20
21 epigenome-wide DNA methylation to that of airway epithelial cells. Our *a priori* hypothesis was
22
23 that methylation profiles from nasal epithelial cells would show a greater similarity than those
24
25 from buccal samples and blood.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Methods

Participants

In order to obtain uncontaminated airway cells, children requiring elective endotracheal intubation were recruited to the study. Six children (four female, two male) aged 5-13 years undergoing elective tonsillectomy or adenotonsillectomy were recruited from the Ear, Nose and Throat clinic at the Royal London Hospital, three of whom had doctor diagnosed asthma. Children with a respiratory condition other than asthma were excluded. Ethical approval was obtained from the National Research Ethics Service Committee London - Riverside (REC 11/LO/1579), and parents or carers consented for study procedures.

Sampling

Samples from 4 tissue sites (airway, nasal, buccal and blood) were obtained. Buccal cells were collected pre-operatively using swabs (Isohelix SK-2S, Cell Projects, Kent, UK). Immediately following intubation, tracheal epithelial cells were collected by performing blind brushings via the endotracheal tube using a 2mm cytology brush (BC-202D-5010, Olympus KeyMed, Essex, UK). This has been shown to be an effective and safe method in children when bronchoscopy is not feasible [18, 19]. Nasal epithelial cells were obtained by directly observed curettage of the inferior nasal turbinates (Rhino-Pro® Nasal Mucosal Curette, Arlington Scientific, Utah, USA). Peripheral blood was obtained by venesection and mixed with EDTA (BD vacutainer, Becton Dickinson, Oxford, UK).

Measurement of DNA methylation

DNA Extraction and Quality control

Upon receipt, buccal swabs were treated as per the Isohelix DNA Isolation Kit DDK-50 protocol Part A, until appropriate numbers for extraction had been received. Nasal mucosal samples and airway brushings were immersed in 0.9% sodium chloride and frozen at -80°C. After thawing, tubes were vortexed to loosen collected cells, probes and brushes were removed, and the resulting cell suspension was then centrifuged at 3,500 RCF for 10 minutes in a bench-top centrifuge to pellet the cells. The supernatant was removed and DNA extraction of buccal swabs, nasal and airway samples was carried out as in the Isohelix protocol (beginning at Part B for buccal swabs), with the optional step 6 included, and the elution volume used was 30µl.

Blood samples were stored at -20° C until extraction. Extraction was performed using a salting out procedure [20]. Briefly, blood samples were defrosted overnight before 40mls of cold water were added to lyse red cells, followed by centrifugation at 3000 RCF for 20 minutes at 4° C to precipitate white cells. Supernatant was discarded and the pellet re-suspended in 40mls solution 1 (0.32M Sucrose, 0.01M Tris-Cl, pH 7.6, 0.005M MgCl₂, 1%TRitonX100, 0.02 Sodium Azide). Samples were centrifuged at 3000 RCF for 20 minutes at 4° C. The supernatant was discarded and the pellet re-suspended in 30mls solution 1. Tubes were centrifuged at 3000 RCF for 20 minutes at 4° C and the pellet re-suspended in 11mls of Solution 2 (0.05M Tris-Cl, pH 8, 0.02M EDTA, pH 8, 2%SDS). Samples were digested with 1mg Proteinase K overnight at 37° C followed by protein precipitation with saturated 4 mls NaCl solution. Protein was precipitated by centrifugation at 3500 RCF for 20min at 4°C and DNA was precipitated with 3x volume 100% Ethanol. DNA quantity and purity were measured using a Nanodrop8000 spectrophotometer (Thermofisher Scientific, Delaware, USA). A Qubit® 2.0 fluorometer

1
2
3 (LifeTechnologies, Paisley, United Kingdom) with the Qubit® dsDNA HS kit was used to
4
5 quantify intact double-stranded DNA. Samples were also checked for RNA contamination by an
6
7 Agilent RNA 6000 Nano kit on the Agilent 2100 Bioanalyser (Agilent Technologies, California,
8
9 USA).

10 11 12 13 14 15 *Bisulphite conversion and Quality control*

16
17 Any sample with less than 500ng total DNA quantity was excluded from further analysis. 500ng
18
19 of DNA were bisulphite converted using the Zymo EZ-96 DNA Methylation Kit D5004 (Zymo
20
21 Research Corp, California, USA). Bisulphite conversion was performed as per manufacturer's
22
23 instructions but with incubation conditions recommended by Illumina for the Infinium®
24
25 Methylation arrays. The resulting quantity of single-stranded, converted sample was measured
26
27 using the Nanodrop with the RNA setting. An initial QC of the bisulfite conversion efficiency
28
29 suggested one of the samples had a particularly low level and hence this was removed before
30
31 normalisation.
32
33
34
35
36
37
38

39 *Illumina Infinium® Methylation 450K arrays*

40
41 Samples with a converted concentration of less than 10ng/μl were excluded from array
42
43 processing. CpG site methylation was analysed at 485,577 sites using the Illumina Infinium
44
45 HumanMethylation 450 Beadchip (450k). This has been validated [21] and recently used to
46
47 demonstrate differential methylation in relation to maternal smoking in pregnancy [8, 9]. 4μl of
48
49 bisulphite converted DNA was used and samples were processed according to the Illumina
50
51 Infinium®HD Meth450 Protocol (Part No 15019522) and scanned on the Illumina iScan (San
52
53
54
55
56
57
58
59
60

1
2
3 Diego, CA). Initial data QC was performed using the Genome Studio analysis package (Illumina
4
5 (San Diego, CA)).
6
7
8
9

10 Statistical analysis

11
12 Raw intensity values extracted from the IDAT files were quantile normalised separately for each
13 colour channel as well as probe type using a custom script. We remapped the genomic sequence
14 of all probes to hg19 and removed those probes which mapped with >90% similarity to more
15 than one genomic location as well as those probes which mapped to either chromosome X or Y.
16 Furthermore we removed those probes which recorded intensity levels around background by
17 using a detection p-value <0.01. This left 441,946 probes which were used for further analysis.
18
19 We calculated beta values using the following equation [21] and used this as a measure of
20 methylation across the 441,946 probes:
21
22
23
24
25
26
27
28
29
30
31

$$32 \quad \beta = \frac{I_m}{I_m + I_u + 100}$$

33
34
35
36
37
38
39 where I_m is the intensity in the methylated channel and I_u is the intensity in the unmethylated
40 channel. To ascertain how similar different tissue types are in terms of their methylation state we
41 performed an unsupervised hierarchical clustering. We first calculated the Euclidean distance (d)
42 between each of the samples by summing the square of the beta differences across all of the
43 probes. Then we clustered the samples using a mean or average linkage. To further identify how
44 close nasal, buccal and blood are in terms of methylation state to airway epithelium we first
45 averaged the beta values of the samples for each tissue and then calculated the Euclidean
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 distance by summing the square of the differences between the average beta value of airway
4
5 epithelium samples and each of the other tissues.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Results

Characteristics of the children are shown in Table 1. One nasal sample and three buccal samples were insufficient for analysis, one blood sample was misplaced, and one sample failed bisulphite conversion, leaving 18 samples (plus 5 duplicate samples, i.e. 23 arrays in total) which were suitable for analysis.

Figure 1 demonstrates the hierarchical clustering of the samples. Taking into account the 441,946 probes for each sample, the diagram represents the difference in methylation at the same site in the DNA sequence between tissues. The greater the distance (represented by height on the diagram), the greater the difference between tissues. This shows that the methylation profile of nasal epithelial cells was most similar to that of airway epithelial cells; the profile of buccal cells was moderately similar; that of blood was least similar. When we quantified this by calculating a standard Euclidean distance of the average methylation beta values for each tissue from the average value for airway epithelial cells, the distances for nasal, buccal and blood were, respectively, 46.1, 72.3, and 91.4. To further evaluate this we carried out a Wilcoxon rank sum test on the Euclidean distances for each sample of nasal, buccal and blood against the average methylation for airway epithelial cells. Both nasal (p-value = 0.004329) and buccal (p-value = 0.009524) were significantly closer to airway epithelial cells than blood.

Figure 2 demonstrates the extent of methylation for the 20,000 most variable probes (defined by having the largest variance in beta values across all of our samples), where yellow represents hypomethylation and blue represents hypermethylation; this gives a visual representation of the difference in methylation across the separate tissues, with blood clearly separate from the

1
2
3 respiratory tract samples. We analysed the blood cell type data for the same 20,000 variable
4 probes analysed in Figure 2. We found that nasal, airway and buccal cells clustered together as
5 before, while whole blood clustered with the various blood subsets (data not shown), confirming
6 that our conclusions above are not affected by blood cell type
7
8
9
10
11
12

13
14
15 Figure 3 demonstrates the hierarchical clustering of the samples when we limited the analyses to
16 children with asthma (n=3). This confirmed a similar hierarchy of methylation profiles across
17 tissues to that seen in all children. Euclidean distances of the average methylation beta values
18 for each tissue from the average value for airway epithelial cells, were 42.5, 84.2 and 92.8 for
19 nasal, buccal and blood respectively.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

Asthma is primarily an epithelial disorder [22] and hence increasing our understanding of epigenetic changes in airway epithelium may shed light on asthma aetiology. In epidemiological studies of the epigenetics of asthma, in which airway epithelial cells cannot be obtained, we therefore need to focus efforts on obtaining the most representative surrogate tissue. In this study we have shown that nasal epithelial cells are a better proxy tissue than buccal cells for making inferences about epigenome-wide methylation patterns in airway epithelial cells, with blood being the worst proxy tissue. Our findings suggest that, whilst studies to date which have measured DNA methylation in blood may have obtained some useful information, future studies of the epigenetics of asthma are likely to prove more informative if buccal cells, or preferably nasal epithelial cells, are collected. We acknowledge, though, that blood DNA methylation profiling may still be informative for understanding epigenetic influences on the immune (atopic) component of asthma, as suggested by previous studies of Th1/Th2 gene methylation [23-25] or possibly in relation to DNA methylation changes induced by exposures that occur *in utero* or are inherited as a result of trans-generational transmission of DNA methylation.

Our conclusion that blood methylation profiles may not be informative for making inferences about DNA methylation in the airway is supported by a recent study of children which compared DNA methylation profiles between airway epithelial cells (AECs) and peripheral blood mononuclear cells (PMBCs)[26]. The authors demonstrated differential methylation of CpG sites in AECs compared with PBMCs, irrespective of disease phenotype, but nasal epithelial cells and buccal cells were not studied. Although we did not separate out PBMCs before DNA extraction in our study, and we did not take into account the differential white cell count in our analysis of DNA methylation in blood, it has recently been shown that polymorphonuclear leucocytes and

1
2
3 PBMCs do not differ in terms of methylation profiles [9]. Whilst it was recently proposed that
4 nasal cell DNA may be a more appropriate and accessible proxy tissue for studying the
5 epigenetics of asthma [27], airway epithelial cells and other proxy tissues were not compared in
6 that study.
7
8
9
10
11

12
13
14
15 A potential limitation of our study is that the sample size was small, as we experienced a number
16 of refusals from parents because of concerns about taking airway brushings and prolonging the
17 anaesthetic time. Despite this, by taking multiple samples from the same subject, we were able to
18 demonstrate that a clear hierarchy of similarity between the methylation profiles of different
19 proxy tissues and that of airway epithelial cells exists within individuals. Furthermore, a similar
20 hierarchy was seen in children with asthma. Our nasal samples were acquired using a small
21 curette which is an uncomfortable procedure when awake. Whilst tolerated well by adults it
22 would be challenging to perform in younger children when not anaesthetised. Soft brushing of
23 the nasal epithelium using a swab, as described by Baccarelli *et al* [27], offers an alternative
24 technique for acquiring nasal epithelial cells in epidemiological studies, and may be more
25 suitable especially in young children.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43
44 In summary, we have shown that the best surrogate tissue for capturing methylation marks
45 relevant to airway epithelium is the nasal epithelium, followed by buccal cells. DNA methylation
46 profiling of blood samples does not accurately reflect the pattern of methylation seen in the
47 airway. Future epidemiological studies of the role of DNA methylation in airways diseases
48 should take this into account, and existing or forthcoming data from studies based on
49 methylation in blood should be interpreted with caution.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Acknowledgements**
4

5
6 We would like to thank the children and families who took part, and the nursing, theatre and
7
8 anaesthetic staff who facilitated sampling.
9

10
11
12 **Competing interests**
13

14
15 No competing interests to declare
16

17
18 **Funding:** Asthma UK, Barts and The London Charity
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Yang IV, Schwartz DA. Epigenetic control of gene expression in the lung. *Am J Respir Crit Care Med* 2011;183:1295-1301.
2. Kabesch M, Michel S, Tost J. Epigenetic mechanisms and the relationship to childhood asthma. *Eur Respir J* 2010;36:950-961.
3. Shaheen SO, Adcock IM. The developmental origins of asthma: Does epigenetics hold the key? *Am J Respir Crit Care Med* 2009;180:690-691.
4. Karmaus W, Ziyab AH, Everson T, et al. Epigenetic mechanisms and models in the origins of asthma. *Curr Opin Allergy Clin Immunol* 2013;13:63-69.
5. Miller RL, Ho SM. Environmental epigenetics and asthma: Current concepts and call for studies. *Am J Respir Crit Care Med* 2008;177:567-573.
6. Morales E, Bustamante M, Vilahur N, et al. DNA hypomethylation at alox12 is associated with persistent wheezing in childhood. *Am J Respir Crit Care Med* 2012;185:937-943.
7. Perera F, Tang WY, Herbstman J, et al. Relation of DNA methylation of 5'-cpg island of acsl3 to transplacental exposure to airborne polycyclic aromatic hydrocarbons and childhood asthma. *PLoS One* 2009;4:e4488.
8. Breton CV, Byun HM, Wenten M, et al. Prenatal tobacco smoke exposure affects global and gene-specific DNA methylation. *Am J Respir Crit Care Med* 2009;180:462-467.
9. Joubert BR, Haberg SE, Nilsen RM, et al. 450k epigenome-wide scan identifies differential DNA methylation in newborns related to maternal smoking during pregnancy. *Environ Health Perspect* 2012;120:1425-1431.
10. Hollingsworth JW, Maruoka S, Boon K, et al. In utero supplementation with methyl donors enhances allergic airway disease in mice. *J Clin Invest* 2008;118:3462-3469.

- 1
2
3 11. Brand S, Kesper DA, Teich R, et al. DNA methylation of th1/th2 cytokine genes affects
4 sensitization and progress of experimental asthma. *J Allergy Clin Immunol* 2012;129:1602-1610.
5
6
- 7 12. Mikhaylova L, Zhang Y, Kobzik L, et al. Link between epigenomic alterations and genome-wide
8 aberrant transcriptional response to allergen in dendritic cells conveying maternal asthma risk. *PLoS One*
9 2013;8:e70387.
10
11
- 12 13. Eckhardt F, Lewin J, Cortese R, et al. DNA methylation profiling of human chromosomes 6, 20
13 and 22. *Nat Genet* 2006;38:1378-1385.
14
15
- 16 14. Yang Y, Haitchi HM, Cakebread J, et al. Epigenetic mechanisms silence a disintegrin and
17 metalloprotease 33 expression in bronchial epithelial cells. *J Allergy Clin Immunol* 2008;121:1393-1399,
18 1399 e1391-1314.
19
20
- 21 15. Bhutani M, Pathak AK, Fan YH, et al. Oral epithelium as a surrogate tissue for assessing smoking-
22 induced molecular alterations in the lungs. *Cancer Prev Res* 2008;1:39-44.
23
24
- 25 16. Lowe R, Gemma C, Beyan H, et al. Buccals are likely to be a more informative surrogate tissue
26 than blood for epigenome-wide association studies. *Epigenetics* 2013;8.
27
28
- 29 17. McDougall CM, Blaylock MG, Douglas JG, et al. Nasal epithelial cells as surrogates for bronchial
30 epithelial cells in airway inflammation studies. *Am J Respir Cell Mol Biol* 2008;39:560-568.
31
32
- 33 18. Doherty GM, Christie SN, Skibinski G, et al. Non-bronchoscopic sampling and culture of bronchial
34 epithelial cells in children. *Clin Exp Allergy* 2003;33:1221-1225.
35
36
- 37 19. Lane C, Burgess S, Kicic A, et al. The use of non-bronchoscopic brushings to study the paediatric
38 airway. *Respir Res* 2005;6:53.
39
40
- 41 20. Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human
42 nucleated cells. *Nucleic Acids Res* 1988;16:1215.
43
44
- 45 21. Sandoval J, Heyn H, Moran S, et al. Validation of a DNA methylation microarray for 450,000 cpg
46 sites in the human genome. *Epigenetics* 2011;6:692-702.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
22. Holgate ST. The sentinel role of the airway epithelium in asthma pathogenesis. *Immunol Rev* 2011;242:205-219.
23. Soto-Ramirez N, Arshad SH, Holloway JW, et al. The interaction of genetic variants and DNA methylation of the interleukin-4 receptor gene increase the risk of asthma at age 18 years. *Clin Epigenetics* 2013;5:1868-7083.
24. White GP, Hollams EM, Yerkovich ST, et al. CpG methylation patterns in the ifngamma promoter in naive t cells: Variations during th1 and th2 differentiation and between atopics and non-atopics. *Pediatr Allergy Immunol* 2006;17:557-564.
25. Kwon NH, Kim JS, Lee JY, et al. DNA methylation and the expression of il-4 and ifn-gamma promoter genes in patients with bronchial asthma. *J Clin Immunol* 2008;28:139-146.
26. Stefanowicz D, Hackett TL, Garmaroudi FS, et al. DNA methylation profiles of airway epithelial cells and pbmcs from healthy, atopic and asthmatic children. *PLoS One* 2012;7:e44213.
27. Baccarelli A, Rusconi F, Bollati V, et al. Nasal cell DNA methylation, inflammation, lung function and wheezing in children with asthma. *Epigenomics* 2012;4:91-100.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Legends for figures

Figure 1: Hierarchical cluster dendrogram of DNA methylation by tissue type. The respiratory tract tissues cluster together, separately from blood. The dendrogram is calculated using agglomerative hierarchical clustering. The root of the dendrogram is at the top of the figure and its y-axis represents the Euclidean distance between clusters at each splitting point. A greater distance indicates greater differences between tissues.

Figure 2: To visually inspect the data we took 20,000 probes with the highest variability across all samples. The rows of the heatmap represent an individual probe while the columns represent an individual sample. The beta value as recorded from the array is plotted from yellow to blue with yellow representing low methylation, green fractional methylation and blue high methylation. In the top part of the heatmap the respiratory tract tissues contain cytosines in the genome that are hypomethylated compared to blood. In the lower half of the heatmap the majority of probes are specifically hypomethylated in blood.

Figure 3: Hierarchical cluster dendrogram of DNA methylation by tissue type restricted to children with asthma.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Table 1: Subject demographics.

<i>Subject</i>	<i>Age</i>	<i>Gender</i>	<i>Ethnicity</i>	<i>Asthma</i>	<i>Atopy</i>	<i>Home ETS exposure</i>	<i>Tissue</i>
1	13	F	South Asian	N	N	N	Na, Aw, Bu
2	10	M	South Asian	Y	N	Y	Na, Aw, Bl
3	5	F	South Asian	N	N	Y	Na*, Aw*, Bu*, Bl*
4	6	F	European	Y	N	Y	Na, Aw, Bl
5	9	F	European- Afro Caribbean	Y	Hayfever	N	Aw*, Bu, Bl
6	5	M	East Asian	N	Eczema	N	Aw, Bl

ETS; environmental tobacco smoke, Aw airway, Na nasal, Bu buccal, Bl blood,

*processed in duplicate

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1: Hierarchical cluster dendrogram of DNA methylation by tissue type. The respiratory tract tissues cluster together, separately from blood. The dendrogram is calculated using agglomerative hierarchical clustering. The root of the dendrogram is at the top of the figure and its y-axis represents the Euclidean distance between clusters at each splitting point. A greater distance indicates greater differences between tissues.

70x52mm (300 x 300 DPI)

Only




Figure 2: To visually inspect the data we took 20,000 probes with the highest variability across all samples. The rows of the heatmap represent an individual probe while the columns represent an individual sample. The beta value as recorded from the array is plotted from yellow to blue with yellow representing low methylation, green fractional methylation and blue high methylation. In the top part of the heatmap the respiratory tract tissues contain cytosines in the genome that are hypomethylated compared to blood. In the lower half of the heatmap the majority of probes are specifically hypomethylated in blood.

70x52mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3: Hierarchical cluster dendrogram of DNA methylation by tissue type restricted to children with asthma.

70x52mm (300 x 300 DPI)

EW Only

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

1
2
3 **DNA methylation profiles between airway epithelium and proxy tissues in**
4
5 **children**
6
7

8
9
10 Author contributions

11
12
13
14 Rossa Brugha screened and recruited children, performed sampling, maintained the
15 sample database and wrote the first draft of the manuscript with Prof Shaheen
16 Robert Lowe performed data analysis, compiled figures for publication, and
17 contributed to the manuscript.
18
19

20
21
22 John Henderson contributed to secure funding, protocol development, and contributed
23 to the manuscript.
24
25

26
27 John Holloway contributed to secure funding, developed the protocol, and contributed
28 to the manuscript.
29
30

31
32 Vardhman Rakyan supervised data analysis, compilation of figures, and contributed
33 to the manuscript.
34
35

36
37 Eva Wozniak performed sample processing and analysis, and contributed to the
38 manuscript.
39
40

41
42 Nadiya Mahmud performed sample processing and analysis, and contributed to the
43 manuscript.
44
45

46
47 Kay Seymour screened and identified patients and was responsible for supervising
48 sampling procedures in theatre, and contributed to the manuscript.
49

50
51 Jonathan Grigg contributed to secure funding, developed the protocol, and contributed
52 to the manuscript.
53

54
55 Seif Shaheen secured funding, supervised the study, and finalised the manuscript. He
56 assumes overall responsibility for the reported findings.
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only