

POINCARÉ DUALITY AND LANGLANDS DUALITY FOR EXTENDED AFFINE WEYL GROUPS

GRAHAM A. NIBLO, ROGER PLYMEN AND NICK WRIGHT

ABSTRACT. In this paper we construct an equivariant Poincaré duality between dual tori equipped with finite group actions. We use this to demonstrate that Langlands duality induces a rational isomorphism between the group C^* -algebras of extended affine Weyl groups at the level of K -theory.

CONTENTS

Introduction	1
1. Statement of Results	3
2. Background	5
2.1. Real Langlands duality	5
2.2. Affine and Extended Affine Weyl groups	7
3. Equivariant Poincaré Duality between $C(T)$ and $C(T^\vee)$	8
3.1. The Poincaré line bundle	8
3.2. Construction of the element \mathcal{Q} in $KK_W(C(T^\vee) \widehat{\otimes} C(T), \mathbb{C})$	10
3.3. The Kasparov product $\mathcal{P} \otimes_{C(T^\vee)} \mathcal{Q}$	14
3.4. The Kasparov product $\mathcal{P} \otimes_{C(T)} \mathcal{Q}$	18
4. Poincaré duality between $C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W)$ and $C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)$	20
4.1. Descent of Poincaré duality	20
4.2. Proof of Theorem 1.1	22
4.3. Proof of Theorem 1.2	23
4.4. The connection with the Baum Connes assembly map	23
5. Langlands Duality and K -theory	26
References	27

INTRODUCTION

Let T be a compact torus and let W be a finite group acting on T with fixed point. We construct a W -equivariant degree 0 Poincaré duality between $C(T)$ and $C(T^\vee)$, where T^\vee denotes the dual torus equipped with the dual action of W .

Moreover we show that there is a non-equivariant Poincaré duality between the crossed product algebras $C(T) \rtimes W$ and $C(T^\vee) \rtimes W$. Indeed we provide a general mechanism to descend equivariant Poincaré duality to Poincaré duality for crossed products. As far as we are aware this does not appear elsewhere in the literature.

The third author was partially supported by EPSRC grant EP/J015806/1.

In the case when W is trivial, our degree 0 duality is connected to the Baum-Connes assembly map in the following way: Let T be a compact torus (equipped with the structure of a Lie group), and let $X^*(T), X_*(T)$ be the groups of characters and cocharacters respectively. By definition the dual torus T^\vee is the torus such that $X^*(T^\vee) = X_*(T)$ and $X_*(T^\vee) = X^*(T)$. Whence the Pontryagin dual of $X_*(T)$ is the torus T^\vee . The Baum-Connes assembly map for $X_*(T)$ gives a degree 0 isomorphism

$$K_*(T) \xrightarrow{\cong} K_*(C^*X_*(T)) \cong K^*(T^\vee).$$

This isomorphism agrees with our Poincaré duality, though this is not immediate from the definition of the two maps, see Section 4.4.

For an isometric action of a group W on a closed Riemannian manifold M^n , Kasparov's Poincaré duality [10], by contrast with our Poincaré duality, provides an isomorphism from $KK_W(C(M), \mathbb{C})$ to $KK_W(\mathbb{C}, C_\tau(M))$ where $C_\tau(M)$ denotes the algebra of continuous sections of the Clifford bundle for the cotangent bundle τ of M . See also Echterhoff *et al*, [7]. If the action is trivial and M is a spin manifold, then the twisting by the Clifford algebra simply induces a dimension shift so Kasparov's Poincaré duality has degree n modulo 2. In the case where M is a torus and W is trivial, this is connected to our Poincaré duality via the Dirac-dual-Dirac method, which addresses the dimension shift. In the equivariant case the group acts nontrivially on the Clifford bundle, so the appearance of this bundle no longer simply gives a dimension shift. Indeed, for example, letting $\mathbb{Z}/2\mathbb{Z}$ act by complex conjugation on the 1-dimensional torus $U(1)$, then $KK_W(\mathbb{C}, C_\tau(U(1)))$ is \mathbb{Z}^3 in dimension 0 and 0 in dimension 1, which agrees with the *unshifted* K -theory group $KK_W(\mathbb{C}, C(U(1)))$.

In this paper in order to describe the KK -cycles defining our Poincaré dualities explicitly, we have given direct proofs of the relevant properties of these cycles and their pairings. As remarked by the referee, it is in principle possible to obtain these elements by combining Kasparov's Poincaré duality elements with the Dirac and dual-Dirac cycles. Providing full details of this reduction to the known results is in itself somewhat complicated and we have opted to give the direct, self-contained argument.

As an application of our Poincaré duality we consider the case where T is the maximal torus in a compact connected semi-simple Lie group and W is the Weyl group. The dual torus is then the maximal torus in the Langlands dual Lie group. In general there is no W -equivariant homeomorphism between the two tori, hence *a priori* one would not expect them to have the same equivariant K -theory. However our Poincaré duality gives a canonical pairing between these two equivariant K -theory groups, and hence ignoring torsion these groups are isomorphic. Moreover our Poincaré duality also provides a canonical pairing between the K -theory of the extended affine Weyl groups of the original Lie group and its Langlands dual. This again yields an isomorphism up to torsion in K -theory, although these discrete groups are not typically isomorphic. In [12] we explore this phenomenon in further detail and give detailed computations of these K -theory groups in a number of cases.

The connection between T -duality and Langlands duality has been studied by Daenzer-van Erp who showed that Langlands duality for complex reductive Lie groups can be implemented by T -dualization for groups whose simple factors are of type A, D or E, [6]. This was generalised by Bunke-Nikolaus, see [5]. The study of T -duality in these papers, involves examining the Lie group viewed as a principal bundle of tori via the action of the maximal torus on the group. Here by contrast we study the Weyl group action on the maximal torus, instead of the maximal torus action on the Lie group. In both cases there is a natural duality arising from Langlands duality of root systems and the possible unification of these two perspectives would provide an interesting future project.

We would like to thank Maarten Solleveld for his helpful comments on the first version of this paper. We would also like to thank the editor Jonathan Rosenberg and the referee, whose careful reading of our paper led to the examination, in Section 4.4, of the relationship between our Poincaré duality and the Baum-Connes conjecture. We are also grateful to the referee for suggesting the simplified proof of Theorem 4.1 presented here.

1. STATEMENT OF RESULTS

Let W be a finite group acting isometrically with a global fixed point on a flat Riemannian torus T , and let \mathfrak{t} denote the universal cover of T . The notation reflects the observation that T can be equipped with the structure of an abelian Lie group with identity at the fixed point, and \mathfrak{t} is then its Lie algebra which inherits a linear isometric action of W . Denote by Γ the lattice in \mathfrak{t} mapping to the identity in T , or equivalently the fundamental group of T . This inherits an action of W from \mathfrak{t} .

Now let T^\vee be the dual torus of T , that is, the group of characters of Γ . We similarly denote by \mathfrak{t}^* the Lie algebra of T^\vee (which is the dual space of \mathfrak{t}) and denote by Γ^\vee the fundamental group of T^\vee . The action of W on T induces dual actions on T^\vee , \mathfrak{t}^* and Γ^\vee .

Let $\mathcal{P} \in KK_W(\mathbb{C}, C(T) \widehat{\otimes} C(T^\vee))$ denote the class of the Poincaré line bundle. To construct our Poincaré duality we will, in Section 3.2, define an element $\mathcal{Q} \in KK_W(C(T^\vee) \widehat{\otimes} C(T), \mathbb{C})$ given by a triple $(L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}, \rho, \mathcal{Q}_0)$, for which \mathcal{P}, \mathcal{Q} is a Poincaré duality pair. The operator is

$$Q_0 = \frac{\partial}{\partial y^j} \otimes \varepsilon^j - 2\pi i y^j \otimes \mathbf{e}_j$$

where $\{\varepsilon^j, \mathbf{e}_j : j = 1, \dots, n\}$ denotes a suitable basis for $\mathfrak{t}^* \times \mathfrak{t}$ acting on a space of spinors \mathcal{S} . The representation ρ is defined by

$$\rho\left(\sum_{\gamma \in \Gamma} a_\gamma e^{2\pi i \langle \eta, \gamma \rangle} \otimes f\right) \xi \otimes s = \sum_{\gamma \in \Gamma} a_\gamma \gamma \cdot (\tilde{f} \xi) \otimes s$$

where γ acts by translation on $L^2(\mathfrak{t})$, \tilde{f} denotes the lift of f to a periodic function on \mathfrak{t} and η denotes a variable in \mathfrak{t}^* .

Theorem 1.1. *Let T be a torus with flat Riemannian metric and T^\vee its dual. Suppose that W is a finite group acting isometrically on T with a global fixed point. The elements \mathcal{P}, \mathcal{Q} define a W -equivariant Poincaré duality in KK -theory from $C(T)$ to $C(T^\vee)$ and there is a ‘descended’ non-equivariant*

Poincaré duality from $C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W)$ to $C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)$. This is natural in the sense that there is a commutative diagram

$$\begin{array}{ccc} KK_W^*(C(T), \mathbb{C}) & \xrightarrow{\cong} & KK_W^*(\mathbb{C}, C(T^\vee)). \\ \downarrow \cong & & \cong \downarrow \\ KK^*(C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W), \mathbb{C}) & \xrightarrow{\cong} & KK^*(\mathbb{C}, C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)) \end{array}$$

where

- the top and bottom maps are induced by the Poincaré dualities,
- the left-hand map is the composition of the W -equivariant Morita equivalence $C(T) \sim C_0(\mathfrak{t}) \rtimes \Gamma$ with the dual Green-Julg isomorphism in K -homology,
- the right-hand map is its dual, i.e. the composition of the Morita equivalence $C(T^\vee) \sim C_0(\mathfrak{t}^*) \rtimes \Gamma^\vee$ with the Green-Julg isomorphism in K -theory.

The vertical maps factor through $KK(C(T) \rtimes W, \mathbb{C})$ on the left and $KK(\mathbb{C}, C(T^\vee) \rtimes W)$ on the right, and these may be identified (by Fourier-Pontryagin duality) with the groups $KK^*(C^*(\Gamma^\vee \rtimes W), \mathbb{C})$ and $KK^*(\mathbb{C}, C^*(\Gamma \rtimes W))$ respectively.

Theorem 1.2. *Let T be a torus with flat Riemannian metric and T^\vee its dual. Suppose that W is a finite group acting isometrically on T with a global fixed point. The Poincaré duality from $C(T)$ to $C(T^\vee)$ descends to give a non-equivariant Poincaré duality as follows.*

$$\begin{array}{ccc} KK_W^*(C(T), \mathbb{C}) & \xrightarrow{\cong} & KK_W^*(\mathbb{C}, C(T^\vee)). \\ \downarrow \cong & & \cong \downarrow \\ KK^*(C^*(\Gamma^\vee \rtimes W), \mathbb{C}) & \xrightarrow{\cong} & KK^*(\mathbb{C}, C^*(\Gamma \rtimes W)) \end{array}$$

where

- the top and bottom maps are induced by the Poincaré dualities,
- the left-hand map is the composition of the W -equivariant Fourier-Pontryagin duality $C(T) \cong C^*(\Gamma^\vee)$ with the dual Green-Julg isomorphism in K -homology,
- the right-hand map is its dual, i.e. the composition of the W -equivariant Fourier-Pontryagin duality $C(T^\vee) \cong C^*(\Gamma)$ with the Green-Julg isomorphism in K -theory.

In Section 4.4 we turn to the question of the relationship between the Baum-Connes assembly map and our Poincaré duality. In particular, we show that the following diagram of isomorphisms commutes.

$$\begin{array}{ccc} KK_{\Gamma \rtimes W}^*(C_0(\mathfrak{t}), \mathbb{C}) & \xrightarrow{\text{Baum-Connes}} & KK^*(\mathbb{C}, C^*(\Gamma \rtimes W)) \\ \downarrow \text{dual Green-Julg} & & \downarrow \text{Morita equivalence} \\ KK^*(C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W), \mathbb{C}) & \xrightarrow{\text{Poincaré duality}} & KK^*(\mathbb{C}, C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)) \end{array}$$

Given the definitions of the maps this is, in some sense surprising since both the Baum-Connes and the dual Green-Julg maps factor through the

descent map, which has target $KK(\mathbb{C}, C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)) \otimes C^*(\Gamma \rtimes W)$. The corresponding square with this latter group in the top left corner as illustrated in Section 4.4, does not commute.

A case of particular interest is provided by the action of a Weyl group W on a torus, provided by a root datum (X^*, R, X_*, R^\vee) . Let $W'_a = X_* \rtimes W$ be the corresponding extended affine Weyl group. The Langlands dual root system (X_*, R^\vee, X^*, R) gives rise to a dual extended affine Weyl group $(W'_a)^\vee = X^* \rtimes W$, which is not usually isomorphic to W'_a . However the Poincaré duality in Theorem 1.2 provides an isomorphism between $K^*(C^*((W'_a)^\vee))$ and $K_*(C^*(W'_a))$.

The Langlands duality between W'_a and $(W'_a)^\vee$ is further amplified by the following theorem.

Corollary 1.3. *Let G be a compact connected semisimple Lie group and G^\vee its Langlands dual, with $W'_a, (W'_a)^\vee$ the corresponding extended affine Weyl groups. Then there is a rational isomorphism*

$$K_*(C^*((W'_a)^\vee)) \cong K_*(C^*(W'_a)).$$

In particular we obtain a duality between affine and extended affine Weyl groups of the following form:

Corollary 1.4. *Let W'_a be the extended affine Weyl group of G , and let W_a, W_a^\vee be the affine Weyl groups of G and its Langlands dual G^\vee . If G is of adjoint type then rationally*

$$K_*(C^*(W_a^\vee)) \cong K_*(C^*(W'_a)).$$

If additionally G is of type $A_n, D_n, E_6, E_7, E_8, F_4, G_2$ then rationally

$$K_*(C^*(W_a)) \cong K_*(C^*(W'_a)).$$

Recall that the extended affine Weyl group W'_a is an extension of W_a by the cyclic group $\pi_1(G)$ so the content of Corollary 1.4 is that, surprisingly, this particular extension does not change the K -theory.

In a companion paper [12] we explore this phenomenon in further detail and give detailed computations of these K -theory groups in a number of cases.

2. BACKGROUND

2.1. Real Langlands duality. Recall that a connected complex reductive Lie group \mathbf{H} is classified by its root datum. That is a 4-tuple (X^*, R, X_*, R^\vee) where X^* is the lattice of characters on a maximal torus in \mathbf{H} , and X_* is the lattice of co-characters, or equivalently the fundamental group of the maximal torus. The set of roots $R \subset X^*$ is in bijection with the reflections in the Weyl group W and in bijection with the set of coroots $R^\vee \subset X_*$. Root data classify connected complex reductive Lie groups, in the sense that two such groups are isomorphic if and only if their root data are isomorphic (in the obvious sense). The Langlands dual of \mathbf{H} , denoted \mathbf{H}^\vee is then the unique connected complex reductive Lie group associated to the dual root datum (X_*, R^\vee, X^*, R) . See [4] for details.

One of the key motivations of this paper is that for extended affine Weyl groups the Baum-Connes correspondence should be thought of as an equivariant duality between maximal tori in a compact connected semisimple Lie group and its real Langlands dual. As in the complex case these are classified by their root data, and we can define the (real) Langlands dual by dualising the root datum as before. Since the real case is not as well known we recall the relationship with the complex case.

For a Lie group G , the *complexification* $G_{\mathbb{C}}$ is a complex Lie group together with a morphism from G , satisfying the universal property that for any morphism of G into a complex Lie group \mathbf{L} there is a unique factorisation through $G_{\mathbb{C}}$.

For T a maximal torus in G , the complexification $\mathbf{S} := T_{\mathbb{C}}$ of T is a maximal torus in $\mathbf{H} := G_{\mathbb{C}}$, and so the dual torus \mathbf{S}^{\vee} is well-defined in the dual group \mathbf{H}^{\vee} . Then T^{\vee} is defined to be the maximal compact subgroup of \mathbf{S}^{\vee} , and satisfies the condition

$$(T^{\vee})_{\mathbb{C}} = \mathbf{S}^{\vee}.$$

The groups X^*, X_* in the root datum are again the groups of characters and co-characters of the torus T respectively. Dually X_*, X^* are the groups of characters and co-characters on the dual torus T^{\vee} , giving the T -duality equation

$$(1) \quad X^*(T^{\vee}) = X_*(T).$$

The torus T^{\vee} is given explicitly by $T^{\vee} = \text{Hom}(X_*(T), \mathbf{U})$. The *Langlands dual* of G , denoted G^{\vee} , is defined to be a maximal compact subgroup of \mathbf{H}^{\vee} containing the torus T^{\vee} .

The process of passing to a maximal compact subgroup is inverse to complexification in the sense that complexifying G^{\vee} recovers \mathbf{H}^{\vee} .

2.1.1. *A table of Langlands dual groups.* Given a compact connected semisimple Lie group G , the product $|\pi_1(G)| \cdot |\mathcal{Z}(G)|$ is unchanged by Langlands duality, i.e. it agrees with the product $|\pi_1(G^{\vee})| \cdot |\mathcal{Z}(G^{\vee})|$. This product is equal to the *connection index*, denoted f , (see [4, IX, p.320]), which is defined in [4, VI, p.240]. The connection indices are listed in [4, VI, Plates I–X, p.265–292].

The following is a table of Langlands duals and connection indices for compact connected semisimple groups:

G	G^{\vee}	f
$A_n = \text{SU}_{n+1}$	PSU_{n+1}	$n + 1$
$B_n = \text{SO}_{2n+1}$	Sp_{2n}	2
$C_n = \text{Sp}_{2n}$	SO_{2n+1}	2
$D_n = \text{SO}_{2n}$	SO_{2n}	4
E_6	E_6	3
E_7	E_7	2
E_8	E_8	1
F_4	F_4	1
G_2	G_2	1

In this table, the simply-connected form of E_6 (resp. E_7) dualises to the adjoint form of E_6 (resp. E_7).

The Lie group G and its dual G^\vee admit a common Weyl group

$$W = N(T)/T = N(T^\vee)/T^\vee.$$

The T -duality equation (1) identifies the action of the Weyl group of T on $X_*(T)$ with the dual action of the Weyl group of T^\vee on $X^*(T^\vee)$.

Remark 2.1. In general, T and T^\vee are *not* isomorphic as W -spaces. For example, let $G = \mathrm{SU}_3$ and take $T = \{(z_1, z_2, z_3) : z_j \in \mathbf{U}, z_1 z_2 z_3 = 1\}$. Then in homogeneous coordinates we have $T^\vee = \{(z_1 : z_2 : z_3) : z_j \in \mathbf{U}, z_1 z_2 z_3 = 1\}$. The Weyl group W is the symmetric group S_3 which acts by permuting coordinates in both cases. Note that the torus T admits three W -fixed points whereas the unique W -fixed point in T^\vee is the identity $(1 : 1 : 1) \in T^\vee$, hence T and T^\vee are not W -equivariantly isomorphic.

The *nodal group* of the torus T is defined to be $\Gamma(T) := \ker(\exp : \mathfrak{t} \rightarrow T)$ and differentiating the action of the Weyl group W we obtain a linear action of W on the Lie algebra \mathfrak{t} which restricts to an action on the nodal group $\Gamma(T)$. Indeed there is a W -equivariant isomorphism $X_*(T) \cong \Gamma(T)$.

We remark that by definition T^\vee is the Pontryagin dual of the nodal group $\Gamma(T)$. Moreover the natural action of W on T^\vee is the dual of the action on $\Gamma(T)$. Hence we have the following:

Lemma 2.2. *Let $\widehat{\Gamma}$ denote the Pontryagin dual of $\Gamma = \Gamma(T)$. Then we have a W -equivariant isomorphism*

$$\widehat{\Gamma} \cong T^\vee.$$

and hence an isomorphism of W - C^* -algebras

$$C^*(\Gamma) \cong C(T^\vee).$$

2.2. Affine and Extended Affine Weyl groups. In this section we will give the definitions of the affine and extended affine Weyl groups of a compact connected semisimple Lie group. As noted in the introduction these are semidirect products of lattices in the Lie algebra \mathfrak{t} of a maximal torus T by the Weyl group W . The affine Weyl group W_a is a Coxeter group while the extended affine Weyl group contains W_a as a finite index normal subgroup and the quotient is the fundamental group of the Lie group G .

Let $p : \widetilde{G} \rightarrow G$ denote the universal cover and let \widetilde{T} be the preimage of T which is a maximal torus in \widetilde{G} . We consider the following commutative diagram.

$$\begin{array}{ccccccc} \Gamma(\widetilde{T}) & \longrightarrow & \mathfrak{t} & \longrightarrow & \widetilde{T} & \longrightarrow & 0 \\ & & \downarrow \iota & & \downarrow \mathrm{id} & & \downarrow p|_{\widetilde{T}} \\ 0 & \longrightarrow & \Gamma(T) & \longrightarrow & \mathfrak{t} & \longrightarrow & T \end{array}$$

By the snake lemma the sequence

$$\begin{array}{ccccccc} \ker(\mathrm{id}) & \rightarrow & \ker(p|_{\widetilde{T}}) & \rightarrow & \mathrm{coker}(\iota) & \rightarrow & \mathrm{coker}(\mathrm{id}) \\ \parallel & & \parallel & & \parallel & & \parallel \\ 0 & & \pi_1(G) & & \Gamma(T)/\Gamma(\widetilde{T}) & & 0 \end{array}$$

is exact, hence $\Gamma(T)/\Gamma(\widetilde{T})$ is isomorphic to $\pi_1(G)$. We thus have a map from $\Gamma(T)$ onto $\pi_1(G)$. The kernel of this map (more commonly denoted $N(G, T)$) is the nodal lattice $\Gamma(\widetilde{T})$ for the torus \widetilde{T} and we have:

Definition 2.3. The *affine Weyl group of G* is

$$W_a(G) = \Gamma(\tilde{T}) \rtimes W$$

and the *extended affine Weyl group of G* is

$$W'_a(G) = \Gamma(T) \rtimes W$$

where W denotes the Weyl group of G .

The following is now immediate.

Lemma 2.4. *Let \tilde{G} denote the universal cover of G and let \tilde{T} denote a maximal torus in \tilde{G} . Then we have*

$$W_a(G) = W'_a(\tilde{G}) = W_a(\tilde{G}).$$

We remark that the extended affine Weyl group $W'_a(G)$ is a split extension of $W_a(G)$ by $\pi_1(G)$.

3. EQUIVARIANT POINCARÉ DUALITY BETWEEN $C(T)$ AND $C(T^\vee)$

We begin by recalling the general framework of Poincaré duality in KK -theory. For \mathfrak{G} - C^* -algebras A, B a Poincaré duality is given by elements $\mathfrak{a} \in KK_{\mathfrak{G}}(B \hat{\otimes} A, \mathbb{C})$ and $\mathfrak{b} \in KK_{\mathfrak{G}}(\mathbb{C}, A \hat{\otimes} B)$ with the property that

$$(2) \quad \begin{aligned} \mathfrak{b} \otimes_A \mathfrak{a} &= 1_B \in KK_{\mathfrak{G}}(B, B) \\ \mathfrak{b} \otimes_B \mathfrak{a} &= 1_A \in KK_{\mathfrak{G}}(A, A). \end{aligned}$$

These then yield isomorphisms between the K -homology of A and the K -theory of B (and vice versa) given by

$$\begin{aligned} \mathfrak{r} &\mapsto \mathfrak{b} \otimes_A \mathfrak{r} \in KK_{\mathfrak{G}}(\mathbb{C}, B) \text{ for } \mathfrak{r} \in KK(A, \mathbb{C}) \\ \mathfrak{\eta} &\mapsto \mathfrak{\eta} \otimes_B \mathfrak{a} \in KK_{\mathfrak{G}}(A, \mathbb{C}) \text{ for } \mathfrak{\eta} \in KK(\mathbb{C}, B). \end{aligned}$$

Throughout this section T will denote a torus with flat Riemannian metric, T^\vee its dual torus and W a finite group acting by isometries on T (and dually on T^\vee). We will construct elements $\mathcal{Q} \in KK_W(C(T^\vee) \hat{\otimes} C(T), \mathbb{C})$ and $\mathcal{P} \in KK_W(\mathbb{C}, C(T) \hat{\otimes} C(T^\vee))$ satisfying 2.

3.1. The Poincaré line bundle. Recall that the Poincaré line bundle over $T \times T^\vee$ is the bundle with total space given by the quotient of $\mathfrak{t} \times T^\vee \times \mathbb{C}$ under the action of Γ defined by $\gamma(x, \chi, z) = (\gamma + x, \chi, \chi(\gamma)z)$. The projection onto the base $T \times T^\vee$ maps the Γ orbit of (x, χ, z) to the Γ orbit of (x, χ) . Here we are identifying elements of T^\vee with characters on Γ . We note that the bundle is W -equivariant with respect to the diagonal action of W on $\mathfrak{t} \times T^\vee$, hence it defines an element in W -equivariant K -theory allowing it to play the role of the element \mathcal{P} in our Poincaré duality.

To place this in the language of KK -theory we consider sections of this bundle, which are given by functions $\sigma : \mathfrak{t} \times T^\vee \rightarrow \mathbb{C}$ such that $\sigma(\gamma + x, \chi) = \chi(\gamma)\sigma(x, \chi)$. They naturally form a module over $C(T \times T^\vee)$ and given two such sections we define $\langle \sigma_1, \sigma_2 \rangle = \overline{\sigma_1} \sigma_2$. We note that this is a Γ periodic function in the first variable, hence the inner product takes values in $C(T \times T^\vee)$, giving the space of sections the structure of a Hilbert module.

We will now give an alternative construction of this Hilbert module. Let $C_c(\mathfrak{t})$ denote the space of continuous compactly supported functions on \mathfrak{t} and equip this with a $C(T) \otimes C(T^\vee)$ -valued inner product defined by

$$\langle \phi_1, \phi_2 \rangle(x, \eta) = \sum_{\alpha, \beta \in \Gamma} \overline{\phi_1(x - \alpha)} \phi_2(x - \beta) e^{2\pi i \langle \eta, \beta - \alpha \rangle}.$$

We remark that the support condition ensures that this is a finite sum, and that it is easy to check that $\langle \phi_1, \phi_2 \rangle(x, \eta)$ is Γ -periodic in x and Γ^\vee -periodic in η .

The space $C_c(\mathfrak{t})$ has a $C(T) \otimes \mathbb{C}[\Gamma]$ -module structure

$$(\phi \cdot (f \otimes [\gamma])) = \phi(x + \gamma) \tilde{f}(x)$$

where we view the function f in $C(T)$ as a Γ -periodic function \tilde{f} on \mathfrak{t} .

Completing $C_c(\mathfrak{t})$ with respect to the norm arising from the inner product, the module structure extends by continuity to give $\overline{C_c(\mathfrak{t})}$ the structure of a $C(T) \widehat{\otimes} C^*(\Gamma) \cong C(T) \widehat{\otimes} C(T^\vee)$ Hilbert module. We denote this Hilbert module by \mathcal{E} and give this the trivial grading.

The group W acts on \mathfrak{t} and hence on $C_c(\mathfrak{t})$ by $(w \cdot \phi)(x) = \phi(w^{-1}x)$. We have

$$(w \cdot (\phi \cdot (f \otimes [\gamma]))) (x) = \phi(w^{-1}x + \gamma) \tilde{f}(w^{-1}x) = ((w \cdot \phi) \cdot (w \cdot f \otimes [w\gamma]))(x)$$

so the action is compatible with the module structure. Now for the inner product we have

$$\begin{aligned} \langle w \cdot \phi_1, w \cdot \phi_2 \rangle(x, \eta) &= \sum_{\alpha, \beta \in \Gamma} \overline{(w \cdot \phi_1)(x - \alpha)} (w \cdot \phi_2)(x - \beta) e^{2\pi i \langle \eta, \beta - \alpha \rangle} \\ &= \sum_{\alpha, \beta \in \Gamma} \overline{\phi_1(w^{-1}x - w^{-1}\alpha)} \phi_2(w^{-1}x - w^{-1}\beta) e^{2\pi i \langle \eta, \beta - \alpha \rangle} \\ &= \sum_{\alpha', \beta' \in \Gamma} \overline{\phi_1(w^{-1}x - \alpha')} \phi_2(w^{-1}x - \beta') e^{2\pi i \langle \eta, w(\beta' - \alpha') \rangle} \\ &= \sum_{\alpha', \beta' \in \Gamma} \overline{\phi_1(w^{-1}x - \alpha')} \phi_2(w^{-1}x - \beta') e^{2\pi i \langle w^{-1}\eta, \beta' - \alpha' \rangle} \\ &= (w \cdot \langle \phi_1, \phi_2 \rangle)(x, \eta). \end{aligned}$$

Hence \mathcal{E} is a W -equivariant Hilbert module.

The identification of the module \mathcal{E} with the sections of the Poincaré line bundle is given by the following analog of the Fourier transform. For each element $\phi \in C_c(\mathfrak{t})$ set

$$\sigma(x, \chi) = \sum_{\gamma \in \Gamma} \phi(x - \gamma) \chi(\gamma).$$

It is routine to verify that $\sigma(x + \delta, \chi) = \chi(\delta) \sigma(x, \chi)$ hence σ is a section of the Poincaré line bundle, and that the W action on $C_c(\mathfrak{t})$ corresponds precisely to the W action on the bundle.

Theorem 3.1. *The triple $(\mathcal{E}, 1, 0)$, where 1 denotes the identity representation of \mathbb{C} on \mathcal{E} , is a W -equivariant Kasparov triple defining an element \mathcal{P} in $KK_W(\mathbb{C}, C(T) \widehat{\otimes} C(T^\vee))$.*

We remark that there is a connection with Fourier-Mukai duality. We recall that Fourier-Mukai duality is given by the map

$$\mathfrak{t} \mapsto p_{2*}(p_1^*\mathfrak{t} \otimes \mathcal{P}),$$

where p_1, p_2 are the projections of $T \times T^\vee$ onto the first and second factors. From the point of view of K -theory the subtlety is to interpret the wrong-way map p_{2*} . This should give a map from the W -equivariant K -theory of $T \times T^\vee$ to the W equivariant K -theory of T^\vee , but to make this well defined we must twist by the Clifford algebra $\mathcal{C}\ell(\mathfrak{t})$. Specifically we can take

$$p_{2*} := [D] \otimes 1_{C(T^\vee)} \in KK_W(C(T \times T^\vee) \otimes \mathcal{C}\ell(\mathfrak{t}), C(T^\vee)),$$

where $[D]$ is the Dirac class in $KK_W(C(T) \otimes \mathcal{C}\ell(\mathfrak{t}), \mathbb{C})$. The Fourier-Mukai map is then given by taking the Kasparov product with the element $p_1^*P i^* p_{2*} = \mathcal{P} p_1^* i^* p_{2*}$ where i is the diagonal inclusion of $T \times T^\vee$ into $(T \times T^\vee)^2$. We note that $p_1^* i^* p_{2*}$ is the tensor product of Kasparov's Poincaré duality element for T (given by its Dirac element) with the identity on $C(T^\vee)$.

3.2. Construction of the element \mathcal{Q} in $KK_W(C(T^\vee) \widehat{\otimes} C(T), \mathbb{C})$. We consider the differential operator Q_0 on \mathfrak{t} with coefficients in the Clifford algebra $\mathcal{C}\ell(\mathfrak{t} \times \mathfrak{t}^*)$ defined using Einstein summation convention by

$$Q_0 = \frac{\partial}{\partial y^j} \otimes \varepsilon^j - 2\pi i y^j \otimes \mathbf{e}_j.$$

Here $\{\mathbf{e}_j = \frac{\partial}{\partial y^j}\}$ is an orthonormal basis for \mathfrak{t} , $\{\varepsilon^j\}$ denotes the dual basis of \mathfrak{t}^* and we regard these as generators of the Clifford algebra $\mathcal{C}\ell(\mathfrak{t} \times \mathfrak{t}^*)$.

We view Q_0 as an unbounded operator on the Hilbert space $L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}$, where \mathcal{S} denotes the space of spinors $\mathcal{S} = \mathcal{C}\ell(\mathfrak{t} \times \mathfrak{t}^*)P$ with P the projection $\prod_j \frac{1}{2}(1 - i\mathbf{e}_j \varepsilon^j)$ in the Clifford algebra. (The space \mathcal{S} is naturally equipped with a representation of $\mathcal{C}\ell(\mathfrak{t} \times \mathfrak{t}^*)$ by left multiplication inducing the action of Q_0 .)

The subtlety in constructing an element in equivariant KK -theory is the need to ensure that P is W -invariant with respect to the diagonal action of W on $\mathfrak{t} \times \mathfrak{t}^*$ and hence that the action of W on $\mathcal{C}\ell(\mathfrak{t} \times \mathfrak{t}^*)$ restricts to a representation on \mathcal{S} . The corner algebra $P\mathcal{C}\ell(\mathfrak{t} \times \mathfrak{t}^*)P$ is $\mathbb{C}P$, which we identify with \mathbb{C} , and this gives \mathcal{S} a canonical inner product given by $\langle aP, bP \rangle = Pa^*bP$.

As a simple example consider the 1-dimensional case. Here $\mathcal{C}\ell(\mathfrak{t} \times \mathfrak{t}^*) = M_2(\mathbb{C})$ and the two generators are $\mathbf{e}_1 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$ and $\varepsilon^1 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. The

projection P is therefore $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ so \mathcal{S} is the space of matrices of the form

$\begin{pmatrix} * & 0 \\ * & 0 \end{pmatrix}$ and the operator is

$$Q_0 = \begin{pmatrix} 0 & -\frac{\partial}{\partial y^1} + 2\pi y^1 \\ \frac{\partial}{\partial y^1} + 2\pi y^1 & 0 \end{pmatrix}.$$

The off-diagonal elements are of course the 1-dimensional annihilation and creation operators.

Returning to the general case we must now construct a representation of $C(T^\vee) \widehat{\otimes} C(T)$ on $L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}$. It suffices to define commuting representations of $C(T^\vee) \widehat{\otimes} 1$ and $1 \widehat{\otimes} C(T)$. The representation of $C(T)$ is the usual pointwise multiplication on $L^2(\mathfrak{t})$ viewing elements of $C(T)$ as Γ -periodic functions on \mathfrak{t} . The representation of $C(T^\vee)$ involves the action of Γ on \mathfrak{t} .

For a an affine isometry of \mathfrak{t} , let L_a be the operator on $L^2(\mathfrak{t})$ induced by the action of a on \mathfrak{t} :

$$(L_a \xi)(y) = \xi(a^{-1} \cdot y).$$

For the function $\eta \mapsto e^{2\pi i \langle \eta, \gamma \rangle}$ in $C(T^\vee)$ we define

$$\rho(e^{2\pi i \langle \eta, \gamma \rangle}) = L_\gamma \otimes 1_{\mathcal{S}}.$$

Identifying $C(T^\vee)$ with $C^*(\Gamma)$ and identifying $L^2(\mathfrak{t})$ with $\ell^2(\Gamma) \otimes L^2(X)$ where X is a fundamental domain for the action of Γ , the representation of the algebra is given by the left regular representation on $\ell^2(\Gamma)$.

Consider the commutators of Q_0 with the representation ρ . For $f \in C(T)$, the operator $\rho(f)$ commutes exactly with the second term $2\pi i y^j \otimes \mathbf{e}_j$ in Q_0 , while, for f smooth, the commutator of $\rho(f)$ with $\frac{\partial}{\partial y^j} \otimes \varepsilon^j$ is given by the bounded operator $\frac{\partial f}{\partial y^j} \otimes \varepsilon^j$. Now for the function $\eta \mapsto e^{2\pi i \langle \eta, \gamma \rangle}$ in $C(T^\vee)$ we have $\rho(e^{2\pi i \langle \eta, \gamma \rangle}) = L_\gamma \otimes 1_{\mathcal{S}}$. This commutes exactly with the differential term of the operator, while

$$L_\gamma(2\pi i y^j) L_\gamma^* = 2\pi i (y^j - \gamma^j)$$

hence the commutator $[L_\gamma \otimes 1_{\mathcal{S}}, 2\pi i y^j \otimes \mathbf{e}_j]$ is again bounded.

We have verified that Q_0 commutes with the representation ρ modulo bounded operators, on a dense subalgebra of $C(T^\vee) \widehat{\otimes} C(T)$. Thus to show that the triple

$$(L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}, \rho, Q_0)$$

is an unbounded Kasparov triple it remains to prove the following.

Theorem 3.2. *The operator Q_0 has compact resolvent. It has 1-dimensional kernel with even grading.*

Proof. In the following argument we will *not* use summation convention. We consider the following operators on $L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}$:

$$\begin{aligned} p_j &= \frac{\partial}{\partial y^j} \otimes \varepsilon^j \\ x_j &= -2\pi i y^j \otimes \mathbf{e}_j \\ q_j &= \frac{1}{2}(1 + 1 \otimes i\mathbf{e}_j \varepsilon^j) \\ A_j &= \frac{1}{2\sqrt{\pi}}(p_j + x_j) \end{aligned}$$

Since A_j anti-commutes with $1 \otimes i\mathbf{e}_j \varepsilon^j$ we have $q_j A_j = A_j(1 - q_j)$, hence we can think of A_j as an off-diagonal matrix with respect to q_j . We write A_j as $a_j + a_j^*$ where $a_j = q_j A_j = A_j(1 - q_j)$ and hence $a_j^* = A_j q_j = (1 - q_j) A_j$. We think of a_j^* and a_j as creation and annihilation operators respectively and we define a number operator $N_j = a_j^* a_j$. The involution $i\varepsilon^j$ intertwines

q_j with $1 - q_j$. We define A'_j, N'_j to be the conjugates of A_j, N_j respectively by $i\varepsilon^j$. Note that

$$A'_j = \frac{1}{2\sqrt{\pi}}(p_j - x_j)$$

and hence

$$A_j^2 = (A'_j)^2 + 2\frac{1}{4\pi}[x_j, p_j] = (A'_j)^2 + 1 \otimes i\mathbf{e}_j\varepsilon^j.$$

We have $N'_j = A'_j(1 - q_j)A'_j = q_j(A'_j)^2$. Thus

$$a_j a_j^* = q_j A_j^2 q_j = q_j A_j^2 = q_j (A'_j)^2 + q_j (1 \otimes i\mathbf{e}_j\varepsilon^j) = N'_j + q_j.$$

Hence the spectrum of $a_j a_j^*$ (viewed as an operator on the range of q_j) is the spectrum of N'_j shifted by 1. However N'_j is conjugate to $N_j = a_j^* a_j$ so we conclude that

$$\mathrm{Sp}(a_j a_j^*) = \mathrm{Sp}(a_j^* a_j) + 1.$$

But $\mathrm{Sp}(a_j a_j^*) \setminus \{0\} = \mathrm{Sp}(a_j^* a_j) \setminus \{0\}$ so we conclude that the spectrum is $\mathrm{Sp}(a_j^* a_j) = \{0, 1, 2, \dots\}$ while $\mathrm{Sp}(a_j a_j^*) = \{1, 2, \dots\}$.

Now since the operators A_j pairwise gradedly commute we have

$$Q_0^2 = 4\pi \sum_j A_j^2 = 4\pi \sum_j a_j^* a_j + a_j a_j^*$$

and noting that the summands commute we see that Q_0^2 has discrete spectrum. To show that $(1 + Q_0^2)^{-1}$ is compact, it remains to verify that $\ker Q_0$ is finite dimensional (and hence that all eigenspaces are finite dimensional). We have

$$\ker Q_0 = \ker Q_0^2 = \bigcap_j \ker A_j^2 = \bigcap_j \ker A_j.$$

Multiplying the differential equation $(p_j + x_j)f = 0$ by $-\exp(\pi(y^j)^2 \otimes i\varepsilon^j \mathbf{e}_j)\varepsilon^j$ we see that the kernel of A_j is the space of solutions of the differential equation

$$\frac{\partial}{\partial y^j} (\exp(\pi(y^j)^2 \otimes i\varepsilon^j \mathbf{e}_j)f) = 0$$

whence for f in the kernel we have

$$f(y^1, \dots, y^n) = \exp(-\pi(y^j)^2 \otimes i\varepsilon^j \mathbf{e}_j) f(y^1, \dots, y^{j-1}, 0, y^{j+1}, \dots, y^n).$$

Since the solutions must be square integrable the values of f must lie in the $+1$ eigenspace of the involution $i\varepsilon^j \mathbf{e}_j$, that is, the range of the projection $1 - q_j$. On this subspace the operator $\exp(-\pi(y^j)^2 \otimes i\varepsilon^j \mathbf{e}_j)$ reduces to $e^{-\pi(y^j)^2}(1 - q_j)$. Since the kernel of Q_0 is the intersection of the kernels of the operators A_j an element of the kernel must have the form

$$f(y) = e^{-\pi|y|^2} \prod_j (1 - q_j) f(0)$$

so the kernel is 1-dimensional. Indeed the product $\prod_j (1 - q_j)$ is the projection P used to define the space of spinors $\mathcal{S} = \mathcal{C}\ell(\mathfrak{t} \times \mathfrak{t}^*)P$, and hence $\prod_j (1 - q_j) f(0)$ lies in the 1-dimensional space $P\mathcal{S} = P\mathcal{C}\ell(\mathfrak{t} \times \mathfrak{t}^*)P$ which has even grading. \square

We have shown that $(L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}, \rho, Q_0)$ defines an unbounded Kasparov triple. It remains to establish W -equivariance.

Let V be a finite dimensional vector space and equip $V \otimes V^*$ with the natural diagonal action of $\mathrm{GL}(V)$. If V is equipped with a non-degenerate symmetric bilinear form g then we can form the Clifford algebra $\mathcal{C}\ell(V)$ and dually $\mathcal{C}\ell(V^*)$. The subgroup $\mathrm{O}(g)$ of $\mathrm{GL}(V)$, consisting of those elements preserving g , acts diagonally on $\mathcal{C}\ell(V) \widehat{\otimes} \mathcal{C}\ell(V^*)$ which we identify with $\mathcal{C}\ell(V \times V^*)$.

We say that an element a of $\mathcal{C}\ell(V \times V^*)$ is *symmetric* if there exists a g -orthonormal¹ basis $\{\mathbf{e}_j : j = 1, \dots, n\}$ with dual basis $\{\varepsilon^j : j = 1, \dots, n\}$ such that a can be written as $p(\mathbf{e}_1 \varepsilon^1, \dots, \mathbf{e}_n \varepsilon^n)$ where $p(x_1, \dots, x_n)$ is a symmetric polynomial.

Proposition 3.3. *For any basis $\{\mathbf{e}_j\}$ of V with dual basis $\{\varepsilon^j\}$ for V^* , the Einstein sum $\mathbf{e}_j \otimes \varepsilon^j$ in $V \otimes V^*$ is $\mathrm{GL}(V)$ -invariant.*

Suppose moreover that V is equipped with a non-degenerate symmetric bilinear form g and that the underlying field has characteristic zero. Then every symmetric element of $\mathcal{C}\ell(V) \widehat{\otimes} \mathcal{C}\ell(V^) \cong \mathcal{C}\ell(V \times V^*)$ is $\mathrm{O}(g)$ -invariant.*

Proof. Identifying $V \otimes V^*$ with endomorphisms of V in the natural way, the action of $\mathrm{GL}(V)$ is the action by conjugation and $\mathbf{e}_j \otimes \varepsilon^j$ is the identity which is invariant under conjugation.

For the second part, over a field of characteristic zero the symmetric polynomials are generated by power sum symmetric polynomials $p(x_1, \dots, x_n) = x_1^k + \dots + x_n^k$, so it suffices to consider

$$\begin{aligned} p(\mathbf{e}_1 \varepsilon^1, \dots, \mathbf{e}_n \varepsilon^n) &= (\mathbf{e}_1 \varepsilon^1)^k + \dots + (\mathbf{e}_n \varepsilon^n)^k \\ &= (-1)^{k(k-1)/2} \left((\mathbf{e}_1)^k (\varepsilon^1)^k + \dots + (\mathbf{e}_n)^k (\varepsilon^n)^k \right). \end{aligned}$$

When k is even, writing $(\mathbf{e}_j)^k = (\mathbf{e}_j^2)^{k/2} = (g_{jj})^{k/2}$ and similarly $(\varepsilon^j)^k = (g^{jj})^{k/2}$, we see that each term $(\mathbf{e}_j)^k (\varepsilon^j)^k$ is 1 since $g_{jj} = g^{jj} = \pm 1$ for an orthonormal basis. Thus $p(\mathbf{e}_1 \varepsilon^1, \dots, \mathbf{e}_n \varepsilon^n) = n(-1)^{k(k-1)/2}$ which is invariant.

Similarly when k is odd we get $(\mathbf{e}_j)^k (\varepsilon^j)^k = \mathbf{e}_j \varepsilon^j$ so

$$p(\mathbf{e}_1 \varepsilon^1, \dots, \mathbf{e}_n \varepsilon^n) = (-1)^{k(k-1)/2} (\mathbf{e}_1 \varepsilon^1 + \dots + \mathbf{e}_n \varepsilon^n).$$

As the sum $\mathbf{e}_j \otimes \varepsilon^j$ in $V \otimes V^*$ is invariant under $\mathrm{GL}(V)$, it is in particular invariant under $\mathrm{O}(g)$, and hence the sum $\mathbf{e}_j \varepsilon^j$ is $\mathrm{O}(g)$ -invariant in the Clifford algebra. \square

Returning to our construction, the projection P is a symmetric element of the Clifford algebra and hence is W -invariant by Proposition 3.3. It follows that \mathcal{S} carries a representation of W . The space $L^2(\mathfrak{t})$ also carries a representation of W given by the action of W on \mathfrak{t} and we equip $L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}$ with the diagonal action of W .

To verify that the representation ρ is W -equivariant it suffices to consider the representations of $C(T)$ and $C(T^\vee)$ separately. As the exponential map

¹We say that $\{\mathbf{e}_j\}$ is g -orthonormal if $g_{jk} = \pm \delta_{jk}$ for each j, k .

$\mathfrak{t} \rightarrow T$ is W -equivariant it is clear that the representation of $C(T)$ on $L^2(\mathfrak{t})$ by pointwise multiplication is W -equivariant.

For $e^{2\pi i \langle \eta, \gamma \rangle} \in C(T^\vee)$ we have $w \cdot (e^{2\pi i \langle \eta, \gamma \rangle}) = e^{2\pi i \langle w^{-1} \cdot \eta, \gamma \rangle} = e^{2\pi i \langle \eta, w \cdot \gamma \rangle}$ thus $\rho(w \cdot (e^{2\pi i \langle \eta, \gamma \rangle})) = L_{w \cdot \gamma} \otimes 1_{\mathcal{S}} = L_w L_\gamma L_{w^{-1}} \otimes 1_{\mathcal{S}}$. Thus the representation of $C(T^\vee)$ is also W -equivariant.

It remains to check that the operator Q_0 is W -equivariant. By definition

$$Q_0 = \frac{\partial}{\partial y^j} \otimes \varepsilon^j - 2\pi i y^j \otimes \mathbf{e}_j.$$

Now by Proposition 3.3 $\frac{\partial}{\partial y^j} \otimes \varepsilon^j = \mathbf{e}_j \otimes \varepsilon^j$ is a $\mathrm{GL}(\mathfrak{t})$ -invariant element of $\mathfrak{t} \otimes \mathfrak{t}^*$ and so in particular it is W -invariant. Writing $y^j = \langle \varepsilon^j, y \rangle$ the W -invariance of the second term again follows from invariance of $\mathbf{e}_j \otimes \varepsilon^j$.

Hence we conclude the following.

Theorem 3.4. *The triple $(L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}, \rho, Q_0)$ constructed above defines an element \mathcal{Q} of $KK_W(C(T^\vee) \widehat{\otimes} C(T), \mathbb{C})$.*

3.3. The Kasparov product $\mathcal{P} \otimes_{C(T^\vee)} \mathcal{Q}$. We will compute the Kasparov product of the Poincaré line bundle $\mathcal{P} \in KK_W(\mathbb{C}, C(T) \widehat{\otimes} C(T^\vee))$ with our inverse $\mathcal{Q} \in KK_W(C(T^\vee) \widehat{\otimes} C(T), \mathbb{C})$ where the product is taken over $C(T^\vee)$ (not $C(T) \widehat{\otimes} C(T^\vee)$).

Recall that \mathcal{P} is given by the Kasparov triple $(\mathcal{E}, 1, 0)$ where \mathcal{E} is the completion of $C_c(\mathfrak{t})$ with the inner product

$$\langle \phi_1, \phi_2 \rangle(x, \eta) = \sum_{\alpha, \beta \in \Gamma} \overline{\phi_1(x - \alpha)} \phi_2(x - \beta) e^{2\pi i \langle \eta, \beta - \alpha \rangle}$$

in $C(T) \widehat{\otimes} C(T^\vee)$. As above \mathcal{Q} is given by the triple $(L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}, \rho, Q_0)$.

To form the Kasparov product we must take that tensor product of \mathcal{E} with $L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}$ over $C(T^\vee)$ and as the operator in the first triple is zero, the operator required for the Kasparov product can be any connection for Q_0 .

We note that the representation ρ is the identity on \mathcal{S} and hence

$$\mathcal{E} \widehat{\otimes}_{C(T^\vee)} (L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}) = (\mathcal{E} \widehat{\otimes}_{C(T^\vee)} L^2(\mathfrak{t})) \widehat{\otimes} \mathcal{S}.$$

Thus we can focus on identifying the tensor product $\mathcal{E} \widehat{\otimes}_{C(T^\vee)} L^2(\mathfrak{t})$. By abuse of notation we will also let ρ denote the representation of $C(T) \widehat{\otimes} C(T^\vee)$ on $L^2(\mathfrak{t})$.

As we are taking the tensor product over $C(T^\vee)$, not over $C(T) \widehat{\otimes} C(T^\vee)$, we are forming the Hilbert module

$$(\mathcal{E} \widehat{\otimes} C(T)) \widehat{\otimes}_{C(T) \widehat{\otimes} C(T^\vee) \widehat{\otimes} C(T)} (C(T) \widehat{\otimes} L^2(\mathfrak{t}))$$

however since the algebra $C(T)$ is unital, it suffices to consider elementary tensors of the form $(\phi \otimes 1) \otimes (1 \otimes \xi)$. Where there is no risk of confusion we will abbreviate these as $\phi \otimes \xi$

Let $\phi_1, \phi_2 \in C_c(\mathfrak{t})$ and let ξ_1, ξ_2 be elements of $L^2(\mathfrak{t})$. Then

$$\langle \phi_1 \otimes \xi_1, \phi_2 \otimes \xi_2 \rangle = \langle 1 \otimes \xi_1, (1 \otimes \rho)(\langle \phi_1, \phi_2 \rangle \otimes 1)(1 \otimes \xi_2) \rangle.$$

The operator $(1 \otimes \rho)(\langle \phi_1, \phi_2 \rangle \otimes 1)$ corresponds to a field of operators

$$\begin{aligned} (1 \otimes \rho)(\langle \phi_1, \phi_2 \rangle \otimes 1)(x) &= \sum_{\alpha, \beta \in \Gamma} \overline{\phi_1(x - \alpha)} \phi_2(x - \beta) \otimes \rho(e^{2\pi i \langle \eta, \beta - \alpha \rangle} \otimes 1) \\ &= \sum_{\alpha, \beta \in \Gamma} \overline{\phi_1(x - \alpha)} \phi_2(x - \beta) \otimes L_\alpha^* L_\beta \end{aligned}$$

and so

$$\begin{aligned} \langle \phi_1 \otimes \xi_1, \phi_2 \otimes \xi_2 \rangle(x) &= \sum_{\alpha, \beta \in \Gamma} \overline{\phi_1(x - \alpha)} \phi_2(x - \beta) \langle L_\alpha \xi_1, L_\beta \xi_2 \rangle \\ &= \left\langle \sum_{\alpha \in \Gamma} \phi_1(x - \alpha) L_\alpha \xi_1, \sum_{\beta \in \Gamma} \phi_2(x - \beta) L_\beta \xi_2 \right\rangle. \end{aligned}$$

We note that $x \mapsto \sum_{\alpha \in \Gamma} \phi_1(x - \alpha) L_\alpha \xi_1$ is a continuous Γ -equivariant (and hence bounded) function from \mathfrak{t} to $L^2(\mathfrak{t})$. Let $C(\mathfrak{t}, L^2(\mathfrak{t}))^\Gamma$ denote the space of such functions equipped with the $C(T)$ module structure of pointwise multiplication in the first variable and gives the pointwise inner product $\langle g_1, g_2 \rangle(x) = \langle g_1(x), g_2(x) \rangle$. We remark that equivariance implies this inner product is a Γ -periodic function on \mathfrak{t} .

The above calculation show that $\mathcal{E} \widehat{\otimes}_{C(T^\vee)} L^2(\mathfrak{t})$ maps isometrically into $C(\mathfrak{t}, L^2(\mathfrak{t}))^\Gamma$ via the map

$$\phi \otimes \xi \mapsto \sum_{\alpha \in \Gamma} \phi(x - \alpha) L_\alpha \xi.$$

Moreover this map is surjective. To see this, note that if ϕ is supported inside a single fundamental domain then for x in that fundamental domain we obtain the function $\phi(x)\xi$. This is extended by equivariance to a function on \mathfrak{t} , and using a partition of unity one can approximate an arbitrary element of $C(\mathfrak{t}, L^2(\mathfrak{t}))^\Gamma$ by sums of functions of this form.

We now remark that $C(\mathfrak{t}, L^2(\mathfrak{t}))^\Gamma$ is in fact isomorphic to the Hilbert module $C(T, L^2(\mathfrak{t}))$ via a change of variables. Given $g \in C(\mathfrak{t}, L^2(\mathfrak{t}))^\Gamma$, let $\tilde{h}(x) = L_{-x}g(x)$. The Γ -equivariance of g ensures that $g(\gamma + x) = L_\gamma g(x)$ whence

$$\tilde{h}(\gamma + x) = L_{-x-\gamma}g(\gamma + x) = L_{-x-\gamma}L_\gamma g(x) = L_{-x}g(x) = \tilde{h}(x).$$

As \tilde{h} is a Γ -periodic function from \mathfrak{t} to $L^2(\mathfrak{t})$ we identify it via the exponential map with the continuous function h from T to $L^2(\mathfrak{t})$ such that $\tilde{h}(x) = h(\exp(x))$. Hence $g \mapsto h$ defines the isomorphism $C(\mathfrak{t}, L^2(\mathfrak{t}))^\Gamma \cong C(T, L^2(\mathfrak{t}))$.

We now state the following theorem.

Theorem 3.5. *The Hilbert module $\mathcal{E} \widehat{\otimes}_{C(T^\vee)} (L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ is isomorphic to $C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ via the map*

$$\phi \otimes (\xi \otimes s) \mapsto \sum_{\alpha \in \Gamma} \phi(x - \alpha) L_{\alpha-x} \xi \otimes s.$$

The representation of $C(T)$ on $L^2(\mathfrak{t})$ induces a representation σ of $C(T)$ on $C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ defined by

$$[\sigma(f)h](\exp(x), y) = f(\exp(x + y))h(\exp(x), y).$$

Here the notation $h(\exp(x), y)$ denotes the value at the point $y \in \mathfrak{t}$ of $h(\exp(x)) \in L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}$.

Proof. We recall that $\mathcal{E} \widehat{\otimes}_{C(T^\vee)} (L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ is isomorphic to $(\mathcal{E} \widehat{\otimes}_{C(T^\vee)} L^2(\mathfrak{t})) \widehat{\otimes} \mathcal{S}$ and we have established that $\mathcal{E} \widehat{\otimes}_{C(T^\vee)} L^2(\mathfrak{t}) \cong C(T, L^2(\mathfrak{t}))$. This provides the claimed isomorphism.

It remains to identify the representation. Given $f \in C(T)$ let $\tilde{f}(x) = f(\exp(x))$ denote the corresponding periodic function on \mathfrak{t} . By definition the representation of $C(T)$ on $\mathcal{E} \widehat{\otimes}_{C(T^\vee)} (L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ takes $\phi \otimes \xi \otimes s$ to $\phi \otimes \tilde{f}\xi \otimes s$. This is mapped under the isomorphism to the Γ -periodic function on \mathfrak{t} whose value at x is

$$\sum_{\alpha \in \Gamma} \phi(x - \alpha) L_{\alpha-x}(\tilde{f}\xi) \otimes s \in L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}.$$

Evaluating this element of $L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}$ at a point $y \in \mathfrak{t}$ we have

$$\sum_{\alpha \in \Gamma} \phi(x - \alpha) \tilde{f}(x - \alpha + y) \xi(x - \alpha + y) \otimes s = \tilde{f}(x + y) \sum_{\alpha \in \Gamma} \phi(x - \alpha) [L_{\alpha-x}\xi](y) \otimes s$$

by Γ -periodicity of \tilde{f} . Thus $\sigma(f)$ pointwise multiplies the image of $\phi \otimes \xi \otimes s$ in $C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ by $\tilde{f}(x + y) = f(\exp(x + y))$ as claimed. \square

We now define an operator \mathbf{Q} on $C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ by

$$(\mathbf{Q}h)(\exp(x)) = Q_0(h(\exp(x)))$$

for $h \in C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$.

Theorem 3.6. *The unbounded operator \mathbf{Q} is a connection for Q_0 in the sense that the bounded operator $\mathbf{F} = \mathbf{Q}(1 + \mathbf{Q}^2)^{-1/2}$ is a connection for $F_0 = Q_0(1 + Q_0^2)^{-1/2}$, after making the identification of Hilbert modules as in Theorem 3.5.*

Proof. Let $Q_x = (L_x \otimes 1_{\mathcal{S}})Q_0(L_{-x} \otimes 1_{\mathcal{S}})$ and correspondingly define

$$F_x = Q_x(1 + Q_x^2)^{-1/2} = (L_x \otimes 1_{\mathcal{S}})F_0(L_{-x} \otimes 1_{\mathcal{S}}).$$

The commutators $[L_x \otimes 1_{\mathcal{S}}, Q_0]$ are bounded (the argument is exactly as for $[L_\gamma \otimes 1_{\mathcal{S}}, Q_0]$ in Section 3.2). It follows (in the spirit of Baaj-Julg, [1]) that the commutators $[L_x \otimes 1_{\mathcal{S}}, F_0]$ are compact. Thus $F_x - F_0$ is a compact operator for all $x \in \mathfrak{t}$.

To show that \mathbf{F} is a connection for F_0 we must show that for $\phi \in \mathcal{E}$, the diagram

$$\begin{array}{ccc} L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S} & \xrightarrow{F_0} & L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S} \\ \phi \otimes \downarrow & & \phi \otimes \downarrow \\ \mathcal{E} \otimes L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S} & & \mathcal{E} \otimes L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S} \\ \cong \downarrow & & \cong \downarrow \\ C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}) & \xrightarrow{\mathbf{F}} & C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}) \end{array}$$

commutes modulo compact operators.

Following the diagram around the right-hand side we have

$$\xi \otimes s \mapsto \sum_{\alpha \in \Gamma} \phi(x - \alpha)(L_{\alpha-x} \otimes 1_{\mathcal{S}})F_0(\xi \otimes s)$$

while following the left-hand side we have

$$\mathbf{F} \left[\sum_{\alpha \in \Gamma} \phi(x - \alpha)(L_{\alpha-x} \otimes 1_{\mathcal{S}})(\xi \otimes s) \right] = \sum_{\alpha \in \Gamma} \phi(x - \alpha)F_0(L_{\alpha-x} \otimes 1_{\mathcal{S}})(\xi \otimes s).$$

As $[F_0, L_{\alpha-x} \otimes 1_{\mathcal{S}}]$ is a compact operator for each x and the sum is finite for each x , the difference between the two paths around the diagram is a function from T to compact operators on $L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}$. It is thus a compact operator from the Hilbert space $L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}$ to the Hilbert module $C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ as required. \square

Theorem 3.7. *The Kasparov product $\mathcal{P} \otimes_{C(T^\vee)} \mathcal{Q}$ is the identity $1_{C(T)}$ in $KK_W(C(T), C(T))$.*

Proof. We define a homotopy of representations of $C(T)$ on $C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ by

$$[\sigma_\lambda(f)h](\exp(x), y) = f(\exp(x + \lambda y))h(\exp(x), y)$$

and note that $\sigma_1 = \sigma$ while σ_0 is simply the representation of $C(T)$ on $C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ by pointwise multiplication of functions on T . It is easy to see that these representations are W -equivariant.

Let f be a smooth function on T and let $h \in C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$. Let $\tilde{f}(x) = f(\exp(x))$ and let $\tilde{h}(x, y) = h(\exp(x), y)$. Then

$$\begin{aligned} & ([\mathbf{Q}, \sigma_\lambda(f)]h)(\exp(x), y) \\ &= \left[\frac{\partial}{\partial y^j} (\varepsilon^j \tilde{f}(x + \lambda y) \tilde{h}(x, y)) - 2\pi i y^j \mathbf{e}_j \tilde{f}(x + \lambda y) \tilde{h}(x, y) \right] \\ & \quad - \left[\tilde{f}(x + \lambda y) \frac{\partial}{\partial y^j} (\varepsilon^j \tilde{h}(x, y)) - \tilde{f}(x + \lambda y) 2\pi i y^j \mathbf{e}_j \tilde{h}(x, y) \right] \\ &= \frac{\partial}{\partial y^j} (\tilde{f}(x + \lambda y)) (\varepsilon^j \tilde{h}(x, y)). \end{aligned}$$

For each λ the operator \mathbf{Q} thus commutes with the representation σ_λ modulo bounded operators on a dense subalgebra of $C(T)$. Hence for each λ $(C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}), \sigma_\lambda, \mathbf{Q})$ defines an unbounded Kasparov triple.

This is true in particular for $\lambda = 1$ and thus $(C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}), \sigma, \mathbf{Q})$ is a Kasparov triple so as the operator in the triple \mathcal{P} is zero while \mathbf{Q} is a connection for Q_0 it follows that $\mathcal{P} \otimes_{C(T^\vee)} \mathcal{Q} = (C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}), \sigma, \mathbf{Q})$ in $KK_W(C(T), C(T))$.

Now applying the homotopy we have $\mathcal{P} \otimes_{C(T^\vee)} \mathcal{Q} = (C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}), \sigma_0, \mathbf{Q})$. Since σ_0 commutes exactly with the operator \mathbf{Q} the representation σ_0 respects the direct sum decomposition of $C(T, L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S})$ as $C(T, \ker(Q_0)) \oplus C(T, \ker(Q_0)^\perp)$. The operator \mathbf{Q} is invertible on the second summand (and commutes with the representation) and hence the corresponding Kasparov triple $(C(T, \ker(Q_0)^\perp), \sigma_0|_{C(T, \ker(Q_0)^\perp)}, \mathbf{Q}|_{C(T, \ker(Q_0)^\perp)})$ is zero in KK -theory.

We thus conclude that $\mathcal{P} \otimes_{C(T^\vee)} \mathcal{Q} = (C(T, \ker(Q_0)), \sigma_0|_{C(T, \ker(Q_0))}, 0)$. Since $\ker Q_0$ is 1-dimensional (Theorem 3.2) the module $C(T, \ker(Q_0))$ is

isomorphic to $C(T)$ and the restriction of σ_0 to this is the identity representation of $C(T)$ on itself. Thus $\mathcal{P} \otimes_{C(T^\vee)} \mathcal{Q} = (C(T), 1, 0) = 1_{C(T)}$. \square

3.4. The Kasparov product $\mathcal{P} \otimes_{C(T)} \mathcal{Q}$. We begin by considering the dual picture, which exchanges the roles of T and T^\vee . There exist elements $\mathcal{Q}^\vee \in KK_W(C(T) \widehat{\otimes} C(T^\vee), \mathbb{C})$ and $\mathcal{P}^\vee \in KK_W(\mathbb{C}, C(T^\vee) \widehat{\otimes} C(T))$ for which the result of the previous section implies $\mathcal{P}^\vee \otimes_{C(T)} \mathcal{Q}^\vee = 1_{C(T^\vee)}$ in $KK_W(C(T^\vee), C(T^\vee))$.

We will show that there is an isomorphism

$$\theta : C(T^\vee) \widehat{\otimes} C(T) \rightarrow C(T) \widehat{\otimes} C(T^\vee)$$

such that $\mathcal{Q} = \theta^* \mathcal{Q}^\vee$ and $\mathcal{P} = \theta_*^{-1} \mathcal{P}^\vee$. This will imply that $\mathcal{P} \otimes_{C(T)} \mathcal{Q} = \mathcal{P}^\vee \otimes_{C(T)} \mathcal{Q}^\vee = 1_{C(T^\vee)}$ in $KK_W(C(T^\vee), C(T^\vee))$ and hence will complete the proof of the Poincaré duality between $C(T)$ and $C(T^\vee)$.

We recall that \mathcal{Q} is represented by the (unbounded) Kasparov triple $(L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}, \rho, Q_0)$ where $\mathcal{S} = \mathcal{C}l(\mathfrak{t} \times \mathfrak{t}^*)P$, for P the projection $P = \prod_j \frac{1}{2}(1 - i\mathbf{e}_j \varepsilon^j)$ and

$$Q_0 = \frac{\partial}{\partial y^j} \otimes \varepsilon^j - 2\pi i y^j \otimes \mathbf{e}_j.$$

For $\gamma \in \Gamma$, $\chi \in \Gamma^\vee$ and correspondingly $e^{2\pi i \langle \eta, \gamma \rangle}$ in $C(T^\vee)$, $e^{2\pi i \langle \chi, x \rangle}$ in $C(T)$, the representation ρ of $C(T^\vee) \widehat{\otimes} C(T)$ is defined by

$$\rho(e^{2\pi i \langle \eta, \gamma \rangle})(\xi \otimes s) = L_\gamma \xi \otimes s, \text{ and } \rho(e^{2\pi i \langle \chi, x \rangle})(\xi \otimes s) = e^{2\pi i \langle \chi, x \rangle} \xi \otimes s.$$

By definition \mathcal{Q}^\vee is represented by the triple $(L^2(\mathfrak{t}^*) \widehat{\otimes} \mathcal{S}^\vee, \rho^\vee, Q_0^\vee)$ where $\mathcal{S}^\vee = \mathcal{C}l(\mathfrak{t}^* \times \mathfrak{t})P^\vee$, for P^\vee the projection $P^\vee = \prod_j \frac{1}{2}(1 - i\varepsilon^j \mathbf{e}_j)$ and

$$Q_0^\vee = \frac{\partial}{\partial \eta_j} \otimes \mathbf{e}_j - 2\pi i \eta_j \otimes \varepsilon^j.$$

For $\gamma \in \Gamma$, $\chi \in \Gamma^\vee$ and correspondingly $e^{2\pi i \langle \eta, \gamma \rangle}$ in $C(T^\vee)$, $e^{2\pi i \langle \chi, x \rangle}$ in $C(T)$, the representation ρ^\vee of $C(T) \widehat{\otimes} C(T^\vee)$ is now defined by

$$\rho^\vee(e^{2\pi i \langle \chi, x \rangle})(\xi^\vee \otimes s^\vee) = L_\chi^\vee \xi^\vee \otimes s^\vee, \text{ and } \rho^\vee(e^{2\pi i \langle \eta, \gamma \rangle})(\xi^\vee \otimes s^\vee) = e^{2\pi i \langle \eta, \gamma \rangle} \xi^\vee \otimes s^\vee.$$

Here L_χ^\vee denotes the translation action of $\chi \in \Gamma^\vee$ on $L^2(\mathfrak{t}^*)$.

In our notation, ε^j is again an orthonormal basis for \mathfrak{t}^* and \mathbf{e}_j is an orthonormal basis for \mathfrak{t} . We can canonically identify $\mathcal{C}l(\mathfrak{t} \times \mathfrak{t}^*)$ with $\mathcal{C}l(\mathfrak{t}^* \times \mathfrak{t})$, and hence think of both \mathcal{S} and \mathcal{S}^\vee as subspaces of this algebra.

We can identify $L^2(\mathfrak{t})$ with $L^2(\mathfrak{t}^*)$ via the Fourier transform: let $\mathcal{F} : L^2(\mathfrak{t}) \rightarrow L^2(\mathfrak{t}^*)$ denote the Fourier transform isomorphism

$$[\mathcal{F}\xi](\eta) = \int_{\mathfrak{t}} \xi(y) e^{2\pi i \langle \eta, y \rangle} dy.$$

It is easy to see that this is W -equivariant.

To identify \mathcal{S} with \mathcal{S}^\vee , let $u \in \mathcal{C}l(\mathfrak{t} \times \mathfrak{t}^*)$ be defined by $u = \varepsilon^1 \varepsilon^2 \dots \varepsilon^n$ when $n = \dim(\mathfrak{t})$ is even and $u = e_1 e_2 \dots e_n$ when n is odd.

Lemma 3.8. *Conjugation by u defines a W -equivariant unitary isomorphism $\mathcal{U} : \mathcal{S} \rightarrow \mathcal{S}^\vee$. For $a \in \mathcal{C}l(\mathfrak{t} \times \mathfrak{t}^*)$ (viewed as an operator on \mathcal{S} by Clifford multiplication) $\mathcal{U}a\mathcal{U}^*$ is Clifford multiplication by uau^* on \mathcal{S}^\vee and in particular $\mathcal{U}\mathbf{e}_j\mathcal{U}^* = \mathbf{e}_j$ while $\mathcal{U}\varepsilon^j\mathcal{U}^* = -\varepsilon^j$.*

Proof. We first note that u respectively commutes and anticommutes with $\mathbf{e}_j, \varepsilon^j$ (there being respectively an even or odd number of terms in u which anticommute with $\mathbf{e}_j, \varepsilon^j$). It follows that $uPu^* = P^\vee$, hence conjugation by u maps \mathcal{S} to \mathcal{S}^\vee .

Denoting by $\pi : \mathbb{C}P \rightarrow \mathbb{C}$ the identification of $\mathbb{C}P$ with \mathbb{C} , the inner product on \mathcal{S} is given by $\langle s_1, s_2 \rangle = \pi(s_1^* s_2)$ while the inner product on \mathcal{S}^\vee is given by $\langle s_1^\vee, s_2^\vee \rangle = \pi(u^*(s_1^\vee)^* s_2^\vee u)$. Thus

$$\langle usu^*, s^\vee \rangle = \pi(u^*(usu^*)^* s^\vee u) = \pi(s^* u^* s^\vee u) = \langle s, u^* s^\vee u \rangle$$

so \mathcal{U}^* is conjugation by u^* which inverts \mathcal{U} establishing that \mathcal{U} is unitary.

We now check that \mathcal{U} is W -equivariant. In the case that \mathfrak{t} is even-dimensional, we note that identifying $\mathcal{C}\ell(\mathfrak{t}^*)$ with the exterior algebra of \mathfrak{t}^* (as a W -vector space), u corresponds to the volume form on \mathfrak{t}^* so $w \cdot u = \det(w)u$. Similarly in the odd dimensional case u corresponds to the volume form on \mathfrak{t} and again the action of w on u is multiplication by the determinant. Thus

$$w \cdot \mathcal{U}(s) = w \cdot (usu^*) = (w \cdot u)(w \cdot s)(w \cdot u^*) = \det(w)^2 u(w \cdot s)u^* = \mathcal{U}(w \cdot s)$$

since $\det(w) = \pm 1$.

Finally for $s^\vee \in \mathcal{S}^\vee$ and $a \in \mathcal{C}\ell(\mathfrak{t} \times \mathfrak{t}^*)$ we have

$$\mathcal{U}a\mathcal{U}^*s^\vee = \mathcal{U}(au^*s^\vee u) = uau^*s^\vee$$

and hence $\mathcal{U}\mathbf{e}_j\mathcal{U}^* = u\mathbf{e}_ju^* = \mathbf{e}_j$, $\mathcal{U}\varepsilon^j\mathcal{U}^* = u\varepsilon^ju^* = -\varepsilon^j$. \square

Since $\mathcal{F} \otimes \mathcal{U}$ is a W -equivariant unitary isomorphism from $L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}$ to $L^2(\mathfrak{t}^*) \widehat{\otimes} \mathcal{S}^\vee$, the triple $(L^2(\mathfrak{t}) \widehat{\otimes} \mathcal{S}, \rho, Q_0)$ representing \mathcal{Q} is isomorphic to the Kasparov triple

$$(L^2(\mathfrak{t}^*) \widehat{\otimes} \mathcal{S}^\vee, (\mathcal{F} \otimes \mathcal{U})\rho(\mathcal{F}^* \otimes \mathcal{U}^*), (\mathcal{F} \otimes u)Q_0(\mathcal{F}^* \otimes \mathcal{U}^*)).$$

Theorem 3.9. *Let $\theta : C(T^\vee) \widehat{\otimes} C(T) \rightarrow C(T) \widehat{\otimes} C(T^\vee)$ be defined by*

$$\theta(g \otimes f) = f \otimes (g \circ \epsilon).$$

where ϵ is the involution on T^\vee defined by $\epsilon(\exp(\eta)) = \exp(-\eta)$. Then $\mathcal{Q} = \theta^* \mathcal{Q}^\vee$ in $KK_W(C(T^\vee) \widehat{\otimes} C(T), \mathbb{C})$.

Proof. We will show that $\rho^\vee \circ \theta = (\mathcal{F} \otimes \mathcal{U})\rho(\mathcal{F}^* \otimes \mathcal{U}^*)$ and $(\mathcal{F} \otimes u)Q_0(\mathcal{F}^* \otimes \mathcal{U}^*) = Q_0^\vee$. We begin with the operator.

The operator Q_0 is given by

$$\frac{\partial}{\partial y^j} \otimes \varepsilon^j - 2\pi i y^j \otimes \mathbf{e}_j.$$

Conjugating the operator $\frac{\partial}{\partial y^j}$ by the Fourier transform we obtain the multiplication by $2\pi i \eta_j$, while conjugating $-2\pi i y^j$ by the Fourier transform we obtain the multiplication by $-2\pi i (\frac{i}{2\pi} \frac{\partial}{\partial \eta_j}) = \frac{\partial}{\partial \eta_j}$. Conjugation by \mathcal{U} negates ε^j and preserves \mathbf{e}_j hence

$$(\mathcal{F} \otimes u)Q_0(\mathcal{F}^* \otimes \mathcal{U}^*) = 2\pi i \eta_j \otimes (-\varepsilon^j) + \frac{\partial}{\partial \eta_j} \otimes \mathbf{e}_j = Q_0^\vee.$$

For the representation, $\rho(e^{2\pi i \langle \chi, x \rangle})$ is multiplication by $e^{2\pi i \langle \chi, x \rangle}$ on $L^2(\mathfrak{t})$ (with the identity on \mathcal{S}) and conjugating by the Fourier transform we get

the translation L_χ^\vee , hence $(\mathcal{F} \otimes \mathcal{U})\rho(e^{2\pi i\langle \chi, x \rangle})(\mathcal{F}^* \otimes \mathcal{U}^*) = \rho^\vee(e^{2\pi i\langle \chi, x \rangle})$. On the other hand $\rho(e^{2\pi i\langle \eta, \gamma \rangle})$ is the translation L_γ and Fourier transforming we get the multiplication by $e^{-2\pi i\langle \eta, \gamma \rangle}$. Thus $(\mathcal{F} \otimes \mathcal{U})\rho(e^{2\pi i\langle \eta, \gamma \rangle})(\mathcal{F}^* \otimes \mathcal{U}^*) = \rho^\vee(e^{2\pi i\langle -\eta, \gamma \rangle})$.

We conclude that $(\mathcal{F} \otimes \mathcal{U})\rho(\mathcal{F}^* \otimes \mathcal{U}^*) = \rho^\vee \circ \theta$ as required. \square

Theorem 3.10. *The Kasparov product $\mathcal{P} \otimes_{C(T)} \mathcal{Q}$ is $1_{C(T^\vee)}$ in the Kasparov group $KK_W(C(T^\vee), C(T^\vee))$.*

Proof. We have $\mathcal{P} \otimes_{C(T^\vee)} \mathcal{Q} = 1_{C(T)}$ in $KK_W(C(T), C(T))$ by Theorem 3.7 while $\mathcal{P}^\vee \otimes_{C(T)} \mathcal{Q}^\vee = 1_{C(T^\vee)}$ in $KK_W(C(T^\vee), C(T^\vee))$ by Theorem 3.7 for the dual group.

By Theorem 3.9 we have $\mathcal{Q}^\vee = (\theta^{-1})^* \mathcal{Q}$ whence

$$1_{C(T^\vee)} = \mathcal{P}^\vee \otimes_{C(T)} \mathcal{Q}^\vee = (\theta^{-1})_* \mathcal{P}^\vee \otimes_{C(T)} \mathcal{Q}.$$

Let $\mathcal{P}' = (\theta^{-1})_* \mathcal{P}^\vee$ in $KK_W(\mathbb{C}, C(T) \widehat{\otimes} C(T^\vee))$. Then

$$\mathcal{P} = \mathcal{P} \otimes_{C(T^\vee)} 1_{C(T^\vee)} = \mathcal{P} \otimes_{C(T^\vee)} (\mathcal{P}' \otimes_{C(T)} \mathcal{Q}).$$

By definition $\mathcal{P}' \otimes_{C(T)} \mathcal{Q} = (\mathcal{P}' \otimes 1_{C(T^\vee)}) \otimes_{C(T) \otimes C(T^\vee)} \mathcal{Q}$ and hence

$$\mathcal{P} = (\mathcal{P} \otimes \mathcal{P}') \otimes_{C(T^\vee) \widehat{\otimes} C(T)} \mathcal{Q}$$

by associativity of the Kasparov product. Here $\mathcal{P} \otimes \mathcal{P}'$ is the ‘external’ product and lives in $KK_W(\mathbb{C}, C(T) \widehat{\otimes} C(T) \widehat{\otimes} C(T^\vee) \widehat{\otimes} C(T^\vee))$, with \mathcal{P} appearing in the first and last factors, and \mathcal{P}' in the second and third. The product with \mathcal{Q} is over the second and last factors. Similarly

$$\mathcal{P}' = \mathcal{P}' \otimes_{C(T)} (\mathcal{P} \otimes_{C(T^\vee)} \mathcal{Q}) = (\mathcal{P}' \otimes \mathcal{P}) \otimes_{C(T) \widehat{\otimes} C(T^\vee)} \mathcal{Q}$$

where \mathcal{P}' now appears as the first and last factors and the product with \mathcal{Q} is over the first and third factors. Up to reordering terms of the tensor product $(\mathcal{P} \otimes \mathcal{P}') \otimes_{C(T^\vee) \widehat{\otimes} C(T)} \mathcal{Q} = (\mathcal{P}' \otimes \mathcal{P}) \otimes_{C(T) \widehat{\otimes} C(T^\vee)} \mathcal{Q}$.

Thus (by commutativity of the external product) $\mathcal{P} = \mathcal{P}' = (\theta^{-1})_* \mathcal{P}^\vee$ and hence $\mathcal{P} \otimes_{C(T)} \mathcal{Q} = 1_{C(T^\vee)}$. This completes the proof. \square

Corollary 3.11. *The elements $\mathcal{Q} \in KK_W(C(T^\vee) \widehat{\otimes} C(T), \mathbb{C})$ and $\mathcal{P} \in KK_W(\mathbb{C}, C(T) \widehat{\otimes} C(T^\vee))$ exhibit a W -equivariant Poincaré duality between the algebras $C(T)$ and $C(T^\vee)$.*

4. POINCARÉ DUALITY BETWEEN $C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W)$ AND $C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)$

4.1. Descent of Poincaré duality. For W a group, a Poincaré duality between two W - C^* -algebras A, B induces a natural family of isomorphisms

$$KK_W(A \widehat{\otimes} D_1, D_2) \cong KK_W(D_1, B \widehat{\otimes} D_2)$$

for W - C^* -algebras D_1, D_2 . In other words the functor $A \widehat{\otimes}$ is left-adjoint to $B \widehat{\otimes}$ on the KK_W category when there is a Poincaré duality from A to B . (The symmetry of Poincaré dualities means that $B \widehat{\otimes}$ is also left-adjoint to $A \widehat{\otimes}$). The element in $KK_W(\mathbb{C}, A \widehat{\otimes} B)$ defining the Poincaré duality is precisely the unit of the adjunction, while the counit is given by the element in $KK_W(B \widehat{\otimes} A, \mathbb{C})$. This categorical view of Poincaré duality appears in [7, 8, 9].

Now let D_1, D_2 be C^* -algebras (without W -action). Let τ denote the trivial-action functor from KK to KK_W , i.e. $\tau D_1, \tau D_2$ are W - C^* -algebras with trivial action of W . In the case that W is a *finite* group a Poincaré duality yields isomorphisms

$$\begin{aligned} KK(A \rtimes W \widehat{\otimes} D_1, D_2) &\cong KK_W(A \widehat{\otimes} \tau D_1, \tau D_2) \\ &\cong KK_W(\tau D_1, B \widehat{\otimes} \tau D_2) \\ &\cong KK(D_1, B \rtimes W \widehat{\otimes} D_2). \end{aligned}$$

The first and last isomorphisms are the dual Green-Julg and Green-Julg isomorphisms respectively, and in categorical terms these amount to the fact that the τ functor is (right and left) adjoint to the descent functor $\rtimes W$, see [11]. We denote the unit and counit by α and β , for the left-adjunction from τ to $\rtimes W$, and by $\widehat{\alpha}$ and $\widehat{\beta}$ for the left adjunction from $\rtimes W$ to τ .

Since this is natural $A \rtimes W \widehat{\otimes}$ is left-adjoint to $B \rtimes W \widehat{\otimes}$, hence there must exist a unit and a counit providing this descended Poincaré duality. We will identify these elements explicitly.

Theorem 4.1. *Let $\mathfrak{a} \in KK_W(B \widehat{\otimes} A, \mathbb{C})$ and $\mathfrak{b} \in KK_W(\mathbb{C}, A \widehat{\otimes} B)$ define a W -equivariant Poincaré duality between W - C^* -algebras A, B , with W finite. Then $\widetilde{\mathfrak{a}} = \mathfrak{T}\mathfrak{r}(\mathfrak{a} \rtimes W)\beta_{\mathbb{C}} \in KK(B \rtimes W \widehat{\otimes} A \rtimes W, \mathbb{C})$ and $\widetilde{\mathfrak{b}} = \alpha_{\mathbb{C}}(\mathfrak{b} \rtimes W)\Delta \in KK(\mathbb{C}, A \rtimes W \widehat{\otimes} B \rtimes W)$ define a Poincaré duality such that the following diagram commutes*

$$\begin{array}{ccc} KK_W^*(A, \mathbb{C}) & \xrightarrow{\quad \mathfrak{b} \widehat{\otimes}_A - \quad} & KK_W^*(\mathbb{C}, B) \\ \downarrow \text{dual Green-Julg} \cong & & \downarrow \text{Green-Julg} \cong \\ KK^*(A \rtimes W, \mathbb{C}) & \xrightarrow{\quad \widetilde{\mathfrak{b}} \widehat{\otimes}_{A \rtimes W} - \quad} & KK^*(\mathbb{C}, B \rtimes W). \end{array}$$

Here $\Delta \in KK((A \widehat{\otimes} B) \rtimes W, A \rtimes W \widehat{\otimes} B \rtimes W)$ is given by the diagonal inclusion of W into $W \times W$ and $\mathfrak{T}\mathfrak{r} \in KK(B \rtimes W \widehat{\otimes} A \rtimes W, (B \widehat{\otimes} A) \rtimes W)$ is dual to this: We define a positive linear map $\text{Tr} : (B \widehat{\otimes} A) \rtimes (W \times W) \rightarrow (B \widehat{\otimes} A) \rtimes W$ by

$$\text{Tr} : (a \otimes b)[w_1, w_2] \mapsto \begin{cases} (a \otimes b)[w_1] & \text{if } w_1 = w_2 \\ 0 & \text{otherwise.} \end{cases}$$

This is a $(B \widehat{\otimes} A) \rtimes W$ -module map and we equip the algebra $(B \widehat{\otimes} A) \rtimes (W \times W)$ with inner product in $(B \widehat{\otimes} A) \rtimes W$ defined by

$$\langle (b \otimes a)[w_1, w_2], (b' \otimes a')[w'_1, w'_2] \rangle_{\text{Tr}} = \text{Tr}([w_1^{-1}, w_2^{-1}](b^* \otimes a^*)(b' \otimes a')[w'_1, w'_2]).$$

The completion of this as a Hilbert module, equipped with the left multiplication representation of $(B \widehat{\otimes} A) \rtimes (W \times W)$ provides the required element $\mathfrak{T}\mathfrak{r} \in KK((B \widehat{\otimes} A) \rtimes (W \times W), (B \widehat{\otimes} A) \rtimes W)$.

To identify the unit and counit $\widetilde{\mathfrak{b}}$ and $\widetilde{\mathfrak{a}}$ one proceeds as follows. The unit $\widetilde{\mathfrak{b}}$ is the image of the identity $1_{A \rtimes W}$ under the isomorphism $KK(A \rtimes W, A \rtimes W) \cong KK(\mathbb{C}, B \rtimes W \widehat{\otimes} A \rtimes W)$. This is the composition of the dual Green-Julg, equivariant Poincaré duality, and Green-Julg maps. The first two yield the Poincaré dual of the unit $\widehat{\alpha}_A$. One must then descend this and pair with the

unit $\alpha_{\mathbb{C}}$. Hence $\tilde{\mathbf{b}} = \alpha_{\mathbb{C}}(b(\widehat{\alpha}_A \otimes 1_B)) \rtimes W = \alpha_{\mathbb{C}}(b \rtimes W)((\widehat{\alpha}_A \otimes 1_B) \rtimes W)$ by naturality of descent. It is not hard to identify $(\widehat{\alpha}_A \otimes 1_B) \rtimes W$ as the element Δ .

Similarly the counit $\tilde{\mathbf{a}}$ is the image of the identity $1_{B \rtimes W}$ under the isomorphism $KK(B \rtimes W, B \rtimes W) \cong KK(A \rtimes W \widehat{\otimes} B \rtimes W, \mathbb{C})$. This is the composition of the Green-Julg, equivariant Poincaré duality, and dual Green-Julg maps, hence $\tilde{\mathbf{a}}$ is obtained by taking the Poincaré dual of the counit β_B , descending, and applying the counit $\widehat{\beta}_{\mathbb{C}}$. We have $\tilde{\mathbf{a}} = ((\widehat{\beta}_B \otimes 1_A)a) \rtimes W \widehat{\beta}_{\mathbb{C}} = ((\widehat{\beta}_B \otimes 1_A) \rtimes W)(a \rtimes W) \widehat{\beta}_{\mathbb{C}}$. A change of variables identifies $(\widehat{\beta}_B \otimes 1_A) \rtimes W$ with $\mathfrak{T}\mathfrak{r}$.

Remark 4.2. Given a Kasparov triple $(\mathcal{E}, 1, D)$ representing \mathbf{b} we can describe explicitly a triple $(\tilde{\mathcal{E}}, \tilde{\alpha}_{\mathbb{C}}, D \otimes 1)$ for $\tilde{\mathbf{b}}$.

The module $\tilde{\mathcal{E}}$ is given by descending \mathcal{E} and inflating the action of W to $W \times W$. Explicitly $\tilde{\mathcal{E}}$ is the completion of $\mathcal{E} \widehat{\otimes} \mathbb{C}[W \times W]$ with respect to the inner product

$$\langle \xi \otimes [w_1, w_2], \xi' \otimes [w'_1, w'_2] \rangle = (w_1^{-1}, w_2^{-1}) \cdot \langle \xi, \xi' \rangle [w_1^{-1} w'_1, w_2^{-1} w'_2].$$

The operator is simply $D \otimes 1$ on $\tilde{\mathcal{E}}$.

The representation $\tilde{\alpha}_{\mathbb{C}}$ of \mathbb{C} on $\tilde{\mathcal{E}}$ takes 1 to the projection corresponding to the trivial representation of W , where W acts diagonally on $\tilde{\mathcal{E}}$ – the unit $\alpha_{\mathbb{C}} \in KK(\mathbb{C}, (\tau\mathbb{C}) \rtimes W)$ is given by inclusion of \mathbb{C} as the trivial representation in $\mathbb{C}[W] = (\tau\mathbb{C}) \rtimes W$.

4.2. Proof of Theorem 1.1. Theorem 1.1 follows from Theorem 4.1 by consideration of the following diagram.

$$\begin{array}{ccc} KK_W^*(C(T), \mathbb{C}) & \xrightarrow{\mathcal{P} \widehat{\otimes}_{C(T)} -} & KK_W^*(\mathbb{C}, C(T^\vee)). \\ \cong \downarrow \text{Morita} & & \cong \downarrow \text{Morita} \\ KK_W^*(C_0(\mathfrak{t}) \rtimes \Gamma, \mathbb{C}) & \xrightarrow{\mathbf{b} \widehat{\otimes}_{C_0(\mathfrak{t}) \rtimes \Gamma} -} & KK_W^*(\mathbb{C}, C_0(\mathfrak{t}^*) \rtimes \Gamma^\vee) \\ \cong \downarrow \text{dual Green-Julg} & & \cong \downarrow \text{Green-Julg} \\ KK^*(C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W), \mathbb{C}) & \xrightarrow{\tilde{\mathbf{b}} \widehat{\otimes}_{C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W)} -} & KK^*(\mathbb{C}, C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)) \end{array}$$

Composition of \mathcal{P} with the Morita equivalences and of \mathcal{Q} with the inverse Morita equivalences, yields a W -equivariant Poincaré duality between $C_0(\mathfrak{t}) \rtimes \Gamma$ and $C_0(\mathfrak{t}^*) \rtimes \Gamma^\vee$ inducing the middle arrow.

To determine the element \mathbf{b} explicitly, recall that \mathcal{P} is given by the Hilbert module of functions $\sigma : \mathfrak{t} \times \mathfrak{t}^* \rightarrow \mathbb{C}$ which are Γ^\vee periodic in the second variable and satisfying

$$\sigma(\gamma + x, \eta) = e^{2\pi i \langle \eta, \gamma \rangle} \sigma(x, \eta).$$

This module is equipped with the inner product

$$\langle \sigma_1, \sigma_2 \rangle(x, \eta) = \overline{\sigma_1(x, \eta)} \sigma_2(x, \eta).$$

The Morita equivalence from $C(T)$ to $C_0(\mathfrak{t}) \rtimes \Gamma$ is given by the completion of $C_c(\mathfrak{t})$ with respect to the inner product

$$\langle \phi_1, \phi_2 \rangle = \sum_{\gamma \in \Gamma} \overline{\phi_1}(\gamma \cdot \phi_2)[\gamma],$$

and similarly for $C(T^\vee)$.

It follows that \mathfrak{b} is given by the Hilbert module completion of $C_c(\mathfrak{t} \times \mathfrak{t}^*)$ with respect to the inner product

$$\langle \theta_1, \theta_2 \rangle = \sum_{(\gamma, \chi) \in \Gamma \times \Gamma^\vee} \overline{\theta_1}((\gamma, \chi) \cdot \theta_2) e^{2\pi i \langle \eta, \gamma \rangle} [(\gamma, \chi)].$$

Applying Theorem 4.1 yields the bottom arrow. Here we identify $(C_0(\mathfrak{t}) \rtimes \Gamma) \rtimes W$ with $C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W)$ and $(C_0(\mathfrak{t}^*) \rtimes \Gamma^\vee) \rtimes W$ with $C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)$. As noted in Remark 4.2 the element $\tilde{\mathfrak{b}}$ has Hilbert module obtained by descending the module and inflating the W action to $W \times W$.

In conclusion we obtain the module by completing $C_c(\mathfrak{t} \times \mathfrak{t}^*) \rtimes (W \times W)$ with respect to the inner product

$$\langle \theta[w_1, w_2], \theta'[w'_1, w'_2] \rangle = (w_1, w_2)^{-1} \cdot \langle \theta, \theta' \rangle [w_1^{-1} w'_1, w_2^{-1} w'_2],$$

where $\langle \theta, \theta' \rangle$ is the inner product on $C_c(\mathfrak{t} \times \mathfrak{t}^*)$ defined above which is equipped with the representation of \mathbb{C} given by the trivial projection in $\mathbb{C}[W]$, where W acts diagonally on all factors.

4.3. Proof of Theorem 1.2. Theorem 1.2 follows from Theorem 4.1 by the consideration of the following diagram.

$$\begin{array}{ccc} KK_W^*(C(T), \mathbb{C}) & \xrightarrow{\mathcal{P} \widehat{\otimes}_{C(T)} -} & KK_W^*(\mathbb{C}, C(T^\vee)). \\ \cong \downarrow \text{Fourier-Pontryagin} & & \cong \downarrow \text{Fourier-Pontryagin} \\ KK_W^*(C^*(\Gamma^\vee), \mathbb{C}) & \xrightarrow{\mathfrak{b} \widehat{\otimes}_{C^*(\Gamma^\vee)} -} & KK_W^*(\mathbb{C}, C^*(\Gamma)) \\ \cong \downarrow \text{dual Green-Julg} & & \cong \downarrow \text{Green-Julg} \\ KK^*(C^*(\Gamma^\vee \rtimes W), \mathbb{C}) & \xrightarrow{\tilde{\mathfrak{b}} \widehat{\otimes}_{C^*(\Gamma^\vee \rtimes W)} -} & KK^*(\mathbb{C}, C^*(\Gamma \rtimes W)) \end{array}$$

Composition of \mathcal{P} and of \mathcal{Q} with the Fourier-Pontryagin isomorphisms yields a W -equivariant Poincaré duality between $C^*(\Gamma^\vee)$ and $C^*(\Gamma)$ inducing the middle arrow. Applying Theorem 4.1 yields the bottom arrow. Here we identify $C^*(\Gamma^\vee) \rtimes W$ with $C^*(\Gamma^\vee \rtimes W)$ and $C^*(\Gamma) \rtimes W$ with $C^*(\Gamma \rtimes W)$.

4.4. The connection with the Baum Connes assembly map.

The Poincaré duality isomorphism appearing in Theorem 1.1

$$KK^*(C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W), \mathbb{C}) \longrightarrow KK^*(\mathbb{C}, C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W))$$

can be identified with the Baum Connes assembly map for the group $\Gamma \rtimes W$ in a sense made explicit by the following diagram. Note that while we have suppressed the indices, these are degree 0 maps of \mathbb{Z}_2 -graded groups.

$$\begin{array}{ccc}
KK_{\Gamma \rtimes W}(C_0(\mathfrak{t}), \mathbb{C}) & \xrightarrow{\text{Baum-Connes}} & KK(\mathbb{C}, C^*(\Gamma \rtimes W)) \\
\downarrow \text{dual Green-Julg} & & \downarrow \text{Morita equivalence} \\
KK(C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W), \mathbb{C}) & \xrightarrow{\text{Poincaré duality}} & KK(\mathbb{C}, C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)) \\
\uparrow \times \widehat{\beta}_{\mathbb{C}} & & \uparrow \times \widehat{\beta}_{\mathbb{C}} \\
KK(C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W), C^*(\Gamma \rtimes W)) & \xrightarrow{\text{P. d.}} & KK(\mathbb{C}, C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W) \otimes C^*(\Gamma \rtimes W))
\end{array}$$

The curved arrow is the descent map. Note that since $\Gamma \rtimes W$ is amenable the full and reduced C^* -algebras agree. The counit $\widehat{\beta}_{\mathbb{C}} \in KK((\tau\mathbb{C}) \rtimes \Gamma \rtimes W, \mathbb{C}) = KK(C^*(\Gamma \rtimes W), \mathbb{C})$ is given by the trivial representation of the group $\Gamma \rtimes W$. This element has the effect of collapsing the coefficients $C^*(\Gamma \rtimes W)$.

The upper and lower Poincaré dualities in the diagram are both provided by Theorem 1.1, in the lower case with the coefficients $C^*(\Gamma \rtimes W)$, and the element inducing the map from K -homology to K -theory is described in detail in Section 4.2.

Clearly the lower square commutes by associativity of the Kasparov product, while the left hand triangle commutes by definition. Therefore, to show that the Baum Connes assembly map corresponds to the upper Poincaré duality it suffices to show that the outer pentagon is commutative.

By definition the assembly map is the composition of descent with a Kasparov product. We will use the notation $\mathbf{A}_{C_0(\mathfrak{t})}$ to denote the relevant element of $KK(\mathbb{C}, C_0(\mathfrak{t}) \rtimes \Gamma \rtimes W)$, which is given by the Hilbert module obtained by completing $C_c(\mathfrak{t})$ with respect to the inner product

$$\langle f, f' \rangle = \sum_{\gamma, w} \overline{f}((\gamma w) \cdot f)[\gamma w].$$

We thus have the following diagram, where the bottom arrow is our Poincaré duality.

$$\begin{array}{ccc}
KK_{\Gamma \rtimes W}(C_0(\mathfrak{t}), \mathbb{C}) & \xrightarrow{\text{Baum-Connes}} & KK(\mathbb{C}, C^*(\Gamma \rtimes W)) \\
\downarrow \text{descent} & \nearrow \mathbf{A}_{C_0(\mathfrak{t}) \times} & \downarrow \text{Morita equivalence} \\
& & KK(\mathbb{C}, C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)) \\
& & \uparrow \times \widehat{\beta}_{\mathbb{C}} \\
KK(C_0(\mathfrak{t}) \rtimes (\Gamma \rtimes W), C^*(\Gamma \rtimes W)) & \longrightarrow & KK(\mathbb{C}, C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W) \otimes C^*(\Gamma \rtimes W))
\end{array}$$

The upper triangle commutes by definition of the assembly map, however, it should be noted that the lower quadrilateral does not commute: the two directions around the quadrilateral collapse different algebras. It is thus not entirely obvious that the outer pentagon itself commutes. However we will

show that the quadrilateral does commute on the image of the descent map so that the outer pentagon commutes as required.

We start with a Kasparov cycle $(H, \rho, T) \in KK_{\Gamma \rtimes W}(C_0(\mathfrak{t}), \mathbb{C})$. Note that since the action of $\Gamma \rtimes W$ on \mathfrak{t} is proper we may, without loss of generality, take T to be exactly invariant and of finite propagation. Now we descend to get $(\mathcal{E}, \hat{\rho}, T \otimes 1)$, where $\mathcal{E} = H \widehat{\otimes} C^*\Gamma \rtimes W$ and $\hat{\rho}$ is a representation defined by $\hat{\rho}(f[g]) = \rho(f)\pi(g) \otimes [g]$, (π denotes the representation of $\Gamma \rtimes W$ on H).

Applying our Poincaré duality, given by the completion of $C_c(\mathfrak{t} \times \mathfrak{t}^*) \rtimes (\Gamma \times \Gamma^\vee)$ described in Section 4.2, along with with representation of \mathbb{C} given by the trivial representation of W , we obtain a Kasparov triple as follows:

Let $H_c = \rho(C_c(\mathfrak{t}))H$. The module in our triple is the completion of $H_c \widehat{\otimes} C_c(\mathfrak{t}^*) \widehat{\otimes} \mathbb{C}[(\Gamma \rtimes W) \times W]$ with respect to the inner product

$$\langle \xi \otimes f[(g, w)], \xi' \otimes f'[(g', w')] \rangle = \sum_{\delta \in \Gamma} \sum_{\chi \in \Gamma^\vee} \langle \xi, \delta \cdot \xi' \rangle [g^{-1} \delta g'] \langle f[w], [\chi] e^{2\pi i \langle \eta, \delta \rangle} f'[w'] \rangle$$

where the last inner product in the formula is taken in the algebra $C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)$ viewed as a module over itself. The representation of \mathbb{C} is once again given by the trivial projection in $\mathbb{C}[W]$ where W acts diagonally on $H_c, C_c(\mathfrak{t}^*), \Gamma \rtimes W$ and W itself. The operator is given by T on H_c and by the identity on the other factors. This is a well defined adjointable operator as we took T to be exactly invariant under the action of $\Gamma \rtimes W$ and of finite propagation.

Applying the element $\widehat{\beta}_{\mathbb{C}}$ reduces this to a module over $C_0(\mathfrak{t}^*) \rtimes (\Gamma^\vee \rtimes W)$ where the inner product is

$$\langle \xi \otimes f[(g, w)], \xi' \otimes f'[(g', w')] \rangle = \sum_{\delta \in \Gamma} \sum_{\chi \in \Gamma^\vee} \langle \xi, \delta \cdot \xi' \rangle \langle f[w], [\chi] e^{2\pi i \langle \eta, \delta \rangle} f'[w'] \rangle.$$

Note that as this no longer depends on g and g' , vectors of the form $\xi \otimes f[(g_1, w)]$ and $\xi \otimes f[(g_2, w)]$ are identified. Thus the module is really a completion of $H_c \widehat{\otimes} C_c(\mathfrak{t}^*) \widehat{\otimes} \mathbb{C}[W]$, which we will denote \mathcal{E}_1 . Once again the representation is provided by the trivial representation of W , and we denote the corresponding projection on \mathcal{E}_1 by p_W . The operator on \mathcal{E}_1 is given by $T \otimes 1 \otimes 1$.

We now trace the other route around the diagram. As before, starting with a Kasparov triple (H, ρ, T) we obtain the descended triple $(\mathcal{E}, \hat{\rho}, T \otimes 1)$. We next apply the element $\mathbf{A}_{C_0(\mathfrak{t})}$ which is given by the completion of $C_c(\mathfrak{t})$ described earlier in this section. We obtain the completion of $H_c \widehat{\otimes} \mathbb{C}[\Gamma \rtimes W]$ with respect to the inner product

$$\langle \xi[g], \xi'[g'] \rangle = \sum_{h \in \Gamma \rtimes W} \langle \xi, h \cdot \xi' \rangle [g^{-1} h g'].$$

The representation of \mathbb{C} is given by the identity while the operator, once again, is given by T on H_c and the identity on the other factor.

The Hilbert module realising the descended Morita equivalence is given by completing the module $C_c(\mathfrak{t}^*) \rtimes W$ with respect to the inner product

$$\langle f[w], f'[w'] \rangle = \sum_{\chi \in \Gamma^\vee} [w^{-1} \bar{f}(\chi \cdot f')] [\chi w']$$

in $C_0(\mathfrak{t}^*) \rtimes W$.

The representation of $C^*(\Gamma \rtimes W)$ on this module is given by the representation of $\Gamma \rtimes W$ where $((\gamma w') \cdot f[w])(\eta) = e^{2\pi i \langle \eta, \gamma \rangle} (w' \cdot f)(\eta)[w'w]$. Hence applying the Morita equivalence we obtain a Kasparov triple where the module is the completion, which we denote by \mathcal{E}_2 , of $H_c \widehat{\otimes} C_0(\mathfrak{t}^*) \widehat{\otimes} \mathbb{C}[W]$ with respect to the inner product

$$\langle \xi \otimes f[w], \xi' \otimes f'[w'] \rangle = \sum_{\delta \in \Gamma} \sum_{u \in W} \sum_{\chi \in \Gamma^\vee} \langle \xi, (\delta u) \cdot \xi' \rangle \langle f[w], e^{2\pi i \langle \eta, \delta \rangle} [\chi u] f'[w'] \rangle,$$

the representation of \mathbb{C} is given by the identity and the operator is given by T on H_c and the identity elsewhere.

To identify this triple with the Kasparov element obtained via the first route, we note that the module \mathcal{E}_2 is isomorphic to the range of the projection p_W on \mathcal{E}_1 . Indeed

$$\langle p_w(\xi \otimes f[w]), p_w(\xi' \otimes f'[w']) \rangle_{\mathcal{E}_1} = \frac{1}{|W|} \langle \xi \otimes f[w], \xi' \otimes f'[w'] \rangle_{\mathcal{E}_2}.$$

This completes the proof.

5. LANGLANDS DUALITY AND K -THEORY

In this section we will consider the K -theory of the affine and extended affine Weyl groups of a compact connected semisimple Lie group.

As remarked in the introduction an extended affine Weyl group and its Langlands dual $(W'_a)^\vee$ need not be isomorphic. For example the extended affine Weyl groups of PSU_3 and its Langlands dual SU_3 are non-isomorphic. However their group C^* -algebras have the same K -theory, see [12].

In this section we will show that this is not a coincidence, indeed passing to the Langlands dual always rationally preserves the K -theory for the extended affine Weyl groups. In particular, where the extended affine Weyl group of the dual of G agrees with the affine Weyl group of G (as for PSU_3) the K -theory for the affine and extended affine Weyl groups of G agrees up to rational isomorphism.

Corollary 1.3. *Let G be a compact connected semisimple Lie group and G^\vee its Langlands dual, with $W'_a, (W'_a)^\vee$ the corresponding extended affine Weyl groups. Then there is a rational isomorphism*

$$K_*(C^*((W'_a)^\vee)) \cong K_*(C^*(W'_a)).$$

Proof. The proof combines the universal coefficient theorem with our Poincaré duality as follows.

We start by writing $W'_a = \Gamma \rtimes W$ and $(W'_a)^\vee = \Gamma^\vee \rtimes W$. By the Green Julg theorem and Fourier-Pontryagin duality:

$$(3) \quad K_*(C^*(W'_a)^\vee) \cong K_*^W(C^*(\Gamma^\vee)) \cong K_*^W(C(T)) = K_W^*(T).$$

Applying the universal coefficient theorem, we have the exact sequence

$$0 \rightarrow \text{Ext}_{\mathbb{Z}}^1(K_W^{*-1}(T), \mathbb{Z}) \rightarrow K_*^W(T) \rightarrow \text{Hom}(K_W^*(T), \mathbb{Z}) \rightarrow 0$$

In particular the torsion-free part of $K_*^W(T)$ agrees with the torsion-free part of $K_W^*(T)$ therefore rationally we have

$$(4) \quad K_W^*(T) \cong K_*^W(T).$$

As in 1.2 we can identify $K_*^W(T) = K_W^*(C(T))$ with $K^*(C^*(W'_a)^\vee)$. The theorem now follows by applying our Poincaré duality from Theorem 1.2 to obtain

$$(5) \quad K^*(C^*(W'_a)^\vee) \cong K_*(C^*(W'_a)).$$

□

In a subsequent paper, [12] we construct the admissible duals for the extended affine Weyl groups of all Lie groups of type A_n , exhibiting these spaces as varieties which decompose as a union of spaces indexed by the representations of the Weyl group. Furthermore we show that the rational isomorphism given above is induced by a homotopy equivalence between the varieties which respects the decomposition. The special case of $SU(n)$ itself was considered by Solleveld in [13].

For the affine Weyl groups we have the following:

Corollary 1.4. *Let W'_a be the extended affine Weyl group of G , and let W_a, W_a^\vee be the affine Weyl groups of G and its Langlands dual G^\vee . If G is of adjoint type then rationally*

$$K_*(C^*(W_a^\vee)) \cong K_*(C^*(W'_a)).$$

If additionally G is of type $A_n, D_n, E_6, E_7, E_8, F_4, G_2$ then rationally

$$K_*(C^*(W_a)) \cong K_*(C^*(W'_a)).$$

Proof. If G is a compact connected semisimple Lie group of adjoint type then its Langlands dual G^\vee is simply connected so $(W'_a)^\vee = W_a^\vee$.

In the case that G is additionally of type $A_n, D_n, E_6, E_7, E_8, F_4, G_2$ the group G^\vee is the universal cover of G and hence $W_a = W_a^\vee$. □

REFERENCES

- [1] S. Baaj and P. Julg, Théorie bivariante de Kasparov et opérateurs non bornés dans les C^* -modules hilbertiens, C.R. Acad. Sci. Paris Sér. I Math. 296 (1983) 875 – 878.
- [2] B. Blackadar, *K-Theory for operator algebras* (Cambridge University Press, 2002.)
- [3] J. Block, Duality and Equivalence of Module Categories in Noncommutative Geometry, CRM Proceedings and Lecture Notes Volume 50, 2010, 311-340
- [4] N. Bourbaki, *Lie groups and Lie algebras* (chapters 4-6, Springer 2002; chapters 7-9, Springer 2008).
- [5] U. Bunke and T. Nikolaus, T-duality via gerby geometry and reductions, Rev. Math. Phys. 27 (2015), no. 5, 1550013, 46
- [6] C.Daenzer and E. van Erp, T-duality for Langlands dual groups, Advances in Theoretical and Mathematical Physics, 18(6), (2014) 1267–1285.
- [7] S. Echterhoff, H. Emerson, H. J. Kim, KK -theoretic duality for proper twisted actions, Math. Ann. (2008) 340:839–873.
- [8] H. Emerson, Lefschetz numbers for C^* -algebras. Canad. Math. Bull. 54 (2011), no. 1, 82–99.

- [9] H. Emerson and R. Meyer, Dualities in equivariant Kasparov theory. New York J. Math. 16 (2010), 245–313.
- [10] G. Kasparov, Equivariant KK-theory and the Novikov conjecture Invent. math. 91 (1988) 147–201.
- [11] R. Meyer, Categorical aspects of bivariant K-theory. K-theory and noncommutative geometry, 1–39, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2008.
- [12] G.A.Niblo, R. Plymen, N.J.Wright, Stratified Langlands duality in the A_n tower, arXiv:1611.05218.
- [13] M.S. Solleveld, Periodic cyclic homology of affine Hecke algebras, PhD Thesis, 2007, FNWI: Korteweg-de Vries Institute for Mathematics (KdVI), <http://hdl.handle.net/11245/1.271180>.

MATHEMATICAL SCIENCES, UNIVERSITY OF SOUTHAMPTON, SO17 1BJ, ENGLAND
E-mail address: `g.a.niblo@soton.ac.uk`, `r.j.plymen@soton.ac.uk`, `wright@soton.ac.uk`