

Strontium concentration, radiogenic ($^{87}\text{Sr}/^{86}\text{Sr}$) and stable ($\delta^{88}\text{Sr}$) strontium isotope systematics in a controlled feeding study.

Lewis^{1*}, J., Pike², A.W.G., Coath¹, C. D. and Evershed³, R. P.

¹*Bristol Isotope Group, School of Earth Sciences, University of Bristol, Wills Memorial Building, Queen's Road, Bristol BS8 1RJ, UK,*

²*Department of Archaeology, University of Southampton, Avenue Campus, Highfield, Southampton, SO17 1BF, UK*

³*Organic Geochemistry Unit, Biogeochemistry Research Centre, School of Chemistry, University of Bristol, Cantock's Close, Bristol, BS8 1TS, UK*

Strontium concentration, radiogenic ($^{87}\text{Sr}/^{86}\text{Sr}$) and stable ($\delta^{88}\text{Sr}$) strontium isotope systematics in a controlled feeding study.

Transhumance and palaeodiet are two central themes in archaeology and using chemical analysis of bones and teeth to reconstruct trends and patterns in diet and mobility has become a cornerstone of bioarchaeology. This study has investigated strontium concentration ([Sr]), radiogenic ($^{87}\text{Sr}/^{86}\text{Sr}$) and stable strontium ($\delta^{88}\text{Sr}$) isotope systematics in a controlled feeding experiment on domestic pigs designed to simulate terrestrial versus marine protein consumption. The results of the radiogenic ($^{87}\text{Sr}/^{86}\text{Sr}$) analysis offer a validation of the strontium isotope methodology. The study confirms that the radiogenic strontium isotope composition of dental enamel does represent the radiogenic strontium isotope composition of the diet. The results of the $\delta^{88}\text{Sr}$ analysis have revealed a distinct shift of 0.322 ± 0.060 ‰ towards isotopically light Sr with trophic level. The magnitude of this shift is consistent with the predictions from the analogous shift observed in calcium isotopes. This is the first time that trophic level fractionation in $\delta^{88}\text{Sr}$ has been identified in a controlled setting. Although still in its infancy, $\delta^{88}\text{Sr}$ analysis has great potential to inform on trophic level systematics, to investigate dietary trends in early life and is potentially useful in examining diagenetic alteration.

Strontium isotopes, palaeodietary reconstruction, biogeochemistry

Introduction

Provenance and palaeodietary reconstruction are two central themes in (bio)archaeology and the use of isotopic techniques to elucidate trends and patterns in these areas has become a cornerstone of bioarchaeological research. Use of the mass fractionation corrected ratio of ^{87}Sr to ^{86}Sr abundances, hereafter radiogenic $^{87}\text{Sr}/^{86}\text{Sr}$ or simply $^{87}\text{Sr}/^{86}\text{Sr}$, in dental enamel to provenance the geographical source of dietary Sr in order to assess mobility in archaeological populations is now an established technique (Bentley, 2006; Ericson, 1985; Price et al., 2002; Sealy et al., 1991). Studies using this technique have a broad scope, ranging from those assessing a few individuals from a single site to answer specific questions (e.g. Taylor et al. 2013; Montgomery, Budd, and Evans 2000) to larger scale regional studies aimed at investigating broader social trends at the population level (e.g. Haverkort et al. 2008; Bentley et al. 2012). There has also been an increasing use of this technique to study the provenance and management strategies of archaeological fauna (Viner et al 2010; Knipper 2011; Stephan et al. 2012).

Strontium isotopes do not as yet have a substantial role in the field of palaeodietary reconstruction. This is principally addressed through analysis of $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\delta^{34}\text{S}$ in bone collagen (DeNiro and Epstein, 1978; Van der Merwe and Vogel, 1978; Chisholm et al., 1982; Schoeninger et al., 1983; Schoeninger and DeNiro, 1984; Nehlich et al. 2011), as well as $\delta^{13}\text{C}$ in bone and tooth apatite structural carbonate ($\delta^{13}\text{C}_{\text{SC}}$) (Lee Thorpe and Van der Merwe, 1991) and, to a lesser extent, by analysis of other stable isotopes in calcified tissues e.g. $\delta^{44}\text{Ca}$ (Skulan and DePaolo, 1999; Reynard et al., 2010). The range of isotopic techniques available for archaeologists studying palaeodiet has recently been expanded by the measurement of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ on the individual amino acids which collagen comprises (Howland et al., 2003; Jim et al., 2003). Amino acid analysis has been shown to be particularly useful in resolving marine

protein consumption in archaeological populations where aridity and the presence of C₄ plants render bulk collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements ineffective (Corr et al., 2005; Styring et al., 2010; Styring et al., 2015).

Analysis of strontium concentrations in calcified tissues, expressed relative to the major element calcium, has been used as a palaeodietary proxy (Sealy, 2001). The technique relies on biogeochemical cycling of strontium from the biosphere into the food chain but exploits the successive decrease in the elemental Sr/Ca ratio of calcified tissues with trophic level as a result of Ca biopurification in the digestive track (Comar et al., 1957; Sillen, 1981; Elias et al., 1982). This has been used to assess the relative proportion of plant and meat components in an individual's diet (Toots and Voorhies, 1965; Burton and Wright, 1995; Burton et al., 1999; Balter et al., 2012).

Initial work on biogeochemical cycling of strontium utilised natural experiments in ecosystems (Price et al., 1985) and small scale laboratory experiments (Price et al., 1986). Much of this work focused on assessing the magnitude and controls on Sr/Ca discrimination with trophic level and this work was subsequently incorporated into $^{87}\text{Sr}/^{86}\text{Sr}$ methodologies for migration studies. Whilst the validity of $^{87}\text{Sr}/^{86}\text{Sr}$ in assigning geographical provenance to individuals has been proven in archaeological and modern studies (e.g. Britton et al., 2009; Tütken et al., 2011) we are aware of no published study to date that has attempted to demonstrate this under controlled conditions.

Strontium isotopes undergo mass-dependent fractionation in nature. Although the magnitude of the fractionation is small compared to light stable isotope systems (e.g. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$), Sr isotopes do fractionate in a mass-dependent manner as a result of small differences in the thermodynamic properties of individual isotopes which are an inherent function of their mass (Urey, 1947). In radiogenic strontium isotope ($^{87}\text{Sr}/^{86}\text{Sr}$)

studies, any mass-dependent fractionation that occurs both in nature and during analysis is corrected for by normalisation to a fixed $^{86}\text{Sr}/^{88}\text{Sr}$ ratio (0.1194, Nier 1938) using an assumed mass-fractionation law. Thus, mass-dependent strontium isotope fractionation does not compromise mobility studies using $^{87}\text{Sr}/^{86}\text{Sr}$.

Measurements of the mass-dependent isotopic fractionation of strontium, $\delta^{88}\text{Sr}$, have been used to study the composition of the early solar system as preserved in meteorites (Patchett, 1980a; Patchett, 1980b; Moynier et al., 2010; Charlier et al., 2012) and used alongside $^{87}\text{Sr}/^{86}\text{Sr}$ as a geological tracer (Pearce et al., 2015; Pearce et al., 2015). $\delta^{88}\text{Sr}$ has been identified as a potential palaeo-thermometer with studies showing variation in the magnitude of $\delta^{88}\text{Sr}$ fractionation between seawater and cold water corals as a function of ocean temperature (Fietzke and Eisenhauer, 2006; Rüggeberg et al., 2008; Krabbenhöft et al., 2010; Vollstaedt et al., 2014; Stevenson et al., 2014). $\delta^{88}\text{Sr}$ has also been measured in fossil apatites (Conodonts) with such samples having the potential to be highly enriched in ^{86}Sr with $\delta^{88}\text{Sr}$ values in excess of -1 ‰ (Neymark et al., 2014).

Strontium isotopes have also been shown to undergo mass-dependent fractionation between soils and plants growing in those soils with plants being enriched in the lighter ^{86}Sr isotope by approximately 0.3 ‰ (de Souza et al., 2010).

Recently, $\delta^{88}\text{Sr}$ has been introduced as a potential palaeodietary tracer for archaeology by Knudson et al. (2010). Building on geological studies of strontium isotope fractionation, and by analogy with calcium isotope fractionation, Knudson et al. posited that Sr should also undergo mass-dependent fractionation with trophic level. Knudson et al. tested this hypothesis on series of Chiribaya-affiliated sites from southern Peru. Their results showed an approximately 0.3 ‰ difference in $\delta^{88}\text{Sr}$

between molluscs and both small and large herbivores confirming the potential for $\delta^{88}\text{Sr}$ as a palaeodietary tracer.

Regarding tracking trophic level with $\delta^{15}\text{N}$, it has been apparent for some time that for a full understanding of the contribution of plant versus meat protein in the diet it is essential to have an accurate estimate of $\Delta^{15}\text{N}_{\text{diet-tissue}}$ (the magnitude of the trophic level shift) (Hedges and Reynard, 2007; O'Connell et al. 2012). Furthermore, it is also necessary to know how $\Delta^{15}\text{N}_{\text{diet-tissue}}$ can vary with dietary protein source (e.g. marine vs. terrestrial) and total amount of protein in the diet. Although $\delta^{88}\text{Sr}$ should track trophic level across the whole diet rather than just the protein component, similarly to $\delta^{13}\text{C}_{\text{SC}}$, the potential for misinterpretation remains unless $\Delta^{88}\text{Sr}_{\text{diet-tissue}}$ can be adequately constrained.

In this study we present strontium concentration ($[\text{Sr}]$), Sr/Ca , $^{87}\text{Sr}/^{86}\text{Sr}$, and $\delta^{88}\text{Sr}$ determinations made on a series of pigs raised in a controlled feeding study in order to (i) demonstrate under controlled conditions a fixed relationship between $^{87}\text{Sr}/^{86}\text{Sr}_{\text{diet}}$ and $^{87}\text{Sr}/^{86}\text{Sr}_{\text{enamel}}$ and (ii) investigate and characterize the fractionation of $\delta^{88}\text{Sr}$ with trophic level.

The feeding experiment

The controlled feeding experiment from which the samples for this study were obtained was designed to assess the effect of marine protein consumption on the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ composition of various pig tissues at the bulk and single amino-acid level. A detailed description of the feeding experiment design and results including growth rates, for the pigs are presented elsewhere (Webb et al., 2016; Webb et al., in press;); only brief summary is given here.

All pigs were raised at Harper Adams University (Shropshire, UK). Pigs in the controlled feeding experiment were each fed exclusively on one of five diets of known

protein source composition ranging from a 100 % marine source (fish meal) to a 100% terrestrial source (soy). The five diets consist of 100 %, 50 %, 25 %, 12.5 % and 0 % marine protein with the remainder of the dietary protein being made up from the terrestrial source. All diets are nutritionally equivalent, have the same amount of total protein and differ only in the source of the dietary protein (Webb et al., 2016, figure 1).

To ensure the pigs reached equilibrium with the diets the experiment was run over two successive generations. The first generation (G1) gilts (young female pigs that have not yet reproduced), 3 per feeding group, were weaned on to the diets and fed on them until sacrifice as adults (>18 months). The gilts were artificially inseminated and the second generation of pigs (G2) fed on the same test diets as their mothers (including sow milk from the same feeding group) from weaning until they were sacrificed as adolescents in two groups at 5 and a half and 6 and half months old respectively

Although the feeding experiment represents a complete range of marine to terrestrial protein diets with individuals of a range of ages and with replicated individuals on the same diet it was not initially conceived to examine the uptake of dietary strontium. Therefore feed ingredients were not specifically selected for their strontium isotope compositions prior to the experiment starting. Teeth from the feeding experiment were initially sampled as a potential $^{87}\text{Sr}/^{86}\text{Sr}$ standard for laser ablation (Lewis et al., 2014). It was hoped that the strontium isotope signature of the terrestrial protein source would differ sufficiently from the marine protein source ($^{87}\text{Sr}/^{86}\text{Sr}$ 0.70918-0.70920, McArthur et al., 2001, $\delta^{88}\text{Sr}$ 0.381 ‰ Fietzke and Eisenhauer, 2006) and that the strontium isotope composition of the pigs on mixed diets would plot between the two end members.

2. Material and methods

2.1 Samples

A range of materials from the feeding experiment have been analysed for [Sr], [Ca], $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}\text{Sr}$ to assess the dietary uptake of strontium. Each of the 5 feeds were analysed as well as drinking water from a borehole at the Harper Adams site, which is common to all feed groups. Tooth enamel was sampled in 5 G1 adults and 14 G2 adolescents with paired dentine measurements made on 5 of the G2 adolescents. In each case the latest erupting lower left molar was sampled, M₃ for G1 and M₁ for G2. One sample of faeces was analysed from each feeding group, four of the faeces samples can be paired to individual G2 pigs.

2.2 Sample preparation

Tooth enamel and dentine samples (10-20 mg) were sawn from whole teeth and weighed into clean perfluoroalkoxy (PFA) beakers and dissolved in 7N HNO₃ (Romil SpA). Approximately 500 ml of drinking water from the site was evaporated to dryness in a PFA beaker and the residue dissolved in 1:1 7N HNO₃/6N HCl (Romil SpA).

Faeces were freeze-dried to remove moisture and picked under a microscope to remove straw, which is used for bedding at the Harper Adams site. Approximately 20 mg of faeces were weighed into PFA beakers. Feed pellets (100-200 mg) were crushed in a mortar and pestle and then weighed into PFA beakers. Due to their high organic contents feed and faeces were digested in a high-pressure asher (HPA-S, Anton Paar). Feed and faeces were pre-digested in 3:1 7N HNO₃/6N HCl then transferred to clean quartz glass vials. The vials were sealed and ashed at 300°C and ~100 bar for three hours. Solutions are then transferred back to PFA beakers. Following digestion all samples were evaporated to dryness and re-dissolved in 3N HNO₃.

2.3 Double-spike isotopic tracer

$\delta^{88}\text{Sr}$ determinations are made by using a double-spike isotope tracer (Dodson, 1963; Dodson, 1970). The double-spike tracer method involves the mixing of samples with a tracer artificially enriched in two isotopes. The advantage of this method is that any isotopic fractionation resulting from the sample preparation, chemical separation and mass spectrometry, which also fractionates the tracer, is corrected for. Consequently, incomplete recovery of the strontium, which may fractionate the isotopes, does not affect the $\delta^{88}\text{Sr}$ determination (Rudge et al., 2009).

The tracer, enriched in ^{84}Sr and ^{87}Sr , was prepared from two ‘single-spike’ solutions, enriched in ^{84}Sr and ^{87}Sr respectively, prepared from isotopically enriched strontium carbonate purchased from Oak Ridge National Laboratory. Solutions of the single-spikes were mixed in such proportions to minimise the uncertainty in the calculated isotope composition of samples in sample-tracer mixtures (Rudge et al., 2009). To back-out the sample composition from a measurement of the mixture, the precise tracer composition must be determined, a procedure called the calibration of the tracer (Coath et al., 2017; Rudge et al., 2009).

The calibration procedure consists of the preparation and analysis of a series of J mixtures of standard and tracer ranging from pure standard to pure tracer. Following Rudge et al., 2009, the instrumental fractionation corrected measurements must lie on a line in ($^{84}\text{Sr}/^{86}\text{Sr}$, $^{87}\text{Sr}/^{86}\text{Sr}$, $^{88}\text{Sr}/^{86}\text{Sr}$)-space passing through the standard composition. The algorithm solves for the J instrumental fractionation parameters and the unit vector in the direction of the line. The solution for the unit vector is critical to accurate measurements whereas the position on the line where the tracer composition lies is much less so, the latter only having a second-order effect for compositions whose fractionation relative to the standard is small. Nevertheless, the actual tracer

composition can be determined as that point on the line lying closest, in a maximum-likelihood sense, from the instrumental fractionation corrected measurement of the pure tracer. Using this method to characterise the tracer, the composition was found to be 48.7815 % ^{84}Sr , 2.5144 % ^{86}Sr , 38.0902 % ^{87}Sr and 10.6139 % ^{88}Sr .

Any isotopic fractionation resulting from dissolution may be corrected for by adding the tracer prior to dissolution. However, Lewis (2015) has shown that no measurable fractionation occurs, even when the high-pressure asher is used to dissolve organic-rich samples. Therefore, the tracer is added to an aliquot of the dissolved solution for $\delta^{88}\text{Sr}$ determinations and further aliquots of the same solution are used for the $[\text{Sr}]$ and $^{87}\text{Sr}/^{86}\text{Sr}$ determinations. Double-spike is added to the sample to give a final spike/sample proportion of approximately 0.3:0.7, which minimises the uncertainty in the calculated sample isotopic composition (Rudge et al., 2009).

2.4 Sample chemistry and mass spectrometry

Strontium separation was accomplished using Sr Spec resin (Horwitz et al., 1992). Approximately 100 μl of pre-cleaned resin was loaded into PFA micro-columns. Between 100 and 500 ng of strontium was loaded on to columns in 0.5 ml of 3N HNO_3 , matrix was eluted in 2 ml of 3N HNO_3 and Sr collected in 1.5 ml of ultrapure water. Separate micro-columns were used for $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}\text{Sr}$ samples.

For $[\text{Sr}]$ and $[\text{Ca}]$ determinations 5 μL of the final dissolved solution was taken, 25 μL of 200 ppb indium solution added as an internal standard and the solution was then diluted to a final volume of 2 ml with 0.3N HNO_3 . Concentrations were measured on a Thermo-Electron Element 2 ICPMS, internally normalised to indium and quantified by external calibration relative to a gravimetrically prepared in-house multi-element standard.

The determinations of $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}\text{Sr}$ were made following the procedure outlined in Lewis et al. (2014), purified Sr was loaded onto rhenium filaments with a TaCl_5 activator (Birck, 1986). Strontium isotope ratios were measured on a Thermo-Finnigan Triton thermal ionisation mass spectrometer. Amplifier gains were calibrated before each analytical session and all isotope ratios were determined in static collection mode with a 4.194 s integration time per cycle and 12 cycles per block, ^{87}Sr beams were corrected for isobaric ^{87}Rb by monitoring ^{85}Rb and using a $^{85}\text{Rb}/^{87}\text{Rb}$ ratio of 2.59265 modified to approximate the effect of instrumental mass bias (Steiger and Jäger, 1977).

For $^{87}\text{Sr}/^{86}\text{Sr}$, data were collected as 30 blocks and corrected for mass fractionation using an exponential mass bias law and a $^{86}\text{Sr}/^{88}\text{Sr}$ of 0.1194 (Nier, 1938; Russell et al., 1978). All data were externally normalised by measuring NIST SRM 987 and assuming a $^{87}\text{Sr}/^{86}\text{Sr}$ ratio of 0.710248 (Avanzinelli et al., 2005). Long-term reproducibility of $^{87}\text{Sr}/^{86}\text{Sr}$ for NIST SRM 987 analyses (not externally normalised) was 0.710232 ± 0.000036 (2 SD, $n = 146$) and a modern seal tooth 0.709180 ± 0.000033 (2 SD, $n = 25$).

For the $\delta^{88}\text{Sr}$ analyses, data were collected as 75 blocks and the sample fractionation solved using the double-spike inversion function provided in the 'Double Spike Toolbox' (Rudge et al., 2009) running under MATLAB. All data were externally normalised to measured NIST SRM 987 and fractionation of the $^{86}\text{Sr}/^{88}\text{Sr}$ reported in permil deviation from NIST SRM 987 with $(^{86}\text{Sr}/^{88}\text{Sr})_{\text{SRM987}} = 0.1194$. Long-term reproducibility of $\delta^{88}\text{Sr}$ for NIST SRM 987 was $-0.0016 \text{ ‰} \pm 0.031 \text{ ‰}$ (2 SD, $n = 87$) and for a modern seal tooth $-0.2383 \text{ ‰} \pm 0.033 \text{ ‰}$ (2 SD, $n = 12$).

3. Results

3.1 Sr concentration and Sr/Ca ratio

For all substrates except drinking water, there is a strong positive correlation (PCC, $R^2 > 0.96$, $p < 0.001$) between the marine protein content of the diet and the [Sr] of the substrate (Fig 2, table 1a). This is consistent with the marine protein source (fish meal) being enriched in Sr relative to the terrestrial protein source (soya) and the remaining non-protein components of the diets which is identical in each feeding group.

The pig feed has low [Sr] ranging from 13 ppm ($\mu\text{g/g}$) in the 100 % terrestrial protein diet to 36 ppm in the 100% marine protein diet. Dental tissues, with [Sr] ranging from 75 ppm to 300 ppm, are concentrated in Sr relative to feed. This is consistent with strontium in mammals being concentrated in the calcified tissues. For each feeding group the [Sr] of the faeces is enriched over the feed and has [Sr] similar to teeth. This is initially surprising given the low [Sr] of the feed however absorption via the gut discriminates against strontium relative to nutrients from the bulk feed hence digestion concentrates strontium in the faeces.

The correlation between [Sr] and the marine protein content of the diet is mirrored in the Sr/Ca ratio (expressed as $\text{Sr/Ca} \times 1000$) of the pig feed and pig dental tissues (PCC, $R^2 > 0.98$, $p < 0.001$, table 1b). However, irrespective of the initial Sr/Ca of the individual diets a consistent decrease was observed in Sr/Ca between the feed and the dental tissues of 0.242 ± 0.044 (2 SD, $n=5$) as a result of calcium biopurification. This value is within the uncertainty of the canonical 5 fold decrease in Sr/Ca with trophic level and in good agreement with the range of values in the literature for this decrease (Elias et al., 1982; Sillen and Kavanagh, 1982; Burton et al., 2003).

There has been some debate regarding the investigation of resource consumption with Sr/Ca ratios (Schoeninger and Peebles, 1981; Burton and Price, 1999). Although

these new data would suggest that marine consumption could be detected with Sr/Ca ratio we do not make this assertion. We consider that the correlation between marine protein content and Sr/Ca ratio is a result of ingredient selection for the experimental feeds. We further note that in our feeds only the protein component of the diet is substituted for a marine source, in cases of marine resource exploitation the whole diet would be from a marine source. Therefore, these results are not relevant in determining if Sr/Ca can be used to detect marine resource consumption.

3.2 Radiogenic strontium ($^{87}\text{Sr}/^{86}\text{Sr}$)

Dental tissues from the feeding experiment have a mean $^{87}\text{Sr}/^{86}\text{Sr}$ of 0.709089 ± 0.000075 (2 SD, n=25) whilst the pig feed and faeces have a mean $^{87}\text{Sr}/^{86}\text{Sr}$ of 0.709056 ± 0.000058 and 0.709055 ± 0.000083 (both 2 SD, n=5) respectively (Fig 2, table 1c). Thus, there is no significant difference in $^{87}\text{Sr}/^{86}\text{Sr}$ between the pig teeth and either the diets or the faeces (Kruskal-Wallis, $p=0.132$). In addition, there is no difference between the G1 or G2 pigs indicating that both generations of the feeding experiment are at equilibrium with the dietary strontium (Mann-Whitney, $p=0.644$). The agreement between dietary $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{87}\text{Sr}/^{86}\text{Sr}$ in dental tissues supports the current use of strontium isotopes in dental enamel as a proxy for dietary strontium.

Across all substrates there is no correlation between a modern marine $^{87}\text{Sr}/^{86}\text{Sr}$ ratio (0.70918-0.70920, McArthur et al., 2001) and the marine protein content of the diets. Two reasons are postulated for this.

I) The bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ of the area where the soya was grown may be similar to that of modern seawater and thus the isotopic difference between the marine and terrestrial protein sources is minimal which is quite possible if the soya was grown in a soil derived from a young carbonate geology. Unfortunately we do not know the

provenance of the soya. Furthermore soya is added as ground soya flour thus we cannot rule out that the soya used in the feeds is from multiple locations, with different bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$, which are then homogenised during processing. This is unsatisfactory from a Sr isotope perspective but was acceptable for the C and N isotope systems the feeding experiment was originally designed to investigate. In this case the terrestrial end member was required to have a C_3 carbon isotope signature (soya is a C_3 plant) in order to be distinct from the marine end member.

II) The mean $^{87}\text{Sr}/^{86}\text{Sr}$ of the non-protein component (principally starch, calcium phosphate, wheat fibre and vitamin/mineral supplements) of the diet is similar to that of seawater and that the contribution of this Sr to the total diet means that the signal from the terrestrial protein source is not well resolved in the dietary mean. Indeed, the above scenarios are not mutually exclusive and that the soya may or may not have a modern seawater $^{87}\text{Sr}/^{86}\text{Sr}$ value but also may be depleted in Sr such that it contributes very little Sr to the dietary average. In either case, both scenarios account for the observation that Sr from the fish meal is detected in the [Sr] but not in $^{87}\text{Sr}/^{86}\text{Sr}$. It is unfortunate in this experiment that the marine and terrestrial diets are so similar in $^{87}\text{Sr}/^{86}\text{Sr}$ that this study offers little insight into how the $^{87}\text{Sr}/^{86}\text{Sr}$ of dental tissues in terrestrial animals is affected by marine resource consumption.

Across all feed groups, G1 and G2 pig teeth have marginally more radiogenic $^{87}\text{Sr}/^{86}\text{Sr}$ than their respective feeds. Although this difference is not statistically significant (see above), this small offset is attributed to the contribution of strontium from the pigs' drinking water, which is more radiogenic than the feeds ($^{87}\text{Sr}/^{86}\text{Sr}$, 0.711572). Taking the mean $^{87}\text{Sr}/^{86}\text{Sr}$ of the feed and the dental tissues, water contributes only 1.4 % of the strontium by mass and, even in the most radiogenic tooth observed, water contributes < 5% of the total strontium. It is also noted that this result is

for a 2.6 ppm Sr water source, which is reasonably concentrated, the mean values for ground waters in England and Wales is 0.64 ppm with a 5 to 95% range of 0.02 ppm to 5 ppm (Shand et al., 2007).

3.3 Stable strontium ($\delta^{88}\text{Sr}$)

The pig feeds and faeces have mean $\delta^{88}\text{Sr}$ of 0.098 ± 0.054 ‰ and 0.127 ± 0.031 ‰ (both 2 SD, n=5) respectively. The pig dental tissues have mean $\delta^{88}\text{Sr}$ of -0.224 ± 0.044 ‰ (2 SD, n=18) (Fig 4, table 1d), which is 0.322 lower than the feed. As with $^{87}\text{Sr}/^{86}\text{Sr}$, there is no significant difference between the G1 and G2 teeth (Mann-Whitney, p=0.412).

The mean $\Delta^{88}\text{Sr}_{\text{diet-teeth}}$ of 0.322 ± 0.060 ‰ (2 SD) is attributed to a trophic level fractionation in strontium isotopes. The magnitude of the observed fractionation is in excellent agreement with predictions from the analogous fractionation which has been observed in $\delta^{44/40}\text{Ca}$. Skulan and DePaolo, (1999) have demonstrated that $\Delta^{44/40}\text{Ca}_{\text{diet-bone}}$ is 1.3 ‰. Based on the mass difference between Ca and Sr isotopes for a 1.3 ‰ shift in $\delta^{44/40}\text{Ca}$ a kinetic fractionation law predicts a 0.314 ‰ fractionation in $\delta^{88/86}\text{Sr}$, which is within the uncertainty of the observed fractionation. This measurement of the magnitude of the fractionation is also consistent with the estimations of de Souza et al. (2010) and Knudson et al. (2010) who have both shown that $\Delta^{88}\text{Sr}$ is approximately 0.3 ‰.

As with $^{87}\text{Sr}/^{86}\text{Sr}$, no trend is observed towards a modern seawater $\delta^{88}\text{Sr}$ (0.381 ‰, Fietzke and Eisenhauer, 2006) with increasing marine protein content of the diet. However, this may not necessarily be expected as the fish meal will itself be fractionated from that of modern seawater and the remainder of the diet and/or the terrestrial protein source have a $\delta^{88}\text{Sr}$ value sufficiently similar to that of the fish meal

that no trend may be observed. Measurements of the individual ingredients that comprise the feeds may elucidate this.

The five samples of faeces have $\delta^{88}\text{Sr}$ within the uncertainty of the feeds. One might expect excreta to be elevated in $\delta^{88}\text{Sr}$ to balance fractionation observed in the calcified tissue. However, there are two points to consider. First, from the [Sr] and Sr/Ca results it is clear that a large proportion of the dietary strontium is not absorbed into the body. Second, by analogy with calcium isotopes, it is suspected that $\delta^{88}\text{Sr}$ fractionation occurs during mineralisation when hydroxyapatite crystals are precipitated. Thus, strontium that is not absorbed from the gut will remain unfractionated and obscure the signal from any isotopically heavy strontium being excreted. Furthermore, if Sr fractionation does occur at the point of mineralisation then the pig urine is a more likely candidate for the isotopic mass balance as the isotopically heavy Sr resulting from the mineralisation will be removed as urine rather than faeces.

4. Discussion

4.1 Defining the local bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ and the minimum $^{87}\text{Sr}/^{86}\text{Sr}$ variation in a population

At the broadest scale this study offers a validation of the archaeological $^{87}\text{Sr}/^{86}\text{Sr}$ method. The excellent agreement between $^{87}\text{Sr}/^{86}\text{Sr}_{\text{diet}}$ and $^{87}\text{Sr}/^{86}\text{Sr}_{\text{enamel}}$ confirm that with respect to $^{87}\text{Sr}/^{86}\text{Sr}$ “you are what you assimilate”. Whilst the validity of this assertion has been proven in both archaeological contexts and modern natural experiments this is the first study where this has been demonstrated under controlled conditions.

Beyond the validation of the $^{87}\text{Sr}/^{86}\text{Sr}$ method this study can also be used to inform on the minimum variation that can be expected from the local $^{87}\text{Sr}/^{86}\text{Sr}$. Numerous methods

have been used to estimate the local bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ range; from measurements of the geology (Budd et al., 2004; Copeland et al., 2010), measurements of plants, soils, and water sources (Bentley et al., 2004; Hodell et al., 2004; Evans et al., 2009; Evans et al., 2010) and the use of paired enamel and dentine samples (Montgomery et al., 2000; Taylor et al., 2013). Of these, no single technique has risen to dominance.

The results from the feeding experiment can be used to answer a more subtle question, namely what is the minimum variation in $^{87}\text{Sr}/^{86}\text{Sr}$ that can be expected from two individuals raised on identical diets in the same location? Given that the five feeds are isotopically identical and all feeding groups have the same water source, each of the 19 individuals can be said to have consumed a single identical diet. Across all feeding groups the G1 and G2 pig teeth have a 2 standard deviation variability of 75 ppm and thus a conservative estimate of 100 ppm for the minimum range could be made, and it could be argued that two individuals that differ by less than this cannot be distinguished on the basis of their $^{87}\text{Sr}/^{86}\text{Sr}$ alone.

However factors outside the controlled setting of this experiment may contribute additional variability to $^{87}\text{Sr}/^{86}\text{Sr}$ such as unknown metabolic differences between individuals, and different taxa or the differences in the quantity of food and water consumed, given that there may be isotopic difference between the two. However, this is likely also to be true in most archaeological contexts, especially where there may be multiple water sources (e.g. spring vs. rainwater). We therefore feel it is unlikely that for archaeological studies two locals will exhibit less variability than the c.100ppm we see in this study.

4.2. The potential for $\delta^{88}\text{Sr}$ in archaeology

The potential for more precisely tracking the dietary source of strontium using $\delta^{88}\text{Sr}$ and

determining the trophic level of marine and terrestrial resources has already been highlighted by Knudson et al. (2010). Here we expand on the range of possibilities for $\delta^{88}\text{Sr}$.

Strontium has no biological function in mammals and is only present as an impurity associated with Ca (Pors Nielsen, 2004). Therefore $\delta^{88}\text{Sr}$ may be robust against the effects of malnutrition and aridity, which have been shown to be problematic for $\delta^{15}\text{N}$, as N is biologically essential (Sealy et al., 1987; Fuller et al., 2005; Hedges and Reynard, 2007). The experimental results, shown in table 1, reveal no trend in magnitude of $\Delta^{88}\text{Sr}_{\text{diet-teeth}}$ with the [Sr] of the diet over the range of [Sr] in the experimental feeds. Thus, $\delta^{88}\text{Sr}$ may be less susceptible to changes in the dietary [Sr] than $\delta^{15}\text{N}$ is to changes in the protein content of the diet, especially in cases of malnutrition. Further, if $\delta^{88}\text{Sr}$ values in faeces can be shown to be similar to that of diet in other animals, especially ruminants, then $\delta^{88}\text{Sr}$ may be less sensitive to manuring which has been shown to increase the $\delta^{15}\text{N}$ baseline (Bogaard et al., 2007).

$\delta^{88}\text{Sr}$ determinations are made on the mineral fraction of tooth enamel and this may have the following advantages.

(i) Tooth enamel survival in the archaeological record is generally robust and therefore $\delta^{88}\text{Sr}$ may be useful when bone collagen survival is poor, such as in very wet or arid environments (Nielsen-Marsh et al., 2007; Smith et al., 2007).

(ii) $\delta^{88}\text{Sr}$ should be representative of the whole diet, similar to $\delta^{13}\text{C}_{\text{SC}}$, thus $\delta^{88}\text{Sr}$ should represent the trophic level of the whole diet (with individual components weighted by their Sr concentrations) and not just the protein portion. It is hoped that $\delta^{88}\text{Sr}$ may be used for a more complete assessment of dietary trophic level, but that $\delta^{88}\text{Sr}$ may prove to be robust against the problems caused by different digestive physiologies

in different animals which give rise to variable trophic level effects in between $\delta^{13}\text{C}_{\text{SC}}$ and $\delta^{13}\text{C}_{\text{diet}}$ (Lee-Thorpe 2008, Passey et al., 2005).

(iii) Measurements of $\delta^{88}\text{Sr}$ on tooth enamel will give information on the dietary trophic level at the time in life when the enamel was being mineralised as enamel does not remodel once mineralised (Budd et al., 2000). This means that $\delta^{88}\text{Sr}$ will give dietary information on early life and through adolescence. Such information may not necessarily be available to archaeologists using $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of bone collagen although recent work on dentine collagen may address this (Beaumont et al., 2013; Sandberg et al., 2014).

(iv) Sequential mineralisation of individual teeth in a dental arcade and the sequential mineralisation of an individual tooth from the tooth crown to root represent a time resolved signal. Inter-tooth measurements of $\delta^{88}\text{Sr}$ in a dental eruption sequence or intra-tooth measurements of $\delta^{88}\text{Sr}$ along the growth axis of an individual tooth, where the crown height permits, could be used to address changes in trophic level within an individual's lifetime. Specifically this might be used to investigate the effects of weaning.

(v) $\delta^{88}\text{Sr}$ could be of potential use is in the detection of diagenesis in bone and enamel. Bone and dentine have been shown to be susceptible to Sr diagenesis whilst enamel is more resistant (Budd et al., 2000; Trickett et al., 2003; Hoppe et al., 2003). One of the critical considerations in accurately detecting diagenesis has been identifying samples where accurate *a priori* assumptions about the biogenic and diagenetic $^{87}\text{Sr}/^{86}\text{Sr}$ ratio can be made. This has mainly limited studies of diagenetic uptake to marine mammals which have been buried in soils with an $^{87}\text{Sr}/^{86}\text{Sr}$ different from that of modern seawater (e.g. Nelson et al., 1986). However, if one considers that $\delta^{88}\text{Sr}$ fractionates with trophic level then there should be a large difference in $\delta^{88}\text{Sr}$ between

the buried bone/tooth and the burial soil, which should be at least two, possibly three, trophic levels. Furthermore, these differences will be systematic and should exist in bones and teeth even when the biogenic and the diagenetic $^{87}\text{Sr}/^{86}\text{Sr}$ are equivalent.

As a counterpoint to the potential benefits of $\delta^{88}\text{Sr}$ there may be some potential concerns for working with $\delta^{88}\text{Sr}$.

(i) The potential of $\delta^{88}\text{Sr}$ relies on understanding and characterising the variation in geological and bioavailable $\delta^{88}\text{Sr}$. To some extent variation in the geological $\delta^{88}\text{Sr}$ within a site catchment may be muted by the same processes of differential weathering of rocks and subsequent preferential leaching of Sr from soils to soil pore waters that reduces the variation in $^{87}\text{Sr}/^{86}\text{Sr}$ (Capo et al., 1998; Sillen et al., 1998; Price et al., 2002). If this is the case, then current methods for determining the local bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$, such as measuring local plant, soils, water and archaeological small fauna, should be sufficient to characterise the local $\delta^{88}\text{Sr}$. However this assumption needs to be tested and more detailed environmental and trophic level studies, particularly using natural experiments would be invaluable.

(ii) Wider scale variation in geological $\delta^{88}\text{Sr}$ may still be problematic when comparing between sites. The presently available studies on geological $\delta^{88}\text{Sr}$ have already demonstrated that $\delta^{88}\text{Sr}$ varies between rock types (Ohno et al., 2008; Charlier et al., 2012; Pearce et al., 2015) and that in rocks derived from oceanic carbonates (e.g. Limestone) variation will be systematic with the temperature of the water that the carbonate was precipitated from (Fietzke and Eisenhauer, 2006; Rüggeberg et al., 2008; Vollstaedt et al., 2014). It may be the case that $\delta^{88}\text{Sr}$ suffers from the same limitation as Sr/Ca and whilst $\delta^{88}\text{Sr}$ may be effective at a single site, comparisons between sites on different geologies may be problematic. Hopefully, careful characterisation of the bioavailable $\delta^{88}\text{Sr}$ can mitigate this.

(iii) Linked to the variation in geological/bioavailable $\delta^{88}\text{Sr}$ signal is the effect that non-local individuals might have on a population. Knudson et al. (2010) correctly point out that $\delta^{88}\text{Sr}$ can be of value in constraining the dietary source of Sr for $^{87}\text{Sr}/^{86}\text{Sr}$ studies. We would add to this that $^{87}\text{Sr}/^{86}\text{Sr}$ is also important in $\delta^{88}\text{Sr}$ studies. Consider, for example, an allochthonous individual buried at an archaeological site. If this individual has been consuming Sr from an area with a different bioavailable $\delta^{88}\text{Sr}$ then this may lead to an incorrect dietary interpretation in the context of the local bioavailable $\delta^{88}\text{Sr}$. Therefore, having an understanding of the provenance of individuals is important where palaeodietary reconstruction is being assessed and where $\delta^{88}\text{Sr}$ is used for this $^{87}\text{Sr}/^{86}\text{Sr}$ and possibly $\delta^{18}\text{O}$ should also be determined. We note that, whilst the use of the double-spike technique to determine $\delta^{88}\text{Sr}$ necessitates the measurement of $^{87}\text{Sr}/^{86}\text{Sr}$, other techniques such as sample standard bracketing, do not necessarily require the measurement of the ^{87}Sr isotope.

5. Conclusion

In this study we have investigated the [Sr], Sr/Ca, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}\text{Sr}$ systematics in a controlled feeding experiment, which was designed to test the effect of marine resource consumption on dietary isotopes.

The results of [Sr] analysis have detected a strong trend in both [Sr] and Sr/Ca with the marine protein content of the diet. However, rather than being a useful indicator of marine resource consumption, we believe that this is a fortuitous result of the feed preparations and is not significant outside of this specific study.

The results of the radiogenic Sr isotope analysis confirm under controlled conditions that the isotopic composition of dental tissues, modified marginally by drinking water, is equivalent to that of dietary strontium. Data from the feeding experiment also constrain the minimum variation between two individual pigs raised on

identical diets. We conclude that in archaeological contexts for two individuals who differ by less than 100 ppm in $^{87}\text{Sr}/^{86}\text{Sr}$ the most parsimonious explanation is that they are from the same geographical provenance and that they cannot be differentiated by this technique alone.

Results from stable strontium isotopes determined by the double-spike isotopic tracer method have identified a 0.322 ± 0.060 ‰ shift in lighter values of $\delta^{88}\text{Sr}$ from strontium in diet to strontium in dental tissues, across all the control diets, as a result of trophic level fractionation. The magnitude of this fractionation is consistent with that which would be predicted from the analogous shift observed elsewhere in calcium isotopes. This is the first time this has been determined in a controlled setting.

Although still in its infancy $\delta^{88}\text{Sr}$ has great potential for use in archaeological isotope investigations. $\delta^{88}\text{Sr}$ may be useful for examining trends in palaeodiet in concert with provenance studies using $^{87}\text{Sr}/^{86}\text{Sr}$ or where collagen preservation is poor and for investigating dietary trends in early life. Furthermore with the demonstration of $\delta^{88}\text{Sr}$ fractionation with trophic level $\delta^{88}\text{Sr}$ may be useful distinguishing biogenic and diagenetic strontium. $\delta^{88}\text{Sr}$ may be key in accurately identifying Sr diagenesis in archaeological enamel.

Acknowledgements

This work was supported by a University of Bristol Tratman Scholarship to Jamie Lewis and the initial feeding experiment carried out under NERC grant [NE/D004535/1] awarded to Richard Evershed. The authors would like to thank Dr Alan Stewart (Harper Adams University) for collecting the drinking water sample. Dr Emily Webb (Bristol) is also thanked for thoughtful discussions on the experimental results.

Figure and table captions

Figure 1. Schematic of the controlled feeding experiment after Webb et al., 2016.

Figure 2. Strontium concentrations for feed, teeth, water and faeces from the five feeding groups in the feeding experiment.

Figure 3. $^{87}\text{Sr}/^{86}\text{Sr}$ in feed, G1 teeth, G2 teeth and faeces from the pig feeding experiment.

Figure 4. $\delta^{88}\text{Sr}$ in feed, water, G1 adult teeth, G2 adolescent teeth and faeces from the pigs in the feeding experiment.

Table 1. Tabulated strontium concentration, $\text{Sr}/\text{Ca} \times 1000$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}\text{Sr}$ results from the feeding experiment.

References

- Avanzinelli R., Boari E., Conticelli S., Francalanci L., Guarnieri L., Perini G., Petrone, Chiara M., Tommasini S. and Ulivi M. 2005. "High Precision Sr, Nd and Pb Isotopic Analyses Using the New Generation Thermal Ionisation Mass Spectrometer ThermoFinnigan Triton-Ti." *Periodico Di Mineralogia* 74: 147–66.
- Balter V., Braga J., Télouk P. and Thackeray J. F. 2012. "Evidence for Dietary Change but Not Landscape Use in South African Early Hominins." *Nature* 489 (7417): 558–60.
- Beaumont J., Gledhill A., Lee-Thorp J. and Montgomery J. 2013. "Childhood Diet: A Closer Examination of the Evidence from Dental Tissues Using Stable Isotope Analysis." *Archaeometry* 55 (2): 277–95.
- Bentley, R. A. 2006. "Strontium Isotopes from the Earth to the Archaeological Skeleton: A Review." *Journal of Archaeological Method and Theory* 13 (3): 135–87.
- Bentley R. A., Bickle P., Fibiger L., Nowell G. M., Dale C. W., Hedges R. E. M., Hamilton J., Wahl J., Francken M., Grupe G., Lenneis E., Teschler-Nicola M., Arbogast R.-M., Hofmann D. and Whittle A. 2012. "Community Differentiation and Kinship among Europe's First Farmers." *Proceedings of the National Academy of Sciences of the United States of America* 109 (24): 9326–30.
- Bentley R. A., Price T. D. and Stephan E. 2004. "Determining the 'local' $^{87}\text{Sr}/^{86}\text{Sr}$ Range for Archaeological Skeletons: A Case Study from Neolithic Europe." *Journal of Archaeological Science* 31 (4): 365–75.
- Birck J. L. 1986. "Precision K-Rb-Sr Isotopic Analysis: Application to Rb-Sr Chronology." *Chemical Geology* 56 (1–2): 73–83.
- Bogaard A., Heaton T. H. E., Poulton P. and Merbach I. 2007. "The Impact of Manuring on Nitrogen Isotope Ratios in Cereals: Archaeological Implications for Reconstruction of

- Diet and Crop Management Practices.” *Journal of Archaeological Science* 34 (3): 335–43.
- Britton K., Grimes V., Dau J. and Richards M. P. 2009. “Reconstructing Faunal Migrations Using Intra-Tooth Sampling and Strontium and Oxygen Isotope Analyses: A Case Study of Modern Caribou (*R. Tarandus Granti*).” *Journal of Archaeological Science* 36 (5): 1163–72.
- Budd P., Millard A. R., Chenery C. A., Lucy S. and Roberts C. 2004. “Investigating Population Movement by Stable Isotope Analysis: A Report from Britain.” *Antiquity*, no. May. Antiquity Publications: 127–141.
- Budd P., Montgomery J. L., Barreiro B. and Thomas R. G. 2000. “Differential Diagenesis of Strontium in Archaeological Human Dental Tissues.” *Applied Geochemistry* 15 (5): 687–94.
- Burton J. H., Price T. D., Cahue L. and Wright L. E. 2003. “The Use of Barium and Strontium Abundances in Human Skeletal Tissues to Determine Their Geographic Origins.” *International Journal of Osteoarchaeology* 13 (1–2): 88–95.
- Burton J. H., Price T. D. and Middleton W. 1999. “Correlation of Bone Ba/Ca and Sr/Ca due to Biological Purification of Calcium.” *Journal of Archaeological Science* 26: 609–16.
- Burton J. H. and Wright L. E. 1995. “Nonlinearity in the Relationship between Bone Sr/Ca and Diet: Paleodietary Implications.” *American Journal of Physical Anthropology* 96 (3): 273–82.
- Burton J. H. and Price T. D. 1999. “Evaluation of Bone Strontium as a Measure of Seafood Consumption.” *International Journal of Osteoarchaeology* 9 (September 1997): 233–36.
- Capo R. C., Stewart B. W. and Chadwick O. A. 1998. “Strontium Isotopes as Tracers of Ecosystem Processes: Theory and Methods.” *Geoderma* 82 (1–3): 197–225.
- Charlier B. L. A., Nowell G. M., Parkinson I. J., Kelley S. P., Pearson D. G. and Burton K. W. 2012. “High Temperature Strontium Stable Isotope Behaviour in the Early Solar System and Planetary Bodies.” *Earth and Planetary Science Letters* 329–330 (May): 31–40.
- Chisholm B. S., Nelson D. E. and Schwarcz H. P. 1982. “Stable-Carbon Isotope Ratios as a Measure of Marine versus Terrestrial Protein in Ancient Diets.” *Science (New York, N.Y.)* 216 (4550): 1131–32.
- Coath, C. D., Elliott, T. and Hin, R. 2017. “Double Spike Inversions for Three Isotope Systems.” *Chemical Geology* 451: 78-89
- Comar C., Russell R. and Wasserman R. 1957. “Strontium-Calcium Movement from Soil to Man.” *Science* 126 (3272): 485–92.
- Copeland S. R., Sponheimer M., Lee-thorp J. A., le Roux P. J., de Ruiter D. J. and Richards M. P. 2010. “Strontium Isotope Ratios in Fossil Teeth from South Africa: Assessing Laser Ablation MC-ICP-MS Analysis and the Extent of Diagenesis.” *Journal of Archaeological Science* 37 (7): 1437–46.
- Corr L. T., Sealy J. C., Horton M. C. and Evershed R. P. 2005. “A Novel Marine Dietary Indicator Utilising Compound-Specific Bone Collagen Amino Acid $\delta^{13}\text{C}$ Values of Ancient Humans.” *Journal of Archaeological Science* 32 (3): 321–30.

- de Souza G. F., Reynolds B. C., Kiczka M. and Bourdon B. 2010. "Evidence for Mass-Dependent Isotopic Fractionation of Strontium in a Glaciated Granitic Watershed." *Geochimica et Cosmochimica Acta* 74 (9): 2596–2614.
- DeNiro M. J. and Epstein S. 1978. "Influence of Diet on the Distribution of Carbon Isotopes in Animals." *Geochimica et Cosmochimica Acta* 42: 495–506.
- Dodson, M.H. 1970. "Simplified Equations for Double-Spiked Isotopic Analyses." *Geochimica et Cosmochimica Acta* 34: 1241–44.
- Dodson, M.H. 1963. "A Theoretical Study of the Use of Internal Standards for Precise Isotopic Analysis by the Surface Ionization Technique: Part I - General First-Order Algebraic Solutions." *Journal of Scientific Instruments* 40: 289–95.
- Elias R., Hirao Y. and Patterson C. 1982. "The Circumvention of the Natural Biopurification of Calcium along Nutrient Pathways by Atmospheric Inputs of Industrial Lead." *Geochimica et Cosmochimica Acta*, 2561–80.
- Ericson J. 1985. "Strontium Isotope Characterization in the Study of Prehistoric Human Ecology." *Journal of Human Evolution* 14 (5): 503–14.
- Evans J. A., Montgomery J. and Wildman G. 2009. "Isotope Domain Mapping of $^{87}\text{Sr}/^{86}\text{Sr}$ Biosphere Variation on the Isle of Skye, Scotland." *Journal of the Geological Society* 166 (4): 617–31.
- Evans J. A., Montgomery J., Wildman G. and Boulton N. 2010. "Spatial Variations in Biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ in Britain." *Journal of the Geological Society* 167 (1): 1–4.
- Fietzke J. and Eisenhauer A. 2006. "Determination of Temperature-Dependent Stable Strontium Isotope ($^{88}\text{Sr}/^{86}\text{Sr}$) Fractionation via Bracketing Standard MC-ICP-MS." *Geochemistry, Geophysics, Geosystems* 7 (8): 1–6.
- Fuller B. T., Fuller J. L., Sage N. E., Harris D. A, O'Connell T. C. and Hedges R. E. M. 2005. "Nitrogen Balance and $\delta^{15}\text{N}$: Why You're Not What You Eat during Nutritional Stress." *Rapid Communications in Mass Spectrometry: RCM* 19: 2497–2506.
- Haverkort C. M., Weber A., Katzenberg M. A., Goriunova O. I., Simonetti A. and Creaser R. A. 2008. "Hunter-Gatherer Mobility Strategies and Resource Use Based on Strontium Isotope $^{87}\text{Sr}/^{86}\text{Sr}$ Analysis: A Case Study from Middle Holocene Lake Baikal, Siberia." *Journal of Archaeological Science* 35 (5): 1265–80.
- Hedges R. E. M. and Reynard L. M. 2007. "Nitrogen Isotopes and the Trophic Level of Humans in Archaeology." *Journal of Archaeological Science* 34 (8): 1240–51.
- Hodell D. A., Quinn R. L., Brenner M. and Kamenov G. 2004. "Spatial Variation of Strontium Isotopes ($^{87}\text{Sr}/^{86}\text{Sr}$) in the Maya Region: A Tool for Tracking Ancient Human Migration." *Journal of Archaeological Science* 31 (5): 585–601.
- Hoppe K. A., Koch P. L. and Furutani T. T. 2003. "Assessing the Preservation of Biogenic Strontium in Fossil Bones and Tooth Enamel." *International Journal of Osteoarchaeology* 13 (1–2): 20–28.
- Horwitz E., Chiarizia R. and Dietz M. L. 1992. "A Novel Strontium-Selective Extraction Chromatographic Resin." *Solvent Extraction and Ion Exchange* 10 (2): 313–36.

- Howland M. R., Corr L. T., Young S. M. M., Jones V., Jim S., Van Der Merwe N. J., Mitchell A. D. and Evershed R. P. 2003. "Expression of the Dietary Isotope Signal in the Compound-Specific $\delta^{13}\text{C}$ Values of Pig Bone Lipids and Amino Acids." *International Journal of Osteoarchaeology* 13 (1–2): 54–65.
- Jim S., Jones V., Copley M. S., Ambrose S. H. and Evershed R. P. 2003. "Effects of Hydrolysis on the $\delta^{13}\text{C}$ Values of Individual Amino Acids Derived from Polypeptides and Proteins." *Rapid Communications in Mass Spectrometry : RCM* 17 (20): 2283–89.
- Knipper, C. 2011. "Die Räumliche Organisation Der Linearbandkeramischen Rinderhaltung : Naturwissenschaftliche Und Archäologische Untersuchungen." *British Archaeological Reports, International Series 2305* (Oxford 2011).
- Knudson K. J., Williams H. M., Buikstra J. E., Tomczak P. D., Gordon G. W. and Anbar A. D. 2010. "Introducing $^{88}\text{Sr}/^{86}\text{Sr}$ Analysis in Archaeology: A Demonstration of the Utility of Strontium Isotope Fractionation in Paleodietary Studies." *Journal of Archaeological Science* 37 (9). 2352–64.
- Krabbenhöft A., Eisenhauer A., Böhm F., Vollstaedt H., Fietzke J., Liebetrau V., Augustin N., Peucker-Ehrenbrink B., Müller M. N., Horn C., Hansen B. T., Nolte N. and Wallmann K. 2010. "Constraining the Marine Strontium Budget with Natural Strontium Isotope Fractionations ($^{87}\text{Sr}/^{86}\text{Sr}$ *, $\delta^{88/86}\text{Sr}$) of Carbonates, Hydrothermal Solutions and River Waters." *Geochimica et Cosmochimica Acta* 74 (14): 4097–4109.
- Lee-thorp J. A. 2008. "On Isotopes and Old Bones." *Archaeometry* 50 (6): 925–50.
- Lee-Thrope J. A. and van der Merwe N. J. 1991. "Aspects of Modern and Fossil Biological Apatites." *Journal of Archaeological Science* 18: 343–54.
- Lewis J., Coath C. D. and Pike A. W. G. 2014. "An Improved Protocol for $^{87}\text{Sr}/^{86}\text{Sr}$ by Laser Ablation Multi-Collector Inductively Coupled Plasma Mass Spectrometry Using Oxide Reduction and a Customised Plasma Interface." *Chemical Geology* 390 (December). 173–81.
- Lewis, J. 2015. "Lying through Your Teeth; Strontium Uptake in Archaeological Enamel." University of Bristol.
- McArthur J. M., Howarth R. J. and Bailey T. R. 2001. "Strontium Isotope Stratigraphy: LOWESS Version 3: Best Fit to the Marine Sr-Isotope Curve for 0–509 Ma and Accompanying Look-up Table for Deriving Numerical Age." *The Journal of Geology* 109 (2): 155–70.
- Montgomery J. L., Budd P. and Evans J. A. 2000. "Reconstructing the Lifetime Movements of Ancient People: A Neolithic Case Study from Southern England." *European Journal of Archaeology* 3 (3): 370–85.
- Moynier F., Agranier A., Hezel D. C. and Bouvier A. 2010. "Sr Stable Isotope Composition of Earth, the Moon, Mars, Vesta and Meteorites." *Earth and Planetary Science Letters* 300 (3–4). 359–66.
- Nehlich O., Fuller B. T., Jay M., Mora A., Nicholson R. A., Smith C. I. and Richards M. P. 2011. "Application of Sulphur Isotope Ratios to Examine Weaning Patterns and Freshwater Fish Consumption in Roman Oxfordshire, UK." *Geochimica et Cosmochimica Acta* 75 (17): 4963–77.

- Nelson B. K., DeNiro M. J., Schoeninger M. J. and De Paolo D. J. 1986. "Effects of Diagenesis on Strontium, Carbon, Nitrogen and Oxygen Concentration and Isotopic Composition of Bone." *Geochimica et Cosmochimica Acta* 50: 1941–49.
- Neymark L. A., Premo W. R., Mel'nikov N. N. and Emsbo P. 2014. "Precise Determination of $\delta^{88}\text{Sr}$ in Rocks, Minerals, and Waters by Double-Spike TIMS: A Powerful Tool in the Study of Geological, Hydrological and Biological Processes." *Journal of Analytical Atomic Spectrometry* 29 (1): 65.
- Nielsen-Marsh C. M., Smith C. I., Jans M. M. E., Nord A., Kars H. and Collins M. J. 2007. "Bone Diagenesis in the European Holocene II: Taphonomic and Environmental Considerations." *Journal of Archaeological Science* 34 (9): 1523–31.
- Nier, A.O. 1938. "The Isotopic Constitution of Strontium, Barium, Bismuth, Thallium and Mercury." *Physical Review* 54 (4): 275–78.
- O'Connell T. C., Kneale C. J., Tasevska N. and Kuhnle G. G. C. 2012. "The Diet-Body Offset in Human Nitrogen Isotopic Values: A Controlled Dietary Study." *American Journal of Physical Anthropology* 149 (3): 426–34.
- Ohno T., Komiya T., Ueno Y., Hirata T. and Maruyama S. 2008. "Determination of $^{87}\text{Sr}/^{86}\text{Sr}$ Mass-Dependent Isotopic Fractionation and Radiogenic Isotope Variation of $^{87}\text{Sr}/^{86}\text{Sr}$ in the Neoproterozoic Doushantuo Formation." *Gondwana Research* 14 (1–2): 126–33.
- Passey B. H., Robinson T. F., Ayliffe L. K., Cerling T. E., Sponheimer M., Dearing M. D., Roeder B. L. and Ehleringer J. R. 2005. "Carbon Isotope Fractionation between Diet, Breath CO₂, and Bioapatite in Different Mammals." *Journal of Archaeological Science* 32 (10): 1459–70.
- Patchett, P. J. 1980a. "Sr Isotopic Fractionation in Allende Chondrules: A Reflection of Solar Nebular Processes." *Earth and Planetary Science Letters* 50: 181–88.
- . 1980b. "Sr Isotopic Fractionation in Ca–Al Inclusions from the Allende Meteorite." *Nature* 283 (5746): 438–41.
- Pearce C. R., Parkinson I. J., Gaillardet J., Charlier B. L. A., Mokadem F. and Burton K. W. 2015. "Reassessing the Stable ($\delta^{88}/^{86}\text{Sr}$) and Radiogenic ($^{87}\text{Sr}/^{86}\text{Sr}$) Strontium Isotopic Composition of Marine Inputs." *Geochimica et Cosmochimica Acta* 157: 125–46.
- Pearce C. R., Parkinson I. J., Gaillardet J., Chetelat B. and Burton K. W. 2015. "Characterising the Stable ($\delta^{88}/^{86}\text{Sr}$) and Radiogenic ($^{87}\text{Sr}/^{86}\text{Sr}$) Isotopic Composition of Strontium in Rainwater." *Chemical Geology* 409: 54–60.
- Pors Nielsen, S. 2004. "The Biological Role of Strontium." *Bone* 35 (3): 583–88.
doi:10.1016/j.bone.2004.04.026.
- Price T. D., Burton J. H. and Bentley R. A. 2002. "The Characterization of Biologically Available Strontium Isotope Ratios for the Study of Prehistoric Migration." *Archaeometry* 44: 117–35.
- Price T. D., Connor M. and Parsen J. 1985. "Bone Chemistry and the Reconstruction of Diet: Strontium Discrimination in White-Tailed Deer." *Journal of Archaeological Science* 12: 419–42.
- Price T. D., Swick R. W. and Chase E. P. 1986. "Bone Chemistry and Prehistoric Diet:

- Strontium Studies of Laboratory Rats.” *American Journal of Physical Anthropology* 70 (3): 365–75.
- Reynard L. M., Henderson G. M. and Hedges R. E. M. 2010. “Calcium Isotope Ratios in Animal and Human Bone.” *Geochimica et Cosmochimica Acta* 74 (13): 3735–50.
- Rudge J. F., Reynolds B. C. and Bourdon B. 2009. “The Double Spike Toolbox.” *Chemical Geology* 265 (3–4): 420–31.
- Rüggeberg A., Fietzke J., Liebetrau V., Eisenhauer A., Dullo W. C. and Freiwald A. 2008. “Stable Strontium Isotopes ($\delta^{88/86}\text{Sr}$) in Cold-Water Corals - A New Proxy for Reconstruction of Intermediate Ocean Water Temperatures.” *Earth and Planetary Science Letters* 269 (3–4): 569–74.
- Russell W., Papanastassiou D. and Tombrello T. 1978. “Ca Isotope Fractionation on the Earth and Other Solar System Materials.” *Geochimica et Cosmochimica Acta* 42: 1075–90.
- Sandberg P. A., Sponheimer M., Lee-Thorp J. and Van Gerven D. 2014. “Intra-Tooth Stable Isotope Analysis of Dentine: A Step toward Addressing Selective Mortality in the Reconstruction of Life History in the Archaeological Record.” *American Journal of Physical Anthropology* 155 (2): 281–93.
- Schoeninger M. J. and DeNiro M. J. 1984. “Nitrogen and Carbon Isotopic Composition of Bone Collagen from Marine and Terrestrial Animals.” *Geochimica et Cosmochimica Acta* 48: 625–39.
- Schoeninger M. J., DeNiro M. and Tauber H. 1983. “Stable Nitrogen Isotope Ratios of Bone Collagen Reflect Marine and Terrestrial Components of Prehistoric Human Diet.” *Science* 220 (4604): 1381–83.
- Schoeninger M. J. and Peebles C. S. 1981. “Effect of Mollusc Eating on Human Bone Strontium Levels.” *Journal of Archaeological Science* 8: 391–97.
- Sealy J. C. 2001. “Body Tissue Chemistry and Palaeodiet.” In *Handbook of Archaeological Sciences*, edited by A.M. Brothwell, D.R. and Pollard, 269–79. Chichester: John Wiley and Sons.
- Sealy J. C., van der Merwe N. J., Sillen A., Kruger F. J. and Krueger H. W. 1991. “ $^{87}\text{Sr}/^{86}\text{Sr}$ as a Dietary Indicator in Modern and Archaeological Bone.” *Journal of Archaeological Science* 18: 399–416.
- Sealy J. C., van der Merwe N. J., Thorp J. A. L. and Lanham J. L. 1987. “Nitrogen Isotopic Ecology in Southern Africa: Implications for Environmental and Dietary Tracing.” *Geochimica et Cosmochimica Acta* 51: 2707–17.
- Shand P., Edmunds W. M., Lawrence A. R., Smedley P. L. and Burke S. 2007. “The Natural (Baseline) Quality of Groundwater in England and Wales.” *BGS Research Report RR/07/06*.
- Sillen, A. 1981. “Strontium and Diet at Hayonim Cave.” *American Journal of Physical Anthropology* 56 (2): 131–37.
- Sillen A., Hall G., Richardson S. and Armstrong R. 1998. “ $^{87}\text{Sr}/^{86}\text{Sr}$ Ratios in Modern and Fossil Food-Webs of the Sterkfontein Valley: Implications for Early Hominid Habitat Preference.” *Geochimica et Cosmochimica Acta* 62 (14): 2463–73.

- Sillen A. and Kavanagh M. 1982. "Strontium and Paleodietary Research: A Review." *American Journal of Physical Anthropology* 25 (S3): 67–90.
- Skulan J. and DePaolo D. J. 1999. "Calcium Isotope Fractionation between Soft and Mineralized Tissues as a Monitor of Calcium Use in Vertebrates." *Proceedings of the National Academy of Sciences of the United States of America* 96 (24): 13709–13.
- Smith C. I. C., Nielsen-Marsh C. M., Jans M. M. E. and Collins M. J. 2007. "Bone Diagenesis in the European Holocene I: Patterns and Mechanisms." *Journal of Archaeological Science* 34 (9): 1485–93.
- Steiger R. and Jäger E. 1977. "Subcommission on Geochronology: Convention on the Use of Decay Constants in Geo- and Cosmochronology." *Earth and Planetary Science Letters* 36: 359–62.
- Stephan E., Knipper C., Schatz K., Price T. D. and Hegner E. 2012. "Strontium Isotopes in Faunal Remains: Evidence of the Strategies for Land Use at the Iron Age Site Eberdingen-Hochdorf (Baden-Württemberg, Germany)." In *Population Dynamics in Prehistory and Early History*, edited by Elke Kaiser, Joachim Burger, and Schier Wolfram, 265–86. Berlin, Boston: De Gruyter.
- Stevenson E. I., Hermoso M., Rickaby R. E. M., Tyler J. J., Minoletti F., Parkinson I. J., Mokadem F. and Burton K. W. 2014. "Controls on Stable Strontium Isotope Fractionation in Coccolithophores with Implications for the Marine Sr Cycle." *Geochimica et Cosmochimica Acta* 128 (December). 225–35.
- Styring A. K., Fraser R. A., Arbogast R.-M., Halstead P., Isaakidou V., Pearson J. A., Schäfer M., Triantaphyllou S., Valamoti S. M., Wallace M., Bogaard A. and Evershed R. P. 2015. "Refining Human Palaeodietary Reconstruction Using Amino Acid $\delta^{15}\text{N}$ Values of Plants, Animals and Humans." *Journal of Archaeological Science* 53: 504–15.
- Styring A. K., Sealy J. C. and Evershed R. P. 2010. "Resolving the Bulk $\delta^{15}\text{N}$ Values of Ancient Human and Animal Bone Collagen via Compound-Specific Nitrogen Isotope Analysis of Constituent Amino Acids." *Geochimica et Cosmochimica Acta* 74 (1). 241–51.
- Taylor G., Tucker K., Butler R., Pike A. W. G., Lewis J., Roffey S., Marter P., Lee O., Wu H. H. T., Minnikin D. E., Besra G. S., Sing P., Cole S. T. and Stewart G. R. 2013. "Detection and Strain Typing of Ancient Mycobacterium Leprae from a Medieval Leprosy Hospital." *PLOS ONE* 8 (4): e62406.
- Toots H. and Voorhies M. R. 1965. "Strontium in Fossil Bones and the Reconstruction of Food Chains." *Science* 149 (3686): 854–55.
- Trickett M., Budd P., Montgomery J. L. and Evans J. A. 2003. "An Assessment of Solubility Profiling as a Decontamination Procedure for the $^{87}\text{Sr}/^{86}\text{Sr}$ Analysis of Archaeological Human Skeletal Tissue." *Applied Geochemistry* 18: 653–58.
- Tütken T., Vennemann T. W. and Pfretzschner H.-U. 2011. "Nd and Sr Isotope Compositions in Modern and Fossil Bones – Proxies for Vertebrate Provenance and Taphonomy." *Geochimica et Cosmochimica Acta* 75 (20): 5951–70.
- Urey, HC. 1947. "The Thermodynamic Properties of Isotopic Substances." *Journal of the Chemical Society*, no. 561–581.

- Van der Merwe N. J. and Vogel J. C. 1978. " $\delta^{13}\text{C}$ Content of Human Collagen as a Measure of Prehistoric Diet in Woodland North America." *Nature* 276 (5690): 815–16.
- Viner S., Evans J. A., Albarella U. and Pearson M. P. 2010. "Cattle Mobility in Prehistoric Britain: Strontium Isotope Analysis of Cattle Teeth from Durrington Walls (Wiltshire, Britain)." *Journal of Archaeological Science* 37 (11). 2812–20.
- Vollstaedt H., Eisenhauer A., Wallmann K., Böhm F., Fietzke J., Liebetrau V., Krabbenhöft A., Farkaš J., Tomašových A., Raddatz J. and Veizer J. 2014. "The Phanerozoic $\delta^{88}\text{Sr}/^{86}\text{Sr}$ Record of Seawater: New Constraints on Past Changes in Oceanic Carbonate Fluxes." *Geochimica et Cosmochimica Acta* 128 (October). 249–65.
- Webb E. C., Stewart A., Miller B., Tarlton J. and Evershed R. P. 2016. "Age Effects and the Influence of Varying Proportions of Terrestrial and Marine Dietary Protein on the Stable Nitrogen-Isotope Compositions of Pig Bone Collagen and Soft Tissues from a Controlled Feeding Experiment." *STAR: Science & Technology of Archaeological Research* 2 (1): 54–66.
- Webb E., Lewis, J., Shain A., Kastrisianaki-Guyton E., Honch N., Stewart A., Miller B., Tarlton J. and Evershed R. P. n.d. "The Influence of Varying Proportions of Terrestrial and Marine Dietary Protein on the Stable Carbon Isotope Compositions of Pig Tissues from a Controlled Feeding Experiment."




Figure 1. Schematic of the controlled feeding experiment after Webb et al., 2016.




Figure 2. Strontium concentrations for feed, teeth, water and faeces from the five feeding groups in the feeding experiment.




Figure 3. $^{87}\text{Sr}/^{86}\text{Sr}$ in feed, G1 teeth, G2 teeth and faeces from the pig feeding experiment.




Figure 4. $\delta^{88}\text{Sr}$ in feed, water, G1 adult teeth, G2 adolescent teeth and faeces from the pigs in the feeding experiment.

a. Strontium concentration and 2 σ uncertainty (ppm)												
Marine Protein %												
0	12.5		25		50		100		Mean	2 SD		
<i>Feed</i>												
12.96	3.55	15.96	3.32	19.37	3.69	26.01	5.72	36.31	3.34			
<i>G2 teeth</i>												
75.95	8.40	104.03	20.79	117.36	19.90	149.01	26.69	299.73	49.18			
<i>Faeces</i>												
76.57	1.53	129.53	2.59	127.58	2.55	204.74	4.09	342.90	6.86			
b. Sr/Ca \times 1000 and 2σ uncertainty												
Marine Protein %												
0	12.5		25		50		100		Mean	2 SD		
<i>Feed</i>												
0.948	0.112	1.352	0.136	1.629	0.155	2.230	0.231	3.233	0.205			
<i>G2 teeth</i>												
0.251	0.064	0.356	0.061	0.384	0.068	0.473	0.060	0.755	0.041			
<i>Sr/Ca_{teeth/diet}</i>												
0.265	0.263		0.236		0.212		0.234		0.242	0.044		
c. $^{87}\text{Sr}/^{86}\text{Sr}$ and 2σ uncertainty in ppm												
Marine Protein %												
0	12.5		25		50		100		Mean			
<i>Feed</i>												
0.709092	12.2	0.709063	10.9	0.709052	19.1	0.709061	13.8	0.709011	19.6	0.709056	58.4	
<i>G1 teeth</i>												
0.709124	12.4	0.709097	9.2	0.709074	6.2	0.709121	7.9	0.709058	16.4			
<i>G2 teeth</i>												

0.709098	72.9	0.709048	17.4	0.709061	12.3	0.709078	5.4	0.709076	59.0	0.709089	75.3
0.709144	8.5	0.709108	13.4	0.709041	6.6	0.709143	6.2	0.709072	5.7		
0.709178	12.4	0.709061	6.5	0.709064	5.7	0.709139	5.6	0.709047	6.3		
		0.709106	10.7	0.709032	5.8			0.709075	4.9		
				0.709094	5.6						
<i>Faeces</i>											
0.709077	6.1	0.709103	9.9	0.709074	17.9	0.709010	12.1	0.709013	8.1	0.709055	83.3
<i>Water</i>											
				0.711572	6.0						

d. $\delta^{88}\text{Sr}$ and 2σ uncertainty in ‰

Marine Protein %

0	12.5	25	50	100	Mean	2 SD					
<i>Feed</i>											
0.115	0.010	0.088	0.026	0.095	0.024	0.132	0.014	0.061	0.009	0.098	0.054
<i>G1 teeth</i>											
-0.255	0.007	-0.225	0.006	-0.254	0.007	-0.232	0.009	-0.190	0.008		
<i>G2 teeth</i>											
-0.228	0.019	-0.211	0.019	-0.256	0.008	-0.227	0.011	-0.260	0.005	-0.224	0.044
-0.207	0.003	-0.206	0.004	-0.231	0.010	-0.200	0.018	-0.231	0.002		
-0.201	0.022	-0.184	0.006	-0.212	0.003	-0.246	0.005	-0.229	0.003		
		-0.199	0.003	-0.241	0.009						
				-0.218	0.004						
$\Delta^{88}\text{Sr}_{\text{diet-teeth}}$											
0.337		0.293		0.331		0.359		0.289		0.322	0.060
<i>Faeces</i>											
0.130	0.015	0.150	0.007	0.112	0.021	0.113	0.005	0.131	0.009	0.127	0.031

<i>Water</i>						
		0.274	0.005			

Table 1. Tabulated strontium concentration, $\text{Sr}/\text{Ca} \times 1000$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}\text{Sr}$ results from the feeding experiment.