

RECEIVED: July 14, 2017

REVISED: November 6, 2017

ACCEPTED: November 9, 2017

PUBLISHED: November 22, 2017

Spontaneous CP violation in multi-Higgs potentials with triplets of $\Delta(3n^2)$ and $\Delta(6n^2)$

Ivo de Medeiros Varzielas,^{a,b} Stephen F. King,^a Christoph Luhn^c and Thomas Neder^{a,d}^a*School of Physics and Astronomy, University of Southampton,
SO17 1BJ Southampton, United Kingdom*^b*CFTP, Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa,
Avenida Rovisco Pais 1, 1049 Lisboa, Portugal*^c*Theoretische Physik 1, Naturwissenschaftlich-Technische Fakultät, Universität Siegen,
Walter-Flex-Straße 3, 57068 Siegen, Germany*^d*AHEP Group, Institut de Física Corpuscular — C.S.I.C./Universitat de València,
Parc Científic de Paterna, C/ Catedrático José Beltrán, 2 E-46980 Paterna (Valencia), Spain
E-mail: ivo.de@udo.edu, king@soton.ac.uk,
christoph.luhn@uni-siegen.de, neder@ific.uv.es*

ABSTRACT: Motivated by discrete flavour symmetry models, we analyse Spontaneous CP Violation (SCPV) for potentials involving three or six Higgs fields (both electroweak doublets and singlets) which fall into irreducible triplet representations of discrete symmetries belonging to the $\Delta(3n^2)$ and $\Delta(6n^2)$ series, including A_4 , S_4 , $\Delta(27)$ and $\Delta(54)$. For each case, we give the potential and find various global minima for different regions of the parameter space of the potential. Using CP-odd basis Invariants that indicate the presence of Spontaneous CP Violation we separate the VEVs into those that do or do not violate CP. In cases where CP is preserved we reveal a CP symmetry of the potential that is preserved by those VEVs, otherwise we display a non-zero CP-odd Invariant. Finally we identify interesting cases where there is Spontaneous Geometrical CP Violation in which the VEVs have geometrical phases.

KEYWORDS: CP violation, Discrete Symmetries, Higgs Physics, Beyond Standard Model

ARXIV EPRINT: [1706.07606](https://arxiv.org/abs/1706.07606)

Contents

1	Introduction	1
2	CP violation	4
2.1	Generalities	4
2.2	Spontaneous CP Violation	7
2.3	Spontaneous CP invariants	9
3	Potentials and Vacuum Expectation Values	12
3.1	Potentials of one triplet	15
3.1.1	$\Delta(3n^2)$ and $\Delta(6n^2)$, with $n > 3$	15
3.1.2	S_4	17
3.1.3	A_4	18
3.1.4	$\Delta(27)$ and $\Delta(54)$	21
3.2	VEVs of potentials of two triplets	24
3.2.1	$\Delta(6n^2)$ with $n > 3$	26
3.2.2	$\Delta(3n^2)$ with $n > 3$	27
3.2.3	S_4	30
3.2.4	A_4	31
3.2.5	$\Delta(54)$	33
3.2.6	$\Delta(27)$	35
4	Spontaneous CP-odd invariants applied	36
4.1	Potentials of one triplet	36
4.1.1	$\Delta(3n^2)$ and $\Delta(6n^2)$ with $n > 3$	36
4.1.2	A_4 and S_4	36
4.1.3	$\Delta(27)$ and $\Delta(54)$	36
4.2	Potentials of two triplets	37
4.2.1	$\Delta(3n^2)$ and $\Delta(6n^2)$ with $n > 3$	37
4.2.2	$\Delta(3n^2)$, $n > 3$ with CP_0	38
4.2.3	$\Delta(3n^2)$, $n > 3$ with CP_{23}	39
4.2.4	$\Delta(3n^2)$, $n > 3$ another CP	40
4.2.5	A_4 and S_4	40
4.2.6	A_4 with CP_0	41
4.2.7	A_4 with CP_{23}	42
4.2.8	$\Delta(27)$ and $\Delta(54)$	42
4.2.9	$\Delta(54)$ with CP_0	43
4.3	Spontaneous Geometrical CP Violation	43
5	Summary of results	46
6	Conclusions	48
A	Technical details from section 2	49

1 Introduction

The discovery of the Higgs boson by the Large Hadron Collider (LHC) [1, 2], indicates that at least one Higgs doublet must be responsible for electroweak symmetry breaking (EWSB). However, there is no special reason why there should be only one Higgs doublet in Nature, and it is entirely possible that there could be additional Higgs doublets, accompanied by further Higgs bosons which could be discovered in the next run of the LHC.

The simplest example is two-Higgs-doublet models (2HDMs) [3–6]. However, 2HDMs generally face severe phenomenological problems with flavour changing neutral currents (FCNCs) and possible charge breaking vacua, and it is common to consider restricted classes of models controlled by various symmetries. In 2HDMs, the full list of possible symmetries of the potential is now known [7–10]. In 2HDMs these symmetries can be conserved or spontaneously violated after the EWSB, depending on the coefficients of the potential. Generalising these results to NHDMs is technically difficult, although there has been some recent progress in this direction [11–14].

The case of three-Higgs-doublet models (3HDMs) is particularly promising for several reasons. To begin with, it is the next simplest example beyond 2HDMs which has been exhaustively studied in the literature. Furthermore, 3HDMs are more tractable than NHDMs, and all possible finite symmetries (but not all continuous ones) have been identified [15]. Finally, and perhaps most intriguingly, 3HDMs may shed light on the flavour problem, namely the problem of the origin and nature of the three families of quarks and leptons, including neutrinos, and their pattern of masses, mixings and CP violation. Typical examples of such models that use discrete symmetries to constrain the structure of mass matrices need several multiplets of scalar fields that also transform under the same symmetry (for reviews, cf. [16–21]). Such models provide a motivation to study multiple SM Higgs singlets (sometimes called “flavons” in this context) as well as electroweak doublets. In the context of flavour models it is natural to consider Higgs doublets or singlets which play the role of “flavons” and form irreducible triplets under some spontaneously broken discrete family symmetry. Motivated by the above considerations, we shall study CP violating potentials with both three and six Higgs doublets and singlets.

CP symmetry, which for a single field is just the combination of particle-antiparticle exchange and space inversion, is presently known to be violated only by the weak interactions involving quarks in the Standard Model (SM) [22]. The origin of the observed SM quark CP violation (CPV) is a natural consequence of three generations of quarks whose mixing is described by a complex CKM matrix. Although the CKM matrix can be parameterised in different ways, it was realised that the amount of CPV in physical processes always depends on a particular weak basis invariant which can be expressed in terms of the quark mass matrices [23]. Although CP is automatically conserved by the Higgs potential of the SM, with more than one Higgs doublet it is possible that the Higgs potential violates CP, providing a new source of CPV [24]. We shall be interested in cases of three and six Higgs doublets and singlets, whose potentials are controlled by various non-Abelian discrete symmetries which admit irreducible triplet representations. In particular we are interested in the cases of such potentials which conserve CP explicitly, but where the vacua

of such potentials may spontaneously break CP. We shall analyse this problem using basis independent CP-odd invariants for the following reason.

As already mentioned in the context of the CKM matrix, the study of CP is a subtle topic because of the basis dependent nature of the phases which control CP violation. Similar considerations also apply to the phases which appear in the parameters of the potentials of multiple scalars. An important tool to assist in determining whether CP is violated or not are basis independent CP-odd invariants (CPIs), whose usefulness has been shown in the SM in addressing CP violation arising from the CKM matrix, sourced from the Yukawa couplings. The first example of the use of such invariants was the Jarlskog invariant [23], which was reformulated in [25] in a form which is generally valid for an arbitrary number of generations. Generalising the invariant approach [25] and applying it to fermion sectors of theories with Majorana neutrinos [26] or with discrete symmetries [27, 28] leads to other relevant CPIs. In extensions of the Higgs sector of the SM, the CP violation arising from the parameters of the scalar potential can be studied in a similar basis invariant way as for the quark sector. For example, in the general two Higgs Doublet Model (2HDM) [24] (see [29] for a recent analysis) a CPI was identified in [30]. More generally, applying the invariant approach to scalar potentials has revealed relevant CPIs [31–33], including for the 2HDM [7, 34]. The basic idea is that if CP is conserved then all CPIs vanish (and vice versa). If any single CPI is non-zero then CP is violated. Finally, CP violating observables all have to be functions of CPIs.

In a recent paper [35] we considered yet more general Higgs potentials and adopted the powerful method of so-called contraction matrices in order to identify and construct new non-trivial CPIs, which we subsequently applied to potentials involving three or six Higgs fields (which can be either electroweak doublets or singlets) which form irreducible triplets under a discrete symmetry [35]. Having translated the well-known technique for constructing CPIs to diagrams and contraction matrices, we applied this formalism to some physically interesting cases which involve three or six Higgs fields which fall into irreducible triplet representations of discrete symmetries belonging to the $\Delta(3n^2)$ and $\Delta(6n^2)$ series, including A_4 , S_4 , $\Delta(27)$ and $\Delta(54)$. We were mainly interested in the question of explicit CP violation for such Higgs potentials, although a simple example of spontaneous CP violation was also discussed. Here we shall be principally concerned with whether those potentials which respect CP can lead to spontaneous CP violation. In other words we extend our formalism by including also Vacuum Expectation Values (VEVs), obtaining Spontaneous CPIs (SCPIs) that are non-vanishing if CP is spontaneously violated (as considered earlier in [31, 32]).

The purpose of this paper, then, is to discuss Spontaneous CP Violation (SCPV) for potentials involving three or six Higgs fields which fall into irreducible triplet representations of discrete symmetries belonging to the $\Delta(3n^2)$ and $\Delta(6n^2)$ series, including A_4 , S_4 , $\Delta(27)$ and $\Delta(54)$. These symmetry groups of the potentials considered in this paper are motivated by the fact that all of them are good candidates for discrete flavour symmetries. It should be noted that the actual symmetry of the potential can be different from the symmetry group imposed and where this distinction is important, it will be discussed. For each case, we write down the potential and find various global minima for different regions

of the parameter space of the potential, as recently summarised in [36]. In principle one could test which CP symmetries are preserved by VEVs, but it can be non-trivial to know all CP symmetries of the potential, we therefore prefer to use invariants. In each case we shall consider CP-odd basis Invariants (CPIs) that indicate SCPV — which we refer to as Spontaneous CP-odd Invariants (SCPIs). This builds on and was enabled by [35], where diagrammatic methods for constructing such invariants were further developed to the point of making them useful for the analysis of complicated example models such as the authors have in mind for this paper. In cases where CP is preserved we give a CP symmetry of the potential that is preserved by those VEVs, otherwise we show a non-zero CP-odd Invariant. In models where CP is violated spontaneously, thanks to the enhanced symmetry at high energies, the number of parameters of the model can be greatly reduced, and thanks to the controlled breaking of CP, the strength of CPV will have to be a function of these parameters too, which relates observable low-energy phenomena to possibly extremely high-energy parameters.

We emphasise that the work here extends the scope of the existing main models of SCPV considered in the literature based on models with 2 Higgs doublets (2HDM) or 3 Higgs doublets (3HDM). We remark that the most general 2HDM has a sufficient number of complex parameters to allow for explicit CPV. If a CP symmetry is imposed on the Lagrangian in the unbroken phase, the explicit CPV disappears and spontaneous CPV becomes possible. However, in both of those cases, when coupled to fermions, the model gives rise to flavour-changing neutral currents. When the latter are forbidden using any kind of symmetry, also both of explicit CPV (i.e. CPV with zero VEV) and SCPV disappear. 3HDMs improve a little on the situation of the 2HDM, as both of explicit CPV and FCNCs can be eliminated separately by CP and flavour-type symmetries, such that SCPV is possible without FCNCs. In particular a general analysis of CP invariants in 3HDMs suitable for SCPV does not exist, and one of the motivations of this work is to extend the discussion of CPIs suitable for these cases. We are also interested in models with 6 Higgs doublets (6HDM), for which the question of SCPV has not been analysed at all, and we extend our analysis to such cases also. Specifically, we expand our analysis by considering also the 6-Higgs potentials invariant under the discrete symmetries A_4 , S_4 , $\Delta(27)$, $\Delta(54)$, $\Delta(3n^2)$ and $\Delta(6n^2)$ with $n > 3$. In very special symmetric cases a phenomenon called geometric CP violation (GCPV) arises which means that values of complex CP-violating phases of VEVs do no longer depend on the parameters of the potentials in the region of the parameter space where they are minima. The only case that was known for a long time was that of Higgs triplets of $\Delta(27)$ [37]. Recently, several other cases had been discussed by [38, 39]. We shall discuss new examples of such GCPV.

It is important to further clarify which relevant results are already available in the literature, and which we need to obtain for the first time. A complete list of possible global minima for the A_4 , S_4 , $\Delta(27)$ and $\Delta(54)$ Higgs potentials with 3 Higgs fields has been obtained by [40], which includes furthermore cases without irreducible triplets such as S_3 (the S_3 is itself studied in great detail in [41]). For the cases where the respective potential is CP conserving in general (e.g. A_4) [40] also verified none of these VEVs spontaneously violate CP. While we analyse cases with 3 Higgs arranged as irreducible triplets, we go

beyond confirming the existing results for the 3-Higgs potentials with SCPIs, employing our methodology to analyse the VEVs and CP properties of other relevant cases. We continue by checking 3-Higgs potentials where a specific CP symmetry is imposed in addition to a discrete symmetry for which the potential is in general CP violating (e.g. the $\Delta(27)$, $\Delta(54)$) — this includes the case where Spontaneous Geometrical CP Violation (SGCPV) was first identified [37].

The layout of the paper is as follows: in section 2 we set our notation and describe how to identify CP violation and spontaneous CP violation within the basis invariant formalism. In section 2.3 we present the spontaneous CP invariants (SCPIs) that we will use throughout the paper. In section 3 we list the discrete symmetry groups and the potentials invariant under them that we consider (with one or two triplets), as well as a list of candidate Vacuum Expectation Values (VEVs) that are global minima of some of those potentials. In sections 4.1, and 4.2 we apply the SCPIs to the potentials invariant under the respective symmetry groups, checking if CP is conserved when all SCPIs we calculate vanish, and otherwise checking which of the VEVs we have found spontaneously violate CP. For any VEVs that do violate CP, we further consider if their phases are geometrical, i.e. if there is Spontaneous Geometrical CP Violation. Section 5 is a summary of our results and section 6 concludes the paper.

2 CP violation

In this section we review the current understanding of CP symmetries in models of several scalar fields.

2.1 Generalities

Usually, CP transformations are thought to act on scalars via complex conjugation of the field itself and parity transformation of the field coordinates, combined with arbitrary unitary basis transformations between fields with equal quantum numbers to form what is then called generalized CP transformations. A multiplet of scalar fields $\varphi = (\varphi_1, \dots, \varphi_n)$ would then be to transform under such a generalized CP transformation as

$$\varphi \mapsto X\varphi^*(x^P) \tag{2.1}$$

with a unitary matrix X and $x^P = (t, -\mathbf{x})$. Recently, [42] re-opened the discussion about how for complex scalars that have no charge associated to an Abelian symmetry, CP transformations without complex conjugation are equivalent to such with complex conjugation, by which only the (generalised) parity part of the transformation plays a role.

In the scalar potentials considered by us, fields are assumed, at least before symmetry breaking, to carry some conserved U(1) charge, as happens automatically for Higgs doublets, and imposed on EW singlets where they are considered, in the latter case primarily to render the potential even. For this reason, we think that at least symmetries of CP-type before the symmetry breaking will involve complex conjugation for the fields considered here.

When defining a model, in principle arbitrary CP-type symmetries can just be imposed onto a model at high energies in addition to all its other symmetries to render it explicitly CP-conserving and in addition maybe constrain it in other ways. If the potential has both pure flavour-type and CP-type symmetries, the notion of the *consistency* of these symmetries exists [43, 44].

By this one means that only certain CP-type symmetries can be imposed onto a model without enlarging the flavour-type symmetries of the potential. In essence, apart from the different physical interpretation, there is no difference between flavour-type and CP transformations, as both just relate different degrees of freedom. Consider a set of scalar fields φ and its complex conjugate, and combine them as

$$\phi = \begin{pmatrix} \varphi \\ \varphi^* \end{pmatrix}. \quad (2.2)$$

When written in this way, pure flavour-type and CP-type transformations act in the following ways:

$$\phi \xrightarrow{\text{flavour}} \begin{pmatrix} \rho & 0 \\ 0 & \rho^* \end{pmatrix} \phi, \text{ and } \phi \xrightarrow{\text{CP}} \begin{pmatrix} 0 & X \\ X^* & 0 \end{pmatrix} \phi. \quad (2.3)$$

Next, consider basis transformations U that at this stage (before symmetry breaking) only connect fields with identical (gauge) charges and thus act in the same way as a flavour transformation on the fields. There are some basis transformations for which $U\rho U^\dagger$ is just another group element ρ' . If the potential already has some CP symmetries, then such basis transformations generate additional CP transformations without changing the flavour type symmetry at all (and can be eliminated in the same way). The CP symmetries generated in this way are of the form

$$\begin{pmatrix} 0 & UXU^T \\ U^*X^*U^\dagger & 0 \end{pmatrix}. \quad (2.4)$$

Such basis transformations had been considered in [45] and have the effect of leaving the potential form-invariant and relate different points in the parameter space of the potential.

However, an arbitrary CP transformation with matrix X , which splits into symmetric and antisymmetric parts, $X = X_s + X_a$, transforms under a flavour basis transformation as follows,

$$UXU^T = UX_sU^T + UX_aU^T = \tilde{X}_s + UX_aU^T =: \tilde{X}, \quad (2.5)$$

where in the second step U can now be chosen such that it diagonalises X_s while UX_aU^T is still antisymmetric. Thus, always a flavour basis exists where at least for one of the pure CP-type symmetries that are not related by flavour transformations or transformations that leave the potential form-invariant the corresponding X can be made the sum of a diagonal and an antisymmetric matrix. In addition, if X was symmetric from the start, it can be made the identity matrix. This basis transformation does not change the size of the overall symmetry of the potential which is generated by all of the above 2×2 block matrices corresponding to flavour and CP transformations. Furthermore, this basis transformation generally does not leave the flavour symmetry or the potential form-invariant. The order-4

CP symmetry discussed in [46] and [42] is of this form of a sum of a diagonal and an antisymmetric matrix.

Arbitrary CP transformations can enlarge the pure flavour-type symmetry. The simplest way in which imposed flavour and CP symmetries combine to a pure flavour-transformation is

$$\begin{pmatrix} 0 & X \\ X^* & 0 \end{pmatrix} \begin{pmatrix} \rho & 0 \\ 0 & \rho^* \end{pmatrix} \begin{pmatrix} 0 & X \\ X^* & 0 \end{pmatrix} = \begin{pmatrix} X\rho^*X^* & 0 \\ 0 & X^*\rho X \end{pmatrix} \quad (2.6)$$

If now $X\rho^*X^*$ is not another element of the imposed flavour group, the actual flavour-type symmetry has been enlarged, and if on the other hand, for every group element g a group element g' exists such that

$$X\rho(g)^*X^* = \rho(g') , \quad (2.7)$$

then X is called consistent, cf. [43, 44].

In addition, it is clear that even powers of CP transformations are flavour transformations and corresponding conditions arise for $(XX^*)^n$.

Next, if the imposed X was symmetric and one can use a basis transformation to go to a basis where $\tilde{X} = 1$, combining CP and flavour transformations as above gives

$$\begin{aligned} \begin{pmatrix} 0 & X \\ X^* & 0 \end{pmatrix} \begin{pmatrix} \rho & 0 \\ 0 & \rho^* \end{pmatrix} \begin{pmatrix} 0 & X \\ X^* & 0 \end{pmatrix} &\rightarrow \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} U & 0 \\ 0 & U^* \end{pmatrix} \begin{pmatrix} \rho & 0 \\ 0 & \rho^* \end{pmatrix} \begin{pmatrix} U^\dagger & 0 \\ 0 & U^T \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ &= \begin{pmatrix} U^*\rho^*U^T & 0 \\ 0 & U\rho U^\dagger \end{pmatrix} \end{aligned} \quad (2.8)$$

while the original flavour transformations become $U\rho U^\dagger$ and the flavour symmetry in the basis where $\tilde{X} = 1$ is generated by all of $U\rho U^\dagger$ and of $U^*\rho^*U^T$. On the other hand, when combined with CP with $X = 1$, the complete original flavour symmetry was generated by all of ρ and of ρ^* .

For CP transformations where no basis exists where X can be made the identity, the same conditions arise, only with an additional matrix \tilde{X} appearing, and the full flavour symmetry in the basis where the symmetric part of X has been made diagonal is generated by all of $U\rho U^\dagger$ and of $\tilde{X}U^*\rho^*U^T\tilde{X}^*$.

Conversely, when one assumes that the flavour symmetry is complete and the CP part of the full symmetry is generated by a single matrix X , and as observables of CPV are invariants under internal basis transformations, this means that *for fixed flavour-type symmetry combined with a single CP generator, the physically different CP generators are of the form of the r.h.s. of eq. (2.5)* (up to changes of the basis of the group matrices that does not extend the flavour-type symmetry).

For models with more than one CP generator, the situation can be more complicated, as in the basis where one of them is of the form of eq. (2.5), the other can still be arbitrary. Unfortunately, this situation has to be outside the scope of this paper.

Later, when example potentials are studied, for chosen flavour symmetries, we still impose various CP symmetries and boldly ignore the question of consistency. If the potential

invariant under the symmetry already is automatically CP invariant under some (general) CP symmetry, then there is no doubt that the CP symmetry is consistent with the flavour symmetry. Otherwise, the potential has explicit CPV in general and if we impose a CP symmetry it necessarily increases the total symmetry content of the potential.¹ In those cases there are situations where the CP symmetries will then enlarge the actual flavour-type symmetry of the potential, and building a model this way is still fine, if one is aware that the full symmetry is always generated by the 2×2 flavour-type and CP-type block matrices. The first potential we discuss here where this is relevant has at least initially a $\Delta(54)$ flavour symmetry and further discussion can be found in the respective section 3.1.4. In all cases we always clearly specify the form of the potential, and when imposing a CP symmetry, what coefficients are made to vanish or become related. In any case, of the (imposed) CP symmetries appearing in table 6 and table 7, all are consistent in the sense of eq. (2.7). However, in the case of $X_{\varphi\varphi'}$ for two triplets of $\Delta(3n^2)$, the CP symmetry is only consistent when considering also the $U(1) \times U(1)$ part of the flavour symmetry, as noted again in section 3.2.2.

2.2 Spontaneous CP Violation

One of the questions we want to investigate in this paper is, when does geometrical CPV arise spontaneously (SGCPV) — what are the conditions on the potential or on the symmetries of the potential? Finding more cases with SGCPV is a step in this direction. In the literature, e.g. [37–39], geometrical CP violation has been defined as the situation when the relative phases of a VEV become geometrical (originally denoted as “calculable” [37]) which is to mean that in the region of parameter space of the potential where this VEV is a (global) minimum, these relative phases do not depend on the model parameters. This criterion has two components, one, that the VEV is geometric, and second, that CP is violated spontaneously by it. Geometric CPV is theoretically appealing because the strength of CP violation is no longer a function simply of arbitrary and at least in the near future unmeasurable parameters and furthermore also as phases arising in geometric CPV are stable against renormalization, as they are protected by a residual symmetry [37].

One good criterion for spontaneous violation of CP that we rely on throughout the paper is when at least one CP-odd invariant that can indicate spontaneous CPV (see next subsection) is non-zero. When CP is violated, then at least one CP-odd invariant is non-zero, but it is not possible at the moment to obtain a complete list of invariants and it is impractical to test all of them in any case. The situation would be simpler, if one had a basis of CP-odd invariants, which is a (small) finite set of CP-odd invariants with the property that if all of them are zero, CP is conserved. Indeed, such a basis of CP-odd invariants is known for the 2HDM, [34], but for the more symmetric potentials we will consider, all invariants from [34] vanish trivially without indicating CP-conservation, because whether a set of invariants forms a basis is model-dependent, which is why in [35] it was found necessary to find additional CP-odd invariants. Finally, if CP is conserved,

¹Note that additional care is required in situations where the total symmetry of the potential becomes continuous, as this may lead to Goldstone bosons arising when the symmetry is broken.

then all CP-odd invariants (which includes all possible basis sets) vanish of course. Again, the strength of CP-odd invariants that indicate spontaneous CP violation is that it is not necessary to systematically know all CP symmetries of the unbroken potential, a task which can be non-trivial.

Without SCPIs, one could also check directly whether a VEV preserves a specific CP symmetry that leaves the potential invariant. This is arguably more direct, but it has the significant drawback that it is not sufficient to find a CP symmetry not preserved by the VEV for there to be CPV, but rather all CP symmetries that left the potential invariant must be broken by the VEV. Finding all inequivalent CP symmetries that leave a potential invariant can be non-trivial, as the above discussion elaborates, and checking each is broken by the particular VEV is also cumbersome. Conversely, if a single CP symmetry that leaves the (unbroken) potential invariant is preserved by the VEV, then CP is conserved (this is a sufficient condition). This makes the direct check very convenient to confirm CP is preserved. We now review this direct condition in a notation that follows [35]. Consider an even scalar potential written in the following standard form,

$$V(\varphi) = Y_b^a \varphi_a \varphi^{*b} + Z_{cd}^{ab} \varphi_a \varphi_b \varphi^{*c} \varphi^{*d}, \quad (2.9)$$

where φ contains as components the components of all fields. Y and Z are tensors that contain all allowed couplings and are subject to possible symmetries acting on φ . In the following only the action of CP transformations is repeated from [35].² Assume that this potential is CP-conserving, so in particular invariant under a set of CP transformations of the fields, such that when one transforms the fields as in

$$\varphi \mapsto X \varphi^*, \quad (2.10)$$

with a unitary matrix X , the potential is unchanged,

$$V(\varphi) = V(X \varphi^*). \quad (2.11)$$

With indices explicitly shown, we write the transformation of eq. (2.10) as³

$$\varphi_a \mapsto X_{aa'} \varphi^{*a'} := \sum_{a'} X_{aa'} (\varphi_{a'})^* \quad (2.12)$$

and for the complex conjugated field as

$$\varphi^{*a} \mapsto X^{*aa'} \varphi_{a'}. \quad (2.13)$$

Note that in this notation, $X^{*aa'} = X^{\dagger a'a}$. Denote the whole set of such X under which a potential is invariant in the unbroken phase by $\mathcal{X} := \{X\}$. In appendix A the traditional argument is repeated in the notation of eq. (2.9) that a VEV $v = \langle \varphi \rangle$ conserves CP if at least for one of the matrices $X \in \mathcal{X}$ holds that

$$v_a = X_{aa'} v^{*a'}. \quad (2.14)$$

²The rules for indices can be summarized as follows: on φ , $(\varphi_a)^* = \varphi^{*a}$, $(\varphi^{*a})^* = \varphi_a$ and on matrices, e.g. a basis transformation U , $(U_a^b)^* = U^{*a}_b$. Consequently, $(U_a^b \varphi_b)^* = U^{*a}_b \varphi^{*b} = (U^\dagger)_b^a \varphi^{*b}$.

³This is a slight improvement on the notation in [35], where CP transformations were written with one upper and one lower index.

Inverting the argument this means that a VEV violates CP if for none of the original CP trasfos in the unbroken phase, eqs. (2.12), (2.13), the previous condition can be fulfilled.⁴

The type of CP transformations without complex conjugation discussed in [42], while they might be a symmetry of the neutral components of scalar fields after symmetry breaking, are in our case not symmetries in the unbroken phase.

2.3 Spontaneous CP invariants

From the standard form of the potential, eq. (2.9), invariants can be built in a model-independent way that are CP-odd by construction. Model-independent here is to mean that these invariants are constructed from the Y and Z tensor in eq. (2.9) and work for every potential that can be put into this standard form.

Under basis transformations, the field multiplet and its conjugate transform as $\varphi \mapsto U\varphi$ with which the Y and Z tensor of the even potential in the standard form of eq. (2.9) transform under basis transformations of the fields and its conjugate as

$$Y_a^b \mapsto U_a^{a'} Y_{a'}^{b'} U_{b'}^b, \quad (2.15)$$

$$Z_{ac}^{bd} \mapsto U_a^{a'} U_c^{c'} Z_{a'c'}^{b'd'} U_{b'}^b U_{d'}^d. \quad (2.16)$$

Additionally, VEVs transform as vectors under basis transformations,

$$v_a \mapsto U_a^{a'} v_{a'} \text{ and } v^{a*} \mapsto U_{a'}^a v^{a*} \quad (2.17)$$

and one can see now that every combination of Y , Z , v and v^* where all indices are correctly contracted forms a basis invariant. Complex conjugation acts on invariants by interchanging upper and lower indices, such that the complex conjugate of a basis invariant can be obtained by interchanging all upper and lower indices of all components. In [35], invariants had been related to diagrams where arrows indicate index contractions and other parts of the diagram tensors and VEVs. In that formalism, an invariant is CP-odd if the diagram does not stay identical when inverting the direction of all arrows, which is the diagrammatic equivalent to complex conjugation. We recall then that those basis invariant contractions can be built into CPIs by subtracting its conjugate, for a given invariant J , $\mathcal{J} = J - J^*$ is the CPI and for those relevant to SCPV we refer to \mathcal{J} as a spontaneous CPI (SCPI). The diagrammatic formalism furthermore allowed for a systematic search for CP-odd invariants. In [35], all inequivalent CPIs up to some order were constructed, including some SCPIs.

Here, all SCPIs that were listed in appendix B.5 of [35] have been evaluated for all example potentials that will be considered in later sections. As a first example, we consider here briefly the most general 2-Higgs Doublet Model (2HDM) potential. In our notation, brackets indicate the $SU(2)_L$ contractions e.g. $(H_1^\dagger H_1)^2 = (h_{1,1}^\dagger h_{1,1} + h_{1,2}^\dagger h_{1,2})^2$, therefore

⁴Recently, in [47], a case was discussed where a CP-type transformation is preserved by the vev, but CP is violated in the interplay of scalar and gauge sector.




Figure 1. The diagram corresponding to the invariant eq. (2.19). (In [35], the precise relation between invariants and diagrams is explained.)

we write the general two-Higgs-doublet (2HDM) potential as

$$\begin{aligned}
 V(H_1, H_2) = & m_1^2 H_1^\dagger H_1 + m_{12}^2 e^{i\theta_0} H_1^\dagger H_2 + m_{12}^2 e^{-i\theta_0} H_2^\dagger H_1 + m_2^2 H_2^\dagger H_2 \\
 & + a_1 (H_1^\dagger H_1)^2 + a_2 (H_2^\dagger H_2)^2 \\
 & + b (H_1^\dagger H_1) (H_2^\dagger H_2) + b' (H_1^\dagger H_2) (H_2^\dagger H_1) \\
 & + c_1 e^{i\theta_1} (H_1^\dagger H_1) (H_2^\dagger H_1) + c_1 e^{-i\theta_1} (H_1^\dagger H_1) (H_1^\dagger H_2) \\
 & + c_2 e^{i\theta_2} (H_2^\dagger H_2) (H_2^\dagger H_1) + c_2 e^{-i\theta_2} (H_2^\dagger H_2) (H_1^\dagger H_2) \\
 & + d e^{i\theta_3} (H_1^\dagger H_2)^2 + d e^{-i\theta_3} (H_2^\dagger H_1)^2. \tag{2.18}
 \end{aligned}$$

In it appear $H_1 = (h_{1,1}, h_{1,2})$ and $H_2 = (h_{2,1}, h_{2,2})$, and the arbitrary coefficients $a_1, a_2, b, b', c_1, c_2, d$. For the 2HDM, all SCPIs in [35] are non-zero although we don't show any as even the smallest expression is rather large. In any case they are not completely meaningful at this stage, as the general 2HDM is explicitly CP violating. When imposing a CP symmetry, the expression becomes meaningful — for example when imposing trivial CP, the expression for the CP-odd invariant

$$\mathcal{J}_{2\text{HDM}} \equiv Z_{a_3 a_7}^{a_1 a_2} Z_{a_1 a_2}^{a_3 a_4} Z_{a_4 a_6}^{a_5 a_6} v_{a_5} v^{*a_7} - \text{c.c.}, \tag{2.19}$$

which of the invariants tested produces the smallest non-zero expressions for the 2HDM, becomes a function of the coefficients in the potential multiplying the VEVs in the following combination:

$$\mathcal{J}_{2\text{HDM}} = F(a_1, a_2, b, b', c_1, c_2, d)[\langle h_{2,1} \rangle \langle h_{1,1} \rangle^* + \langle h_{2,2} \rangle \langle h_{1,2} \rangle^* - \text{h.c.}] \tag{2.20}$$

i.e. it correctly identifies that SCPV depends in this case on the relative phase between the VEVs of H_1 and H_2 (note that charge preserving VEVs correspond to $\langle h_{1,1} \rangle = \langle h_{2,1} \rangle = 0$). The diagram corresponding to this invariant is shown in figure 1. The same quadratic VEV dependence factors out also when calculating the $J_1^{(3,2)}$ SCPI that has 2 pairs of VEVs, as the full expression is a more complicated quadratic function of the VEVs multiplying $[\langle h_{2,1} \rangle \langle h_{1,1} \rangle^* + \langle h_{2,2} \rangle \langle h_{1,2} \rangle^* - \text{h.c.}]$. For the more symmetric potentials considered in the remainder, the above simple invariant vanishes trivially, i.e. without indicating CP-conservation. We will not consider the 2HDM further and will instead use as our examples 3HDM and 6HDM potentials that are invariant under discrete symmetries.

For all potentials besides the 2HDM that we consider as examples, of the SCPIs listed in [35] only $\mathcal{J}_1^{(3,2)}$ is non-zero. For this reason we drop the subscript, referring to it as




Figure 2. The diagram corresponding to the invariant $J^{(3,2)}$.




Figure 3. The diagram corresponding to the invariant $J^{(3,3)}$.

$\mathcal{J}^{(3,2)}$ (we keep the superscript to distinguish from another SCPI which we will use). The contraction $J^{(3,2)}$ of Z tensors and VEVs v out of which $\mathcal{J}^{(3,2)}$ is formed, is in terms of the Z tensor and VEVs given by

$$J^{(3,2)} \equiv Z_{a_4 a_5}^{a_1 a_2} Z_{a_2 a_6}^{a_3 a_4} Z_{a_7 a_8}^{a_5 a_6} v_{a_1} v_{a_3} v^{*a_7} v^{*a_8}. \quad (2.21)$$

The diagram corresponding to this invariant is shown in figure 2, where it also be seen that it is CP-odd as the diagram with inverted arrows cannot be made identical to the original diagram, however the dots (= Z tensors) and crosses (= VEVs) are moved around. Given that only one of the previously constructed SCPIs is non-trivial for the potentials we are interested in, the question arises if there is another SCPI which gives non-trivial results. Searching for SCPIs with an additional Z tensor, i.e. considering invariants of type $J_i^{(4,2)}$, an explicit (but not systematic) search did not yield a positive answer. On the other hand, an invariant $J^{(3,3)}$ with an additional pair of VEVs revealed another interesting SCPI which turned out to be sufficient for our purposes (and therefore we also do not use a subscript to label it):

$$J^{(3,3)} \equiv Z_{a_1 a_2}^{a_7 a_8} Z_{a_3 a_4}^{a_5 a_9} Z_{a_5 a_6}^{a_1 a_3} v_{a_7} v_{a_8} v_{a_9} v^{*a_2} v^{*a_4} v^{*a_6}. \quad (2.22)$$

The diagram corresponding to this invariant is shown in figure 3.

The results of evaluating both these SCPIs for the various potentials, CP symmetries and VEVs is summarized in section 5, including table 6 and table 7. As shown in the following sections where different potentials are considered in detail, $J_1^{(3,3)}$ provides us with additional relevant information that complements what we learn from $\mathcal{J}^{(3,2)}$. For example, from our examples, the simplest 3 field potential that for arbitrary VEV values (which not necessarily can be obtained from the potential) gives a non-vanishing expression for $\mathcal{J}_1^{(3,3)}$

is invariant under S_4 , $V_{S_4}(\varphi)$. The SCPI $J^{(3,2)}$ gives a vanishing expression independently of the VEVs considered, whereas the expression for $\mathcal{J}^{(3,3)}$ is in general:

$$\mathcal{J}^{(3,3)} = c(c^2 - s^2) \left[|v_1|^2 v_1^2 (v_2^{*2} + v_3^{*2}) + |v_2|^2 v_2^2 (v_3^{*2} + v_1^{*2}) + |v_3|^2 v_3^2 (v_1^{*2} + v_2^{*2}) - \text{h.c.} \right]. \quad (2.23)$$

Note that this doesn't mean that it is possible to have SCPV in this potential, as when plugging in any of the possible VEVs [40], $\mathcal{J}^{(3,3)}$ does indeed vanish, as shown in section 4.1.2 and summarised in section 5.

3 Potentials and Vacuum Expectation Values

In this paper we are mainly interested in three- and six-Higgs-doublet models, where fields form one or two irreducible triplet representations of some discrete symmetry group. The most complicated potentials that will be considered in this paper then describe six scalar $SU(2)_L$ doublets (or if only $U(1)_{em}$ -preserving minima are considered, equivalently, $SU(2)_L$ singlets), and finding (global) minima can be a non-trivial task.

The aim of this section is to arrive at a list (not necessarily exhaustive, but complete enough for further analysis) of possible VEVs of potentials of two scalar triplets of A_4 , S_4 , $\Delta(27)$, $\Delta(54)$, and $\Delta(3n^2)$ and $\Delta(6n^2)$ with $n > 3$.

First, the potentials of one triplet will be re-analysed, after which the potentials of two triplets are examined. As the aim in the next section will be to study geometrical CPV, potentials are made CP-conserving via various additional CP symmetries.

We identify classes of potentials by the symmetry imposed by us. This differs from [40], where potentials were identified by their full symmetry, both of flavour- and CP-type, which was enabled by a classification of those symmetries in [15]. The full symmetry of the potential can be different from the imposed symmetry, e.g. because cutting off the potential at renormalisable operators can enlarge the symmetry group.

The full symmetry of the potential or of a part of it is what will be relevant in the following, and it will hopefully always be made clear which exact symmetry is under discussion.

For $SU(2)_L$ singlets, always an additional $U(1)$ symmetry was imposed to force the potential to be even. In addition, to not break $U(1)_{em}$, in a basis where at least one VEV component of one doublet is zero, also the same component of the other doublets has to be zero. When in a term in a potential of singlet fields the flavour indices allow for various ways of contracting $SU(2)_L$ indices, then if without loss of generality, the bottom component of all doublets is set to zero, such terms in doublet potentials will coincide again.

For the aforementioned reasons the possible vacuum expectation values that don't break $U(1)_{em}$ of a doublet potential with some symmetry will be identical to those of the corresponding singlet potential made even by an additional $U(1)$. Some of these potentials had been considered in [40], where also the full list of global VEVs was obtained via a geometric method.

When searching for the minima of a potential, one is not just interested in the minimum at some position in the parameter space of the potential, but one would ideally like to know what the minima of the potential are in every corner of its parameter space.

Minima that are related by symmetries of the potential have the same energy by definition and such sets of related VEVs are called orbits, of which a potential can have more than one. Notably, when the symmetry of the potential is increased (of flavour- or CP-type), this has the effect of merging orbits. Beyond this it should be noted that the full symmetry of the VEV set is larger than that of the potential. The set of VEVs is also invariant under transformations under which the potential was just form-invariant, cf. [45], as such transformations connect orbits. In other words, while the symmetries of the potential are valid at every point of the whole space of the parameters of the potential, the set of VEVs additionally contains information about all potentials that differ by the values of their parameters but not by their symmetries.

In the notation of section 2.2, classical minima of the potential are solutions of

$$0 = \frac{\partial V}{\partial v_i} = Y_b^i v^{*b} + 2Z_{cd}^{ib} v_b v^{*c} v^{*d} \quad (3.1)$$

and the corresponding conjugated equation, as well as the condition that the matrix of all second derivatives is positive definite. The mixed derivative by VEV and conjugated VEV would e.g. be

$$\frac{\partial^2 V}{\partial v_i \partial v^{*j}} = Y_j^i + 4Z_{jd}^{ib} v_b v^{*d}. \quad (3.2)$$

An alignment is basically, in accord with the literature, a VEV where one does not care about its absolute length but only about its direction. The linear symmetries in consideration can in any case not constrain the length of the VEV, as the overall sizes of quadratic and quartic terms are entirely unrelated. At the end of the day one is interested in interesting alignments, especially such that violate CP spontaneously or even geometrically. In a following section, CP violation will be discussed in greater detail, this section focuses on obtaining VEVs of the potentials considered in this paper.

New VEVs are generated by analysing which degrees of freedom become physical when going from a potential with high symmetry to one with lower symmetry. Consider a potential

$$V = V_G + V_H \text{ with } H \subset G. \quad (3.3)$$

All alignments v_i of potentials generally fall into orbits $\{H \cdot v_i\}$. If one considers the subpotential V_G separately, it will have minima that again fall into orbits $\{G \cdot v_i\}$. Next, consider V_H as terms that explicitly break the symmetry of the potential from G to H . By this, the orbits of V_G are now split into several orbits organized by H , sometimes only a finite number of orbits, occasionally there can be a new continuous parameter distinguishing them. In this case the full potential can be minimized in this new parameter, which is often possible analytically.

First, it is thus tested for potentials with one triplet of scalars, which orbits can be obtained from the orbits of more symmetric potentials, just by applying the symmetries of the less symmetric potential to the orbits of the more symmetric one. In this way, degrees of freedom that were unphysical in the more symmetric potential can become physical and after minimizing just the part of the potential that depends on these new degrees of freedom, one obtains as global minima exactly those that had also been obtained

in [40]. One could thus conjecture that for certain relations between symmetry groups, also relations between the global minima of potentials with these symmetries exist which might be made more precise.

After that, the same method is applied to obtain some of the (conjectured) global minima of potentials of two triplets of scalars from those of potentials that are just the sum of potentials of one triplet without cross-terms. Again this simply shows which degrees of freedom become physical due to the reduced symmetry. Still, many new interesting global minima are obtained, many of which are geometric and many of which will turn out to be CP violating and especially geometrically CP violating, which constitute the first examples of this phenomenon outside the original $\Delta(27)$ model and [38, 39]. The cases found here are in particular the first new examples of geometric CPV with six scalars.

Generally, a VEV is defined as being geometric when in the region of the parameter space of the potential where it is a (global) minimum, its direction but not its length depend on the potential parameters, or if the potential is expressed in terms of Y and Z tensors,

$$v = |v|(Y, Z)\hat{v}, \quad (3.4)$$

where $|v|$ is the length of the VEV and the only part that depends on the potential parameters, and \hat{v} is the normalised direction that does not depend on them. This is equivalent to

$$\frac{\partial \hat{v}}{\partial Y_b^a} = 0 \text{ and } \frac{\partial \hat{v}}{\partial Z_{cd}^{ab}} = 0. \quad (3.5)$$

A VEV is primarily a solution of the minimisation conditions, in the notation of eq. (2.9), given by

$$\frac{\partial V}{\partial \phi_i} \rightarrow Y_b^i v^* + 2Z_{cd}^{ib} v_b v^{*c} v^{*d} \equiv 0 \text{ and } \frac{\partial V}{\partial \phi^{*j}} \rightarrow Y_j^a v_j + 2Z_{jd}^{ab} v_a v_b v^{*d}. \quad (3.6)$$

It is normally not possible to easily arrive at a closed form of the VEV, but it is implicitly defined as solution of the minimisation conditions. This allows to use the theorem about the derivatives of implicitly defined functions to arrive at a relation involving $\partial \hat{v} / \partial Y$ and $\partial \hat{v} / \partial Z$. Call a vector of the potential parameters \vec{z} . If $f(|v|, \hat{v}, \vec{z})$ is the implicit function defining $|v|$ and \hat{v} , i.e. the minimisation condition, then an expression for $d\hat{v} / d\vec{z}$, which is identical to $\partial \hat{v} / \partial \vec{z}$, as \hat{v} only depends on \vec{z} , can be obtained via the differential of f :

$$df = \frac{\partial f}{\partial \vec{z}} d\vec{z} + \frac{\partial f}{\partial |v|} d|v| + \frac{\partial f}{\partial \hat{v}} d\hat{v}, \quad (3.7)$$

from which follows

$$\frac{\partial \hat{v}}{\partial \vec{z}} = - \left(\frac{\partial f}{\partial \hat{v}} \right)^{-1} \left(\frac{\partial f}{\partial \vec{z}} + \frac{\partial f}{\partial |v|} \frac{\partial |v|}{\partial \vec{z}} \right). \quad (3.8)$$

While the above is interesting and the most direct line of thought, we did not manage to extract any practical criteria on the potential or its symmetries and do not follow it further in this work.

We had recently listed the minima obtained in the following in a short note, [36], but without such a detailed discussion of the derivation as given here and without considering the effect of CP symmetries on the minima of the potentials of two triplets.

3.1 Potentials of one triplet

3.1.1 $\Delta(3n^2)$ and $\Delta(6n^2)$, with $n > 3$

We start with a discussion of the simplest potential considered in this paper: that of one triplet of $\Delta(3n^2)$, which is identical to the potential of one triplet of $\Delta(6n^2)$.

In the notation from [48], triplets of $\Delta(3n^2)$ can be enumerated as $\mathbf{3}_{(k,l)}$, where $k, l = 0, 1, \dots, n-1$. The conjugate of $\mathbf{3}_{(k,l)}$ is $\mathbf{3}_{(-k,-l)}$. For triplets in the same irrep.,

$$\mathbf{3}_{(k,l)} \otimes \mathbf{3}_{(k,l)} = [\mathbf{3}_{(2k,2l)} + \mathbf{3}_{(-k,-l)}]_s + [\mathbf{3}_{(-k,-l)}]_a. \quad (3.9)$$

Here, s and a are the symmetric and antisymmetric combinations.

This potential had not been analysed in [15, 40], as the large discrete symmetry is so constraining that the renormalisable potential ends up having a continuous symmetry: apart from the $U(1)$ symmetry that make the potentials even in ϕ , this potential has additional continuous symmetries, cf. eq. (3.21). The potential is

$$V_{\Delta(6n^2)}(\varphi) = V_{\Delta(3n^2)}(\varphi) \equiv V_0(\varphi) \quad (3.10)$$

where, as it will appear in subsequent potentials, we define

$$V_0(\varphi) \equiv -m_\varphi^2 \sum_i \varphi_i \varphi^{*i} + r \left(\sum_i \varphi_i \varphi^{*i} \right)^2 + s \sum_i (\varphi_i \varphi^{*i})^2. \quad (3.11)$$

If each component of the triplet is an $SU(2)_L$ doublet,

$$H = (h_{1\alpha}, h_{2\beta}, h_{3\gamma}), \quad (3.12)$$

one additional invariant appears, due to the two different ways to perform the $SU(2)_L$ contraction on the discrete symmetry invariant $(\sum_i \varphi_i \varphi^{*i})^2$, when the φ are replaced by Higgs doublets,⁵

$$\sum_{i,j,\alpha,\beta} \left[r_1(h_{i\alpha} h^{*i\alpha})(h_{j\beta} h^{*j\beta}) + r_2(h_{i\alpha} h^{*i\beta})(h_{j\beta} h^{*j\alpha}) \right]. \quad (3.13)$$

Here we highlight the $SU(2)_L$ indices to clarify the distinct $SU(2)_L$ contractions. Then we have

$$V_{\Delta(6n^2)}(H) = V_{\Delta(3n^2)}(H) \equiv V_0(H) \quad (3.14)$$

and we define $V_0(H)$ in analogy with eq. (3.11):

$$\begin{aligned} V_0(H) = & -m_h^2 \sum_{i,\alpha} h_{i\alpha} h^{*i\alpha} + \sum_{i,j,\alpha,\beta} \left[r_1(h_{i\alpha} h^{*i\alpha})(h_{j\beta} h^{*j\beta}) + r_2(h_{i\alpha} h^{*i\beta})(h_{j\beta} h^{*j\alpha}) \right] \\ & + s \sum_{i,\alpha,\beta} (h_{i\alpha} h^{*i\alpha})(h_{i\beta} h^{*i\beta}). \end{aligned} \quad (3.15)$$

⁵Since the doublet **2** of $SU(2)_L$ is a pseudoreal representation, it is also possible to combine $(h_{i\alpha} h_{j\beta} \epsilon^{\alpha\beta})(h^{*i\gamma} h^{*j\delta} \epsilon_{\gamma\delta})$ using the antisymmetric ϵ tensor. However, such a term is not independent of the two terms in eq. (3.13) as can be easily seen in an explicit calculation or by noting that **2** \times **2** = **1** + **3** which entails only two independent $SU(2)_L$ invariant quartic terms.

The potential eq. (3.11) can be minimized analytically and one obtains for $m_\varphi \neq 0$ three classes of non-zero VEVs that are internally related by symmetry transformations and representatives of those classes are

$$v_1 \cdot (1, 0, 0), \quad v_2 \cdot (1, 1, 0), \quad v_3 \cdot (1, 1, 1), \quad (3.16)$$

where

$$v_1^2 = \frac{m_\varphi^2}{2r + 2s}, \quad v_2^2 = \frac{m_\varphi^2}{4r + 2s}, \quad v_3^2 = \frac{m_\varphi^2}{6r + 2s}. \quad (3.17)$$

It can be tested that for all of the alignments listed previously, regions in the parameter space exist such that for potentials with parameters in that region, the corresponding alignment is the global minimum of the potential.

The potential in eq. (3.11) decays into two parts, one, namely the two invariants with parameters m_φ and r , invariant under all of $U(3)$, and a second part, consisting of the invariant with parameter s , invariant under $((U(1) \times U(1)) \rtimes S_3) \times U(1) =: \Delta(6\infty^2) \times U(1)$, where the $U(1)^2$ factor in brackets arises from sending n to infinity in $\Delta(6n^2)$, and the third $U(1)$ was imposed to keep the potential even:

$$V_0 = V_{U(3)} + \delta V_{\Delta(6\infty^2) \times U(1)}, \quad (3.18)$$

where

$$V_{U(3)} = -m_\varphi^2 \sum_i \varphi_i \varphi^{*i} + r \left(\sum_i \varphi_i \varphi^{*i} \right)^2, \quad (3.19)$$

and

$$\delta V_{\Delta(6\infty^2) \times U(1)} = s \sum_i (\varphi_i \varphi^{*i})^2, \quad (3.20)$$

from eq. (3.11). Occasionally, the label on a potential will refer to the imposed symmetry and occasionally, but only when talking about parts of potentials, to the full symmetry of that part of the potential. It is hoped that it is mostly clear from context which symmetry is meant.

The alignments of $V_{U(3)}$ all fall into one large orbit, represented e.g. by $(1, 0, 0)$, connected by arbitrary unitary transformations. The effect of the second, less symmetric, part of the potential is that this big orbit decays into several orbits in which now not yet phases but the direction of the VEV becomes physical.

To each of the alignments in eq. (3.16) quite a big orbit is attached, the members of which are related by the full symmetries of the potential, both such of flavour and CP type. The flavour symmetries of V_0 are generated by

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} e^{i\alpha} & 0 & 0 \\ 0 & e^{i\alpha} & 0 \\ 0 & 0 & e^{i\alpha} \end{pmatrix}, \begin{pmatrix} e^{i\beta} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{-i\beta} \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\gamma} & 0 \\ 0 & 0 & e^{-i\gamma} \end{pmatrix}, \quad (3.21)$$

where α, β, γ are arbitrary phases. Additionally, the potential is automatically invariant under canonical CP transformations, which we refer to as CP_0 . For one triplet, this

corresponds to an X matrix

$$X_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (3.22)$$

Note that the alignments in eq. (3.16) all conserve canonical CP (CP_0). The orbits of alignments of this potential are

$$\left\{ \begin{pmatrix} e^{i\eta} \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ e^{i\eta} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ e^{i\eta} \end{pmatrix} \right\}, \left\{ \begin{pmatrix} e^{i\eta} \\ e^{i\zeta} \\ 0 \end{pmatrix}, \text{permut.} \right\}, \left\{ \begin{pmatrix} e^{i\eta} \\ e^{i\zeta} \\ e^{i\theta} \end{pmatrix}, \text{permut.} \right\}. \quad (3.23)$$

Now one could wonder into what kind of orbits the above alignments of eq. (3.23) are bundled by applying a smaller symmetry group than $\Delta(6n^2) \times U(1)$. In other words, one asks, which phases and permutations of the alignments of the one-triplet potential of $\Delta(6n^2)$, that were unphysical under the symmetries of eq. (3.21) would no longer be unphysical, assuming for a moment that they could actually be minima of a potential with a smaller symmetry.

3.1.2 S_4

The potential of one triplet of S_4 is

$$V_{S_4}(\varphi) = V_0(\varphi) + \delta V_{S_4 \times U(1)}(\varphi) \quad (3.24)$$

with $V_0(\varphi)$ from eq. (3.11) and

$$\begin{aligned} \delta V_{S_4 \times U(1)}(\varphi) = & b (\varphi_1 \varphi_1 \varphi^{*3} \varphi^{*3} + \varphi_2 \varphi_2 \varphi^{*1} \varphi^{*1} + \varphi_3 \varphi_3 \varphi^{*2} \varphi^{*2}) \\ & + b (\varphi^{*1} \varphi^{*1} \varphi_3 \varphi_3 + \varphi^{*2} \varphi^{*2} \varphi_1 \varphi_1 + \varphi^{*3} \varphi^{*3} \varphi_2 \varphi_2). \end{aligned} \quad (3.25)$$

Note that b is real. The abbreviations cycl. to denote the cyclic permutations, and h.c. to indicate the hermitian conjugate allow us to write the potential of eq. (3.25) (and subsequent potentials) in a more compact way:

$$\delta V_{S_4 \times U(1)}(\varphi) = b [(\varphi_1 \varphi_1 \varphi^{*3} \varphi^{*3} + \text{cycl.}) + \text{h.c.}], \quad (3.26)$$

$$V_{S_4}(\varphi) = V_0(\varphi) + b [(\varphi_1 \varphi_1 \varphi^{*3} \varphi^{*3} + \text{cycl.}) + \text{h.c.}]. \quad (3.27)$$

The $SU(2)_L$ doublet version is similar:

$$V_{S_4}(H) = V_0(H) + \sum_{\alpha, \beta} b \left[(h_{1\alpha} h_{1\beta} h^{*3\alpha} h^{*3\beta} + \text{cycl.}) + \text{h.c.} \right]. \quad (3.28)$$

The flavour symmetries of the S_4 potentials, eq. (3.27) and (3.28) for $SU(2)$ singlets and doublets respectively, are generated by

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} e^{i\alpha} & 0 & 0 \\ 0 & e^{i\alpha} & 0 \\ 0 & 0 & e^{i\alpha} \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad (3.29)$$

and in addition, the potential is automatically invariant under CP_0 . Under these symmetries, the elements of the orbits of the potential of one triplet of $\Delta(6n^2)$, eq. (3.23), fall into the following orbits:

$$\begin{pmatrix} e^{i\eta} \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} e^{i\eta} \\ e^{i\zeta} \\ 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 \\ e^{i\zeta'} \\ 0 \end{pmatrix} \text{ with } \zeta' \in [0, \pi] \quad (3.30)$$

and

$$\begin{pmatrix} e^{i\eta} \\ e^{i\zeta} \\ e^{i\theta} \end{pmatrix} \rightarrow \begin{pmatrix} 1 \\ e^{i\zeta'} \\ e^{i\theta'} \end{pmatrix} \text{ with } \zeta'' \in [0, \pi] \text{ and } \theta' \in [0, 2\pi]. \quad (3.31)$$

What has happened so far is just that phases that were unphysical for $\Delta(6n^2)$ can now be physical. Minimizing the parts of the potential that depend on these potentially physical phases for the various orbits yields the following possible global minima of just the phase-dependent part, $\delta V_{S_4 \times U(1)}$,

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} \pm 1 \\ \omega \\ \omega^2 \end{pmatrix}, \begin{pmatrix} 1 \\ i \\ 0 \end{pmatrix}, \quad (3.32)$$

which have been rephased into the form as they appear in [40], and where we have defined

$$\omega \equiv e^{i2\pi/3}. \quad (3.33)$$

The above alignments obtained in this way are, maybe surprisingly, in agreement with [40].

Two questions arise now: firstly, does e.g. the potential of one triplet of S_4 have global VEVs that don't arise from making unphysical phases in $V_{\Delta(6n^2)}$ physical in $\delta V_{S_4 \times U(1)}$, which would be in contradiction to [40], at least for the potential of one triplet of S_4 ? With respect to the first question, for S_4 or any of the one-triplet potentials this is not the case. Secondly, what is the situation when going from $\Delta(6n^2)$ to $\Delta(54)$, $\Delta(27)$ or A_4 , or from S_4 to A_4 , or more interestingly for the actual topic of this paper, from potentials of two copies of triplets without cross-terms to such with cross terms? We address this second question in section 3.2.

3.1.3 A_4

The irreps. of A_4 are three one-dimensional representations and a real triplet. Two triplet products decompose as

$$\mathbf{3} \otimes \mathbf{3} = (\mathbf{1}_0 + \mathbf{1}_1 + \mathbf{1}_2 + \mathbf{3})_s + \mathbf{3}_a, \quad (3.34)$$

where subscripts s and a , denote symmetric and antisymmetric combinations respectively. Throughout, we use the [49] basis for A_4 products.

We charge φ under a $U(1)$ symmetry (or a discrete subgroup) forbidding terms such as $\varphi_i \varphi_i$ or $\varphi_i \varphi_i \varphi_i$, for a more direct generalisation to cases where the triplets transform

under the SM gauge group. The renormalisable scalar potential invariant under A_4 and such a $U(1)$ is then:

$$V_{A_4}(\varphi) = V_0(\varphi) + [c(\varphi_1\varphi_1\varphi^{*3}\varphi^{*3} + \text{cycl.}) + \text{h.c.}] , \quad (3.35)$$

with $V_0(\varphi)$ as defined in eq. (3.11), and where c is complex in general.

This potential is automatically invariant under the CP symmetry with:

$$X_{23} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} , \quad (3.36)$$

even for arbitrary complex coefficient c .

For $SU(2)_L$ doublets, the potential is similar

$$V_{A_4}(H) = V_0(H) + \sum_{\alpha, \beta} \left[c \left(h_{1\alpha} h_{1\beta} h^{*3\alpha} h^{*3\beta} + \text{cycl.} \right) + \text{h.c.} \right] . \quad (3.37)$$

This potential is also invariant under a CP transformation that involves swapping the second and third component in flavour space while keeping $SU(2)_L$ contractions unchanged, i.e. $h_{2\alpha} \rightarrow h^{*3\alpha}$ etc.:

$$X_{23}^H = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \otimes \delta_\beta^\alpha . \quad (3.38)$$

Therefore, CP is conserved automatically for this potential and all possible explicit CPIs necessarily vanish.

One notes that the potential of one triplet of A_4 , eqs. (3.35) and (3.37) for $SU(2)$ singlets and doublets respectively, is an extension of the potential of one triplet of $\Delta(3n^2)$ (in eqs. (3.10), (3.11) and eq. (3.14), (3.15)) by a term that is invariant only under $A_4 \times U(1)$:

$$V_{A_4} = V_0 + \delta V_{A_4 \times U(1)} . \quad (3.39)$$

The full flavour-type symmetries of the full potential V_{A_4} are generated by

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} , \begin{pmatrix} e^{i\alpha} & 0 & 0 \\ 0 & e^{i\alpha} & 0 \\ 0 & 0 & e^{i\alpha} \end{pmatrix} , \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} , \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} , \quad (3.40)$$

and in addition the potential has the CP symmetry generated by X_{23} . Under these symmetries, the elements of the orbits of the potential of one triplet of $\Delta(6n^2)$, eq. (3.23), fall into the following orbits:

$$\begin{pmatrix} e^{i\eta} \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} , \begin{pmatrix} e^{i\eta} \\ e^{i\zeta} \\ 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 \\ e^{i\zeta'} \\ 0 \end{pmatrix} \text{ with } \zeta' \in [0, \pi] \quad (3.41)$$

and

$$\begin{pmatrix} e^{i\eta} \\ e^{i\zeta} \\ e^{i\theta} \end{pmatrix} \rightarrow \begin{pmatrix} 1 \\ e^{i\zeta'} \\ e^{i\theta'} \end{pmatrix} \text{ with } \zeta'' \in [0, \pi] \text{ and } \theta' \in [0, 2\pi]. \quad (3.42)$$

So far this has been straightforward and one has learned that out of the orbits of the potential invariant under $\Delta(6n^2)$, up to two phases can become physical, which now have to be determined by minimizing the parts of the potential that depend on them, which happens to be exactly $\delta V_{A_4 \times U(1)}$. For example, plugging the alignment $(1, e^{i\zeta'}, 0)$ into $\delta V_{A_4 \times U(1)}$, one obtains

$$\delta V_{A_4 \times U(1)}[(1, e^{i\zeta'}, 0)] = ce^{2i\zeta'} + c^*e^{-2i\zeta'}, \quad (3.43)$$

which when minimizing this in ζ' of course yields $\zeta' = -\text{Arg}(c)/2 \bmod \pi$.

For the alignment candidate $(1, e^{i\zeta''}, e^{i\theta'})$ one obtains after a only slightly longer calculation for the phases which minimize $\delta V_{A_4 \times U(1)}$ and after eliminating phase constellations that are related by symmetries of the potential, eq. (3.40),

$$(\zeta'', \theta') = (0, 0), (5\pi/3, \pi/3), (2\pi/3, 4\pi/3), \quad (3.44)$$

which correspond to alignments of the type $(1, 1, 1)$ and both of $(\pm 1, \omega, \omega^2)$, such that with eq. (3.41) the list of possible global alignments from [50, 51] and [40] for one triplet of A_4 is completed. All of the alignments found above, in summary,

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} \pm 1 \\ \omega \\ \omega^2 \end{pmatrix}, \begin{pmatrix} 1 \\ e^{i\zeta'} \\ 0 \end{pmatrix} \quad (3.45)$$

have the same energy for V_0 and are just further differentiated by $\delta V_{A_4 \times U(1)}$, as is well known in the literature.

There are two ways in which one can reach the symmetries of the A_4 potential from the symmetries of the $\Delta(6n^2)$ potential, namely directly and via the symmetries of the S_4 potential.

The potential of one triplet of S_4 is actually contained in the potential of one triplet of A_4 :

$$V_{A_4} = V_0 + \delta V_{S_4 \times U(1)} + \delta V'_{A_4 \times U(1)} \quad (3.46)$$

with $\delta V_{S_4 \times U(1)}$ from eq. (3.26) (with b real and the h.c. inside the bracket) and

$$\delta V'_{A_4 \times U(1)} = (c - b) (\varphi_1 \varphi_1 \varphi^{*3} \varphi^{*3} + \varphi_2 \varphi_2 \varphi^{*1} \varphi^{*1} + \varphi_3 \varphi_3 \varphi^{*2} \varphi^{*2}) + \text{h.c.} \quad (3.47)$$

Note that while b is real, c is complex and in this expression the h.c. is outside of the bracket. Following the same path, namely writing out the orbits of the VEVs of the S_4 potential and reducing them by the full symmetries of the A_4 potential, one recovers the first three VEVs in eq. (3.45). However, there is no way of obtaining $(1, e^{i\zeta'}, 0)^T$ from the $S_4 \times U(1)$ orbit of $(1, i, 0)^T$: the full symmetry of the potential invariant under S_4 is such that the orbit of $(1, i, 0)^T$ does not contain $(1, e^{i\zeta'}, 0)$. Interestingly, this A_4 VEV is not geometric.

3.1.4 $\Delta(27)$ and $\Delta(54)$

As in the A_4 case, we start with a single triplet of SM singlets and later consider up to two $\Delta(27)$ triplets of $SU(2)_L$ doublets.

$\Delta(27)$ has irreps. **3**, its conjugate **$\bar{3}$** , and nine one-dimensional irreps. The product of triplets decomposes as

$$\mathbf{3} \otimes \mathbf{3} = (\bar{\mathbf{3}} + \bar{\mathbf{3}})_s + \bar{\mathbf{3}}_a, \quad (3.48)$$

s and a are the symmetric and antisymmetric combinations, respectively.

In the basis used in [52, 53] The renormalisable scalar potential or one triplet is ,

$$V_{\Delta(27)}(\varphi) = V_0(\varphi) + [d(\varphi_1\varphi_1\varphi^{*2}\varphi^{*3} + \text{cycl.}) + \text{h.c.}] . \quad (3.49)$$

Apart from $d \in \mathbb{C}$, the coefficients are real. Note that this potential has no automatic CP symmetry. In the literature, for the potential eq. (3.49), a variety of CP symmetries have been discussed, starting with 12 CP transformations that had been found to be consistent with $\Delta(27)$ in [54]. The enumeration of the X matrices here follows that paper. The full discrete flavour symmetry of the potential invariant under $\Delta(27)$ is $\Delta(54)$ and in this case, pairs of X matrices become related under the enlarged symmetry. In the following, the X matrices are listed by the relation they enforce on the parameters of the potential. After that, basis transformations that relate these CP symmetries, and their role are discussed.

- $\text{Arg}(d) = 0$

$$X_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{or} \quad X_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad (3.50)$$

- $\text{Arg}(d) = 4\pi/3$

$$X_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \omega \end{pmatrix} \quad \text{or} \quad X_8 = \begin{pmatrix} \omega & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad (3.51)$$

- $\text{Arg}(d) = 2\pi/3$

$$X_3 = X_2^* = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \omega^2 \end{pmatrix} \quad \text{or} \quad X_9 = X_8^* = \begin{pmatrix} \omega^2 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad (3.52)$$

- $s = (d + d^*) = 2\text{Re}(d)$

$$X_4 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \quad \text{or} \quad X_5 = X_4 X_1 = X_4^* = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{pmatrix} \quad (3.53)$$

- $s = -\text{Re}(d) - \sqrt{3}\text{Im}(d)$

$$X_6 = \frac{-i}{\sqrt{3}} \begin{pmatrix} 1 & \omega & \omega \\ \omega & \omega & 1 \\ \omega & 1 & \omega \end{pmatrix} \quad \text{or} \quad X_{10} = X_6 X_1 = \frac{-i}{\sqrt{3}} \begin{pmatrix} 1 & \omega & \omega \\ \omega & 1 & \omega \\ \omega & \omega & 1 \end{pmatrix} \quad (3.54)$$

- $s = -\text{Re}(d) + \sqrt{3}\text{Im}(d)$

$$X_7 = X_6^* = \frac{i}{\sqrt{3}} \begin{pmatrix} 1 & \omega^2 & \omega^2 \\ \omega^2 & \omega^2 & 1 \\ \omega^2 & 1 & \omega^2 \end{pmatrix} \quad \text{or} \quad X_{11} = X_7 X_1 = \frac{i}{\sqrt{3}} \begin{pmatrix} 1 & \omega^2 & \omega^2 \\ \omega^2 & 1 & \omega^2 \\ \omega^2 & \omega^2 & 1 \end{pmatrix} \quad (3.55)$$

We note that X_0 was defined already in eq. (3.22) and X_1 (matching [54]) was defined as X_{23} in eq. (3.36).

As all of the above X matrices are symmetric, there are basis transformations U_i such that in the new basis the matrix X_i becomes the identity, and for X_2 and X_3 , these also leave the potential form invariant, as was realised in [45],

$$U_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \omega^2 \end{pmatrix} \quad \text{and} \quad U_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \omega \end{pmatrix}. \quad (3.56)$$

There are of course also basis transformations from X_4, X_5, X_6 to X_0 , but these do not leave the potential form-invariant. The basis transformation e.g. from X_4 to X_0 is

$$U_4 = \begin{pmatrix} 0 & 0 & 1 \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{1+\sqrt{3}}{\sqrt{6+2\sqrt{3}}} & \frac{1}{\sqrt{3+\sqrt{3}}} \\ 0 & \frac{\sqrt{3}-1}{\sqrt{6-2\sqrt{3}}} & \frac{1}{\sqrt{3-\sqrt{3}}} \end{pmatrix} \begin{pmatrix} -(-1)^{3/4} & 0 & 0 \\ 0 & -i & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (3.57)$$

For $SU(2)_L$ doublets, the potential is

$$V_{\Delta(27)}(H) = V_0(H) + \sum_{\alpha, \beta} \left[d \left(h_{1\alpha} h_{1\beta} h^{*2\alpha} h^{*3\beta} + \text{cycl.} \right) + \text{h.c.} \right]. \quad (3.58)$$

For one triplet, the corresponding potentials invariant under are $\Delta(54)$ identical to those invariant under $\Delta(27)$ [55]

$$V_{\Delta(54)}(\varphi) = V_{\Delta(27)}(\varphi), \quad (3.59)$$

$$V_{\Delta(54)}(H) = V_{\Delta(27)}(H). \quad (3.60)$$

The potential eq. (3.58) has been analysed in [40] and is the 3HDM potential with the largest of the purely discrete symmetries (apart for the $U(1)$ that arises from the potential being even) for which CP may be spontaneously violated. In addition, this was the case where so-called geometric violation was first discovered, see also section 4.1.3.

For generic values of the potential parameters, the potential violates CP explicitly. Additional CP symmetries may be imposed that force the potential to be explicitly CP-conserving. In [40] the two types of CP symmetry that are normally considered consistent with the flavour-type symmetry of the potential are analysed. In [54], 12 CP symmetries are listed as being consistent with $\Delta(27)$, but if considered in the $\Delta(54)$ context this number reduces to 6. For example, CP_0 and the CP_{23} (the CP associated with X_{23}) become related by $\Delta(54)$. In this case, of the 6 remaining, 3 restrict the phase of parameter d in the

potential, the remaining 3 additionally enforces an additional relation on the parameters, such as $2s = (d + d^*)$ if CP with X matrix

$$X_4 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}, \quad (3.61)$$

is imposed. This second CP-type transformation (like X_4) enlarges the flavour-type symmetry of the potential to $\Sigma(36)$. It should be noted that the three CP symmetries that restrict the phase of d are related by basis-transformations that leave the potential form-invariant, cf. [45]. In addition, we found that the three CP symmetries that enforce $2s = d + d^*$ are also related with each other via such basis transformations, the simplest ones being for connecting X_4 with X_{10} and X_{11} , respectively,

$$U_{10} = \sqrt{-i} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega \end{pmatrix} \text{ and } U_{11} = \sqrt{-i} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}. \quad (3.62)$$

Furthermore, as X_4 of eq. (3.61) is also symmetric, in principle a basis transformation exists, such that in the new basis X_4 would be diagonal, however this basis transformation does not leave the potential form-invariant.

The full flavour-type symmetries of an even potential of one triplet of $\Delta(27)$ or one triplet of $\Delta(54)$, both produce identical renormalisable potentials, are generated by

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} e^{i\alpha} & 0 & 0 \\ 0 & e^{i\alpha} & 0 \\ 0 & 0 & e^{i\alpha} \end{pmatrix}, \begin{pmatrix} e^{i2\pi/3} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{-i2\pi/3} \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i2\pi/3} & 0 \\ 0 & 0 & e^{-i2\pi/3} \end{pmatrix}. \quad (3.63)$$

Using these, the orbits of one triplet of $\Delta(6n^2)$, eq. (3.23) become

$$(1, 0, 0), (1, e^{i\beta'}, 0), (1, e^{i\beta''}, e^{i\gamma'}) \quad (3.64)$$

with $\beta', \beta'', \gamma' \in [0, 2\pi/3]$. The CP symmetries, whether CP_0 or the CP symmetry associated with X_4 do at this point not further constrain the potentially physical phases. The phases appearing here can now be physical within $\Delta(54)$ and need to be determined by minimizing the parts of the potential that depend on them. Curiously, the phase dependent part of $V_{\Delta(54)}$,

$$\delta V_{\Delta(54) \times U(1)} = [d(\varphi_1 \varphi_1 \varphi^{*2} \varphi^{*3} + \text{cycl.}) + \text{h.c.}] \quad (3.65)$$

yields simply zero for the alignment $(1, e^{i\beta'}, 0)$, which already means that the phase β' remains unphysical. $(1, 1, 0)$ could still be a local minimum of the potential, as it already is a possible global minimum of V_0 . When minimizing $\delta V_{\Delta(54) \times U(1)}$ with $(1, e^{i\beta''}, e^{i\gamma'})$, one obtains the usual alignments of $(1, 1, 1)$, $(1, 1, \omega)$, $(1, \omega, \omega)$ (or for the last one equivalently $(1, 1, \omega^2)$). Evaluating the potential at those alignments explains the fate of the $(1, 1, 0)$ alignment as a global minimum: $\delta V_{\Delta(54) \times U(1)}$ always attains a negative value for at least one of the alignments $(1, 1, 1)$, $(1, 1, \omega)$, $(1, 1, \omega^2)$, for any value of $\text{Arg}(d)$. However, as

this scales with the absolute size of d as well, $(1, 1, 0)$ could still be quite a metastable minimum. In any case, again, taking VEV candidates just from the VEV orbits of the $\Delta(6n^2)$ potential produces the full list of global minima that were obtained (painstakingly) by [40], in summary:

$$(1, 0, 0), (1, 1, 1), (1, 1, \omega), (1, 1, \omega^2). \quad (3.66)$$

To also repeat the effect of CP transformations from [40]: with CP_0 , the last two VEVs become related by symmetry and no longer represent different breaking patterns. With the type of CP that extends the flavour symmetry to $\Sigma(36)$, with matrix X_4 , eq. (3.61), again the last two VEVs in eq. (3.66) become part of the same orbit. In addition also the first two VEVs in eq. (3.66) become part of the same orbit (separate from the last two VEVs).

3.2 VEVs of potentials of two triplets

Typically, realistic models of flavour require more than just one triplet flavon. We therefore consider potentials involving two physically different flavon fields φ and φ' which both transform under a triplet representation of the symmetry groups we consider. Similarly we also consider 6HDMs with the $SU(2)_L$ doublets transforming as two triplets of the discrete symmetries.

Potentials of two triplets consist of three parts: two sets of terms that each only couple components of one triplet to each other on one hand, and cross terms that couple components of different triplets to each other,

$$V(\varphi, \varphi') = V(\varphi) + V'(\varphi') + V_c(\varphi, \varphi'). \quad (3.67)$$

If the two triplets transform identically under the symmetry, then $V(\varphi)$ and $V'(\varphi')$ will be functionally identical.

The orbits of minima of the potentials of single triplets are known completely for the flavour symmetries in consideration. Similarly to potentials of one triplet, one can now analyse which degrees of freedom of VEVs that are unphysical by the symmetry of the two single-triplet potentials $V(\varphi) + V'(\varphi')$ can become physical by reducing the symmetry either by introducing cross-terms between triplets, $V_c(\varphi, \varphi')$, or by considering a whole potential invariant only under a smaller symmetry group.

As before, only the alignments of the VEVs are shown and for two triplets, this does not mean that the VEVs v_1, v_2 of both triplets have to have the same length. Instead, arbitrary lengths of both triplets are allowed and possible and generally $|v_1| \neq |v_2|$.

In the following subsections, we will consider potentials of two triplets, starting with those invariant under $\Delta(3n^2)$ and $\Delta(6n^2)$ with $n > 3$ and then continuing to those with smaller symmetries, all of which share certain common terms (this was already the case for the potentials with a single triplet invariant under these groups). For this reason, it is

useful to define for the $\Delta(3n^2)$ symmetries

$$\begin{aligned}
V_1(\varphi, \varphi') = & +\tilde{r}_1 \left(\sum_i \varphi_i \varphi^{*i} \right) \left(\sum_j \varphi'_j \varphi'^{*j} \right) + \tilde{r}_2 \left(\sum_i \varphi_i \varphi'^{*i} \right) \left(\sum_j \varphi'_j \varphi^{*j} \right) \\
& + \tilde{s}_1 \sum_i (\varphi_i \varphi^{*i} \varphi'_i \varphi'^{*i}) \\
& + \tilde{s}_2 (\varphi_1 \varphi^{*1} \varphi'_2 \varphi'^{*2} + \varphi_2 \varphi^{*2} \varphi'_3 \varphi'^{*3} + \varphi_3 \varphi^{*3} \varphi'_1 \varphi'^{*1}) \\
& + i \tilde{s}_3 [(\varphi_1 \varphi'^{*1} \varphi'_2 \varphi^{*2} + \text{cycl.}) - (\varphi^{*1} \varphi'_1 \varphi'^{*2} \varphi_2 + \text{cycl.})]. \tag{3.68}
\end{aligned}$$

Note that in this definition, the term multiplied by \tilde{r}_1 contains the term multiplied by \tilde{s}_2 as well as the term obtained from the latter by interchanging φ with φ' ,

$$(\varphi'_1 \varphi'^{*1} \varphi_2 \varphi^{*2} + \varphi'_2 \varphi'^{*2} \varphi_3 \varphi^{*3} + \varphi'_3 \varphi'^{*3} \varphi_1 \varphi^{*1}), \tag{3.69}$$

which is not included separately in \tilde{s}_2 .

Earlier, when considering a potential of one triplet of $SU(2)_L$ doublets, the only difference was that the term with coefficient r split into two different invariants corresponding to two different possible $SU(2)_L$ contractions, cf. eq. (3.13). Similarly, the potential of two triplets of SM doublets,

$$H = (h_{1\alpha}, h_{2\beta}, h_{3\gamma}), \quad H' = (h'_{1\alpha}, h'_{2\beta}, h'_{3\gamma}), \tag{3.70}$$

can be obtained from the corresponding potential of singlets, eq. (3.68). In the first two parts of the potential, $V_0(\varphi)$ and $V_0(\varphi')$, as earlier, there are two different ways of $SU(2)_L$ -contracting the invariants with coefficients r and r' . In the potential $V_1(\varphi, \varphi')$, there are two possible ways of $SU(2)_L$ -contracting for each invariant and therefore this part of the potential becomes in its $SU(2)_L$ doublet version

$$\begin{aligned}
V_1(H, H') = & \sum_{i,j,\alpha,\beta} \left[\tilde{r}_{11} h_{i\alpha} h'^{*i\alpha} h'_{j\beta} h'^{*j\beta} + \tilde{r}_{12} h_{i\alpha} h'^{*j\alpha} h'_{j\beta} h^{*i\beta} \right] \\
& + \sum_{i,j,\alpha,\beta} \left[\tilde{r}_{21} h_{i\alpha} h'^{*i\alpha} h'_{j\beta} h^{*j\beta} + \tilde{r}_{22} h_{i\alpha} h^{*j\alpha} h'_{j\beta} h'^{*i\beta} \right] \\
& + \sum_{i,\alpha,\beta} \left[\tilde{s}_{11} h_{i\alpha} h^{*i\alpha} h'_{i\beta} h'^{*i\beta} + \tilde{s}_{12} h_{i\alpha} h'^{*i\alpha} h'_{i\beta} h^{*i\beta} \right] \\
& + \sum_{\alpha,\beta} \left[\tilde{s}_{21} (h_{1\alpha} h^{*1\alpha} h'_{2\beta} h'^{*2\beta} + \text{cycl.}) + \tilde{s}_{22} (h_{1\alpha} h'^{*2\alpha} h'_{2\beta} h^{*1\beta} + \text{cycl.}) \right] \\
& + i \tilde{s}_{31} \sum_{\alpha,\beta} [(h_{1\alpha} h'^{*1\alpha} h'_{2\beta} h^{*2\beta} + \text{cycl.}) - (h^{*1\alpha} h'_{1\alpha} h'^{*2\beta} h_{2\beta} + \text{cycl.})] \\
& + i \tilde{s}_{32} \sum_{\alpha,\beta} [(h_{1\alpha} h^{*2\alpha} h'_{2\beta} h'^{*1\beta} + \text{cycl.}) - (h^{*1\alpha} h_{2\alpha} h'^{*2\beta} h'_{1\beta} + \text{cycl.})]. \tag{3.71}
\end{aligned}$$

As it is useful for all of the $\Delta(6n^2)$ symmetries, we define also

$$\begin{aligned}
V_2(\varphi, \varphi') = & \tilde{r}_1 \left(\sum_i \varphi_i \varphi^{*i} \right) \left(\sum_j \varphi'_j \varphi'^{*j} \right) + \tilde{r}_2 \left(\sum_i \varphi_i \varphi'^{*i} \right) \left(\sum_j \varphi'_j \varphi^{*j} \right) \\
& + \tilde{s}_1 \sum_i (\varphi_i \varphi^{*i} \varphi'_i \varphi'^{*i}), \tag{3.72}
\end{aligned}$$

and the $SU(2)_L$ doublet version

$$\begin{aligned} V_2(H, H') = & \sum_{i,j,\alpha,\beta} \left[\tilde{r}_{11} h_{i\alpha} h^{*i\alpha} h'_{j\beta} h^{*j\beta} + \tilde{r}_{12} h_{i\alpha} h'^{*j\alpha} h'_{j\beta} h^{*i\beta} \right] \\ & + \sum_{i,j,\alpha,\beta} \left[\tilde{r}_{21} h_{i\alpha} h'^{*i\alpha} h'_{j\beta} h^{*j\beta} + \tilde{r}_{22} h_{i\alpha} h^{*j\alpha} h'_{j\beta} h'^{*i\beta} \right] \\ & + \sum_{i,\alpha,\beta} \left[\tilde{s}_{11} h_{i\alpha} h^{*i\alpha} h'_{i\beta} h'^{*i\beta} + \tilde{s}_{12} h_{i\alpha} h'^{*i\alpha} h'_{i\beta} h^{*i\beta} \right]. \end{aligned} \quad (3.73)$$

We note that one can obtain V_2 from V_1 by imposing

$$\tilde{s}_2 = \tilde{s}_3 = 0, \quad (3.74)$$

for $SU(2)_L$ singlets and, for $SU(2)_L$ doublets, $s_{22} = \tilde{s}_{31} = \tilde{s}_{32} = 0$.

As previously for one-triplet-potentials, minima of several doublets that conserve $U(1)_{em}$ can be identified with the minima of the same number of singlets.

By restricting the potential with $U(1)$ symmetries for each of the scalar fields, the mixed terms appearing are limited to the form $\varphi \varphi' \varphi^* \varphi'^*$. For $SU(2)_L$ doublets, it is even sufficient to impose e.g. a Z_3 symmetry with non-trivial charge for only one of the two triplets of Higgs doublets to distinguish them.

3.2.1 $\Delta(6n^2)$ with $n > 3$

The potentials of two triplets under $\Delta(6n^2)$ of singlets and doublets have the form

$$V_{\Delta(6n^2)}(\varphi, \varphi') = V_0(\varphi) + V'_0(\varphi') + V_2(\varphi, \varphi'), \quad (3.75)$$

$$V_{\Delta(6n^2)}(H, H') = V_0(H) + V'_0(H') + V_2(H, H'), \quad (3.76)$$

where $V'_0(\varphi')$ has the same functional form as $V_0(\varphi)$ with different coefficients $m'_{\varphi'}$, r' , s' and depends on φ' . $V_0(\varphi)$, $V_0(H)$ were defined in eqs. (3.11) and (3.15), $V_2(\varphi, \varphi')$ and $V_2(H, H')$ were defined in eqs. (3.72) and (3.73). These $\Delta(6n^2)$ potentials (with $n > 3$) conserve CP explicitly. As the parameters are all real, one of the CP symmetries is CP_0 .

The orbits of minima of a potential of one triplet of $\Delta(6n^2)$ are (cf. eq. (3.23))

$$\left\{ \begin{pmatrix} e^{i\eta} \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ e^{i\eta} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ e^{i\eta} \end{pmatrix} \right\}, \left\{ \begin{pmatrix} e^{i\eta} \\ e^{i\zeta} \\ 0 \end{pmatrix}, \text{permut.} \right\}, \left\{ \begin{pmatrix} e^{i\eta} \\ e^{i\zeta} \\ e^{i\theta} \end{pmatrix}, \text{permut.} \right\}. \quad (3.77)$$

Without cross terms $V_2(\varphi, \varphi')$, both triplets can transform independently under $\Delta(6n^2)$ and thus have independent orbits. Thus, every member of each orbit may now be combined with every other member of each orbit, but only the symmetries of the full potential of two triplets including cross terms can be used to eliminate unphysical phases. The symmetries of the full potential are generated by the simultaneous transformations of both triplets under $\Delta(6\infty^2)$ and in addition separate $U(1)$ phases acting on each triplet,

$$\begin{pmatrix} e^{i\alpha} & & \\ & e^{i\alpha} & \\ & & e^{i\alpha} \end{pmatrix} \oplus \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix}, \text{ and } \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix} \oplus \begin{pmatrix} e^{i\alpha'} & & \\ & e^{i\alpha'} & \\ & & e^{i\alpha'} \end{pmatrix}, \quad (3.78)$$

as well as an overall canonical CP transformation (CP_0). With these symmetries one obtains the following reduction of combinations of orbits of VEVs of a potential of one triplet of $\Delta(6n^2)$:

$$(e^{i\eta}, 0, 0)|(e^{i\eta'}, 0, 0) \rightarrow (1, 0, 0)|(1, 0, 0) \quad (3.79)$$

$$(e^{i\eta}, 0, 0)|(0, e^{i\eta'}, 0) \rightarrow (1, 0, 0)|(0, 1, 0) \quad (3.80)$$

$$(e^{i\eta}, 0, 0)|(e^{i\eta'}, e^{i\zeta'}, 0) \rightarrow (1, 0, 0)|(1, 1, 0) \quad (3.81)$$

$$(e^{i\eta}, 0, 0)|(0, e^{i\eta'}, e^{i\zeta'}) \rightarrow (1, 0, 0)|(0, 1, 1) \quad (3.82)$$

$$(e^{i\eta}, 0, 0)|(e^{i\eta'}, e^{i\zeta'}, e^{i\theta'}) \rightarrow (1, 0, 0)|(1, 1, 1) \quad (3.83)$$

$$(e^{i\eta}, e^{i\zeta}, 0)|(0, e^{i\zeta'}, e^{i\theta'}) \rightarrow (1, 1, 0)|(0, 1, 1) \quad (3.84)$$

$$(e^{i\eta}, e^{i\zeta}, 0)|(e^{i\eta'}, e^{i\zeta'}, 0) \rightarrow (1, 1, 0)|(1, e^{i\zeta'}, 0) \quad (3.85)$$

$$(e^{i\eta}, e^{i\zeta}, 0)|(e^{i\eta'}, e^{i\zeta'}, e^{i\theta'}) \rightarrow (1, 1, 0)|(1, e^{i\zeta'}, 1) \quad (3.86)$$

$$(e^{i\eta}, e^{i\zeta}, e^{i\theta})|(e^{i\eta'}, e^{i\zeta'}, e^{i\theta'}) \rightarrow (1, 1, 1)|(1, e^{i\zeta'}, e^{i\theta'}). \quad (3.87)$$

Note that at this stage orbit pairs that arise from interchanging first and second triplet are redundant, which is why e.g. $(1, 1, 0)|(1, 0, 1)$ is not listed. The remaining phases are determined by minimizing the parts of the potential that depend on them and one obtains for both of $(1, 1, 0)|(1, e^{i\zeta'}, 0)$ and $(1, 1, 0)|(1, e^{i\zeta'}, 1)$ that $\zeta' = 0$ for $r'_2 > 0$ and $\zeta' = \pi$ for $r_2 < 0$, leading to the following alignments:

$$(1, 1, 0)|(1, \pm 1, 0) \text{ and } (1, 1, 0)|(1, \pm 1, 1) \quad (3.88)$$

where different sign choices correspond to different orbits. For $(1, 1, 1)|(1, e^{i\zeta'}, e^{i\theta'})$ minimizing the part of the potential depending on ζ' and θ' produces for $r_2 < 1$ the orbit

$$(1, 1, 1)|(1, 1, 1) \quad (3.89)$$

and for $r_2 > 0$ the two orbits

$$(1, 1, 1)|(1, \omega, \omega^2) \text{ and } (1, 1, 1)|(1, \omega^2, \omega). \quad (3.90)$$

To summarize all possible pairs we list them in table 1.

3.2.2 $\Delta(3n^2)$ with $n > 3$

Next, consider the potential of two triplets of $\Delta(3n^2)$ with $n > 3$,

$$V_{\Delta(3n^2)}(\varphi, \varphi') = V_0(\varphi) + V'_0(\varphi') + V_1(\varphi, \varphi'), \quad (3.91)$$

with $V_0(\varphi)$ and $V_1(\varphi, \varphi')$ defined in eqs. (3.11) and (3.68) for singlets and

$$V_{\Delta(3n^2)}(H, H') = V_0(H) + V'_0(H') + V_1(H, H') \quad (3.92)$$

for doublets, where $V_0(H)$ and $V_1(H, H')$ were defined in eqs. (3.15) and (3.71).

$(1, 0, 0) (1, 0, 0)$
$(1, 0, 0) (0, 1, 0)$
$(1, 0, 0) (1, 1, 0)$
$(1, 0, 0) (0, 1, 1)$
$(1, 0, 0) (1, 1, 1)$
$(1, 1, 0) (0, 1, 1)$
$(1, 1, 0) (1, \pm 1, 0)$
$(1, 1, 0) (1, \pm 1, 1)$
$(1, 1, 1) (1, 1, 1)$
$(1, 1, 1) (1, \omega, \omega^2)$
$(1, 1, 1) (1, \omega^2, \omega)$

Table 1. List of VEV directions for the potential of two triplets of $\Delta(6n^2)$, $V_{\Delta(6n^2)}(\varphi, \varphi')$.

The orbits of VEVs of the corresponding potential of one triplet are identical to that of one triplet of $\Delta(6n^2)$, cf. eq. (3.77). The difference to the previous potential lies in the fact that the full symmetries of $V_{\Delta(3n^2)}$ only allow for cyclic permutations, i.e. only

$$\begin{pmatrix} 1 & & \\ & 1 & \\ 1 & & \end{pmatrix} \oplus \begin{pmatrix} 1 & \\ & 1 \\ 1 & \end{pmatrix} \quad (3.93)$$

with otherwise identical phase symmetries, i.e. all simultaneous phase symmetries arising from $\Delta(3n^2)$, and eq. (3.78).

This potential has no automatic CP symmetry. Possible CP symmetries are overall CP_0

$$X_0^{\varphi\varphi'} = \begin{pmatrix} X_0 & 0 \\ 0 & X_0 \end{pmatrix}, \quad \text{where} \quad X_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (3.94)$$

and simultaneous CP_{23} , i.e. a CP symmetry acting on both triplets with

$$X_{23}^{\varphi\varphi'} = \begin{pmatrix} X_{23} & 0 \\ 0 & X_{23} \end{pmatrix}, \quad \text{where} \quad X_{23} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \quad (3.95)$$

Besides overall CP_0 and CP_{23} , we will also consider a CP symmetry, $CP_{\varphi\varphi'}$, with

$$X_{\varphi\varphi'} = \begin{pmatrix} X_\varphi & 0 \\ 0 & X_{\varphi'} \end{pmatrix}, \quad \text{with} \quad X_\varphi = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \quad X_{\varphi'} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix}, \quad (3.96)$$

which was found in [35] for this potential.⁶ Under CP_0 , all coefficients of the potential become real such that $\tilde{s}_3 = 0$. For CP_{23} , $\tilde{s}_2 = 0$. The CP symmetry with $X_{\varphi\varphi'}$, eq. (3.96), relates parameters of the potential via

$$\tilde{s}_3 = \tilde{r}_2 i(\omega - \omega^2). \quad (3.97)$$

As $\omega = e^{2\pi i/3}$, we get $\tilde{s}_3 = -\sqrt{3}\tilde{r}_2$. The CP transformation where the roles of the explicit matrices in eq. (3.96) are exchanged enforces $\tilde{s}_3 = \sqrt{3}\tilde{r}_2$. The effect of the CP symmetries on the VEV orbits will be discussed at the end of this subsection.

Again, arbitrary members of the one-triplet orbits can be combined to pairs which are combined into orbits under the symmetry of the full potential. Compared to the potential of two triplets of $\Delta(6n^2)$, due to the missing permutation generator in $\Delta(3n^2)$, several orbits split. However, again at this stage it does not make difference to interchange the first and second triplet. For this reason apart from eqs. (3.79)–(3.87), only one new combination survives:

$$(e^{i\eta}, 0, 0) | (e^{i\eta'}, 0, e^{i\zeta'}) \rightarrow (1, 0, 0) | (1, 0, 1). \quad (3.98)$$

Furthermore, the orbits that contained continuous degrees of freedom now need to be minimized for the more complicated potential invariant under $\Delta(3n^2)$. This is still analytically possible and one obtains

$$(1, 1, 0) | (1, e^{i\zeta'}, 0) \text{ and} \quad (3.99)$$

$$(1, 1, 0) | (1, e^{i\zeta'}, 1) \quad (3.100)$$

with

$$\zeta' = \arctan(\tilde{r}_2/\tilde{s}_3) \quad (3.101)$$

a function of \tilde{s}_3 and \tilde{r}_2 in contrast to the situation with a $\Delta(6n^2)$ symmetry, where $\zeta' = 0, \pi$, depending on the value of \tilde{r}_2 . The orbit with two phases again results in

$$(1, 1, 1) | (1, 1, 1), \quad (3.102)$$

$$(1, 1, 1) | (1, \omega, \omega^2), \quad (3.103)$$

$$(1, 1, 1) | (1, \omega^2, \omega), \quad (3.104)$$

depending on the values of \tilde{r}_2 and \tilde{s}_3 .

To summarize all possible pairs before applying CP symmetries we list them in table 2.

Next, consider the effects of the various CP symmetries on the orbits. For CP_0 , nothing happens to the orbits with real representatives. $\tilde{s}_3 = 0$, so ζ' becomes $\pm\pi/2$ and thus $\zeta' = \pm i$. Furthermore, the orbits $(1, 1, 1) | (1, \omega, \omega^2)$ and $(1, 1, 1) | (1, \omega^2, \omega)$ merge. For

⁶This CP symmetry is not consistent with the $\Delta(3n^2)$ part of the flavour symmetry acting on both triplets simultaneously. However, under the action of the two independent $U(1)$ s of each triplet, any different overall phases appearing in the consistency condition for each triplet can be absorbed, which makes this CP transformation consistent with the full flavour symmetry of the potential. The same is true for the related CP transformation where X_φ and $X_{\varphi'}$ have been interchanged.

(1, 0, 0) (1, 0, 0)
(1, 0, 0) (0, 1, 0)
(1, 0, 0) (1, 1, 0)
(1, 0, 0) (1, 0, 1)
(1, 0, 0) (0, 1, 1)
(1, 0, 0) (1, 1, 1)
(1, 1, 0) (0, 1, 1)
(1, 1, 0) (1, $e^{i\zeta'}$, 0)
(1, 1, 0) (1, $e^{i\zeta'}$, 1)
(1, 1, 1) (1, 1, 1)
(1, 1, 1) (1, ω, ω^2)
(1, 1, 1) (1, ω^2, ω)

Table 2. List of VEV directions for the potential of two triplets of $\Delta(3n^2)$, $V_{\Delta(3n^2)}(\varphi, \varphi')$.

CP_{23} , $(1, 0, 0)|(1, 0, 1)$ and $(1, 0, 0)|(1, 1, 0)$ merge. ζ' remains free and nothing happens to the orbits with ω s. Lastly, for the CP symmetry with $X^{\varphi\varphi'}$, $\tilde{s}_3 = \sqrt{3}\tilde{r}_2$, thus $\zeta' = \pm\pi/6$. Additionally, $(1, 1, 1)|(1, \omega, \omega^2)$ and $(1, 1, 1)|(1, 1, 1)$ merge. For the CP symmetry similar to that with $X^{\varphi\varphi'}$ but with block matrices interchanged, $(1, 1, 1)|(1, \omega^2, \omega)$ merges with $(1, 1, 1)|(1, 1, 1)$.

3.2.3 S_4

The S_4 potential for one triplet was

$$V_{S_4}(\varphi) = V_0(\varphi) + b \left[(\varphi_1 \varphi_1 \varphi^{*3} \varphi^{*3} + \text{cycl.}) + \text{h.c.} \right], \quad (3.105)$$

with real b . For two triplets we use eq. (3.72), and write

$$\begin{aligned} V_{S_4}(\varphi, \varphi') &= V_0(\varphi) + V'_0(\varphi') + V_2(\varphi, \varphi') \\ &+ b \left[(\varphi_1 \varphi_1 \varphi^{*3} \varphi^{*3} + \text{cycl.}) + \text{h.c.} \right] + b' \left[(\varphi'_1 \varphi'_1 \varphi'^{*3} \varphi'^{*3} + \text{cycl.}) + \text{h.c.} \right] \\ &+ \tilde{b} \left[(\varphi_1 \varphi'_1 \varphi^{*3} \varphi'^{*3} + \text{cycl.}) + \text{h.c.} \right]. \end{aligned} \quad (3.106)$$

For $SU(2)$ doublets, we use eq. (3.73) and write

$$\begin{aligned} V_{S_4}(H, H') &= V_0(H) + V'_0(H') + V_2(H, H') \\ &+ \sum_{\alpha, \beta} b \left[\left(h_{1\alpha} h_{1\beta} h^{*3\alpha} h^{*3\beta} + \text{cycl.} \right) + \text{h.c.} \right] \\ &+ \sum_{\alpha, \beta} b' \left[\left(h'_{1\alpha} h'_{1\beta} h'^{*3\alpha} h'^{*3\beta} + \text{cycl.} \right) + \text{h.c.} \right] \\ &+ \sum_{\alpha, \beta} \tilde{b}_1 \left[\left(h_{1\alpha} h^{*3\alpha} h'_{1\beta} h'^{*3\beta} + \text{cycl.} \right) + \text{h.c.} \right] \\ &+ \sum_{\alpha, \beta} \tilde{b}_2 \left[\left(h_{1\alpha} h'^{*3\alpha} h'_{1\beta} h^{*3\beta} + \text{cycl.} \right) + \text{h.c.} \right]. \end{aligned} \quad (3.107)$$

$(1, 0, 0) (1, 0, 0)$
$(1, 0, 0) (0, 1, 0)$
$(1, 0, 0) (1, 0, i)$
$(1, 0, 0) (0, 1, i)$
$(1, 0, 0) (1, 1, 1)$
$(1, 0, 0) (1, \omega^2, \omega)$
$(1, 0, i) (1, 0, \pm i)$
$(1, 0, i) (1, i, 0)$
$(1, 0, i) (1, 1, 1)$
$(1, 0, i) (1, \omega^2, \pm \omega)$
$(1, 1, 1) (1, 1, \pm 1)$
$(1, 1, 1) (1, \pm \omega^2, \omega)$
$(1, \omega^2, \omega) (1, \omega^2, \pm \omega)$
$(1, \omega^2, \omega) (1, -\omega, -\omega^2)$
$(1, \omega^2, \omega) (1, \omega, \omega^2)$

Table 3. List of VEV directions for the potential of two triplets of S_4 , $V_{S_4}(\varphi, \varphi')$.

All S_4 -invariant potentials listed here, even the two triplet cases, conserve CP explicitly: the parameters, including $b, b', \tilde{b}, \tilde{b}_1, \tilde{b}_2$ are all real, so one of the CP symmetries is CP_0 .

For one triplet of S_4 , the full symmetries of an even potential are generated by eq. (3.29), and notably, this potential has an automatic CP symmetry. The orbits are (and again plus permutations)

$$\left\{ \begin{pmatrix} \pm e^{i\alpha} \\ 0 \\ 0 \end{pmatrix} \right\}, \left\{ \begin{pmatrix} (-1)^k e^{i\alpha} \\ (-1)^l e^{i\alpha} \\ (-1)^{k+l} e^{i\alpha} \end{pmatrix} \right\}, \left\{ \begin{pmatrix} (-1)^k e^{i\alpha} \\ \omega(-1)^l e^{i\alpha} \\ \omega^2(-1)^{k+l} e^{i\alpha} \end{pmatrix} \right\}, \left\{ \begin{pmatrix} -(-1)^k e^{i\alpha} \\ \omega(-1)^l e^{i\alpha} \\ \omega^2(-1)^{k+l} e^{i\alpha} \end{pmatrix} \right\}, \left\{ \begin{pmatrix} 0 \\ \pm e^{i\alpha} \\ \pm ie^{i\alpha} \end{pmatrix} \right\}, \quad (3.108)$$

where the last set stands for separate orbits for each combination of signs. These combine to the VEVs pairs listed in table 3, where the sign choices correspond to separate orbits (again only orbit representatives).

3.2.4 A_4

In the case of two A_4 triplets distinguished by additional symmetries so that the total symmetry is $A_4 \times U(1) \times U(1)'$, the potential includes a total of seven independent mixed quartic invariants of the form $\varphi \varphi' \varphi^* \varphi'^*$.

The A_4 symmetric renormalisable potential takes the following explicit form, with V_0 as defined in eq. (3.11), and using eq. (3.68) we write

$$\begin{aligned} V_{A_4}(\varphi, \varphi') = & V_0(\varphi) + V'_0(\varphi') + V_1(\varphi, \varphi') + \\ & + [c(\varphi_1 \varphi_1 \varphi^{*3} \varphi'^{*3} + \text{cycl.}) + \text{h.c.}] + [c'(\varphi'_1 \varphi'_1 \varphi'^{*3} \varphi'^{*3} + \text{cycl.}) + \text{h.c.}] \\ & + [\tilde{c}(\varphi_1 \varphi'_1 \varphi^{*3} \varphi'^{*3} + \text{cycl.}) + \text{h.c.}]. \end{aligned} \quad (3.109)$$

The potential is explicitly CP violating [35] but it is interesting to consider the CP symmetry where one imposes X_{23} of eq. (3.36) on both triplets, i.e. the block matrix $X_{23}^{\varphi\varphi'}$ in eq. (3.95). This CP symmetry constrains the potential such that $\tilde{s}_2 = 0$, which forces all explicit CPIs to vanish as expected from the presence of a CP symmetry [35].

Furthermore, applying instead the trivial CP symmetry CP_0 forces $\tilde{s}_3 = 0$ and all complex parameters (c, c', \tilde{c}) to be real.

For the $SU(2)_L$ version we use eq. (3.71) and, for the remainder of the potential A_4 potential with two triplets, only the invariant with coefficient \tilde{c} from eq. (3.109) needs to be doubled:

$$\sum_{\alpha, \beta} \left[\tilde{c}_1 (h_{1\alpha} h^{*3\alpha} h'_{1\beta} h'^{*3\beta} + \text{cycl.}) + \tilde{c}_2 (h_{1\alpha} h'^{*3\alpha} h'_{1\beta} h^{*3\beta} + \text{cycl.}) + \text{h.c.} \right]. \quad (3.110)$$

We therefore write

$$\begin{aligned} V_{A_4}(H, H') = & V_0(H) + V'_0(H') + V_1(H, H') \\ & + \sum_{\alpha, \beta} \left[c \left(h_{1\alpha} h_{1\beta} h^{*3\alpha} h^{*3\beta} + \text{cycl.} \right) + c' \left(h'_{1\alpha} h'_{1\beta} h'^{*3\alpha} h'^{*3\beta} + \text{cycl.} \right) + \text{h.c.} \right] \\ & + \sum_{\alpha, \beta} \left[\tilde{c}_1 (h_{1\alpha} h^{*3\alpha} h'_{1\beta} h'^{*3\beta} + \text{cycl.}) + \tilde{c}_2 (h_{1\alpha} h'^{*3\alpha} h'_{1\beta} h^{*3\beta} + \text{cycl.}) + \text{h.c.} \right]. \end{aligned} \quad (3.111)$$

We note that due to $SU(2)_L$ not allowing cubic invariants of H and/or H' , it is sufficient to use a Z_3 symmetry to distinguish the A_4 triplets.⁷

It is possible to impose a CP symmetry with

$$X_{23}^{HH'} = \begin{pmatrix} X_{23} & 0 \\ 0 & X_{23} \end{pmatrix} \otimes \delta_\beta^\alpha, \quad (3.112)$$

which, similarly to previous examples, restricts the coefficients in the potential, namely

$$\tilde{s}_{21} = \tilde{s}_{22} = 0. \quad (3.113)$$

Imposing, alternatively, the canonical CP symmetry CP_0 leads to $\tilde{s}_{31} = \tilde{s}_{32} = 0$ as well as $c, c', \tilde{c}_1, \tilde{c}_2 \in \mathbb{R}$.

The full symmetries of an even potential acting on one triplet are generated by eq. (3.40),

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} e^{i\alpha} & 0 & 0 \\ 0 & e^{i\alpha} & 0 \\ 0 & 0 & e^{i\alpha} \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad (3.114)$$

and the orbits of one-triplet VEVs are, again with permutations, where now only cyclic permutations are allowed and no longer, as for S_4 , all possible permutations, and in addition

⁷The potential invariant under a Z_2 [56] would additionally allow for invariants of the form $h_{i\alpha} h'^{*i\alpha} h_{j\beta} h'^{*j\beta}$ and $h_{i\alpha} h'^{*i\beta} h_{j\beta} h'^{*j\alpha}$ where the conjugated fields are both related to H' .

$(1, 0, 0) (1, 0, 0)$
$(1, 0, 0) (0, 1, 0)$
$(1, 0, 0) (1, e^{i\zeta'}, 0)$
$(1, 0, 0) (0, 1, e^{i\zeta'})$
$(1, 0, 0) (e^{i\zeta'}, 0, 1)$
$(1, 0, 0) (1, 1, 1)$
$(1, 0, 0) (1, \omega, \omega^2)$
$(1, e^{i\zeta}, 0) (1, \pm e^{i\zeta'}, 0)$
$(1, e^{i\zeta}, 0) (0, 1, e^{i\zeta'})$
$(1, e^{i\zeta}, 0) (e^{i\zeta'}, 0, 1)$
$(1, e^{i\zeta}, 0) (1, \pm 1, 1)$
$(1, e^{i\zeta}, 0) (1, \pm \omega, \omega^2)$
$(1, 1, 1) (1, 1, \pm 1)$
$(1, 1, 1) (1, \omega, \pm \omega^2)$
$(1, \omega, \omega^2) (1, \omega, \pm \omega^2)$

Table 4. List of VEV directions for the potential of two triplets of A_4 , $V_{A_4}(\varphi, \varphi')$.

β is an arbitrary phase,

$$\left\{ \begin{pmatrix} \pm e^{i\alpha} \\ 0 \\ 0 \end{pmatrix} \right\}, \left\{ \begin{pmatrix} (-1)^k e^{i\alpha} \\ (-1)^l e^{i\alpha} \\ (-1)^{k+l} e^{i\alpha} \end{pmatrix} \right\}, \left\{ \begin{pmatrix} (-1)^k e^{i\alpha} \\ \omega(-1)^l e^{i\alpha} \\ \omega^2(-1)^{k+l} e^{i\alpha} \end{pmatrix} \right\}, \left\{ \begin{pmatrix} -(-1)^k e^{i\alpha} \\ \omega(-1)^l e^{i\alpha} \\ \omega^2(-1)^{k+l} e^{i\alpha} \end{pmatrix} \right\}, \left\{ \begin{pmatrix} 0 \\ \pm e^{i\alpha} \\ \pm e^{i\alpha+i\beta} \end{pmatrix} \right\}. \quad (3.115)$$

The alignment classes listed after this paragraph arise in this case. Note that in the following, the phases ζ and ζ' are not arbitrary, but are fixed by the one-triplet parts of the two-triplet potential, as by eq. (3.43). Where two sign choices are given, they correspond to separate orbits.

table 4 contains the list of VEVs for $V_{A_4}(\varphi, \varphi')$. Minima $(1, 0, 0)|(e^{i\zeta'}, 1, 0)$ correspond to a rephasing of the triplets and lie in the same orbit as $(1, 0, 0)|(1, e^{i\zeta'}, 0)$. Similarly, $(1, 0, 0)|(0, e^{i\zeta'}, 1)$ and $(1, 0, 0)|(1, 0, e^{i\zeta'})$ are already included in the orbits above.

Under trivial CP, orbits with phases merge with their complex conjugates. In particular, as $c, c' = 0$, $\zeta, \zeta' = 0$.

3.2.5 $\Delta(54)$

Next, consider potentials of two triplets of $\Delta(54)$. Using eq. (3.72) we write

$$\begin{aligned} V_{\Delta(54)}(\varphi, \varphi') &= V_0(\varphi) + V'_0(\varphi') + V_2(\varphi, \varphi') \\ &+ [d(\varphi_1 \varphi_1 \varphi^{*2} \varphi^{*3} + \text{cycl.}) + \text{h.c.}] + [d'(\varphi'_1 \varphi'_1 \varphi'^{*2} \varphi'^{*3} + \text{cycl.}) + \text{h.c.}] \\ &+ \tilde{d}_1 [(\varphi_1 \varphi'_1 \varphi^{*2} \varphi'^{*3} + \text{cycl.}) + (\varphi_1 \varphi'_1 \varphi^{*3} \varphi'^{*2} + \text{cycl.})] + \text{h.c..} \end{aligned} \quad (3.116)$$

Finally, setting

$$\tilde{d}_{21} = \tilde{d}_{11}, \quad \tilde{d}_{22} = \tilde{d}_{12}, \quad \tilde{s}_{21} = \tilde{s}_{22} = \tilde{s}_{31} = \tilde{s}_{32} = 0, \quad (3.117)$$

we obtain

$$\begin{aligned}
V_{\Delta(54)}(H, H') &= V_0(H) + V'_0(H') + V_2(H, H') \\
&+ \sum_{\alpha, \beta} \left[d \left(h_{1\alpha} h_{1\beta} h^{*2\alpha} h^{*3\beta} + \text{cycl.} \right) + d' \left(h'_{1\alpha} h'_{1\beta} h'^{*2\alpha} h'^{*3\beta} + \text{cycl.} \right) + \text{h.c.} \right] \\
&+ \sum_{\alpha, \beta} \left[\tilde{d}_{11} (h_{1\alpha} h^{*2\alpha} h'_{1\beta} h'^{*3\beta} + \text{cycl.}) + \tilde{d}_{12} (h_{1\alpha} h'^{*3\alpha} h'_{1\beta} h^{*2\beta} + \text{cycl.}) + \text{h.c.} \right] \\
&+ \sum_{\alpha, \beta} \left[\tilde{d}_{11} (h_{1\alpha} h^{*3\alpha} h'_{1\beta} h'^{*2\beta} + \text{cycl.}) + \tilde{d}_{12} (h_{1\alpha} h'^{*2\alpha} h'_{1\beta} h^{*3\beta} + \text{cycl.}) + \text{h.c.} \right].
\end{aligned} \tag{3.118}$$

The full symmetries of an even potential of one triplet of $\Delta(54)$ were generated by eq. (3.63). Again, the potential of two triplets of $\Delta(54)$ has no automatic CP symmetries. The potential of two triplets of $\Delta(54)$ could be reached in several ways: the single-triplet parts of the potential each consist of a $\Delta(6n^2)$ part and a $\Delta(54)$ part, while the same is true for the cross-terms. With eq. (3.67) we can write

$$V_{\Delta(54)}(\varphi, \varphi') = V_{\Delta(54)}(\varphi) + V'_{\Delta(54)}(\varphi') + V_{\Delta(54),c}(\varphi, \varphi'), \tag{3.119}$$

where each part splits into the part that is symmetric under the larger group and the part that is specific to $\Delta(54)$, e.g. for the cross-terms

$$V_{\Delta(54),c}(\varphi, \varphi') = V_{\Delta(6n^2),c}(\varphi, \varphi') + V'_{\Delta(54),c}(\varphi, \varphi'). \tag{3.120}$$

The orbits of VEVs of a potential of one triplet of $\Delta(54)$ are

$$\begin{aligned}
&\left\{ \begin{pmatrix} \omega^k e^{i\alpha} \\ 0 \\ 0 \end{pmatrix}, \text{perm.} \right\}, \left\{ \begin{pmatrix} \omega^k e^{i\alpha} \\ \omega^l e^{i\alpha} \\ \omega^{2k+2l} e^{i\alpha} \end{pmatrix}, \text{perm.} \right\}, \\
&\left\{ \begin{pmatrix} \omega^k e^{i\alpha} \\ \omega^l e^{i\alpha} \\ \omega^{2k+2l+1} e^{i\alpha} \end{pmatrix}, \text{perm.} \right\}, \left\{ \begin{pmatrix} \omega^k e^{i\alpha} \\ \omega^l e^{i\alpha} \\ \omega^{2k+2l+2} e^{i\alpha} \end{pmatrix}, \text{perm.} \right\}.
\end{aligned} \tag{3.121}$$

Here, there are no permutations that are not reproduced by some combination of (k, l) .

The full phase symmetries of two triplets are generated by the direct sum of the generators in eq. (3.63) except for the potential having two separate phase symmetries for each triplet, equivalently to eq. (3.78). Again combining members of single triplet orbits and using the symmetries of the full potential to reduce the orbits yields the VEVs in table 5.

Here, the last line stands for several orbits distinguished by the value of i' and i . As all of $k, k', l, l', i, i' = 0, 1, 2$, especially pairs like $(1, \omega, 1) | (1, 1, \omega^2)$ occur, where the phase differences appearing between first and second triplet had previously been unphysical. Nevertheless, as these orbits had only been obtained by reducing the orbits of triplets that already had a $\Delta(54)$ symmetry instead of $\Delta(6n^2)$, the above list is probably incomplete, similarly to the A_4 reduction of S_4 orbits at the end of section 3.1.3.

The effects of imposing CP_0 are analysed in section 4.2.9.

(1, 0, 0) (1, 0, 0)
(1, 0, 0) (0, 1, 0)
(1, 0, 0) (1, 1, 1)
(1, 0, 0) (1, 1, ω)
(1, 0, 0) (1, 1, ω^2)
(1, 1, ω^i) ($\omega^{k'-k}$, $\omega^{l'-l}$, $\omega^{2k'+2l'-2k-2l+i'}$)

Table 5. List of VEV directions for the potential of two triplets of $\Delta(54)$, $V_{\Delta(54)}(\varphi, \varphi')$.

3.2.6 $\Delta(27)$

Using eq. (3.68), the potential of two triplets of $\Delta(27)$ even in both triplets is

$$\begin{aligned} V_{\Delta(27)}(\varphi, \varphi') &= V_0(\varphi) + V'_0(\varphi') + V_1(\varphi, \varphi') \\ &+ [d(\varphi_1\varphi_1\varphi^{*2}\varphi^{*3} + \text{cycl.}) + \text{h.c.}] + [d'(\varphi'_1\varphi'_1\varphi'^{*2}\varphi'^{*3} + \text{cycl.}) + \text{h.c.}] \\ &+ [\tilde{d}_1(\varphi_1\varphi'_1\varphi^{*2}\varphi'^{*3} + \text{cycl.}) + \text{h.c.}] + [\tilde{d}_2(\varphi_1\varphi'_1\varphi^{*3}\varphi'^{*2} + \text{cycl.}) + \text{h.c.}]. \end{aligned} \quad (3.122)$$

Only d , d' , \tilde{d}_1 and \tilde{d}_2 are generally complex, the other coefficients are real. We note also that the $\Delta(54)$ potential (eq. (3.116)) is a particular case of this one, obtained by setting

$$\tilde{s}_2 = \tilde{s}_3 = 0, \quad \tilde{d}_1 = \tilde{d}_2. \quad (3.123)$$

For $SU(2)_L$ doublets, using eq. (3.71), the potential is:

$$\begin{aligned} V_{\Delta(27)}(H, H') &= V_0(H) + V'_0(H') + V_1(H, H') + \\ &+ \sum_{\alpha, \beta} \left[d \left(h_{1\alpha} h_{1\beta} h^{*2\alpha} h^{*3\beta} + \text{cycl.} \right) + d' \left(h'_{1\alpha} h'_{1\beta} h'^{*2\alpha} h'^{*3\beta} + \text{cycl.} \right) + \text{h.c.} \right] \\ &+ \sum_{\alpha, \beta} \left[\tilde{d}_{11} (h_{1\alpha} h^{*2\alpha} h'_{1\beta} h'^{*3\beta} + \text{cycl.}) + \tilde{d}_{12} (h_{1\alpha} h'^{*3\alpha} h'_{1\beta} h^{*2\beta} + \text{cycl.}) + \text{h.c.} \right] \\ &+ \sum_{\alpha, \beta} \left[\tilde{d}_{21} (h_{1\alpha} h^{*3\alpha} h'_{1\beta} h'^{*2\beta} + \text{cycl.}) + \tilde{d}_{22} (h_{1\alpha} h'^{*2\alpha} h'_{1\beta} h^{*3\beta} + \text{cycl.}) + \text{h.c.} \right]. \end{aligned} \quad (3.124)$$

The VEVs of one triplet of $\Delta(27)$ are identical to that of one triplet of $\Delta(54)$, the VEV pairs generated in this way are almost identical for two triplets of $\Delta(27)$ to the above ones of two triplets of $\Delta(54)$. The only effect of the missing permutation in $\Delta(27)$ with respect to $\Delta(54)$ is that several orbits split and that in addition to table 5, the following pairs represent new independent orbits:

$$(1, 0, 0)|(1, 1, \omega) \quad (3.125)$$

$$(1, 0, 0)|(1, \omega, 1) \quad (3.126)$$

$$(1, 0, 0)|(1, 1, \omega^2) \quad (3.127)$$

$$(1, 0, 0)|(1, \omega^2, 1). \quad (3.128)$$

4 Spontaneous CP-odd invariants applied

Recall that for $SU(2)$ doublets we consider only VEVs that preserve $U(1)_{em}$. By an $SU(2)$ gauge transformation one can without loss of generality assume that for each doublet one component is zero. Due to this, the conclusions about presence or absence of CPV are the same as for the respective potentials of $SU(2)$ singlets, and the SCPI expressions are slightly simpler. For this reason, in this section we refer only to the potentials of (φ) or (φ, φ') .

4.1 Potentials of one triplet

4.1.1 $\Delta(3n^2)$ and $\Delta(6n^2)$ with $n > 3$

For the potential of one triplet of $\Delta(6n^2)$ or $\Delta(3n^2)$, eq. (3.11), the CP-odd invariants $\mathcal{J}^{(3,2)}$ and $\mathcal{J}^{(3,3)}$ vanish independently of the triplet VEV. As the phases of the allowed VEVs are unphysical, one can always find a CP symmetry that is present in the potential and preserved by the particular VEV. There is no possibility for SCPV.

4.1.2 A_4 and S_4

Even for A_4 and S_4 , the one triplet cases eq. (3.35) and eq. (3.27) are fairly trivial, as $\mathcal{J}^{(3,2)}$ vanishes before introducing any VEV and $\mathcal{J}^{(3,3)}$ vanishes after introducing each of the VEVs found. In each case, a CP symmetry preserved by the respective VEV has been found — in particular all the VEVs preserve X_{23} , except $(0, 1, e^{i\alpha})$, which preserves a rephased version of X_{23} (alternatively, one can perform an unphysical global rephasing of the VEV such that it also preserves X_{23}).

4.1.3 $\Delta(27)$ and $\Delta(54)$

For the discrete symmetries $\Delta(27)$ and $\Delta(54)$, the single triplet potentials eq. (3.49) and eq. (3.59) are the same. This potential features the first known case of SGCPV.

In [35], already the value of $J^{(3,2)}$ was found for $V_{\Delta(27)}(\varphi) = V_{\Delta(54)}(\varphi)$. For completeness, we repeat it here:

$$\begin{aligned} \mathcal{J}^{(3,2)} = & \frac{1}{4}(d^{*3} - d^3)(|v_1|^4 + |v_2|^4 + |v_3|^4 - 2|v_1|^2|v_2|^2 - 2|v_1|^2|v_3|^2 - 2|v_2|^2|v_3|^2) \\ & + \left[\frac{1}{2}(dd^{*2} - 2d^*s^2 + d^2s)(v_2v_3v_1^{*2} + v_1v_3v_2^{*2} + v_1v_2v_3^{*2}) - \text{h.c.} \right]. \end{aligned} \quad (4.1)$$

When imposing CP_0 (trivial CP, which in this case constrains $\text{Arg}(d) = 0$) or any of the other 5 CP transformations listed by [54] which constrain $\text{Arg}(d)$ to be 0, $2\pi/3$ or $4\pi/3$, the expression simplifies to

$$\mathcal{J}_{CP_0}^{(3,2)} = \frac{1}{2}[dd^{*2} - 2d^*s^2 + d^2s] [v_2v_3v_1^{*2} + v_1v_3v_2^{*2} + v_1v_2v_3^{*2}] - \text{h.c.} \quad (4.2)$$

As described previously [37, 55, 57, 58], the potential admits minima such as the complex VEV $(1, \omega, \omega^2)$, which is however not CP violating when the potential had trivial CP imposed. It can be confirmed easily that the expression above vanishes for this case —

it preserves not CP_0 , but the product of CP_0 with one of the group elements of $\Delta(27)$ (indeed, $(1, \omega, \omega^2)$ is in the same VEV orbit as $(1, 1, 1)$, which is real and preserves CP_0). Instead, the SGCPV VEV $(\omega, 1, 1)$ can be inserted into the expression and gives (for CP_0 making $d = d^*$):

$$\mathcal{J}_{CP_0}^{(3,2)}[v(\omega, 1, 1)] = \frac{3}{2}d(d-s)(d+2s)(\omega-\omega^2)v^4 \quad (4.3)$$

For the CP transformation X_3 forcing $\text{Arg}(d) = 2\pi/3$ (see eq. (3.52)), d is complex, the results are that the VEVs $(0, 0, 1)$ and $(\omega, 1, 1)$ preserve some subset of the CP symmetries, so again no SCPV occurs. Instead, the real VEV $(1, 1, 1)$ and in the same orbit, the complex VEV $(1, \omega, \omega^2)$ show SGCPV [45], as indicated by the SCPI giving $\mathcal{J}_1^{(3,2)} \propto (\omega - \omega^2)$.

The other 6 CP transformations that can be applied to the potential [54], such as X_4 (see eq. (3.53)), don't constrain the phase of d , but rather relate the parameters d and s such that the SCPI simplifies to:

$$\mathcal{J}_{X_4}^{(3,2)} = \frac{1}{4}(d^{*3} - d^3)(|v_1|^4 + |v_2|^4 + |v_3|^4 - 2|v_1|^2|v_2|^2 - 2|v_1|^2|v_3|^2 - 2|v_2|^2|v_3|^2). \quad (4.4)$$

The SCPI reveals that for these CP symmetries, SCPV is independent of the phases of the VEV. We verified that the known VEVs for the CP_0 symmetric potential, including real VEVs such as $(0, 0, 1)$, $(1, 1, 1)$ and complex ones such as $(\omega, 1, 1)$ are still candidate VEVs of the X_4 symmetric potential and all violate CP spontaneously, as indicated by the SCPI not vanishing, e.g.:

$$\mathcal{J}_{X_4}^{(3,2)}[v(1, 0, 0)] = \frac{1}{4}(d^{*3} - d^3)v^4 \quad (4.5)$$

4.2 Potentials of two triplets

4.2.1 $\Delta(3n^2)$ and $\Delta(6n^2)$ with $n > 3$

The useful SCPI $\mathcal{J}^{(3,2)}$ also takes a non-zero expression for the potential of two triplets of $\Delta(3n^2)$, eq. (3.91) for $SU(2)_L$ singlets (the respective potential for doublets can be found in eq. (3.92)).

$$\begin{aligned} \mathcal{J}^{(3,2)} = & -\frac{1}{16}\tilde{s}_2[\tilde{r}_2(-4s-4s'+2\tilde{s}_1-\tilde{s}_2+3\tilde{r}_2)-\tilde{s}_3^2]W_{CP_0} \\ & -\frac{1}{8}i\tilde{s}_3[\tilde{s}_3^2-3\tilde{r}_2^2]W_{CP_{23}} \\ & -\frac{1}{16}i\tilde{s}_2\tilde{s}_3[-4s-4s'+2\tilde{s}_1-\tilde{s}_2](v'_1v_2v_1^*v_2^*+v'_2v_3v_2^*v_3^*+v_1v'_3v_1^*v_3^*+\text{h.c.}) \end{aligned} \quad (4.6)$$

where we define VEV dependences for convenience:

$$W_{CP_{23}} \equiv [(|v'_1|^2|v_2|^2 + |v'_2|^2|v_3|^2 + |v'_3|^2|v_1|^2) - (|v_1|^2|v'_2|^2 + |v_2|^2|v'_3|^2 + |v_3|^2|v'_1|^2)] \quad (4.7)$$

and

$$W_{CP_0} \equiv [(v_1v_1^*v'_2v_2^* + v_2v_2^*v'_3v_3^* + v_3v_3^*v'_1v_1^*) - \text{h.c.}] \quad (4.8)$$

which is similar to the remaining VEV dependence apart from the relative minus sign with respect to the hermitian conjugate.

In contrast, for $\Delta(6n^2)$ with two triplets, shown in eq. (3.75), $\mathcal{J}_1^{(3,2)}$ vanishes independently of the VEVs, and $\mathcal{J}_1^{(3,3)}$ takes a non-vanishing expression in general which vanishes when inserting the VEV candidates we found. In each case, all the VEV candidates for the two triplets preserve a CP symmetry (some combination of CP_0 or CP_{23}), that is automatically present in the general $V_{\Delta(6n^2)}(\varphi, \varphi')$ potential - therefore this potential does not admit SCPV.

4.2.2 $\Delta(3n^2)$, $n > 3$ with CP_0

For $V_{\Delta(3n^2)}$, $n > 3$, the non-zero $\mathcal{J}_1^{(3,2)}$ expression becomes meaningful as a measure of SCPV when imposing a CP symmetry on the potential. If we choose CP_0 , then $\tilde{s}_3 = 0$ and the SCPI becomes:

$$\mathcal{J}_{CP_0}^{(3,2)} = -\frac{1}{16}\tilde{r}_2\tilde{s}_2[-4s - 4s' + 2\tilde{s}_1 - \tilde{s}_2 + 3\tilde{r}_2]W_{CP_0} \quad (4.9)$$

One can note from

$$W_{CP_0} \equiv [(v_1 v'_1 v'_2 v_2^* + v_2 v'_2 v'_3 v_3^* + v_3 v'_3 v'_1 v_1^*) - \text{h.c.}] \quad (4.10)$$

that the expression vanishes if the VEVs of the two triplets do not have matching entries, such as $v(1, 0, 0)|v'(0, 1, 1)$, where v and v' are real here and throughout the rest of the paper. This suggests that those types of VEVs will not show SCPV in this case, regardless of complex phases, confirming that only the relative phase across the same component of the two triplets is physical for this potential. In contrast, a non-zero result is sufficient to show that the VEV pair does SCPV, and the three classes of VEVs that can SCPV in this case are those that in general depend at least on one phase, such as

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1, 1, 0)|v'(1, e^{i\zeta'}, 0)] = \frac{1}{8}i\tilde{r}_2\tilde{s}_2(3\tilde{r}_2 - 4s + 2\tilde{s}_1 - \tilde{s}_2 - 4s')\sin(\zeta')v^2v'^2 \quad (4.11)$$

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1, 1, 0)|v'(1, e^{i\zeta'}, 1)] = \frac{1}{8}i\tilde{r}_2\tilde{s}_2(3\tilde{r}_2 - 4s + 2\tilde{s}_1 - \tilde{s}_2 - 4s')\sin(\zeta')v^2v'^2 \quad (4.12)$$

In this notation, the square brackets denote the pair of VEV considered (including the v , v' normalisation), in order to more easily track which VEVs are being plugged in. Note that the ordering of the VEVs can affect the results, e.g.

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1, 1, 0)|v'(1, e^{i\zeta'}, 1)] = -\mathcal{J}_{CP_0}^{(3,2)}[v(1, e^{i\zeta'}, 1)|v'(1, 1, 0)]. \quad (4.13)$$

We note also that for $\tilde{s}_3 = 0$, $\zeta' = \pi/2$ and therefore the VEV of the second triplet becomes $(1, i, 0)$ and $(1, i, 1)$ respectively (cf. eq. (3.101)). As the CP violating phases are fixed to geometric values, they are geometrical and these cases are considered SGCPV. The third class that is also very interesting is

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1, 1, 1)|v'(1, \omega, \omega^2)] = -\frac{3}{16}\tilde{r}_2\tilde{s}_2(3\tilde{r}_2 - 4s + 2\tilde{s}_1 - \tilde{s}_2 - 4s')(\omega - \omega^2)v^2v'^2 \quad (4.14)$$

as the phases of the VEVs take special values ω and ω^2

When the VEV pair representatives take the special values of phases that minimize the potential in certain regions of parameter space, namely one of the triplets aligns in the

directions with one entry being i , or in the $(1, \omega, \omega^2)$ direction, the SCPI reveals cases with SGCPV. This is interesting as these are the first reported cases of SGCPV in potentials with 6 fields (arranged here as two triplets), the potential is relatively simple due to the symmetry and the special ω, ω^2 phases appear for a symmetry that is not $\Delta(27)$ or $\Delta(54)$ (the $n = 3$ cases).

4.2.3 $\Delta(3n^2)$, $n > 3$ with CP_{23}

When CP_{23} (2-3 swap CP) is imposed on $V_{\Delta(3n^2)}(\varphi, \varphi')$, the respective SCPI $\mathcal{J}_1^{(3,2)}$ expression becomes particularly simple. CP_{23} for two triplets swaps the 2-3 components of both triplets, i.e. $X_{23}^{\varphi\varphi'}$ in eq. (3.95), which leads to $\tilde{s}_2 = 0$:

$$\mathcal{J}_{CP_{23}}^{(3,2)} = -\frac{1}{8}i\tilde{s}_3 [\tilde{s}_3^2 - 3\tilde{r}_2^2] W_{CP_{23}} \quad (4.15)$$

where the VEV dependence

$$W_{CP_{23}} \equiv [(|v'_1|^2|v_2|^2 + |v'_2|^2|v_3|^2 + |v'_3|^2|v_1|^2) - (|v_1|^2|v'_2|^2 + |v_2|^2|v'_3|^2 + |v_3|^2|v'_1|^2)] \quad (4.16)$$

reveals SPCV occurs (or not) regardless of the phases of the VEVs. The three representatives that give non-zero results are proportional to $v^2v'^2$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1, 0, 0)|v'(0, 1, 0)] \neq 0 \quad (4.17)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1, 0, 0)|v'(1, 0, 1)] \neq 0 \quad (4.18)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1, 1, 0)|v'(0, 1, 1)] \neq 0 \quad (4.19)$$

featuring three cases with SGCPV. This may appear peculiar given all the phases are zero, but they are geometrical and therefore these cases fall under the definition of SGCPV — the (vanishing) phases of the VEVs are stable under small variations of the potential parameters, and there is SPCV.

For the type of VEV where the phase ζ' appears with a physical non-trivial value, CP is spontaneously violated, even though both invariants we are using vanish:

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1, 1, 0)|v'(1, e^{i\zeta'}, 0)] = \mathcal{J}_{CP_{23}}^{(3,3)}[v(1, 1, 0)|v'(1, e^{i\zeta'}, 0)] = 0 \quad (4.20)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1, 1, 0)|v'(1, e^{i\zeta'}, 1)] = \mathcal{J}_{CP_{23}}^{(3,3)}[v(1, 1, 0)|v'(1, e^{i\zeta'}, 1)] = 0 \quad (4.21)$$

As a systematic search for further SCPIs is beyond the scope of this work, instead we have excluded the possibility of X matrices that are simultaneously a symmetry of the potential with general $\tilde{s}_3 \neq 0$ and are preserved by the VEVs. The first condition fails for CP symmetries that transform the triplets differently, whereas the second cannot be fulfilled if both triplets transform identically under the CP symmetry. In fact, we were able to find a SCPI which indeed gives a non-zero result for $v(1, 1, 0)|v'(1, e^{i\zeta'}, 1)$, but as this is the only instance in this paper where it is important, we haven't mentioned it previously. The relevant SCPI is:

$$J_{CP_{23}}^{(3,4)} = Z_{a_5 a_7}^{a_1 a_2} Z_{a_6 a_8}^{a_3 a_4} Z_{a_9 a_{10}}^{a_5 a_6} v_{a_1} v_{a_2} v_{a_3} v_{a_4} v^{*a_7} v^{*a_8} v^{*a_9} v^{*a_{10}} \quad (4.22)$$




Figure 4. The diagram corresponding to the invariant eq. (4.22).

with diagram in figure 4 and we have

$$\mathcal{J}_{CP_{23}}^{(3,4)}[v(1,1,0)|v'(1,e^{i\zeta'},1)] \propto (2\sin\zeta' - \sin 2\zeta'). \quad (4.23)$$

4.2.4 $\Delta(3n^2)$, $n > 3$ another CP

In [35], explicit CP violation was studied and another class of CP symmetry was found that made the potential CP conserving by relating the parameters \tilde{s}_3 and \tilde{r}_2 in 3 similar ways. Of these, we denoted in eq. (3.96)):

$$X_{\varphi\varphi'} = \begin{pmatrix} X_\varphi & 0 \\ 0 & X_{\varphi'} \end{pmatrix}, \quad \text{with} \quad X_\varphi = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \quad X_{\varphi'} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix}, \quad (4.24)$$

which we repeat here as this CP symmetry relates the parameters as

$$\tilde{s}_3 = \tilde{r}_2 i(\omega - \omega^2), \quad (4.25)$$

i.e. $\tilde{s}_3 = -\sqrt{3}\tilde{r}_2$. This CP symmetry makes the explicit CPIs vanish [35] and simplifies the SCPI we are considering such that the first two lines of eq. (4.6) vanish. Indeed, when this is inserted into the SCPI (considering that $2\omega = i(\sqrt{3} + i)$):

$$\mathcal{J}_{X_{\varphi\varphi'}}^{(3,2)} = \frac{1}{8} \tilde{r}_2 \tilde{s}_2 [4s + 4s' - 2\tilde{s}_1 + \tilde{s}_2] [\omega(v_1 v_1' v_2' v_2^* + v_2 v_2' v_3' v_3^* + v_3 v_3' v_1' v_1^*) - \text{h.c.}], \quad (4.26)$$

the SCPI becomes similar to the CP_0 case but there are relevant factors of ω and ω^2 which change the VEV dependence crucially (note this is not ωW_{CP_0} , so it is not just an overall multiplicative factor). Indeed, the cases with VEVs featuring powers of ω that were SGCPV for CP_0 are CP conserving for this case and instead, the pair of VEVs $(1,1,1)|(1,1,1)$ is SGCPV. Further, the classes of VEVs with ζ' , which gets fixed to $\zeta' = -\pi/6$ due to the CP symmetry, also gives a non-zero value for the SCPI and are therefore more cases with SGCPV.

4.2.5 A_4 and S_4

The general expression of $\mathcal{J}^{(3,2)}$ for the $V_{A_4}(\varphi, \varphi')$ potential with two triplets (listed in eq. (3.109)) is non-zero. We don't show it as it is rather long, and strictly speaking it carries no physical meaning as the potential is explicitly CPV. The expression simplifies

sufficiently for relevant cases, when specific CP symmetries are imposed, as shown below as we test each of the twenty two (22) VEV pairs from section 3.2.4.

In contrast, $V_{S_4}(\varphi, \varphi')$ (seen in eq. (3.106)) is a particular case of $V_{A_4}(\varphi, \varphi')$ that automatically preserves both CP_0 and CP_{23} . The general expression of $\mathcal{J}^{(3,2)}$ for $V_{S_4}(\varphi, \varphi')$ vanishes, although this does not happen with the SCPI $\mathcal{J}^{(3,3)}$ for $V_{S_4}(\varphi, \varphi')$. When testing the twenty (20) VEV pairs from section 3.2.3 we found the SCPI $\mathcal{J}^{(3,3)}$ always vanishes, which is understood as the VEVs either preserve CP_0 (the real VEVs) or CP_{23} . In this potential there is no SCPV.

Given this, of the many SCPIs we calculated for the discrete groups A_4 and S_4 , it is only interesting to look in more detail to $V_{A_4}(\varphi, \varphi')$.

4.2.6 A_4 with CP_0

When CP_0 (trivial CP) is imposed on $V_{A_4}(\varphi, \varphi')$, the respective SCPI $\mathcal{J}^{(3,2)}$ becomes:

$$\mathcal{J}_{CP_0}^{(3,2)} = C_{A_4} W_{CP_0} \quad (4.27)$$

with the coefficient dependence

$$C_{A_4} = \frac{1}{16} \tilde{s}_2 (\tilde{r}_2 (-3\tilde{r}_2 + 4s - 2\tilde{s}_1 + \tilde{s}_2 + 4s') - 4\tilde{c}(c + c')) \quad (4.28)$$

and the same VEV dependence, W_{CP_0} in eq. (4.8), that appeared in the SCPI in eq. (4.9), of the analogous $\Delta(3n^2)$ case invariant under CP_0 . This is not completely unexpected, as the SCPI should vanish regardless of the discrete symmetry being A_4 or $\Delta(3n^2)$ with $n > 3$, for VEVs that preserve CP_0 (note though that for the $\Delta(27)$ case, the functional dependence is different).

An analysis of the expression reveals that for this SCPI, only relative phases between the same component of the two triplets appear. This was the case for the $\Delta(3n^2)$ potential, but for A_4 there are actually some additional physical phases, which the more complicated SCPI $\mathcal{J}^{(3,3)}$ is sensitive to.

Using $\mathcal{J}^{(3,2)}$ we can confirm that the following VEV pairs SCPV, being proportional to $v^2 v'^2$:

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1, e^{i\zeta}, 0) | v'(1, \pm e^{i\zeta'}, 0)] \neq 0 \quad (4.29)$$

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1, e^{i\zeta}, 0) | v'(1, \pm 1, 1)] \neq 0 \quad (4.30)$$

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1, e^{i\zeta}, 0) | v'(1, \omega, \pm\omega^2)] \neq 0 \quad (4.31)$$

and specifically

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1, 1, 1) | v'(1, \omega, \omega^2)] = 3C_{A_4}(\omega - \omega^2)v^2 v'^2 \quad (4.32)$$

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1, 1, 1) | v'(1, \omega, -\omega^2)] = C_{A_4}(\omega - \omega^2)v^2 v'^2. \quad (4.33)$$

The most interesting cases are arguably these latter ones where VEVs only have geometrical phases and therefore SGCPV, although one notes they are essentially the same cases obtained for the slightly simpler $\Delta(3n^2)$ potential.

For the cases where the pair of VEVs returns a vanishing expression we confirm if CP is not broken by identifying a CP symmetry preserved by the VEVs (such as CP_0 itself, for any real VEVs).

4.2.7 A_4 with CP_{23}

When CP_{23} (2-3 swap CP) is imposed on $V_{A_4}(\varphi, \varphi')$, the respective SCPI $\mathcal{J}_1^{(3,2)}$ expression becomes:

$$\mathcal{J}_{CP_{23}}^{(3,2)} = \frac{1}{8} [i\tilde{s}_3 (2\tilde{c}^*(c+c') + 2\tilde{c}(c^*+c^*) + 3\tilde{r}_2^2) + 2\tilde{r}_2(\tilde{c}^*(c-c') - c^*\tilde{c} + c^*\tilde{c}) - i\tilde{s}_3^3] W_{CP_{23}}. \quad (4.34)$$

Note this is the same VEV dependence, $W_{CP_{23}}$ in eq. (4.7), that does not depend on the phases of the VEVs and which appeared in the SCPI in eq. (4.15), of the analogous $\Delta(3n^2)$ potential invariant under CP_{23} . This is not completely unexpected, as the SCPI should vanish, regardless of the discrete symmetry being A_4 or $\Delta(3n^2)$ with $n > 3$, for VEVs that preserve CP_{23} .

We use this SCPI to check the pairs of VEVs for this potential. We display here only the non-vanishing ones, which are always proportional to $v^2v'^2$, and are:

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,0,0)|v'(0,1,0)] \neq 0 \quad (4.35)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,0,0)|v'(1,e^{i\zeta},0)] \neq 0 \quad (4.36)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,0,0)|v'(0,1,e^{i\zeta})] \neq 0 \quad (4.37)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,0,0)|v'(e^{i\zeta},0,1)] \neq 0 \quad (4.38)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,e^{i\zeta},0)|v'(0,1,e^{i\zeta'})] \neq 0 \quad (4.39)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,e^{i\zeta},0)|v'(e^{i\zeta'},0,1)] \neq 0 \quad (4.40)$$

The SCPI vanishes for the pair $(1,1,1)|(1,\omega,\omega^2)$ regardless of the physical complex phases (this is clear from the SCPI expression, which depends only on the absolute values). It must do so, as the VEV pair preserves CP_{23} . In contrast, $[(1,0,0)|(0,1,0)]$ is a SGCPV pair of VEVs as the vanishing phases are geometrical.

Furthermore, for A_4 we have cases entirely analogous to the two VEV pairs that CPV, even though the SCPIs $\mathcal{J}^{(3,2)}$, $\mathcal{J}^{(3,3)}$ vanish:

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,1,0)|v'(1,e^{i\zeta'},0)] = \mathcal{J}_{CP_{23}}^{(3,3)}[v(1,1,0)|v'(1,e^{i\zeta'},0)] = 0 \quad (4.41)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,1,0)|v'(1,e^{i\zeta'},1)] = \mathcal{J}_{CP_{23}}^{(3,3)}[v(1,1,0)|v'(1,e^{i\zeta'},1)] = 0 \quad (4.42)$$

The conclusions are the same as in $\Delta(3n^2)$, as there is no CP symmetry that is both preserved by these VEVs (this condition requires X matrices transforming the two triplets differently) and is a symmetry of the potential with $\tilde{s}_3 \neq 0$ (this condition requires X matrices that transform the two triplets equally).

4.2.8 $\Delta(27)$ and $\Delta(54)$

For two triplets the potentials $V_{\Delta(27)}(\varphi, \varphi')$ and $V_{\Delta(54)}(\varphi, \varphi')$ are different, as seen in eq. (3.122) and eq. (3.116) respectively. Nonetheless, the VEV candidates are the same, and the SCPI results are similar. We found several new VEV pairs, many of which are cases with SGCPV.

It turns out the SCPI expressions are slightly more complicated in the $\Delta(27)$ case, but they vanish whenever the corresponding expression in $\Delta(54)$ vanishes, and they are proportional to $(\omega - \omega^2) = i\sqrt{3}$ when the corresponding SCPI expression in $\Delta(54)$ has that dependence. For this reason, we display the SCPI results only for $\Delta(54)$.

4.2.9 $\Delta(54)$ with CP_0

The last 3 representative VEV pairs from section 3.2.5 can give non-zero results with SGCPV. In this case we show some cases that make the SCPI vanish, for clarity:

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,1,1)|v'(1,\omega,\omega^2)] = 0 \quad (4.43)$$

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,\omega,\omega^2)|v'(1,\omega,\omega^2)] = 0. \quad (4.44)$$

The non-vanishing cases include

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,0,0)|v'(1,1,\omega)] = -\mathcal{J}_1^{(3,2)}[v(1,0,0)|v'(1,1,\omega^2)] \quad (4.45)$$

$$= \frac{3}{16} \left(2\tilde{d}_1^2(d' + s') + 4\tilde{d}_1\tilde{r}_2s' + d'(8(d' - s')(d' + 2s') - \tilde{s}_1(2\tilde{r}_2 + \tilde{s}_1)) \right) (\omega - \omega^2)v'^4$$

and

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,1,\omega)|v'(1,\omega,\omega^2)] = -\mathcal{J}_1^{(3,2)}[v(1,1,\omega^2)|v'(1,\omega,\omega^2)] \quad (4.46)$$

$$= \frac{3}{16} \left(8d^3 + 8d^2s + d \left(2\tilde{d}_1^2 - \tilde{s}_1(2\tilde{r}_2 + \tilde{s}_1) - 16s^2 \right) + 2\tilde{d}_1s(\tilde{d}_1 + 2\tilde{r}_2) \right) (\omega - \omega^2)v^4$$

Note the dependence is either on v^4 or v'^4 exclusively in these cases (and not on $v^2v'^2$). We recall for $\Delta(27)$ and $\Delta(54)$, SCPV is already possible in the single triplet potentials. This contrasts with the A_4 and $\Delta(3n^2)$ cases, where only the two triplet potentials admit SCPV — with only two triplets enabling SCPV, the non-vanishing SCPIs for those cases depend always on $v^2v'^2$. In $\Delta(54)$ we also found VEV pairs for which the SCPI has the $v^2v'^2$ dependence, one such instance is $(1,1,\omega)|(1,1,\omega^2)$ which when plugged into the SCPI has all three dependences, v^4 , v'^4 and $v^2v'^2$.

4.3 Spontaneous Geometrical CP Violation

We have found several new cases of SGCPV. Through these new cases we hope to advance the understanding of what the conditions are for it to occur.

Clearly, as part of the definition, all of the cases found that show SGCPV have geometrical phases inside the VEVs. We also demonstrate in this section that these VEVs preserve a different geometrical CP symmetry (e.g. real VEVs that violate CP_{23} but would have preserved CP_0). We conjecture that this is a requirement for SGCPV to occur. It is important to stress that the CP symmetries that these VEVs would have preserved are themselves geometrical, having only geometrical phases that do not depend on the parameters of the potential. For cases with SCPV that is not geometric one can find CP symmetries that would have been preserved by the respective VEVs, but those CP symmetries will also not be geometrical: the phases inside the VEVs and the associated CP

symmetries the VEVs would have preserved depend on the parameters of the potential, which run with energy scale.

This usefulness of this conjecture is illustrated by our examples. In cases where it is possible to identify distinct geometrical CP symmetries that can be imposed to constrain the parameters of a specific potential, that potential can then give rise to SGCPV, depending of course on the VEVs allowed by the respective minimisation conditions.

We divide our cases mainly in VEVs of one or two triplets where the imposed CP symmetry is CP_0 and cases where the imposed CP symmetry is CP_{23} .

In the latter case, where CP_{23} is imposed, we found three cases with SGCPV for $\Delta(3n^2)$ with $n > 3$:

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,0,0)|v'(0,1,0)] \neq 0 \quad (4.47)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,0,0)|v'(1,0,1)] \neq 0 \quad (4.48)$$

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,1,0)|v'(0,1,1)] \neq 0 \quad (4.49)$$

and one similar case for A_4 :

$$\mathcal{J}_{CP_{23}}^{(3,2)}[v(1,0,0)|v'(0,1,0)] \neq 0. \quad (4.50)$$

It occurs always in situations with two triplets where the VEVs have the geometrical phase of zero (i.e. the VEVs are real), in pairs that simultaneously violate CP_{23} and the cyclic permutations of CP_{23} (which we may refer to as CP_{31} and CP_{12}) that are also CP symmetries of the potential when CP_{23} is imposed, due to the flavour symmetries we are considering. Indeed, it is understandable why there is no SGCPV of this type for the respective one triplet potentials, as a real single triplet VEV cannot violate all 3 of those CP symmetries. We note though that the real VEV pairs that SGCPV in these cases would have preserved CP_0 , which is not however a symmetry of the potentials (otherwise there wouldn't be SCPV).

In the cases where CP_0 is imposed, which include also the original SGCPV cases with one triplet of $\Delta(27)$ (or $\Delta(54)$), we note also that the VEVs with geometrical phases that have SGCPV would have preserved some other geometrical CP symmetry⁸ that is not a symmetry of the potential. In the one triplet case, take:

$$\mathcal{J}_{CP_0}^{(3,2)}[v(\omega,1,1)] = \frac{3}{2}d(d-s)(d+2s)(\omega - \omega^2)v^4 \quad (4.51)$$

this VEV would have preserved a variant of the X_2 , X_3 CP transformations (which is consistent):

$$X'_2 = \begin{pmatrix} \omega & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (4.52)$$

⁸By extension of the definition of “calculable” phases [37], which we refer to as geometrical phases heres, we refer to a geometrical CP symmetry as one that is independent of the parameters of the potential.

The SGCPV VEVs in $\Delta(3n^2)$:

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,1,0)|v'(1,i,0)] = \frac{1}{8}i\tilde{r}_2\tilde{s}_2(3\tilde{r}_2 - 4s + 2\tilde{s}_1 - \tilde{s}_2 - 4s')v^2v'^2 \quad (4.53)$$

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,1,0)|v'(1,i,1)] = \frac{1}{8}i\tilde{r}_2\tilde{s}_2(3\tilde{r}_2 - 4s + 2\tilde{s}_1 - \tilde{s}_2 - 4s')v^2v'^2 \quad (4.54)$$

would have preserved a CP with diagonal matrix $diag(1, 1, 1, 1, -1, 1)$,⁹ and

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,1,1)|v'(1,\omega,\omega^2)] = -\frac{3}{16}\tilde{r}_2\tilde{s}_2(3\tilde{r}_2 - 4s + 2\tilde{s}_1 - \tilde{s}_2 - 4s')(\omega - \omega^2)v^2v'^2 \quad (4.55)$$

would have preserved CP_{23} . CP_{23} would also be preserved by the first of the SGCPV pair of VEVs in A_4 :

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,1,1)|v'(1,\omega,\omega^2)] = 3C_{A_4}(\omega - \omega^2)v^2v'^2 \quad (4.56)$$

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,1,1)|v'(1,\omega,-\omega^2)] = C_{A_4}(\omega - \omega^2)v^2v'^2 \quad (4.57)$$

The second pair, due to the minus sign, would have preserved the CP symmetry with a block matrix with the regular X_{23} for the first triplet and for the second triplet:

$$X_{(-23)} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}. \quad (4.58)$$

The new SGCPV VEVs we find in $\Delta(54)$ (and $\Delta(27)$) are essentially two triplet versions of the one triplet case:

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,0,0)|v'(1,1,\omega)] \neq 0 \quad (4.59)$$

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,1,\omega)|v'(1,\omega,\omega^2)] \neq 0 \quad (4.60)$$

$$\mathcal{J}_{CP_0}^{(3,2)}[v(1,1,\omega)|v'(1,1,\omega^2)] \neq 0 \quad (4.61)$$

and for each case one can find X matrices of the geometrical CP symmetries that would have been preserved. With one triplet the other case with SGCPV that we presented is

$$\mathcal{J}_{X_4}^{(3,2)}[v(1,0,0)] = \frac{1}{4}(d^{*3} - d^3)v^4 \quad (4.62)$$

and this VEV would have preserved CP_0 . Finally, the other CP symmetry we studied for two triplets of $\Delta(3n^2)$ with $n > 3$ led to the following cases of SGCPV:

$$\mathcal{J}_{CP_{\varphi\varphi'}}^{(3,2)}[v(1,1,0)|v'(1,e^{-i\pi/6},0)] \neq 0 \quad (4.63)$$

$$\mathcal{J}_{CP_{\varphi\varphi'}}^{(3,2)}[v(1,1,0)|v'(1,e^{-i\pi/6},1)] \neq 0 \quad (4.64)$$

$$\mathcal{J}_{CP_{\varphi\varphi'}}^{(3,2)}[v(1,1,1)|v'(1,1,1)] \neq 0. \quad (4.65)$$

Of these, the last VEV pair would clearly have preserved CP_0 , and the other VEVs preserve a CP with diagonal matrix $diag(1, 1, 1, 1, e^{i\pi/3}, 1)$ featuring only geometrical phases (0 and $\pi/3$).

⁹This CP symmetry would not be consistent with the flavour symmetry of the potential, but if imposed it changes the potential only by setting $\tilde{r}_2 = \tilde{s}_3 = 0$.

5 Summary of results

In this section we summarise the results presented throughout section 4, in two tables. Table 6 lists the potentials with one triplet of the discrete symmetries and table 7 the potentials with two triplets. For both tables, the leftmost column notes what is the imposed discrete symmetry and, in cases where it is also imposed, the CP symmetry. The next two columns show the two SCPIs we calculated and distinguish whether the expression vanishes in general (before plugging in the VEVs) — note that in some of the cases, despite the expression not vanishing in general, it vanishes for all the VEVs. The fourth column lists if a CP symmetry is present, listing the matrix associated with the CP symmetry of one triplet (for two triplets, this means it is the diagonal 2×2 block matrix with the same transformation for both triplets) — this column is relevant because in many cases where there was no CP symmetry imposed, there is nevertheless a CP symmetry present for the potential that is invariant under the imposed discrete flavour symmetry. The last column notes whether there is Spontaneous CP Violation. In this last column we note 2 categories when there is no imposed CP symmetry — the potential is either CP conserving, and all of these cases have no SCPV either,¹⁰ or alternatively the potential is explicitly CP violating and therefore we denote the SCPV column with NA for Not Applicable. The final possibility is when a CP symmetry is imposed, and it turned out that for all the cases analysed with these discrete symmetries, whenever there was the possibility of SCPV, there were some (but not all) VEVs that have SGCPV, and therefore this column is marked as “S(G)CPV”, because that potential can have regular SCPV or SGCPV depending on the VEVs (we note though that this is probably due to the discrete symmetries we analyse being special, as many potentials, such as the 2HDM with imposed CP symmetry, can have SCPV but don’t have SGCPV).

In summary:

- We present new results for $V_{\Delta(3n^2)}(H) = V_{\Delta(6n^2)}(H)$.
- Of the 1 triplet potentials we studied, only $V_{\Delta(27)}(H) = V_{\Delta(54)}(H)$ has non-trivial CP properties, which includes cases with Spontaneous Geometrical CP Violation.
- All the 2 triplet results we present are new.
- Of the 2 triplets potentials we studied, all that can have SCPV, also show explicit CPV.
- For the potentials we studied where SCPV can be found, some VEVs exist with SGCPV (this is a peculiarity of the symmetries we considered, as it is not true in general).
- We have found several new cases of SGCPV beyond the few known cases up to now, finding also the first cases of SGCPV for 6HDMs.
- We formulate a conjecture relating SGCPV with geometrical CP symmetries.

¹⁰These results support a conjecture from [40], proven for specific cases in [59].

	$J^{(3,2)}$	$J^{(3,3)}$	CP	CP type	SCPV
$\mathbf{3}_{\Delta(3n^2)=\Delta(6n^2)}$	0	0	X_0 , eq. (3.22); X_{23} , eq. (3.36)	Automatic	No
$\mathbf{3}_{A_4}$	0	*	X_{23} , eq. (3.36)	Automatic	No
$\mathbf{3}_{S_4}$	0	*	X_0 , eq. (3.22); X_{23} , eq. (3.36)	Automatic	No
$\mathbf{3}_{\Delta(27)=\Delta(54)}$	*	*	None	NA	NA
$\mathbf{3}_{\Delta(27)=\Delta(54)}, X_0$	*	*	X_0 , eq. (3.22)	Imposed	S(G)CPV
$\mathbf{3}_{\Delta(27)=\Delta(54)}, X_4$	*	*	X_4 , eq. (3.53)	Imposed	S(G)CPV

Table 6. Summary of the value of $J^{(3,2)}$, $J^{(3,3)}$, CP symmetry transformations and whether SCPV is possible, for the 1 triplet scalar potentials analysed. The * denotes that a SCPI is non-zero for arbitrary VEVs (it may still vanish for the potential's VEVs), NA stands for Not Applicable (SCPV is not meaningful in cases without a CP symmetry).

	$J^{(3,2)}$	$J^{(3,3)}$	CP	Type of CP	SCPV
$2 \times \mathbf{3}_{\Delta(3n^2)}$	*	*	None	NA	NA
$2 \times \mathbf{3}_{\Delta(3n^2)}, CP_0$	*	*	$X_0^{\varphi\varphi'}$, eq. (3.94);	Imposed	S(G)CPV
$2 \times \mathbf{3}_{\Delta(3n^2)}, CP_{23}$	*	*	$X_{23}^{\varphi\varphi'}$, eq. (3.95);	Imposed	S(G)CPV
$2 \times \mathbf{3}_{\Delta(3n^2)}, CP_{\varphi\varphi'}$	*	*	$X_{\varphi\varphi'}$, eq. (3.96)	Imposed	S(G)CPV
$2 \times \mathbf{3}_{\Delta(6n^2)}$	0	*	$X_0^{\varphi\varphi'}$, eq. (3.94) $X_{23}^{\varphi\varphi'}$, eq. (3.95)	Automatic	No
$2 \times \mathbf{3}_{A_4}$	*	*	None	NA	NA
$2 \times \mathbf{3}_{A_4}, CP_0$	*	*	$X_0^{\varphi\varphi'}$, eq. (3.94)	Imposed	S(G)CPV
$2 \times \mathbf{3}_{A_4}, CP_{23}$	*	*	$X_{23}^{\varphi\varphi'}$, eq. (3.95)	Imposed	S(G)CPV
$2 \times \mathbf{3}_{S_4}$	0	*	$X_0^{\varphi\varphi'}$, $X_{23}^{\varphi\varphi'}$	Automatic	No
$2 \times \mathbf{3}_{\Delta(27)}$	*	*	None	NA	NA
$2 \times \mathbf{3}_{\Delta(27)}, CP_0$	*	*	$X_0^{\varphi\varphi'}$, eq. (3.94)	Imposed	S(G)CPV
$2 \times \mathbf{3}_{\Delta(54)}$	*	*	None	NA	NA
$2 \times \mathbf{3}_{\Delta(54)}, CP_0$	*	*	$X_0^{\varphi\varphi'}$, eq. (3.94)	Imposed	S(G)CPV

Table 7. Summary of the value of $J^{(3,2)}$, $J^{(3,3)}$, CP symmetry transformations and whether SCPV is possible, for the 2 triplet scalar potentials analysed. The * denotes the SCPI is non-zero for arbitrary VEVs (it may still vanish for the potential's VEVs), NA stands for Not Applicable (SCPV is not meaningful in cases without a CP symmetry).

6 Conclusions

We have analysed 3 and 6 Higgs scalar potentials invariant under discrete symmetries $\Delta(3n^2)$ and $\Delta(6n^2)$ with $n = 2$ (A_4, S_4), $n = 3$ ($\Delta(27), \Delta(54)$) and $n > 3$. For these potentials, we have presented their VEVs and considered whether they can have Spontaneous CP Violation, and if so, if it can be Spontaneous Geometrical CP Violation.

Concerning the minimisation of such complicated potentials, the strategy we have followed in this paper may be summarised as follows. We started with more symmetric cases, which can be minimised analytically, where each of the minima of these potentials is associated with an orbit, which is generated by the representative VEV by the symmetry group. For the less symmetric potentials, we have analysed how these orbits are broken up and which phases can become physical as a result of reducing the original symmetry. We find that for potentials involving one triplet, this method reproduces perfectly the minima found by a previous thorough analysis done by other authors. We then proceeded to construct the orbits of potentials of two triplets, again starting from more symmetric cases, which in this case means the potentials of two triplets without cross-terms. We then used the conjectured minima in the subsequent analysis of CPV.

With respect to Spontaneous CP Violation, using the basis invariant formalism, we presented two Spontaneous CP-odd Invariants and used these to confirm cases where there is Spontaneous CP Violation. With our methods, we have confirmed several existing results in the literature for the 3 Higgs potentials, i.e. involving one triplet under the specified symmetries of $A_4, S_4, \Delta(27)$ and $\Delta(54)$. Beyond that, we present new results for the potentials invariant under $\Delta(3n^2)$ and $\Delta(6n^2)$ with $n > 3$ and those with 6 Higgs, i.e. involving two triplets under the above symmetries. Of the potentials considered, for those that were automatically CP invariant we found no VEVs that spontaneously violate CP — all the SCPIs vanish and we found for each VEV at least one preserved CP symmetry. This is in line with what was the case for 3 Higgs [40] and the more general conjecture proven in the particular case of rephasing symmetries in [59].

For potentials that in general have explicit CP violation, however, we considered imposing different CP symmetries and then checking which (if any) VEVs spontaneously violate CP. After revisiting the well-known case for the 3 Higgs potential invariant under both $\Delta(27)$ and trivial CP (where SGCPV was first identified) we also found SGCPV in the 6 Higgs potentials invariant under $A_4, \Delta(27), \Delta(54)$ and $\Delta(3n^2)$ with $n > 3$. These new cases were found by finding a non-zero SCPI and checking that the VEVs in question have geometrical phases. We have proposed the following conjecture: SGCPV appears when there are different geometrical CP symmetries that can be imposed on a potential, and the VEV violates the imposed geometrical CP symmetry but would have preserved a different geometrical CP symmetry.

Finally we remark that the results in this paper may be applied to physical CP violating processes involving three and six-Higgs doublet potentials controlled by classes of non-Abelian discrete symmetries.

Acknowledgments

The authors thank Igor Ivanov and Renato Fonseca for useful discussions.

IdMV acknowledges funding from Fundação para a Ciência e a Tecnologia (FCT) through the contract IF/00816/2015 and partial support from the HARMONIA project under contract UMO-2015/18/M/ST2/00518 (2016-2019). This project has received funding from the European Union’s Seventh Framework Programme for research, technological development and demonstration under grant agreement no PIEF-GA-2012-327195 SIFT. This work was partially supported by Fundação para a Ciência e a Tecnologia (FCT, Portugal) through the project CFTP-FCT Unit 777 (UID/FIS/00777/2013) which is partially funded through POCTI (FEDER), COMPETE, QREN and EU.

The work of CL is supported by the Deutsche Forschungsgemeinschaft (DFG) within the Research Unit FOR 1873 “Quark Flavour Physics and Effective Field Theories”.

SFK acknowledges the STFC Consolidated Grant ST/L000296/1 and the European Union’s Horizon 2020 Research and Innovation programme under Marie Skłodowska-Curie grant agreements Elusives ITN No. 674896 and InvisiblesPlus RISE No. 690575.

Work supported by MINECO grants FPA2014-58183-P, Multidark CSD2009-00064, and the PROMETEOII/2014/084 grant from Generalitat Valenciana.

A Technical details from section 2

In this appendix, first the argument leading to eq. (2.14) is given. After that, additionally a related relation between VEVs is derived. From each allowed CP transformation, conditions on Y and Z follow,

$$Y_b^a = Y_{a'}^{b'} X_{b'b} X^{*a'a} \quad (\text{A.1})$$

and

$$Z_{cd}^{ab} = Z_{a'b'}^{c'd'} X_{c'c} X_{d'd} X^{*a'a} X^{*b'b}. \quad (\text{A.2})$$

When the potential “acquires” a VEV, i.e. when the parameters of the potential are chosen such that the energy is classically minimized at a value of the field different from zero,

$$\langle \phi \rangle = v \neq 0, \quad (\text{A.3})$$

then first the fields and with them the potential can be expanded around this VEV:

$$\varphi \mapsto v + \varphi \quad (\text{A.4})$$

and

$$V(\phi) \equiv V(v, \varphi) = Y_b^a (v + \varphi)_a (v + \varphi)^{*b} + Z_{cd}^{ab} (v + \varphi)_a (v + \varphi)_b (v + \varphi)^{*c} (v + \varphi)^{*d}. \quad (\text{A.5})$$

This potential can be reordered by the number of fields φ , i.e.

$$\begin{aligned}
V(v, \varphi) &= Y_b^a v_a v^{*b} + Y_b^a v_a \varphi^{*b} + Y_b^a \varphi_a v^{*b} + Y_b^a \varphi_a \varphi^{*b} \\
&\quad + Z_{cd}^{ab} v_a v_b v^{*c} v^{*d} + \dots + Z_{cd}^{ab} \varphi_a \varphi_b \varphi^{*c} \varphi^{*d} \\
&= Y_b^a v_a v^{*b} + Z_{cd}^{ab} v_a v_b v^{*c} v^{*d} \\
&\quad + (Y_b^a v^{*b} + \dots + Z_{cd}^{ab} v_b v^{*c} v^{*d}) \varphi_a \\
&\quad + \dots \\
&\quad + Z_{cd}^{ab} \varphi_a \varphi_b \varphi^{*c} \varphi^{*d}.
\end{aligned} \tag{A.6}$$

The coefficients of this potential are now combinations of Y , Z , and v . This potential is CP-conserving in the new degrees of freedom, φ , if a CP trafo

$$\varphi \mapsto \tilde{X} \varphi^* \tag{A.7}$$

exists, such that the minimized potential is invariant,

$$V(v, \varphi) = V(v, \tilde{X} \varphi^*). \tag{A.8}$$

From this, conditions on the coefficients of the expanded potential follow. For the coefficient quartic in φ , this happens to be the original condition, only with X replaced with \tilde{X} ,

$$Z_{cd}^{ab} \tilde{X}_a^{a'} \tilde{X}_b^{b'} \tilde{X}_c^{*c} \tilde{X}_{d'}^{*d} = Z_{c'd'}^{*a'b'}, \tag{A.9}$$

while for the trilinear coefficient in φ , a new condition arises,

$$Z_{cd}^{ab} v_a \tilde{X}_b^{b'} \tilde{X}_{c'}^{*c} \tilde{X}_{d'}^{*d} = Z_{c'd'}^{*e b'} v_e \tag{A.10}$$

together with the CP-conjugate of this condition. (Similarly for lower powers of φ , only with more complicated conditions that involve combinations of Y and Zvv .)

Can there be \tilde{X} that are not in $\mathcal{X} = \{X\}$? First, as the condition for CP-conservation in the quartic terms in ϕ and φ are identical, and all CP transformations of ϕ were assumed to be known, there cannot be any more general CP transformations under which the whole potential is invariant than these known ones. In other words, all candidates for CP transformations of φ , \tilde{X} are in $\mathcal{X} = \{X\}$.

Using the invariance of Z on the l.h.s. of eq. (A.10) leads to

$$Z_{c'd'}^{*a'b'} X_{a'}^{\dagger a} v_a = Z_{c'd'}^{*a'b'} v_{a'} \tag{A.11}$$

from which with $X_{a'}^{\dagger a} v_a = (v_{a'})^*$ follows that the potential conserves CP if for at least one of the original X matrices holds that

$$v_a = X_{aa'} v^{*a'}. \tag{A.12}$$

Which conditions follow on VEVs or on relations between VEVs from an existing CP symmetry with unitary matrix X at high energy? Transform the potential with said CP transformation to obtain

$$V = Y_b^a X_{aa'} X^{*bb'} \phi^{*a'} \phi_{b'} + Z_{cd}^{ab} X_{aa'} X_{bb'} X^{*cc'} X^{*dd'} \phi^{*a'} \phi^{*b'} \phi_{c'} \phi_{d'}. \tag{A.13}$$

From this potential, minimisation conditions can be derived, and from

$$\frac{\partial V}{\partial \phi_i} = 0 \quad (\text{A.14})$$

follows

$$0 = Y_i^a X_{aa'} v^{*a'} + 2Z_{id}^{ab} X_{aa'} v^{*a'} X_{bb'} v^{*b'} X^{*dd'} v_{d'}. \quad (\text{A.15})$$

Comparing this with the CP-conjugate of the untransformed minimisation condition, i.e. the condition obtained by taking the derivative of V by ϕ^{*i} ,

$$0 = Y_i^a v^a + 2Z_{id}^{ab} v_a v_b v^{*d}, \quad (\text{A.16})$$

one finds that v_a and $u_a := X_{aa'} v^{*a'}$ fulfill exactly the same sets of equations. This means that the solution sets, $\{v_a\}$ and $\{X_{aa'} v^{*a'}\}$ are identical:

$$\{v_a\} = \{X_{aa'} v^{*a'}\} =: \mathcal{V}. \quad (\text{A.17})$$

Note that this does *not* mean that for every v_a holds that $v_a = X_{aa'} v^{*a'}$, but instead, as X is unitary, the only thing that can be said is that for symmetric X for every $v_a \in \mathcal{V}$ exactly one $u_a \in \mathcal{V}$ exists, such that

$$v_a = X_{aa'} u^{*a'} \quad (\text{A.18})$$

and longer chains for non-symmetric X matrices.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](#)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] ATLAS collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[arXiv:1207.7214](#)] [[INSPIRE](#)].
- [2] CMS collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[arXiv:1207.7235](#)] [[INSPIRE](#)].
- [3] J.F. Gunion, H.E. Haber, G.L. Kane and S. Dawson, *The Higgs Hunter’s Guide*, *Front. Phys.* **80** (2000) 1 [[INSPIRE](#)].
- [4] G.C. Branco, P.M. Ferreira, L. Lavoura, M.N. Rebelo, M. Sher and J.P. Silva, *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516** (2012) 1 [[arXiv:1106.0034](#)] [[INSPIRE](#)].
- [5] S. Chang, S.K. Kang, J.-P. Lee, K.Y. Lee, S.C. Park and J. Song, *Two Higgs doublet models for the LHC Higgs boson data at $\sqrt{s} = 7$ and 8 TeV*, *JHEP* **09** (2014) 101 [[arXiv:1310.3374](#)] [[INSPIRE](#)].
- [6] I.P. Ivanov, *Building and testing models with extended Higgs sectors*, *Prog. Part. Nucl. Phys.* **95** (2017) 160 [[arXiv:1702.03776](#)] [[INSPIRE](#)].

- [7] S. Davidson and H.E. Haber, *Basis-independent methods for the two-Higgs-doublet model*, *Phys. Rev. D* **72** (2005) 035004 [*Erratum ibid. D* **72** (2005) 099902] [[hep-ph/0504050](#)] [[INSPIRE](#)].
- [8] M. Maniatis, A. von Manteuffel and O. Nachtmann, *CP violation in the general two-Higgs-doublet model: A geometric view*, *Eur. Phys. J. C* **57** (2008) 719 [[arXiv:0707.3344](#)] [[INSPIRE](#)].
- [9] C.C. Nishi, *Physical parameters and basis transformations in the Two-Higgs-Doublet model*, *Phys. Rev. D* **77** (2008) 055009 [[arXiv:0712.4260](#)] [[INSPIRE](#)].
- [10] I.P. Ivanov, *Minkowski space structure of the Higgs potential in 2HDM. II. Minima, symmetries and topology*, *Phys. Rev. D* **77** (2008) 015017 [[arXiv:0710.3490](#)] [[INSPIRE](#)].
- [11] I.P. Ivanov, V. Keus and E. Vdovin, *Abelian symmetries in multi-Higgs-doublet models*, *J. Phys. A* **45** (2012) 215201 [[arXiv:1112.1660](#)] [[INSPIRE](#)].
- [12] I.P. Ivanov and C.C. Nishi, *Properties of the general NHDM. I. The orbit space*, *Phys. Rev. D* **82** (2010) 015014 [[arXiv:1004.1799](#)] [[INSPIRE](#)].
- [13] I.P. Ivanov, *Properties of the general NHDM. II. Higgs potential and its symmetries*, *JHEP* **07** (2010) 020 [[arXiv:1004.1802](#)] [[INSPIRE](#)].
- [14] R.N. Mohapatra and C.C. Nishi, *Absence of Spontaneous CP-violation in Multi-Higgs Doublet Extension of MSSM*, *Phys. Rev. D* **84** (2011) 095023 [[arXiv:1108.1769](#)] [[INSPIRE](#)].
- [15] I.P. Ivanov and E. Vdovin, *Classification of finite reparametrization symmetry groups in the three-Higgs-doublet model*, *Eur. Phys. J. C* **73** (2013) 2309 [[arXiv:1210.6553](#)] [[INSPIRE](#)].
- [16] G. Altarelli and F. Feruglio, *Discrete Flavor Symmetries and Models of Neutrino Mixing*, *Rev. Mod. Phys.* **82** (2010) 2701 [[arXiv:1002.0211](#)] [[INSPIRE](#)].
- [17] H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada and M. Tanimoto, *Non-Abelian Discrete Symmetries in Particle Physics*, *Prog. Theor. Phys. Suppl.* **183** (2010) 1 [[arXiv:1003.3552](#)] [[INSPIRE](#)].
- [18] W. Grimus and P.O. Ludl, *Finite flavour groups of fermions*, *J. Phys. A* **45** (2012) 233001 [[arXiv:1110.6376](#)] [[INSPIRE](#)].
- [19] S.F. King and C. Luhn, *Neutrino Mass and Mixing with Discrete Symmetry*, *Rept. Prog. Phys.* **76** (2013) 056201 [[arXiv:1301.1340](#)] [[INSPIRE](#)].
- [20] S.F. King, A. Merle, S. Morisi, Y. Shimizu and M. Tanimoto, *Neutrino Mass and Mixing: from Theory to Experiment*, *New J. Phys.* **16** (2014) 045018 [[arXiv:1402.4271](#)] [[INSPIRE](#)].
- [21] S.F. King, *Models of Neutrino Mass, Mixing and CP-violation*, *J. Phys. G* **42** (2015) 123001 [[arXiv:1510.02091](#)] [[INSPIRE](#)].
- [22] M. Kobayashi and T. Maskawa, *CP Violation in the Renormalizable Theory of Weak Interaction*, *Prog. Theor. Phys.* **49** (1973) 652 [[INSPIRE](#)].
- [23] C. Jarlskog, *Commutator of the Quark Mass Matrices in the Standard Electroweak Model and a Measure of Maximal CP-violation*, *Phys. Rev. Lett.* **55** (1985) 1039 [[INSPIRE](#)].
- [24] T.D. Lee, *A Theory of Spontaneous T Violation*, *Phys. Rev. D* **8** (1973) 1226 [[INSPIRE](#)].
- [25] J. Bernabeu, G.C. Branco and M. Gronau, *CP Restrictions On Quark Mass Matrices*, *Phys. Lett. B* **169** (1986) 243 [[INSPIRE](#)].

- [26] G.C. Branco, L. Lavoura and M.N. Rebelo, *Majorana Neutrinos and CP Violation in the Leptonic Sector*, *Phys. Lett. B* **180** (1986) 264 [[INSPIRE](#)].
- [27] G.C. Branco, I. de Medeiros Varzielas and S.F. King, *Invariant approach to CP in family symmetry models*, *Phys. Rev. D* **92** (2015) 036007 [[arXiv:1502.03105](#)] [[INSPIRE](#)].
- [28] G.C. Branco, I. de Medeiros Varzielas and S.F. King, *Invariant approach to \mathcal{CP} in unbroken $\Delta(27)$* , *Nucl. Phys. B* **899** (2015) 14 [[arXiv:1505.06165](#)] [[INSPIRE](#)].
- [29] V. Keus, S.F. King, S. Moretti and K. Yagyu, *CP Violating Two-Higgs-Doublet Model: Constraints and LHC Predictions*, *JHEP* **04** (2016) 048 [[arXiv:1510.04028](#)] [[INSPIRE](#)].
- [30] A. Mendez and A. Pomarol, *Signals of CP-violation in the Higgs sector*, *Phys. Lett. B* **272** (1991) 313 [[INSPIRE](#)].
- [31] L. Lavoura and J.P. Silva, *Fundamental CP-violating quantities in a $SU(2) \times U(1)$ model with many Higgs doublets*, *Phys. Rev. D* **50** (1994) 4619 [[hep-ph/9404276](#)] [[INSPIRE](#)].
- [32] F.J. Botella and J.P. Silva, *Jarlskog-like invariants for theories with scalars and fermions*, *Phys. Rev. D* **51** (1995) 3870 [[hep-ph/9411288](#)] [[INSPIRE](#)].
- [33] G.C. Branco, M.N. Rebelo and J.I. Silva-Marcos, *CP-odd invariants in models with several Higgs doublets*, *Phys. Lett. B* **614** (2005) 187 [[hep-ph/0502118](#)] [[INSPIRE](#)].
- [34] J.F. Gunion and H.E. Haber, *Conditions for CP-violation in the general two-Higgs-doublet model*, *Phys. Rev. D* **72** (2005) 095002 [[hep-ph/0506227](#)] [[INSPIRE](#)].
- [35] I. de Medeiros Varzielas, S.F. King, C. Luhn and T. Neder, *CP-odd invariants for multi-Higgs models: applications with discrete symmetry*, *Phys. Rev. D* **94** (2016) 056007 [[arXiv:1603.06942](#)] [[INSPIRE](#)].
- [36] I. de Medeiros Varzielas, S.F. King, C. Luhn and T. Neder, *Minima of multi-Higgs potentials with triplets of $\Delta(3n^2)$ and $\Delta(6n^2)$* , [arXiv:1704.06322](#) [[INSPIRE](#)].
- [37] G.C. Branco, J.M. Gerard and W. Grimus, *Geometrical T Violation*, *Phys. Lett. B* **136** (1984) 383 [[INSPIRE](#)].
- [38] I. de Medeiros Varzielas, *Geometrical CP-violation in multi-Higgs models*, *JHEP* **08** (2012) 055 [[arXiv:1205.3780](#)] [[INSPIRE](#)].
- [39] I.P. Ivanov and L. Lavoura, *Geometrical CP-violation in the N-Higgs-doublet model*, *Eur. Phys. J. C* **73** (2013) 2416 [[arXiv:1302.3656](#)] [[INSPIRE](#)].
- [40] I.P. Ivanov and C.C. Nishi, *Symmetry breaking patterns in 3HDM*, *JHEP* **01** (2015) 021 [[arXiv:1410.6139](#)] [[INSPIRE](#)].
- [41] D. Emmanuel-Costa, O.M. Ogreid, P. Osland and M.N. Rebelo, *Spontaneous symmetry breaking in the S_3 -symmetric scalar sector*, *JHEP* **02** (2016) 154 [*Erratum ibid. 08 (2016) 169*] [[arXiv:1601.04654](#)] [[INSPIRE](#)].
- [42] A. Aranda, I.P. Ivanov and E. Jiménez, *When the C in CP does not matter: anatomy of order-4 CP eigenstates and their Yukawa interactions*, *Phys. Rev. D* **95** (2017) 055010 [[arXiv:1608.08922](#)] [[INSPIRE](#)].
- [43] F. Feruglio, C. Hagedorn and R. Ziegler, *Lepton Mixing Parameters from Discrete and CP Symmetries*, *JHEP* **07** (2013) 027 [[arXiv:1211.5560](#)] [[INSPIRE](#)].
- [44] M. Holthausen, M. Lindner and M.A. Schmidt, *CP and Discrete Flavour Symmetries*, *JHEP* **04** (2013) 122 [[arXiv:1211.6953](#)] [[INSPIRE](#)].

- [45] M. Fallbacher and A. Trautner, *Symmetries of symmetries and geometrical CP-violation*, *Nucl. Phys. B* **894** (2015) 136 [[arXiv:1502.01829](https://arxiv.org/abs/1502.01829)] [[INSPIRE](#)].
- [46] I.P. Ivanov and J.P. Silva, *CP-conserving multi-Higgs model with irremovable complex coefficients*, *Phys. Rev. D* **93** (2016) 095014 [[arXiv:1512.09276](https://arxiv.org/abs/1512.09276)] [[INSPIRE](#)].
- [47] M. Ratz and A. Trautner, *CP violation with an unbroken CP transformation*, *JHEP* **02** (2017) 103 [[arXiv:1612.08984](https://arxiv.org/abs/1612.08984)] [[INSPIRE](#)].
- [48] C. Luhn, S. Nasri and P. Ramond, *The flavor group $\Delta(3n^2)$* , *J. Math. Phys.* **48** (2007) 073501 [[hep-th/0701188](https://arxiv.org/abs/hep-th/0701188)] [[INSPIRE](#)].
- [49] E. Ma and G. Rajasekaran, *Softly broken A_4 symmetry for nearly degenerate neutrino masses*, *Phys. Rev. D* **64** (2001) 113012 [[hep-ph/0106291](https://arxiv.org/abs/hep-ph/0106291)] [[INSPIRE](#)].
- [50] R. de Adelhart Toorop, F. Bazzocchi, L. Merlo and A. Paris, *Constraining Flavour Symmetries At The EW Scale I: The A_4 Higgs Potential*, *JHEP* **03** (2011) 035 [*Erratum ibid.* **01** (2013) 098] [[arXiv:1012.1791](https://arxiv.org/abs/1012.1791)] [[INSPIRE](#)].
- [51] R. de Adelhart Toorop, F. Bazzocchi, L. Merlo and A. Paris, *Constraining Flavour Symmetries At The EW Scale II: The Fermion Processes*, *JHEP* **03** (2011) 040 [[arXiv:1012.2091](https://arxiv.org/abs/1012.2091)] [[INSPIRE](#)].
- [52] I. de Medeiros Varzielas, S.F. King and G.G. Ross, *Neutrino tri-bi-maximal mixing from a non-Abelian discrete family symmetry*, *Phys. Lett. B* **648** (2007) 201 [[hep-ph/0607045](https://arxiv.org/abs/hep-ph/0607045)] [[INSPIRE](#)].
- [53] E. Ma, *Neutrino Mass Matrix from Delta(27) Symmetry*, *Mod. Phys. Lett. A* **21** (2006) 1917 [[hep-ph/0607056](https://arxiv.org/abs/hep-ph/0607056)] [[INSPIRE](#)].
- [54] C.C. Nishi, *Generalized CP symmetries in $\Delta(27)$ flavor models*, *Phys. Rev. D* **88** (2013) 033010 [[arXiv:1306.0877](https://arxiv.org/abs/1306.0877)] [[INSPIRE](#)].
- [55] I. de Medeiros Varzielas and D. Emmanuel-Costa, *Geometrical CP-violation*, *Phys. Rev. D* **84** (2011) 117901 [[arXiv:1106.5477](https://arxiv.org/abs/1106.5477)] [[INSPIRE](#)].
- [56] I. de Medeiros Varzielas, O. Fischer and V. Maurer, *A_4 symmetry at colliders and in the universe*, *JHEP* **08** (2015) 080 [[arXiv:1504.03955](https://arxiv.org/abs/1504.03955)] [[INSPIRE](#)].
- [57] I. de Medeiros Varzielas, D. Emmanuel-Costa and P. Leser, *Geometrical CP-violation from Non-Renormalisable Scalar Potentials*, *Phys. Lett. B* **716** (2012) 193 [[arXiv:1204.3633](https://arxiv.org/abs/1204.3633)] [[INSPIRE](#)].
- [58] G. Bhattacharyya, I. de Medeiros Varzielas and P. Leser, *A common origin of fermion mixing and geometrical CP-violation and its test through Higgs physics at the LHC*, *Phys. Rev. Lett.* **109** (2012) 241603 [[arXiv:1210.0545](https://arxiv.org/abs/1210.0545)] [[INSPIRE](#)].
- [59] G.C. Branco and I.P. Ivanov, *Group-theoretic restrictions on generation of CP-violation in multi-Higgs-doublet models*, *JHEP* **01** (2016) 116 [[arXiv:1511.02764](https://arxiv.org/abs/1511.02764)] [[INSPIRE](#)].