

Title: Both rare and common species make unique contributions to functional diversity in an ecosystem unaffected by human activities

Short title: Rarity and uniqueness at hydrothermal vents

Authors: Abbie S. A. Chapman¹, Verena Tunnicliffe², and Amanda E. Bates^{1,3}

Affiliations: ¹Ocean and Earth Science, University of Southampton Waterfront Campus, National Oceanography Centre, European Way, Southampton, UK SO14 3ZH. ²Department of Biology and School of Earth and Ocean Sciences, University of Victoria, P.O. Box 3020, Victoria, B.C., Canada V8W 3N5. ³Memorial University of Newfoundland, Department of Ocean Sciences, St. John's, NL, Canada, A1C 5S7.

Email: Abbie S. A. Chapman – asac1g09@soton.ac.uk
Verena Tunnicliffe – verenat@uvic.ca
Amanda E. Bates – bates.amanda@gmail.com

ACKNOWLEDGEMENTS

We would like to thank Maia Tsurumi and Jean Marcus for providing their sample data for use in this paper. We also thank S. Beaulieu, J. Copley, S. Hourdez, A. Metaxas, and A. Warén for their expert advice for trait scoring. Thanks also to T. Bird, R. Cooke, J. Lefcheck, S. Southgate, R. Stainthorp, C. Waldock, and the sFDvent working group, for valuable discussions that improved this work. Support from D. Chapman, L. Chapman, L. Chapman, and S. Chapman was also greatly appreciated. ASAC is a PhD candidate funded by the SPITFIRE Doctoral Training Partnership (supported by the Natural Environmental Research

Council, grant number: NE/L002531/1) and the University of Southampton. VT is sponsored through the Canada Research Chair Programme.

ABSTRACT

Aim Rare species typically contribute more to functional diversity than common species.

However, humans have altered the occupancy and abundance patterns of many species - the basis upon which we define ‘rarity’. Here, we use a globally unique dataset from hydrothermal vents – an untouched ecosystem – to test whether rare species over-contribute to functional diversity.

Location Juan de Fuca Ridge hydrothermal vent fields, Northeast Pacific Ocean.

Methods We first conduct a comprehensive review to set up expectations for the relative contributions of rare and common species to functional diversity. We then quantify the rarity and commonness of 37 vent species with relevant trait information to assess the relationship between rarity and functional distinctiveness - a measure of the uniqueness of the traits of a species relative to traits of coexisting species. Next, we randomly assemble communities to test whether rare species over-contribute to functional diversity in artificial assemblages ranging in species richness. Then, we test whether biotic interactions influence functional diversity contributions by comparing the observed contribution of each species to a null expectation. Finally, we identify traits driving functional distinctiveness using a distance-based redundancy analysis.

Results Across functional diversity metrics and species richness levels, we find that both rare and common species can contribute functional uniqueness. Some species always offer unique trait combinations, and these species host bacterial symbionts and provide habitat complexity. Moreover, we find that contributions of species to functional diversity may be influenced by biotic interactions.

Main Conclusions Our findings show that many common species make persistent, unique contributions to functional diversity. Thus, it is key to consider whether the abundance and occupancy of species have been reduced, relative to historical baselines, when interpreting the contributions of rare species to functional diversity. Our work highlights the importance of testing ecological theory in ecosystems unaffected by human activities for the conservation of biodiversity.

Keywords: conservation, diversity, functional distinctiveness, biological trait, human impact, hydrothermal vent, rarity, review, species richness, unique trait combination.

INTRODUCTION

Rare species, in having small populations, restricted geographic ranges, and, often, narrow environmental niches, are more vulnerable than common species to disturbance, environmental change, and competitive exclusion (Gaston, 1994; Rabinowitz, 1981). These rare species are often considered unique, endemic, and threatened - a combination that puts them at a higher risk of extinction than their common counterparts (Gaston, 1994, 2003; Gaston & Fuller, 2008; Hartley & Kunin, 2003; Margules & Pressey, 2000; Pimm, Jones, & Diamond, 1988). Yet, in possessing the characteristics that define rarity (low occupancy, abundance, and biomass), rare species may also contribute less than common species to ecosystem functioning processes (Grime, 1998; Smith & Knapp, 2003).

Functional traits are characteristics of a species affecting its contribution to the functioning of, and fitness within, an ecosystem (e.g., body size, trophic level). They can support empirical approaches when evaluating the respective roles of rare and common species in communities (Violle et al., 2017). Studies assessing the contributions of rare species to functional diversity

encompass numerous methods, scales, and systems, which we comprehensively review in Table S1.1, provided in the Supporting Information. Seven of the eight studies that focus on contributions of rare species to functional diversity (Table S1.1) showed that rare species contributed more to functional diversity than expected given small abundances, occupancies, or ranges. Thus, the majority of evidence suggests that rare species contribute disproportionately to communities by offering functional uniqueness and, therefore, supporting diverse ecosystem functions (Bracken & Low, 2012; Ellingsen, Hewitt, & Thrush, 2007; Jain et al., 2014; Leitão et al., 2016; Mouillot et al., 2013; Smith & Knapp, 2003). Rare species introduce functional redundancy by sharing traits with other species (a phenomenon that becomes more likely as species richness increases) and, consequently, may provide insurance and resilience for an ecosystem under different, future environmental conditions (Jain et al., 2014; Mouillot et al., 2013; Walker, Kinzig, & Langridge, 1999; Yachi & Loreau, 1999).

Most studies reporting over-contribution of rare species to functional diversity focus on ecosystems that humans have fundamentally altered (e.g., through fishing and aquaculture, tourism, and logging), including: coral reefs, rainforests, marine soft sediments, and alpine meadows (Mora et al., 2011; Vitousek, Mooney, Lubchenco, & Melillo, 1997; and see Table S1.1). By contrast, deep-sea hydrothermal vent communities thrive thousands of metres below the sea surface, without light, in warm, mineral-rich fluids that emerge from the seafloor, supporting diverse microbial communities – the primary producers in this system – and fauna highly adapted to these environments (e.g., Figure 1c; Ramirez-Llodra et al., 2010). Energy in deep-ocean hydrothermal environments is provided via chemosynthesis, where microorganisms use the reduced compounds in vent fluid and inorganic carbon to form organic matter (e.g., Cavanaugh, McKiness, Newton, & Stewart, 2006 and Ramirez-Llodra et al., 2010). Hydrothermal vents therefore provide a rare opportunity to assess the relative contributions of rare and common species to functional diversity in a chemosynthetic

environment with which only scientists have interacted, and thus human impacts are minimal, relative to terrestrial and shallow water systems (Ramirez-Llodra et al., 2011).

Here, we take advantage of a globally unique dataset of hydrothermal vent macrofaunal samples from one biogeographic region to test the hypothesis that rare species over-contribute to functional diversity. We focus on abundance, occupancy, and geographic extent as the main facets of rarity, in concordance with other rarity-oriented studies that use a functional trait approach, based on Rabinowitz' forms of rarity (Hartley & Kunin, 2003; Jain et al., 2014; Rabinowitz, 1981; Violle et al., 2017). We artificially assemble communities ranging in species richness from four to thirty-seven. For this range in species richness, we test the relative contribution of each species to local functional diversity in both artificially assembled and observed communities. Next, for the species that are functionally redundant with increasing species richness, we test whether species contributions differ from our null expectation as a result of community assembly processes. Finally, we identify the traits driving functional distinctiveness – a measure of the uniqueness of the trait values of a species, relative to the traits of all other species in a community (Grenié, Denelle, Tucker, Munoz, & Violle, 2017; Violle et al., 2017). Our work highlights differences in how species contribute to functional diversity in ecosystems that have not yet been reshaped by humans and, henceforth, a need to develop and test ecological theory in unaltered systems.

METHODS

Measuring rarity and commonness using abundance, occupancy, and geographic extent

This study focussed on invertebrate species assemblages ('communities') living more than 1,000 metres below the sea surface, at hydrothermal vents on basalt rocks along the Juan de Fuca Ridge tectonic plate boundary in the Northeast Pacific Ocean (Figure 1). Samples include sites in the Endeavour Marine Protected Area (Figure 1b). Sixty-three tubeworm

samples (44 tubeworm grabs, 10 suction samples, and nine samples compiled from both) were taken from 47 basalt-hosted sites between 1986 and 2001 (as described in detail in: Marcus, Tunnicliffe, & Butterfield, 2009; Tsurumi & Tunnicliffe 2001, 2003; Tunnicliffe et al., 1997; and Tunnicliffe 2000). Based on these samples, and for each of 37 species, we quantified the: (i) maximum relative abundance, to capture dominance potential and to discount sampling variability inherent in grab and suction samples (see Figure 1c); (ii) occupancy (the number of samples within which a species occurred); and (iii) geographic extent (the number of vent fields where each species occurred). We also computed a combined rarity index for each species (calculated as outlined in Leitão et al. (2016) but without log transformation), to enable cross-ecosystem comparability. The values for all four rarity metrics, for each of the 37 species, are provided in Table S1.2.

We include all macrofauna (≥ 1 mm in size) with species-level taxonomic identities and, thus, with fully assigned trait information (see species list in Table S1.2, verified using the World Register of Marine Species (Horton et al., 2017)). *Ridgeia piscesae* individuals (worms living inside tubes, aggregating to form bushes attached to the basalt rock substratum) were not included in this dataset, as they form the substratum for the sampled communities (e.g., see Figure 1c), in the same way that corals are often excluded from datasets as they provide habitat for reef fauna. The collector's curve in Figure 1a illustrates that sampling effort was sufficient to capture most of the Juan de Fuca Ridge vent species pool.

Species traits

We assembled a matrix of species traits based on a comprehensive literature search and expert knowledge of these species (see Table S1.2). We selected five species-level traits that should reflect adaptations to rarity or commonness, as well as the functional contribution of a species to an ecosystem: relative adult mobility, maximum (recorded) body size, trophic level

indicator, nutritional symbionts (present or absent), and forms a three-dimensional structure (yes or no). The justification and description of our traits are provided in Table S1.2.

Testing whether rare species over-contribute to functional diversity in vent ecosystems

To test the hypothesis that rare species over-contribute to functional diversity, we compared the functional distinctiveness of each of the 37 species to their rarity (abundance, occupancy, geographic extent, and a combined rarity index). Functional distinctiveness was computed using the ‘funrar’ and ‘cluster’ packages in R (Grenié et al., 2017; Maechler, Rousseeuw, Struyf, Hubert, & Hornik, 2017; R Core Team 2017; Violle et al., 2017). We also assessed whether functional distinctiveness was related to species richness, as functional diversity and taxonomic richness often have a strong linear relationship in other systems, detected using commonly used functional diversity metrics (e.g., functional richness – FRic, functional dispersion - FDis, and Rao’s quadratic entropy (Laliberté, Legendre, & Shipley, 2014); see Figure S1.1). We then created artificial communities, applying a random subsampling approach, to test the hypothesis that rare species over-contribute to functional diversity in vent communities ranging from four species (the minimum observed species richness on the local scale) to 37 species (the maximum, regional-scale species pool). We assembled 1,000 communities per level of species richness (four to thirty-seven), sampling randomly without replacement to create a presence-absence matrix. This matrix was used to compute the proportion of artificially assembled communities within which each species would contribute a novel unique trait combination (UTC - i.e. the combination of traits a species possesses is not found in other species it coexists with; see equation below); this proportion was plotted against species richness using a LOESS line of best fit alongside the proportion calculated using observed (sampled) data (displayed as points). Supporting R (R Core Team, 2017) script is provided in Appendix S2. For the full set of N species, we calculate the proportion of communities (artificially assembled or sampled) within which a species i ($i = 1, \dots, N$) makes a unique trait combination contribution. This proportion, $p_{species\ i}$, is the number of UTCs in a

community when species i is included (UTC_{total}) minus the number of UTCs when species i is excluded from the community (UTC_{-i}), divided by the total number of communities species i is present in, C :

$$p_{species\ i} = \frac{\Sigma (UTC_{total} - UTC_{-i})}{C}.$$

We used the Unique Trait Combination (UTC) metric to quantify species contributions to functional diversity to capture the redundancy and uniqueness of combinations of traits from a multifunctional perspective. This approach was selected as Mouillot et al. (2013) show that species with distinct trait combinations are more likely to support vulnerable ecosystem functions than species with commonly observed trait combinations. The UTC metric was computed using the ‘mvfd’ function of the ‘multirich’ R package (Keyel & Wiegand, 2016; R Core Team, 2017). This metric does not share limitations with the commonly used convex-hull based ‘FRic’ functional richness metric (of particular relevance, given the low richness of vent communities) (Cornwell, Schilke, & Ackerly, 2006; Laliberté et al., 2014; Villéger, Mason, & Mouillot, 2008; Schmera, Podani & Erős, 2009); the UTC metric can accommodate the limited number of continuous measurements available for relatively inaccessible and expensive to sample deep-sea hydrothermal vent species by incorporating categorical trait values (Keyel & Wiegand, 2016).

Species were placed in one of three groups (‘always unique’, ‘redundant with richness’, or ‘rapidly redundant’) according to the shape of the relationship between species richness and the proportion of communities within which their trait combination was unique in 1,000 randomly assembled artificial communities. We tested whether community assembly processes and biotic interactions could be shaping the functional uniqueness of the species that we identified as becoming increasingly redundant with richness (for species observed in at least 10 samples – an arbitrary threshold selected to minimize sampling effort bias). We compared the proportion of sampled communities in which each of the species in the

‘redundant with richness’ group observed in at least 10 samples contributed a UTC to a null expectation established using our randomly assembled community data. We used a binomial test to determine whether the proportion contributed by sampled species differed significantly from the proportion species would be expected to contribute based on the outcomes of the randomly assembled samples. Richness levels were binned for the tests as, despite using one of the world’s most complete hydrothermal vent sample datasets, we had insufficient samples to conduct the test for each richness level. Bins are highlighted as dotted boxes in Figure 4 (with further details and binomial test results presented in Table S1.3). We selected the minimum probability value for the null expectation in each bin to be conservative.

Finally, to assess the trait similarity among the species in the ‘always unique’ group, we used a distance-based redundancy analysis (db-RDA; Legendre & Anderson 1999). We adapted the db-RDA with an input of a species by trait matrix as our explanatory matrix and a species by functional distinctiveness matrix as our response. Given the nature of our functional trait data (i.e., some semi-quantitative traits), we used the Gower distance metric for the db-RDA (Gower & Legendre 1986; Laliberté & Legendre 2010). The db-RDA is appropriate because our data did not meet assumptions of Euclidean distances and normality (Legendre & Anderson 1999; Legendre & Legendre 1998).

RESULTS

In 63 hydrothermal vent community samples, 27 species were relatively rare (if we consider a species with a rarity index value of < 0.5 to be rare, as the index is scaled from 0 to 1; Figure S1.2) and 10 relatively common (> 0.5 rarity index value; Figure S1.2). In these 63 samples, some of the most rare and common species contributed to functional distinctiveness. All three measures of rarity, as well as a combined rarity index, hold the same flat relationship with distinctiveness (Figure 2); therefore, this outcome does not support a hypothesis of

increasing contribution with rarity (Figure 2a; Table S1.4 contains the outputs of the linear models presented in Figure 2b-e). Nonetheless, as species richness increased in sampled communities, functional distinctiveness decreased (Figure 3), supporting the expected pattern of increasing functional redundancy with richness; when number of species increases, the probability of a new species contributing to functional distinctiveness decreases because there is a higher chance that the traits of a new species already exist in another species in the community. These results were not simply an outcome of trait selection or scoring methodology, as tested and discussed in Appendices S3 and S4.

Given the strong relationship between species richness and functional distinctiveness, we further tested whether rare and common species both contribute to functional distinctiveness across a range of species richness levels, spanning beyond the maximum sample richness (21) to that of the regional pool (37). We accounted for the contributions made by species given known co-occurrences, rather than contributions to the overall regional pool - which can include species that, in reality, never co-occur. Considering the independent contributions of each species (Figure 4), we found that 12 species, including eight rare and two very common species, contributed unique trait combinations at all levels of richness ('always unique'). Twelve species contributed uniqueness at low levels of richness but became increasingly redundant with richness ('redundant with richness'), while 13 became rapidly redundant at lower richness ('rapidly redundant'). For the most part, the groupings highlighted in Figure 4 comprised a mix of taxonomic classes, but all the scaleworm polychaetes (in the phylum Annelida) fell only within the 'rapidly redundant' group. Each group is described in more detail in Table S1.5.

The proportion of communities in which eight, well-sampled species in the 'redundant with richness' group made unique contributions to the assemblages in which they were sampled was compared to the random expectation. The worms, *Paralvinella pandorae* and

Amphisamytha carldarei, and snails, *Depressigyra globulus* and *Provanna variabilis*, contributed a unique trait combination (UTC) in fewer communities than would be expected by chance (Figure 4 and Table S1.3). By contrast, the worm, *Paralvinella palmiformis*, and sea spider, *Sericosura verenae*, contributed a UTC in more communities than expected (Figure 4 and Table S1.3). The contributions of two worms, *Protomystides verenae* and *Parougia wolfi*, did not differ significantly from random (see Table S1.3).

Two key traits emerged as driving functional distinctiveness in the ‘always unique’ group: hosting nutritional symbionts and forming a three-dimensional structure (Figure 5, and see Table S1.6). These two traits also shaped the functional distinctiveness of species in the ‘redundant with richness’ and ‘rapidly redundant’ groups, influencing the relative distances between all species (Tables S1.5 and S1.6). The direction of influence differs in these traits for species in the ‘rapidly redundant’ group, though, as these species do not form 3D structures or have nutritional symbionts (Table S1.5).

DISCUSSION

Here, we show that rare species do not contribute significantly more to the functional distinctiveness of vent communities than more common species. This finding is supported at all levels of species richness (from the minimum sampled richness – four species – to the 37 species regional pool). Our results contradict those of most research into rarity-functional diversity relationships to date, which, instead, demonstrate that rare species over-contribute to functional diversity (a review of the research on this topic is presented in Table S1.1). For example, in coral reefs, tropical rainforests, and alpine meadows, the most distinct trait combinations are supported by rare species (Mouillot et al., 2013). In addition, in a removal experiment conducted on a rocky shore community, rare species had bottom-up influences on the diversity and abundance of consumers (Bracken & Low, 2012).

269

270 Despite diverging from the expectation set in other trait-based studies, our findings are
271 supported by ecological theory. If rare species are specialists, adapted to specific
272 environmental conditions (hence their small geographic range, for example), we expect rarer
273 species to have traits that reflect this specialisation. These specialist traits would be dissimilar
274 to the traits of species occupying other habitats and niches, enabling specialist rare species to
275 persist with low occupancy and abundance (Gaston 1994; Rabinowitz 1981). At the same
276 time, common species are expected to thrive in a wider range of habitats and environmental
277 conditions, and have more generalist traits enabling them to do so; yet, common species must
278 also possess unique traits (or combinations of traits) to successfully outcompete other species
279 for space and resources, attain high abundances, maintain broad geographic ranges, and
280 occupy many habitats (Gaston, 2010; Gaston, 2011; Tilman, 1999).

281

282 While our results are supported by ecological theory, there remain several explanations as to
283 why our results differ from previous studies. These include: (1) sampling method (i.e., at
284 vents, remotely operated equipment enables the retrieval of intact assemblages comprising
285 diverse taxa, rather than the taxon-specific ‘community’ data collected for some studies
286 described in Table S1.1); (2) trait selection (e.g., see Lefcheck, Bastazini, & Griffin, 2014)
287 and scoring methodology (as the number of modalities per trait can influence trait space and,
288 thus, indices computed using a multidimensional trait space volume; Lefcheck et al., 2014);
289 (3) species richness of the study system, which can shape functional diversity by influencing
290 the potential for functional redundancy; and (4) the natural and anthropogenic processes
291 affecting ecosystem stability and function. We discuss these possible explanations in more
292 detail in Appendix S3 and argue (with support in Appendix S4) why each was unlikely to
293 explain the novel result that common species can contribute to functional diversity.

294

After considering alternative explanations, we propose that our results may, instead, differ from the expectation set in shallow marine and terrestrial ecosystems because vents are unique in being a relatively untouched ecosystem on Earth. Species abundance and occupancy patterns at vents have not yet been markedly altered by human activities (Ramirez-Llodra et al., 2011). For example, Juan de Fuca Ridge predator species *Buccinum thermophilum*, a snail, and *Nereis piscesae*, a worm, are redundant (in terms of the traits we selected) with other species in nearly every community in which they occur. It is likely that relatively large predatory species like these would be unique if incorporated in trait-based studies of ecosystems within which humans have removed many large, mobile predators (e.g., by hunting). Conversely, in Juan de Fuca vent communities, we find many relatively large predators that are mobile and carnivorous (Bergquist et al., 2007; Kelly, Metaxas, & Butterfield, 2007), making these traits - and thus these snail and worm species – functionally redundant in these systems.

At vents and in other remote ecosystems, rarity and commonness are the result of various abiotic and biotic processes shaping abundance, occupancy, and geographic range extent. In contrast, in ecosystems such as coral reefs and tropical rainforests, human activities have caused species losses and gains, and have fundamentally altered species abundances, occupancies, and geographic range dimensions (e.g., see Inger et al., 2015). Thus, when we study ‘rare’ species in human-altered systems, we might be including species that were once common but appear low in abundance, extent, or occupancy at the time of study as a result of human action (Gaston, 2008; Gaston & Fuller, 2008); this would affect our perceived contributions of ‘rare’ species to functional diversity. For example, *Dipturus batis* (common skate) is a demersal marine species that has been reduced in number by human activities (Gaston & Fuller, 2008), and would be considered ‘rare’ if studied today, as opposed to several decades prior.

It could therefore be argued that ecological research more widely would benefit from studying undisturbed systems like hydrothermal vents, as results such as those presented here have conservation and management implications. Rare species are often the focus of conservation strategies because they may be more prone to extinction (e.g., a species low in abundance could be lost altogether with the loss of several individuals) (Gaston, 1994; Kruckeberg & Rabinowitz, 1985; Margules & Pressey, 2000). However, we have shown that common species also influence functional diversity in vent systems; thus, common species losses may also have important implications for ecosystem functioning and stability (e.g. see Lyons, Brigham, Traut, & Schwartz, 2005 and Tilman et al., 1997 for discussions of relationships between functional diversity and ecosystem functioning). For instance, if a common species facilitates a rare species, reducing the number of individuals of this common species will likely have a knock-on effect on the rare species (Gaston, 2011; Lindenmayer et al., 2011). For hydrothermal vent communities specifically, as deep-sea mining is likely to affect these presently undisturbed assemblages, conservation approaches must move rapidly to assess effects of species loss (Van Dover et al., 2017). Given our findings at the Juan de Fuca Ridge, we propose that conservation planning incorporating functional assessments that include the roles of common species may be particularly effective at vents, as well as in other ecosystems, where rare species are more often the focus at present.

Furthermore, our work advances understanding of the ecology of hydrothermal vent communities in identifying features of these communities that clarify assemblage structures and key species roles (summarised in Table S1.7). We identify two traits underpinning contributions to relative functional distinctiveness: ‘possessing nutritional symbionts’ and ‘forming a three-dimensional (3D) structure’. Possessing nutritional, bacterial symbionts enables a host species to access chemical energy and accumulate biomass in the same way that corals rely on zooxanthellae to reach high biomass in a low-productivity tropical ocean. Common species hosting nutritional symbionts will have direct access to primary productivity

(Cavanaugh et al., 2006; Stewart, Newton, & Cavanaugh, 2005) and, as space can be limited at vents, this often translates to increased abundance and biomass. At the same time, rare species might also benefit from hosting nutritional symbionts, to thrive with limited resources or compete with common species, especially if the symbiont-host association requires conditions that differ from competitive dominants.

In addition to, or instead of, having nutritional symbionts, for some vent species that have limited access to tolerable conditions in high temperature or fluid flux gradients, the ability to form 3D structures may enable them to modulate flow and/or access specific habitats (Bates, Lee, Tunnicliffe, & Lamare, 2010; Bates, Tunnicliffe, & Lee, 2005; Kelly & Metaxas, 2008; Tsurumi & Tunnicliffe, 2003). Common *Lepetodrilus fucensis* limpets access and alter fluid flow patterns by stacking vertically (see Bates et al., 2005), like epiphytes on forest trees. Such species that form 3D structures likely also play a key role as ecosystem engineers, increasing habitat complexity and facilitating colonisation by other species. In other ecosystems, common species are typically the engineers (Gaston, 2011). Also common are the limpets and tube-forming polychaetes that form 3D structures at Juan de Fuca Ridge vents, enabling other species to colonise the augmented surface area. Their structures can modify local fluid flow patterns and provide surfaces for microbial colonisation, thereby increasing food resources for grazers. Additionally, the functionally distinct rare species that form 3D structures potentially act as ‘cornerstone species’ in small and limited niches, playing an important role in structuring the local space (as defined and observed in rocky intertidal communities in Bracken & Low 2012). For example, the most distinct rare species (bivalves *Calyptogena starobogatovi* and *Idas washingtonius*) usually inhabit other chemosynthetic environments; but, in our tubeworm bush samples, their shells provide stable habitats for settlers of other species, even when the bivalves themselves are no longer alive.

Coexistence theory helps to explain the unique trait combination (UTC) contributions of these vent species. Kraft et al. (2008) tested coexistence theories (e.g., neutral and niche-based models) in Amazonian forest trees using functional traits. They compared measures of community-trait structure to a null expectation of random assembly, with habitat filtering deemed to be taking place if the range of observed trait values was smaller than that of randomly assembled communities. In our study, the bristle worm *Paralvinella pandorae* offered a redundant trait combination more times than expected based on random assembly; the opposite was true for its congener, *Paralvinella palmiformis*. Indeed, these two alvinellid worms are competitors, with *Paralvinella pandorae* the inferior competitor because of its narrower trophic and space requirements (Levesque, Juniper, & Marcus, 2003; Marcus et al., 2009; Tunnicliffe et al., 1997). Meanwhile, the worm, *Amphisamytha carldarei*, and snails, *Depressigyra globulus* and *Provanna variabilis*, make unique contributions to significantly fewer communities than we would expect by chance. The contributions of these species might, therefore, be affected by habitat filtering (in addition to, or instead of, competitive interactions). In contrast, the sea spider, *Sericosura verenae*, makes a unique contribution in more communities than in the null expectation, so its contributions might be affected by niche differentiation.

We used a novel approach to test whether the proportion of communities each species contributed a UTC to was significantly different from a random expectation. Trait-based studies of rarity and functional diversity have previously used a regional pool based approach, thereby incorporating species that, in reality, never co-occur. Our method may, in cases with representative sampling, provide a tool to identify species that are limited in functional distinctiveness by other co-occurring species. Thus, our approach could be used to test for community assembly mechanisms – a common goal in trait-based ecology (Ackerly & Cornwell, 2007; de Bello, 2012; de Bello et al., 2012) – alongside analyses of other influential processes, such as larval dispersal, colonisation, and suitability of the abiotic environment

(Kraft et al., 2015; MacArthur & Levins, 1967). Our approach could also prove a useful ecological tool for investigating more general relationships between rarity, species richness, functional distinctiveness, and co-occurrence patterns in communities.

In conclusion, our understanding of many ecological processes, biodiversity patterns, and resilience are based on systems that are significantly altered by human activities. Using sampled and artificial (randomly assembled) hydrothermal vent communities, we show that rare and common species both offer functional distinctiveness, with contributions of species shaped by traits important for chemosynthesis, ecosystem engineering, and physico-chemical tolerance. Furthermore, functional uniqueness can be constrained by biotic interactions, such as competition, habitat filtering, and niche differentiation. Our findings offer new perspectives on rarity, commonness, distinctiveness, and redundancy; thus, we suggest that hydrothermal vent habitats and other relatively untouched environments offer unique windows into ecology, conservation, and biodiversity theory. Ultimately, here we highlight a need to test ecological hypotheses in Earth's remaining untouched systems, to facilitate our ecological understanding of the systems that we, as humans, have already altered.

REFERENCES

- Ackerly, D. D., & Cornwell, W. K. (2007). A trait-based approach to community assembly: partitioning of species trait values into within- and among-community components. *Ecology Letters*, 10, 135-145. doi:10.1111/j.1461-0248.2006.01006.x
- Bates, A. E., Lee, R. W., Tunnicliffe, V., & Lamare, M. D. (2010). Deep-sea hydrothermal vent animals seek cool fluids in a highly variable thermal environment. *Nature Communications*, 1(14), 1-6. doi:10.1038/ncomms1014

429 Bates, A. E., Tunnicliffe, V., & Lee, R. W. (2005). Role of thermal conditions in habitat
 430 selection by hydrothermal vent gastropods. *Marine Ecology Progress Series*, 305, 1-15.
 431

432 Beaulieu, S. (2015). *InterRidge Global Database of Active Submarine Hydrothermal Vent*
 433 *Fields: prepared for InterRidge, Version 3.4.*
 434

435 Bergquist, D.C., Eckner, J.T., Urcuyo, Istvan A., Cordes, E.E., Hourdez, S., Macko, S.A., &
 436 Fisher, C. R. (2007). Using stable isotopes and quantitative community characteristics to
 437 determine a local hydrothermal vent food web. *Marine Ecology Progress Series*, 330, 49-65.
 438

439 Bracken, M. E. S., & Low, N. H. N. (2012). Realistic losses of rare species disproportionately
 440 impact higher trophic levels. *Ecology Letters*, 15, 461-467. doi:10.1111/j.1461-
 441 0248.2012.01758.x
 442

443 Cavanaugh, C. M., McKiness, Z. P., Newton, I. L. G., & Stewart, F. J. (2006). Marine
 444 Chemosynthetic Symbioses *The Prokaryotes* (3rd Edition ed., pp. 475-507). New York:
 445 Springer.
 446

447 Chadwick, W. W., Clague, D. A., Embley, R. W., Perfit, M. R., Butterfield, D. A., Caress, D.
 448 W., . . . Bobbitt, A. M. (2013). The 1998 eruption of Axial Seamount: New insights on
 449 submarine lava flow emplacement from high-resolution mapping. *Geochemistry, Geophysics,*
 450 *Geosystems*, 14, 3939-3968. doi:10.1002/ggge.20202
 451

452 Coleman, B. D., Mares, M. A., Willis, M. R., & Hsieh, Y. (1982). Randomness, area and
 453 species richness. *Ecology*, 63, 1121-1133.
 454

455 Cornwell, W. K., Schilck, D. W., & Ackerly, D. D. (2006). A trait-based test for habitat
 456 filtering: convex hull volume. *Ecology*, 87(6), 1465-1471.

457

458 de Bello, F., Price, J.N., Münkemüller, T., Liira, J., Zobel, M., Thuiller, W., . . . Pärtel, M.
 459 (2012). Functional species pool framework to test for biotic effects on community assembly.
 460 *Ecology*, 93, 2263-2273.

461

462 de Bello, F. (2012). The quest for trait convergence and divergence in community assembly:
 463 are null-models the magic wand? *Global Ecology and Biogeography*, 21, 312-317.
 464 doi:10.1111/j.1466-8238.2011.00682.x

465

466 Ellingsen, K. E., Hewitt, J. E., & Thrush, S. F. (2007). Rare species, habitat diversity and
 467 functional redundancy in marine benthos. *Journal of Sea Research*, 58, 291-301.
 468 doi:10.1016/j.seares.2007.10.001

469

470 Esri, GEBCO, NOAA, Geographic, National, DeLorme, HERE, . . . and other contributors.
 471 (2012). *Ocean Basemap*.

472

473 Gaston, K. J. (1994). *Rarity* (1st ed.). Netherlands: Springer.

474

475 Gaston, K. J. (2003). *The Structure and Dynamics of Geographic Ranges*. Oxford, UK:
 476 Oxford University Press.

477

478 Gaston, K. J. (2010). Valuing Common Species. *Science*, 327(5962), 154-155.

479

480 Gaston, K. J., & Fuller, R. A. (2008). Commonness, population depletion and conservation
 481 biology. *Trends in Ecology and Evolution*, 23(1), 14-19. doi:10.1016/j.tree.2007.11.001

482

483 Gaston, Kevin J. (2008). Biodiversity and extinction: the importance of being common.

484 *Progress in Physical Geography*, 32(1), 73-79. doi:10.1177/0309133308089499

485

486 Gaston, Kevin J. (2011). Common Ecology. *BioScience*, 61, 354-362.

487 doi:10.1525/bio.2011.61.5.4

488

489 Gower, J. C., & Legendre, P. (1986). Metric and Euclidean Properties of Dissimilarity

490 Coefficients. *Journal of Classification*, 3, 5-48.

491

492 Grenié, M., Denelle, P., & Tucker, C. (2017). funrar: Functional Rarity Indices Computation

493 (Version R package version 1.0.3): R package version 1.0.3. Retrieved from [http://CRAN.R-](http://CRAN.R-project.org/package=funrar)

494 [project.org/package=funrar](http://CRAN.R-project.org/package=funrar)

495

496 Grenié, M., Denelle, P., Tucker, C. M., Munoz, F., & Violle, C. (2017). funrar: An R package

497 to characterize functional rarity. *Diversity and Distributions*, 1-7. doi:10.1111/ddi.12629

498

499 Grime, J. P. (1998). Benefits of plant diversity to ecosystems: immediate, filter and founder

500 effects. *Journal of Ecology*, 86, 902-910.

501

502 Hartley, S., & Kunin, W. E. (2003). Scale Dependence of Rarity, Extinction Risk, and

503 Conservation Priority. *Conservation Biology*, 17, 1559-1570.

504

505 Horton, T., Kroh, A., Bailly, N., Boury-Esnault, N., Brandão, S. N., Costello, M. J., . . .

506 Zeidler, W. (2017). World Register of Marine Species (WoRMS). Retrieved 2017-07-28,

507 from WoRMS Editorial Board <http://www.marinespecies.org>

508

509 Inger, R., Gregory, R., Duffy, J. P., Stott, I., Voříšek, P., & Gaston, K. J. (2015). Common
 510 European birds are declining rapidly while less abundant species' numbers are rising. *Ecology*
 511 *Letters*, 18, 28-36. doi:10.1111/ele.12387
 512
 513 Jain, M., Flynn, D. F., Prager, C. M., Hart, G. M., Devan, C. M., Ahrestani, F. S., . . . Naeem,
 514 S. (2014). The importance of rare species: a trait-based assessment of rare species
 515 contributions to functional diversity and possible ecosystem function in tall-grass prairies.
 516 *Ecology and Evolution*, 4, 104-112. doi:10.1002/ece3.915
 517
 518 Kelly, N., & Metaxas, A. (2008). Diversity of invertebrate colonists on simple and complex
 519 substrates at hydrothermal vents on the Juan de Fuca Ridge. *Aquatic Biology*, 3, 271-281.
 520 doi:10.3354/ab00085
 521
 522 Kelly, N., Metaxas, A., & Butterfield, D. (2007). Spatial and temporal patterns of
 523 colonization by deep-sea hydrothermal vent invertebrates on the Juan de Fuca Ridge, NE
 524 Pacific. *Aquatic Biology*, 1, 1-16. doi:10.3354/ab00001
 525
 526 Keyel, A. C., & Wiegand, K. (2016). Validating the use of unique trait combinations for
 527 measuring multivariate functional richness. *Methods in Ecology and Evolution*, 1-8.
 528 doi:10.1111/2041-210x.12558
 529
 530 Kraft, N. J. B., Valencia, R., & Ackerly, D. D. (2008). Functional Traits and Niche-Based
 531 Tree Community Assembly in an Amazonian Forest. *Science*, 322, 580-582.
 532
 533 Kraft, N. J. B., Adler, P. B., Godoy, O., James, E. C., Fuller, S., Levine, J. M., & Fox, J.
 534 (2015). Community assembly, coexistence and the environmental filtering metaphor.
 535 *Functional Ecology*, 29, 592-599. doi:10.1111/1365-2435.12345

536

537 Kruckeberg, A. R., & Rabinowitz, D. (1985). Biological aspects of endemism in higher
538 plants. *Annual Review of Ecology and Systematics*, 16, 447-479.

539

540 Laliberté, E., & Legendre, P. (2010). A distance-based framework for measuring functional
541 diversity from multiple traits. *Ecology*, 91, 299-305.

542

543 Laliberté, E., Legendre, P., & Shipley, B. (2014). FD: measuring functional diversity from
544 multiple traits, and other tools for functional ecology (Version R package version 1.0-12): R
545 package version 1.0-12.

546

547 Lefcheck, J. S., Bastazini, V. A. G., & Griffin, J. N. (2014). Choosing and using multiple
548 traits in functional diversity research. *Environmental Conservation*, 1-4.

549 doi:10.1017/s0376892914000307

550

551 Legendre, P., & Anderson, M. J. (1999). Distance-based redundancy analysis: Testing
552 multispecies responses in multifactorial ecological experiments. *Ecological Monographs*,
553 69(1), 1-24.

554

555 Legendre, P., & Legendre, L. (1998). *Numerical Ecology* (Second English ed. Vol. 24).
556 Amsterdam: Elsevier Science B. V.

557

558 Leitão, R. P., Zuanon, J., Villéger, S., Williams, S. E., Baraloto, C., Fortunel, C., . . .

559 Mouillot, D. (2016). Rare species contribute disproportionately to the functional structure of
560 species assemblages. *Proceedings of the Royal Society B*, 283, 1-9.

561 doi:10.1098/rspb.2016.0084

562

Levesque, C., Juniper, S. K., & Marcus, J. (2003). Food resource partitioning and competition among alvinellid polychaetes of Juan de Fuca Ridge hydrothermal vents. *Marine Ecology Progress Series*, 246, 173-182.

Lindenmayer, D. B., Wood, J. T., McBurney, L., MacGregor, C., Youngentob, K., & Banks, S. C. (2011). How to make a common species rare: A case against conservation complacency. *Biological Conservation*, 144, 1663-1672. doi:10.1016/j.biocon.2011.02.022

Lyons, K. G., Brigham, C. A., Traut, B. H., & Schwartz, M. W. (2005). Rare Species and Ecosystem Functioning. *Conservation Biology*, 19, 1019-1024. doi:10.1111/j.1523-1739.2005.00106.x

MacArthur, R., & Levins, R. (1967). The Limiting Similarity, Convergence, and Divergence of Coexisting Species. *The American Naturalist*, 101, 377-385.

Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M., & Hornik, K. (2017). cluster: Cluster Analysis Basics and Extensions (Version R package version 2.0.6.): R package version 2.0.6.

Marcus, J., Tunnicliffe, V., & Butterfield, D. A. (2009). Post-eruption succession of macrofaunal communities at diffuse flow hydrothermal vents on Axial Volcano, Juan de Fuca Ridge, Northeast Pacific. *Deep-Sea Research II*, 56, 1586-1598. doi:10.1016/j.dsr2.2009.05.004

Margules, C.R., & Pressey, Robert L. (2000). Systematic conservation planning. *Nature*, 405, 243-253.

Mora, C., Aburto-Oropeza, O., Ayala Bocos, A., Ayotte, P. M., Banks, S., Bauman, A. G., . . .
 Zapata, F. A. (2011). Global Human Footprint on the Linkage between Biodiversity and
 Ecosystem Functioning in Reef Fishes. *PLoS Biology*, 9(4), e1000606.
 doi:10.1371/journal.pbio.1000606

Mouillot, D., Bellwood, D. R., Baraloto, C., Chave, J., Galzin, R., Harmelin-Vivien, M., . . .
 Thuiller, W. (2013). Rare Species Support Vulnerable Functions in High-Diversity
 Ecosystems. *PLoS Biology*, 11(5), e1001569. doi:10.1371/journal.pbio.1001569

Newman, K. R., Nedimović, M. R., Canales, J. P., & Carbotte, S. M. (2011). Evolution of
 seismic layer 2B across the Juan de Fuca Ridge from hydrophone streamer 2-D travelttime
 tomography. *Geochemistry, Geophysics, Geosystems*, 12(5), 1-24. doi:10.1029/2010gc003462

Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., ...
 Wagner, H. (2017). vegan: Community Ecology Package. (Version R package version 2.4-3):
 R package version 2.4-3.

Pimm, S. L., Jones, H. L., & Diamond, J. (1988). On the Risk of Extinction. *The American
 Naturalist*, 132, 757-785.

R Core Team. (2017). R: A Language and Environment for Statistical Computing. Vienna,
 Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org>

Rabinowitz, D. (1981). Seven forms of rarity. In H. Synge (Ed.), *The Biological Aspects of
 Rare Plant Conservation* (pp. 205-217): John Wiley & Sons Ltd.

Ramirez-Llodra, E., Brandt, A., Danovaro, R., De Mol, B., Escobar, E., German, C. R., . . .
 Vecchione, M. (2010). Deep, diverse and definitely different: unique attributes of the world's
 largest ecosystem. *Biogeosciences*, 7, 2851-2899. doi:10.5194/bg-7-2851-2010

Ramirez-Llodra, E., Tyler, P. A., Baker, M. C., Bergstad, O. A., Clark, M. R., Escobar, E., . . .
 Van Dover, C. L. (2011). Man and the Last Great Wilderness: Human Impact on the Deep
 Sea. *PLoS ONE*, 6(7), e22588. doi:10.1371/journal.pone.0022588

Schmera, D., Podani, J., & Erős, T. (2009). Measuring the contribution of community
 members to functional diversity. *Oikos*, 118, 961-971. doi:10.1111/j.1600-0706.2009.17076.x

Smith, M. D., & Knapp, A. K. (2003). Dominant species maintain ecosystem function with
 non-random species loss. *Ecology Letters*, 6, 509-517.

Stewart, F. J., Newton, I. L., & Cavanaugh, C. M. (2005). Chemosynthetic endosymbioses:
 adaptations to oxic-anoxic interfaces. *Trends in Microbiology*, 13, 439-448.
 doi:10.1016/j.tim.2005.07.007

Tilman, D. (1999). The ecological consequences of changes in biodiversity: a search for
 general principles. *Ecology*, 80, 1455-1474.

Tilman, D., Knops, J., Wedin, D., Reich, P., Ritchie, M., & Siemann, E. (1997). The
 Influence of Functional Diversity and Composition on Ecosystem Processes. *Science*, 277,
 1300-1302.

640 Tsurumi, M., & Tunnicliffe, V. (2001). Characteristics of a hydrothermal vent assemblage on
641 a volcanically active segment of Juan de Fuca Ridge, northeast Pacific. *Canadian Journal of*
642 *Fisheries and Aquatic Sciences*, 58, 530-542. doi:10.1139/cjfas-58-3-530
643

644 Tsurumi, M., & Tunnicliffe, V. (2003). Tubeworm-associated communities at hydrothermal
645 vents on the Juan de Fuca Ridge, northeast Pacific. *Deep-sea Research I*, 50, 611-629.
646 doi:10.1016/s0967-0637(03)00039-6
647

648 Tunnicliffe, V. (2000). *A documentation of biodiversity characteristics of the hydrothermal*
649 *vent assemblages at High Rise Ventfield, Endeavour Segment, Juan de Fuca Ridge*. Contract
650 Report to Department of Fisheries and Oceans, 53pp.
651

652 Tunnicliffe, V., Embley, R.W., Holden, J.F., Butterfield, David A., Massoth, Gary J., &
653 Juniper, S. K. (1997). Biological colonization of new hydrothermal vents following an
654 eruption on Juan de Fuca Ridge. *Deep-sea Research I*, 44, 1627-1644.
655

656 Van Dover, C. L., Ardron, J. A., Escobar, E., Gianni, M., Gjerde, K. M., Jaeckel, A., . . .
657 Weaver, P. P. E. (2017). Biodiversity loss from deep-sea mining. *Nature Geoscience*, 10, 464-
658 465. doi:10.1038/ngeo2983
659

660 VanderBeek, B. P., Toomey, D. R., Hooft, E. E. E., & Wilcock, W. S. D. (2016).
661 Segmentation of mid-ocean ridges attributed to oblique mantle divergence. *Nature*
662 *Geoscience*, 9, 636-642. doi:10.1038/ngeo2745
663

664 Villéger, S., Mason, N. W. H., & Mouillot, D. (2008). New multidimensional functional
665 diversity indices for a multifaceted framework in functional ecology. *Ecology*, 89, 2290-2301.
666 doi:10.1890/07-1206.1

667

668 Violle, C., Thuiller, W., Mouquet, N., Munoz, F., Kraft, N. J. B., Cadotte, M. W., . . .

669 Mouillot, D. (2017). Functional Rarity: The Ecology of Outliers. *Trends in Ecology and*

670 *Evolution*, 32, 356-367. doi:10.1016/j.tree.2017.02.002

671

672 Vitousek, Peter M., Mooney, H. A., Lubchenco, J., & Melillo, J. M. (1997). Human

673 Domination of Earth's Ecosystems. *Science*, 277, 494-499.

674

675 Walker, B., Kinzig, A., & Langridge, J. (1999). Plant Attribute Diversity, Resilience, and

676 Ecosystem Function: The Nature and Significance of Dominant and Minor Species.

677 *Ecosystems*, 2, 95-113.

678

679 Yachi, S., & Loreau, M. (1999). Biodiversity and ecosystem productivity in a fluctuating

680 environment: The insurance hypothesis. *Proceedings of the National Academy of Sciences*

681 *USA*, 96, 1463-1468.

682

683 **BIOSKETCH**

684

685 Abbie Chapman is a PhD Candidate at the University of Southampton

686 (www.abbiechapman.com). She is researching the biodiversity of hydrothermal vents across

687 the globe using a functional trait approach and is co-PI (with Amanda Bates) of the ‘sFDvent’

688 working group, building the first global functional trait database for vent species. Verena

689 Tunnicliffe is a Canada Research Chair in Deep Ocean Research based at the University of

690 Victoria. Verena focuses on the characteristics that enable populations to survive in extreme

691 habitats and primarily researches deep-sea ecosystems such as hydrothermal vents and

692 submarine volcanoes. Amanda Bates' research quantifies biodiversity in time and space, and

693 relates emergent patterns to environmental drivers and physiological tolerances of species.

694 Her aim is to transform physiological data into predictive tools that incorporate both exposure

and species' sensitivities to global change drivers. In doing so, she generates novel theories and approaches for managing and conserving marine living resources.

Author contributions: ASAC and AEB designed the study; VT provided sample data; all authors contributed trait data; ASAC and AEB undertook the analyses, with outcomes verified by VT; ASAC drafted the manuscript; and all authors contributed substantially to revisions.

DATA ACCESSIBILITY

Should the manuscript be accepted, the data supporting the results will be archived in an appropriate public repository such as Dryad or Figshare and the data DOI will be included at the end of the article.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

Appendix S1: Tables: S1.1 - review of trait-based studies assessing the contributions of rare species to functional diversity; S1.3 - binomial test outcomes; S1.4 - linear model parameter estimates associated with Figure 2; S1.5 - species traits for the simulated groups shown in Figure 4; S1.6 – identification of traits affecting relative functional distinctiveness shown in Figure 5; S1.7 - features of hydrothermal vent communities identified using a trait-based approach; Figure S1.1 - relationship between functional diversity and taxonomic richness, using four metrics of functional diversity; and Figure S1.2 – relative commonness of each of the 37 species in this study.

Table S1.2: Species traits matrix used in this study with a supporting key and rationale, references, and species list.

Appendix S2: R script for computing the proportion of communities each species contributes a UTC in, using observed (sample) data and artificially assembled communities, and a distance-based redundancy analysis using traits.

Appendix S3: Discussion of alternative explanations for the differences between our results and those of previous studies, conducted in other systems (e.g., those reviewed in Table S1.1).

Appendix S4: Assessing the robustness of methods and results presented in our manuscript, with supporting Figures (S4.2 – S4.6).

Table S4.9: Species traits matrix used in Appendix S4, with supporting key and references.

FIGURES

Figure 1: Overview of sampling effort and collection methods used to obtain macrofaunal data: (a) collector's curve (with 95% confidence intervals shown in grey) for the regional species pool compiled by combining all species sampled and using the Coleman et al. (1982) method; (b) location map, placing the hydrothermal vents ('sites') sampled along the Juan de Fuca Ridge (with 'n' the number of vents sampled at a particular vent field (labelled in bold), preceding the years of sampling, and the ridge marked with dashed lines); (c) image showing tubeworm grab sampling with a submersible claw - a method commonly used to sample communities hosted by tubeworms like *Ridgeia piscesae*. The suction sample hose supplements the grab to retrieve animals on the substratum under the bushes, and mobile species escaping the grab. Image about 120 cm across. Bathymetry in (a) was sourced from Esri et al. (2012), vent field locations from sample records and the InterRidge Vents Database

(Beaulieu, 2015) (note that I.D. is International District), and the Juan de Fuca Ridge was drawn using information in Newman et al. (2011), Chadwick et al. (2013), and VanderBeek et al. (2016).

Figure 2: The relationship between rarity and functional distinctiveness in Juan de Fuca Ridge vent communities. (a) The expected linear relationship between rarity and distinctiveness, whereby more common species offer less functional distinctiveness than rare species. (b) The observed relationship between maximum relative abundance of each species and functional distinctiveness relative to all other species in the community. (c) The relationship between occupancy (number of samples within which the species occurs) and functional distinctiveness for each species. (d) The relationship between geographic extent (number of vent fields within which the species was observed) and functional distinctiveness of each species. (e) The relationship between the Rarity Index (per Leitão et al. (2016) without log transformation) and functional distinctiveness. Note that the relationships shown in (b), (c), (d), and (e) are relatively flat, contrary to the expectation presented in (a), suggesting that rare and common species contribute functional distinctiveness. Model outputs to support panels (b), (c), (d) and (e) are provided in Table S1.4.

Figure 3: The relationship between taxonomic richness per sample and mean (across all species present in a sample) functional distinctiveness per sample, as computed using the ‘distinctiveness’ function of the ‘funrar’ functional rarity package (Grenié et al., 2017). The linear relationship was fitted using a linear model (formula: mean FDistinct ~ richness) and is a significant linear relationship, with slope -0.005 and p-value <0.05 (adjusted R-squared value: 0.4157; F-statistic: 37.29 on 1 and 50 degrees of freedom).

Figure 4: The proportion of communities within which each species makes a unique trait combination (UTC) contribution. Each panel in this figure shows the proportion of

communities to which a species (named at the top of each panel) makes a unique contribution (y axis), relative to species richness (x axis). Solid lines in each panel depict the relationships identified using artificial, randomly assembled communities of four to thirty-seven species in richness (mean values based on communities that were randomly assembled 1,000 times per level of richness - see Methods). Circles in each panel are observations from sample data, with circle size relative to the number of samples with the given result. Dotted boxes, shown on some central panels, encompass the richness bins used in the binomial tests to see whether the observed UTC contribution made by a species significantly differed from the null, random expectation (shown in the artificial community line). Each panel is shaded according to the groups listed in the legend that are assigned based on the shape of the relationship between species richness and UTC contributions revealed in the randomly assembled communities. Cartoon inserts illustrate the taxonomic group of each species, as outlined in the Legend. Species names have been shortened, but are given in full in Table S1.2. The species presented on this figure combine to represent the regional species pool – the 37 taxa identified to species level for basalt samples from the Juan de Fuca Ridge.

Figure 5: Distance-based redundancy analysis (db-RDA) of Gower distances showing the functional distinctiveness of each of the species relative to one another and the potential traits driving distinctiveness differences (labelled in bold next to arrows, with trait names shortened as follows: NUT – Nutritional Symbionts, FOR – Forms a 3D Structure, BOD –Maximum Body Size (mm), MOB – Relative Adult Mobility, and TRO – Trophic Level Indicator). ‘Nutritional Symbionts’ and ‘Forms a 3D Structure’ were significant traits in the db-RDA model, based on an ANOVA by terms with 200 permutations (a permutation test used to assess the significance of constraints for each trait separately (Oksanen et al., 2017); see Table S1.6). Functional distinctiveness is colour coded, with the two most functionally distinct species (*Lepetodrilus fucensis* and *Calypptogena starobogatovi*) shown in black and the point colours becoming lighter with decreasing functional distinctiveness. The species

803 labelled on this plot belong to the ‘always unique’ group (see Figure 4), though the functional
804 distinctiveness of all 37 species was driven by the same traits. Species names shortened in
805 this Figure are: ALVIS – *Munidopsis alvisca*, AXIAL – *Amphiduropsis axialensis*, ENDEA –
806 *Pardalisca endeavouri*, FUCEN – *Lepetodrilus fucensis*, GLOBO – *Ophryotrocha*
807 *globopalpata*, GLOBU – *Hyalogyrina globularis*, RIDGE – *Sphaerosyllis ridgensis*, STARO
808 – *Calyptogena starobogatovi*, SULFI – *Paralvinella sulfincola*, VALEN – *Thermanemertes*
809 *valens*, VENTI – *Sericosura venticola*, and WASHI – *Idas washingtonius*.









