




Contents lists available at ScienceDirect

Australian Critical Care

journal homepage: www.elsevier.com/locate/aucc

Review Paper

Effect of organisational factors on the variation in incidence of delirium in intensive care unit patients: A systematic review and meta-regression analysis

Paul Rood, RN, MSc, CCRN ^{a, b, *}
 Getty Huisman - de Waal, PhD ^b
 Hester Vermeulen, RN, PhD ^b
 Lisette Schoonhoven, PhD, FEANS ^{c, d}
 Peter Pickkers, MD, PhD ^a
 Mark van den Boogaard, RN, PhD, CCRN ^a

^a Department of Intensive Care Medicine, Radboud University Medical Center, Nijmegen, The Netherlands

^b Radboud Institute for Health Sciences, Radboud University Medical Center, Nijmegen, The Netherlands

^c Faculty of Health Sciences, University of Southampton, Southampton, United Kingdom

^d National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care (Wessex), United Kingdom

ARTICLE INFORMATION

Article history:

Received 21 June 2017

Received in revised form

5 January 2018

Accepted 1 February 2018

Keywords:

Delirium
 Critical care
 Incidence
 Organisational factors
 Systematic review
 Meta-analysis

ABSTRACT

Background: Delirium occurs frequently in intensive care unit (ICU) patients and is associated with numerous deleterious outcomes. There is a large variation in reported delirium occurrence rates, ranging from 4% to 89%. Apart from patient and treatment-related factors, organisational factors could influence delirium incidence, but this is currently unknown.

Objective: To systematically review delirium incidence and determine whether or not organisational factors may contribute to the observed delirium incidence in adult ICU patients.

Methods: Systematic review of prospective cohort studies reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement. Included articles were independently assessed by two researchers. Quality of the articles was determined using the Strengthening of Reporting of Observational Studies in Epidemiology checklist. Subsequently, apart from patient characteristics, a meta-regression analysis was performed on available organisational factors, including hospital type, screening method and screening frequency.

Data Sources: PubMed, Embase, CINAHL, and Cochrane Library databases were searched from inception to 27 January 2017, without language limitation.

Results: A total of 9357 articles were found, of which 19 articles met the inclusion criteria and were considered as true delirium incidence studies. The articles were of good methodological quality (median [interquartile range] 32/38 [30–35] points), published between 2005 and 2016, originated from 17 countries. A total of 9867 ICU patients were included. The incidence rate of delirium varied between 4% and 55%, with a mean \pm standard deviation of $29 \pm 14\%$. Data relating to three organisational factors were included in the studies, but they were not significantly associated with the reported delirium incidence: hospital type (p 0.48), assessment methods (p 0.41), and screening frequency (p 0.28).

Conclusions: The mean incidence of delirium in the ICU was 29%. The organisational factors found including methods of delirium assessment, screening frequency, and hospital type were not related to the reported ICU delirium incidence.

© 2018 Australian College of Critical Care Nurses Ltd. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author at: Department of Intensive Care Medicine, Radboud Institute for Health Sciences, Radboud University Medical Center, P.O. Box 9101, Internal post 710, 6500 HB, Nijmegen, The Netherlands.

E-mail address: paul.rood@radboudumc.nl (P. Rood).

<https://doi.org/10.1016/j.aucc.2018.02.002>

1036-7314/© 2018 Australian College of Critical Care Nurses Ltd. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Delirium is a serious problem in the intensive care unit (ICU), as it is associated with numerous short-term adverse events such as

increased duration of mechanical ventilation¹ and length of stay.^{2–4} Also, delirium is associated with long-term adverse effects such as persistent cognitive decline^{5,6} and increased 6-month mortality.⁷ Delirium occurs frequently in the ICU; a recent meta-analysis found that delirium occurs in approximately one-third of ICU patients,⁸ but a large variation is reported. Depending whether incidence, defined as a new onset of delirium after ICU admission, or prevalence, which also includes patients who were already delirious before ICU admission, is estimated,⁹ occurrence rates vary between 4% and 89%.^{10,11} The reason for this large variability is currently not fully understood.

From a research perspective, the collection of implementation data is essential for program evaluations,¹² and multilevel causal factors are described to impact implementation outcomes, such as patient, provider, and organisational factors.¹³ First, several patient-related risk factors are clearly associated with the development of delirium, such as respiratory failure, a history of cognitive impairments, and urgent ICU admission.^{14–17} Although patient-related risk factors account for a considerable part of the variability, they do not explain all of it.^{8,18–24} Second, delirium incidence is also influenced by the provider through the ICU treatment. The current international delirium guideline emphasises that timely management of the cause is essential to reduce the delirium incidence, severity, and duration. Also aspects of care such as sedation management and early mobilisation are considered to influence the development of delirium. Third, organisational factors may contribute to delirium. These include the selection of the screening method (currently either the Confusion Assessment Method for the Intensive Care Unit [CAM-ICU] or the Intensive Care Delirium Screening Checklist [ICDSC] are recommended, of which the performance is limited,^{25,26} even when performed by experts²⁷), daily screening frequency (as delirium may not be detected if screening frequency is too low because of its fluctuating course during the day), and hospital type (as variations in different types of hospitals may account for differences). Also, strategies used for education of staff and screening compliance as the measured delirium incidence may differ because of to fluctuating course of delirium and the adequate recognition of delirium symptoms.

Recent reviews have provided insight into patient- and provider-related risk factors,^{8,15,28,29} but did not address organisational factors. As there is empirical support that the level of implementation may affect the outcome in prevention programs,¹² we feel that these should be clarified so they can be incorporated when collecting and analysing data. As the incidence of delirium can be influenced by ICU treatment, it is important to discern incidence from prevalence figures. Subsequently, the association of organisational factors with ICU delirium incidence needs to be clarified.

Therefore, the aim of our study was to systematically review delirium incidence and determine whether or not organisational factors may contribute to the observed delirium incidence in adult ICU patients.

2. Material and methods

A systematic review and meta-regression analysis was performed, according to the Cochrane Handbook for systematic reviews³⁰ and reported according to the steps of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement.³¹ The selection of articles, data extraction, and methodological quality assessment were performed independently by two reviewers (PR and MvdB). Included articles were assessed for methodological quality using the “Strengthening of Reporting of Observational Studies in Epidemiology” (STROBE) statement.³²

2.1. Eligibility criteria

Prospective cohort studies were included in which the ICU delirium incidence was determined in adult medical, surgical, or mixed ICU patients (18 years or older). Delirium incidence was defined as a new onset of delirium during ICU admission without delirium before ICU admission, diagnosed by a positive delirium screening using a validated screening tool or as a reported medical diagnosis. Articles were excluded when only delirium prevalence (i.e. delirium before ICU admission was not an exclusion criterion) was reported, when the focus was on other ICU subpopulations, or if no full-text article was available. To ensure the quality of reported delirium incidence rates, only prospective cohort studies designed to study delirium incidence were included.

2.2. Search

A systematic literature search was conducted in PubMed, Embase, CINAHL and Cochrane Library databases. References of included articles were searched for additional relevant articles. Databases were searched through combining “Delirium,” “Intensive Care Unit,” and “Incidence”, as well as relevant synonyms. The complete strategy is provided in an online supplemental. Languages were not limited during the search. Articles published from database inception until January 27th, 2017, were included. Data management was performed using Endnote X8 (Thomson Reuters).

2.3. Study selection

Eligibility of articles was independently assessed by screening title and abstract by two researchers (PR and MB) using the inclusion criteria. Eligible articles were obtained in full-text by the first author; if not possible, we planned to contact the authors of the article. Articles that were irretrievable would have been excluded from further analysis. Both turned out not to be necessary, as all articles could be obtained. After independent full-text assessment of the eligible articles, discrepancies were discussed. In case of disagreement, a third researcher (HV) was asked to make a final judgement.

2.4. Methodological quality

The selected articles were screened for methodological quality using the “STROBE statement”.³² Because an index test was not found in most of the reviewed articles, this tool was deemed most suitable for critical quality appraisal of the included articles. It allowed for structured and transparent assessment of bias and applicability of primary diagnostic accuracy articles. Each domain was assessed in terms of risk of bias, and a rating per item was given (2 = present, 1 = partially present, 0 = not present, NA. = not applicable). Afterwards sum scores were calculated. A maximum score of 38 points could be obtained. The lower limit for inclusion in the review was set at 70% of achievable points.

2.5. Data collection

Data extraction was performed by the primary researcher (PR) using a standardised data extraction form containing patients characteristics and treatment and organisational factors. For treatment factors, we aimed to gather data on delirium treatment algorithms, as well as analgesia, sedative, and sleep enhancement strategies.^{19,33} For organisational factors, articles were searched for factors regarding country and continent, hospital type, implementation strategies, staff knowledge, motivation, and screening compliance.^{34,35}

2.6. Statistics

Extracted measures were reported using descriptive statistics: Patient characteristics, ICU treatment, and organisational factors as available in all included articles were extracted. To determine the association of organisational factors with the reported delirium incidences, a meta-regression analysis was performed. Owing to heterogeneity and a limited number of included articles, we performed a univariate analysis using a random effects model. Mean \pm standard deviation, or median (interquartile range) are reported, depending on the normality of the distribution of data. Data were analysed using IBM SPSS Statistics 22 and R statistics 3.2.4.³⁶ The standard error of the incidence was calculated for each study as $(\text{Incidence} - (1 - \text{Incidence})/N)^2$. A *p*-value of ≤ 0.05 was considered statistically significant.

3. Results

3.1. Study selection

The search yielded a total of 9357 articles, of which after removal of duplicates 8178 unique articles remained. Based on analysing title and abstract, 8060 articles were excluded, leaving 118 articles. Another 71 articles were excluded as only a conference abstract was available, and 28 articles were excluded because incidence rates were not mentioned in the full-text article. Finally, in total, 19 articles were included in this meta-regression analysis (Fig. 1).

A total of 9867 ICU patients were included in the articles. Sample size varied from 80 to 4450 patients. All articles were published

between 2005 and 2016. Nine studies were conducted in a general ICU population, four only included medical and six only surgical patients (Table 1).

3.2. Organisational risk factors

In 16 articles, a screening tool validated for the ICU was used (eleven studies used the CAM-ICU,³ five used the ICDSC³⁷). In two studies, a general delirium screening tool (Delirium rating scale-revised-98,³⁸ Nursing Delirium Symptom Checklist³⁹) was used, and in one study the formal diagnostic criteria according to the Diagnostic Statistical Manual IV–Third Revision⁴⁰ were applied. Nine articles originated from Europe, six from Asia, two from Northern America, one from Southern America, and one from Oceania. Sixteen studies were performed in one or more university hospitals and three were performed in general hospitals. Nine articles reported the number of ICU beds, which varied between 8 and 33 beds. Daily screening frequency varied between 1 and 3 times a day, the time window in which the first, follow-up, and last delirium screening was performed varied greatly. In 10 studies, the delirium screening was performed by ICU nurses, in five by ICU physicians, in two studies by dedicated research personnel, and in two studies by psychiatrists (Table 2).

3.3. Methodological quality

Of the 19 included articles, STROBE quality assessment yielded a median score of 32^{30–35} of the possible 38 points. All articles obtained a quality score over 70% and were considered of sufficient methodological quality for inclusion in the analysis. Most elements were reported in the majority of the articles, except for bias (mentioned in 13% of articles) and funding (mentioned in 50% of articles). An overview of the individual assessment scores is provided in the online supplement 2.

3.4. Results of individual articles

The mean delirium incidence rate was $29 \pm 14\%$; the lowest incidence found was 4%, and the highest was 55%. The samples that were included were heterogeneous. Overall, most studies included older patients, who originated from medical, surgical, or mixed ICU populations, and more males than females were included. A broad variation was observed in admission type, severity of illness, length of stay, and mortality rates (Table 1).

3.5. Associations of ICU treatment and organisational factors

Using a meta-regression analysis, we assessed the association of organisational factors to reported delirium incidence rates. No significant differences in delirium incidence were found between university hospitals (median 28%^{21–35} in 9463 patients) and non-university hospitals 33%^{20–46} in 404 patients ($p = 0.48$). Related to the screening method used to detect delirium, five different methods were described. In 11 studies that used the CAM-ICU,⁴¹ a median incidence of 24%^{21–31} was found in 3767 patients. In five studies that used the ICDSC,³⁷ a median incidence of 32%^{11–42} was found in 5654 patients. Three non-ICU specific methods were used: (i) one study used the Delirium rating scale-revised-98³⁸ and found an incidence of 24% in 140 patients; (ii) one study used a psychiatric interview according to the Diagnostic Statistical Manual IV–Third Revision⁴⁰ criteria and found an incidence of 55% in 142 patients; and (iii) one study used the Nursing Delirium Symptom Checklist³⁹ and found an incidence of 44.5% in 164 patients. The incidence of delirium did not differ significantly between screening methods ($p = 0.28$). Finally, the influence of the daily screening




Fig. 1. Flow diagram of study selection.

Table 1
Patient-related characteristics of included articles.

Reference	Delirium incidence	N	Age	Population	Male (N, %)	Urgent admission (N, %)	Severity of illness score	Mechanically ventilated (N, %)	LOS-ICU	LOS hospital	Mortality rate (N, %)
Ayllon Garrido et al. ⁶⁰	41.3%	112	63 (±18)	Mixed	67 (60)	–	APACHE II 14 (±7)	47 (42)	3 [2–8]	–	–
Bilge et al. ⁶¹	18.4%	250	60 (±13)	Surgical	D 21 (46) ND 121 (59)	–	ASA scor. (I/II/III/IV) D 0/10/26/10 ND 4/107/83/10	–	–	–	–
Kanova et al. ⁶²	31.2%	125	64 (±21)	Mixed	87 (61%)	–	APACHE II 12 (±14)	61 (43)	3 (±6)	–	–
Lahariya et al. ⁶³	9.3%	309	D 62 (±14); ND 57 (±13)	Medical	D 51 (63) ND 30 (37)	D 78 (96) ND 205(90)	APACHE II D 15 (±5) ND 7 (±3)	–	D 2 (±2) ND 2 (±1)	–	D 22 (27) ND 2 (0.8)
Limpawattana et al. ⁶⁴	22.2%	99	D 79 (±7); ND 75 (±7)	Medical	D 24 (40) ND 22 (55)	–	APACHE II D 24 (±1) ND 19 (±1)	D 30 (68) ND 18 (33)	–	–	–
Mori et al. ¹⁴	46.3%	149	D 65 [53–80]; ND 54 [30–68]	Mixed	D 42 (61) ND 49 (61)	–	SAPS III D 58 (16) ND 45(17)	–	D 11 [0–27] ND 4 [0–8]	–	D 16 (23) ND 9 (11)
Norkiene et al. ⁶⁵	13.3%	87	D 68 (±10); ND 65 (±11)	Surgical	–	0 (0)	EUROSCORE II D 2.4 (±1.4) ND 2.0 (±1.4)	87 (100)	D 5 (±2) ND 3 (±1)	–	–
Ouimet et al. ⁶⁶	31.8%	820	D 65 (±14); ND 63 (±15)	Mixed	D 147 (61) ND 303 (58)	–	APACHE II D 18 (±8) ND 14 (±8)	645 (79)	D 12 (±12) ND 4 (±4)	D 18 (±16) ND 13 (±19)	D 49 (20) ND 54 (10)
Page et al. ⁶⁷	31.0%	80	D 70 [56–76]; ND 73 [60–77]	Mixed	D 16 (72) ND 34 (69)	D 22 (45) ND 27 (55)	APACHE II D 21 [17–30] ND 15 [11–20]	D 17 (63) ND 10 (37)	–	–	D 8 (36) ND 5 (10)
Peterson et al. ⁶⁸	24.4%	614	53 (±18)	Medical	309 (50)	–	APACHE II 20 (±9)	298 (49)	–	–	–
Pipankaporn et al. ¹¹	3.6%	4450	D 65 (±16); ND 62 (±17)	Surgical	D 102 (63) ND 2505 (58)	D 79 (66) ND 987 (29)	APACHE II D 16 [12–23] ND 10 [7–15]	D 141 (87) ND 2635 (62)	D 8 [5–19] ND 2 [1–4]	D 22 [14–34] ND 15 [9–26]	D 38 (24) ND 349 (8)
Roberts et al. ⁶⁹	45.0%	185	D 64; ND 60	Mixed	D 52 (62) ND 56 (55)	–	APACHE II D 21 ND 17	–	D 10 (±7) ND 7 (±5)	D 23(±16) ND 19 (±12)	D 10 (12) ND 10 (10)
Sabol et al. ¹⁷	20.8%	250	65 (±10)	Surgical	171 (68)	–	EUROSCORE II 2.6 (±2.7)	250 (100)	5 (±3)	11 (±7)	4 (2)
Sharma et al. ⁹	24.4%	140	D 50 (±19); ND 37 (±14)	Mixed	D 43 (57) ND 29 (45)	–	APACHE II D 20 (±6) ND 15 (±6)	D 60 (80) ND 34 (52)	D 8 [5–13] ND 5 [3–6]	–	D 23 (31) ND 0 (0)
Shi et al. ⁷⁰	44.5%	164	D 73 (±8); ND 66 (±11)	Surgical	D 44 (60) ND 54 (59)	–	APACHE II D 13 (±4) ND 7 (±4)	D 32 (44) ND 32 (35)	D 1 [1–14] ND 1 [1–6]	D 18 [7–74] ND 13 [3–48]	D 5 (7) ND 1 (1)
Smulter et al. ⁴²	54.9%	142	D 77 (±5); ND 76 (±4)	Surgical	D 54 (69) ND 38 (59)	–	–	142 (100)	D 1.1 (±1.1) ND 0.76 (±0.23)	–	–
Svenningsen et al. ⁷¹	40.2%	139	D 63; ND 64	Mixed	78 (56)	–	–	–	D 21 ND 14	–	D 8 (20) ND 4 (7)
Van den Boogaard et al. ⁷²	25.5%	1613	61 (±14)	Mixed	792 (66)	526 (44)	APACHE II D 18 (±6) ND 13 (±5)	D 363 (88) ND 903 (75)	D 6 [2–13] ND 1 [1–2]	D 20 [10–20] ND 7 [5–14]	D ND 40 (3)
Woien et al. ⁷³	23.4%	139	55 (±15)	Medical	87 (63)	–	SAPS II 42 (±16)	138 (99)	5 (range 1–53)	–	25 (18)

D = delirious patients, ND = non-delirious patients; LOS = length of stay; ICU = intensive care unit.

Illness severity scores: APACHE-II score = Acute Physiology and Chronic Health Evaluation-II score⁷⁵, ASA Score = American Society of Anaesthesiologists score⁷⁴, SAPS II/III = Simplified Acute Physiology Score⁷⁷, EUROSCORE II = European System for Cardiac Operative Risk Evaluation II⁷⁶.

Data are expressed as number and percentage (N/%), mean (±standard deviation) or median [interquartile range], unless mentioned otherwise.

Table 2
Organisational characteristics of included articles.

Reference	Incidence	Year	N	Type center	Beds	Country	Screening method	Daily screening frequency	Screening by	Comments	Quality
Ayllon Garrido et al. ⁶⁰	41.3%	2007	112	UMC	8	Spain	ICDSC	1	Nurses	Only patients admitted >72 h were included. Patients were assessed once. Patients who required post-operative mechanical ventilation were excluded.	34/38
Bilge et al. ⁶¹	18.4%	2015	250	UMC	–	Turkey	CAM-ICU	1	Physicians		36/38
Kanova et al. ⁶²	31.2%	2015	125	UMC	–	Czech Republic	CAM-ICU	1	Physicians	Prevalence 29%, incidence defined as first positive score >24 after admission. First CAM-ICU was performed within 48 h after admission. Readmitted patients and patients/family who refused cooperation were excluded from analysis. After 5 days the screening, frequency was reduced.	30/38
Lahariya et al. ⁶³	9.3%	2014	309	UMC	22	India	CAM-ICU; DSM-IV-TR	1	Clinicians—psychiatrists		31/38
Limpawattana et al. ⁶⁴	22.2%	2016	99	UMC	–	Thailand	CAM-ICU	1	Nurses	33/38	
Mori et al. ¹⁴	46.3%	2016	149	UMC	17	Brazil	CAM-ICU	2	Nurses	34/38	
Norkiene et al. ⁶⁵	13.3%	2013	87	UMC	–	Lithuania	ICDSC	3	Clinicians	Screening was started 24 h after surgery, repeated every 8 h during ICU stay (max 5 days).	30/38
Ouimet et al. ⁶⁶	31.8%	2007	820	UMC	16	Canada	ICDSC	1	Clinicians	Prevalence 39.6%. Range between SICUs: 0–13.9%. Exclusion: <24 h stay, neurosurgical and cardiac. Inclusion of 185/2568 patients admitted due to exclusion when ICU LOS <36 h or hospital stay >96 h prior to ICU admission.	29/38
Page et al. ⁶⁷	31.0%	2009	80	Gen	8	United Kingdom	CAM-ICU	2	Nurses		27/38
Peterson et al. ⁶⁸	24.4%	2006	614	UMC	14	USA	CAM-ICU	2	Nurses		35/38
Pipanmekaporn et al. ¹¹	3.6%	2015	4450	UMC	–	Thailand	ICDSC	1	Nurses		31/38
Roberts et al. ⁶⁹	45.0%	2005	185	Gen	–	Australia/ New Zealand	ICDSC	2	Researchers or trained professionals	31/38	
Sabol et al. ¹⁷	20.8%	2015	250	UMC	–	Slovakia	CAM-ICU	2	Nurses	Delirium assessment was performed until day 7 post-operative or the disappearance of delirious symptoms for 2 consecutive days. Patients were assessed pre-operatively and on day 1 and day 4 post-operatively. Patients were assessed >48 h after ICU admission until discharged.	32/38
Sharma et al. ⁹	24.4%	2012	140	UMC	8	India	DRS-R-98	1	Psychiatrists		35/38
Shi et al. ⁷⁰	44.5%	2010	164	UMC	–	China	Nu-DESC	3	Physicians and nurses		29/38
Smulter et al. ⁴²	54.9%	2013	142	UMC	–	Sweden	DSM-IV-TR	1	Trained research nurses	32/38	
Svenningsen et al. ⁷¹	40.2%	2011	139	UMC	28	Denmark	CAM-ICU	2	Nurses	28/38	
Van den Boogaard et al. ⁷²	25.5%	2012	1613	UMC	33	The Netherlands	CAM-ICU	3	Nurses	35/38	
Woien et al. ⁷³	23.4%	2013	139	Gen	–	Norway	CAM-ICU	3	Nurses	35/38	

LOS = length of stay; ICU = intensive care unit; CAM-ICU = Confusion Assessment Method for the Intensive Care Unit; ICDSC = Intensive Care Delirium Screening Checklist; DRS-R-98 = Delirium rating scale-revised-98; DSM-IV-TR = Diagnostic Statistical Manual IV–Third Revision; Nu-DESC = Nursing Delirium Symptom Checklist; SICUs = Surgical Intensive Care Units.

frequency was assessed. A median incidence of 26%^{14–38} was found in 6447 patients in nine studies in which patients were screened for delirium once daily, 34%^{23–46} in 1417 patients in six studies which screened twice daily, and 27%^{6–47} in 2003 patients in the studies which screened three times daily. No significant differences were observed ($p = 0.41$) (Table 3).

4. Discussion

In this meta-analysis we found 19 good quality articles (median [interquartile range] score 32 [30–35] of possible 38 points), published between 2005 and 2016, originated from 17 countries. A total of 9867 ICU patients were included. All articles specifically described delirium incidence. A mean ICU delirium incidence of 29 ± 14 was found, ranging from 4%^{11–55},⁴² which is in concordance with previous reviews.^{8,19} Organisational factors were scarcely reported. The variation in reported ICU delirium incidence could not be explained by hospital type, screening methods used, or daily screening frequency.

Several studies have been conducted to examine patient- and treatment-related factors that contribute to the development of delirium in the ICU, like need for mechanical ventilation (odds ratio [OR] = 7.0; 95% confidence interval [CI] = 3.4–14.8), need for physical restraints (OR = 33.8; 95% CI = 11.2–102.4), and use of sedatives (OR = 13.66; 95% CI = 7.2–26.1).⁴³ Based on these risk factors, delirium prediction models accurately predict if a patient will develop delirium or not in 77% of the patients.^{16,44} Although these patient- and treatment-related factors may account for a considerable part of the variability, they do not fully explain the observed delirium incidence.¹⁹ Identifying potentially modifiable organisational factors that might contribute to enhanced delirium recognition or occurrence may improve delirium management.¹³ In our meta-analysis, only three organisational factors could be extracted from the articles. First, the hospital type could have contributed to the delirium incidence found because of differences in severity of illness. Second, the daily screening frequency might have added to the sensitivity of the observed incidence owing to the known fluctuating nature of delirium during the day.³³ Third, the used assessment methods might have influenced the incidence found because the validity and psychometric properties of some screening instruments remain controversial, and differences in delirium incidence have been reported when both the CAM-ICU and the ICDSC are used simultaneously.²⁵ However, based on our findings, no significant association with delirium incidence can be concluded.

Delirium is found in up to 89% of critically ill patients.¹⁰ However, we showed these highest estimates reflect prevalence rather

than incidence. These high prevalence figures probably reflect onset of delirium before ICU admission. For clinical practice, it is important to have insight in the incidence of ICU delirium, so that effects of interventions and quality improvement programs, for example decreasing sedation-management^{33,45} or improving early mobilisation,⁴⁶ can be assessed. Future studies should strive to discriminate between ICU-acquired “incidence” and pre-ICU-acquired “prevalence” figures and also incorporate organisational and treatment aspects as robust data on effectiveness of specific non-organisational are scarce.⁴⁷

Although not significantly associated with the delirium incidence, a broad consensus on which occurrence measure, diagnostic instrument, and which screening algorithm to use might help to optimise delirium treatment and reduce its incidence. The current international Society of Critical Care Medicine (SCCM) delirium treatment ICU guideline⁴⁸ recommends the use of a standardised validated ICU delirium screening instrument. Both the CAM-ICU³ and the ICDSC³⁷ are recommended despite their methodological differences in assessment of delirium (testing patient performance versus observation by the professional).⁴⁹ In our meta-analysis, we did not find that the assessment tools used explained differences in delirium incidence. Potentially, because of the difference in assessment methods, a tool that combines testing patient performance with observation by a professional may provide even better screening performances and insight in the exact proportion of delirium in the ICU, but this has not been tested.

Some limitations of our study should be addressed. First, the number of organisational factors that could be analysed was limited to only three, as these were the only three that are currently reported. There might be other important organisational factors, such as structure and process factors like implementation and educational strategies, staff knowledge, motivation, and screening compliance, which may influence delirium incidence.^{34,35} However, these are difficult to quantify and therefore will be challenging to assess to what extent they may influence delirium incidence. Delirium is a multifactorial syndrome,⁵⁰ which is often the result of several interacting processes.¹⁹ It is vital to have good insight in causal linkages of the organisational processes and the outcomes of care by perceiving insights in these processes.^{51,52} Although, we did not find these associations, this does not exclude that other not reported organisational factors, such as delirium education, implementation strategies, staff knowledge, motivation, and screening compliance, may attribute to the observed delirium incidence as they are known to affect other outcomes in the ICU, such as infections.^{53,54}

In our study, we found articles of good methodological quality, assessed by the STROBE criteria.³² Therefore, the data we found are likely to be valid. Notably, however, the lowest delirium incidence rate (4%) was found in the largest study.¹¹ Although this study fulfilled the criteria for a good methodological quality, the low reported incidence contrasts to previous results⁸ and may be the result of a suboptimal screening regime, as in this national study some participating centres found incidences of 0%, which is unlikely to be accurate, considering the severity of illness (Acute Physiology and Chronic Health Evaluation-II) in the population as well as previous incidences found in large meta-analyses.^{8,23} Because this national study was not only primarily focused on delirium incidence but also on all other adverse events and outcomes associated with critically ill surgical patients,⁵⁵ we hypothesised that this would probably affect their reported delirium incidence rates. However, the results of this study did not statistically affect our reported mean delirium incidence of 29%.

It is possible that we were not able to detect an effect of organisational factors on delirium incidence because important

Table 3
Meta-regression analysis^a.

Variable	Estimates	Standard error	p-value
<i>Type of hospital</i>			
University	0.279	0.035	0.48
Non-university	0.332	0.065	
<i>Daily screening frequency</i>			
1	0.260	0.053	0.41
2	0.343	0.045	
3	0.266	0.064	
<i>Screening method</i>			
CAM-ICU	0.262	0.031	0.28
ICDSC	0.268	0.080	
Other methods	0.412	0.090	

CAM-ICU= Confusion Assessment Method for the Intensive Care Unit; ICDSC = Intensive Care Delirium Screening Checklist.

^a Forest plots available in an online supplement.

organisational factors were simply not addressed. To enable uniform and comparable research addressing all potentially important aspects of delirium, it may be useful to compose a minimal data set for delirium research, which is already used within critical care and other fields of health care,^{56,57} and might be useful for defining consensus-based patient and nursing sensitive outcomes, such as screening compliance and education type e-learning and hands on learning.⁵⁸ This data set might have most potential when added to the current ICU delirium management guidelines³³ and ICU delirium research agenda.⁵⁹ Second, Because of the limited number of studies which specifically reported delirium incidence in the context of organisational factors, we could not perform a multivariate regression analysis. This analysis would have enabled us to assess the associations between the extracted factors related to the delirium incidence found. Although optimal recognition of delirium does not reduce its burden on patients directly, it might help to provide optimal delirium care which may reduce its burden on patients. Further research may align the understanding of the variation in the occurrence of delirium in the ICU, through which lessons may be learned to minimise the burden of delirium in ICU patients.

5. Conclusions

This systematic review showed that 29% of patients develop delirium during their ICU stay. We were unable to demonstrate that organisational factors including methods of delirium assessment, screening frequency, and hospital type were related to the reported ICU delirium incidence.

Acknowledgements

The authors thank Michail Belias (PhD student biostatistics at the Radboud University Medical Center, Department for Health Evidence) for his assistance performing the meta-regression analysis.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.aucc.2018.02.002>.

References

- [1] Lat I, McMillian W, Taylor S, Janzen JM, Papadopoulos S, Korth L, et al. The impact of delirium on clinical outcomes in mechanically ventilated surgical and trauma patients. *Crit Care Med* 2009;37:1898–905.
- [2] Ely EW, Gautam S, Margolin R, Francis J, May L, Speroff T, et al. The impact of delirium in the intensive care unit on hospital length of stay. *Intensive Care Med* 2001;27:1892–900.
- [3] Ely EW, Inouye SK, Bernard GR, Gordon S, Francis J, May L, et al. Delirium in mechanically ventilated patients: validity and reliability of the confusion assessment method for the intensive care unit (CAM-ICU). *J Am Med Assoc* 2001;286:2703–10.
- [4] McNicoll L, Pisani MA, Zhang Y, Ely EW, Siegel MD, Inouye SK. Delirium in the intensive care unit: occurrence and clinical course in older patients. *J Am Geriatr Soc* 2003;51:591–8.
- [5] Girard TD, Jackson JC, Pandharipande PP, Pun BT, Thompson JL, Shintani AK, et al. Delirium as a predictor of long-term cognitive impairment in survivors of critical illness. *Crit Care Med* 2010;38:1513–20.
- [6] Jackson JC, Hart RP, Gordon SM, Shintani A, Truman B, May L, et al. Six-month neuropsychological outcome of medical intensive care unit patients. *Crit Care Med* 2003;31:1226–34.
- [7] Ely EW, Shintani A, Truman B, Speroff T, Gordon SM, Harrell Jr FE, et al. Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *Jama* 2004;291:1753–62.
- [8] Salluh JF, Wang H, Schneider EB, Nagaraja N, Yenokyan G, Damluji A, et al. Outcome of delirium in critically ill patients: systematic review and meta-analysis. 2015.
- [9] Sharma A, Malhotra S, Grover S, Jindal SK. Incidence, prevalence, risk factor and outcome of delirium in intensive care unit: a study from India. *Gen Hosp Psychiatr* 2012;34:639–46.
- [10] Ely EW, Girard TD, Shintani AK, Jackson JC, Gordon SM, Thomason JW, et al. Apolipoprotein E4 polymorphism as a genetic predisposition to delirium in critically ill patients. *Crit Care Med* 2007;35:112–7.
- [11] Pipanmekaporn T, Chittawatanarat K, Chaiwat O, Thawitsri T, Wacharasint P, Kongsayreepong S. Incidence and risk factors of delirium in multi-center Thai surgical intensive care units: a prospective cohort study. *J Intensive Care Med* 2015;3:53.
- [12] Durlak JA, DuPre EP. Implementation matters: a review of research on the influence of implementation on program outcomes and the factors affecting implementation. *Am J Community Psychol* 2008;41:327–50.
- [13] Chaudoir SR, Dugan AG, Barr CH. Measuring factors affecting implementation of health innovations: a systematic review of structural, organizational, provider, patient, and innovation level measures. *Implement Sci* 2013;8:22.
- [14] Mori S, Takeda JR, Carrara FS, Cohrs CR, Zane SS, Whitaker IY. Incidence and factors related to delirium in an intensive care unit. *Rev Esc Enferm USP* 2016;50:587–93.
- [15] Zaal IJ, Devlin JW, Peelen LM, Slooter AJ. A systematic review of risk factors for delirium in the ICU. *Crit Care Med* 2015;43:40–7.
- [16] Wassenaar A, van den Boogaard M, van Achterberg T, Slooter AJ, Kuiper MA, Hoogendoorn ME, et al. Multinational development and validation of an early prediction model for delirium in ICU patients. *Intensive Care Med* 2015;41:1048–56.
- [17] Sabol F, Bily B, Artemiou P, Kolesar A, Torok P, Bilecova-Rabajdova M, et al. Incidence and risk factors of delirium in patients after cardiac surgery: modifiable and non-modifiable factors. *Cor Vasa* 2015;57:e168–75.
- [18] Gossett AN, Slooter AJ, Boere PR, Zaal IJ. Risk factors for delirium after on-pump cardiac surgery: a systematic review. *Crit Care (London, England)* 2015;19:346.
- [19] Reade MC, Finfer S. Sedation and delirium in the intensive care unit. *N Engl J Med* 2014;370:444–54.
- [20] Karnatovskaia LV, Johnson MM, Benzo RP, Gajic O. The spectrum of psychocognitive morbidity in the critically ill: a review of the literature and call for improvement. *J Crit Care* 2015 Feb;30(1):130–7.
- [21] Kalabalik J, Brunetti L, El-Srougy R. Intensive care unit delirium: a review of the literature. *J Pharm Pract* 2014;27:195–207.
- [22] Awissi DK, Lebrun G, Coursin DB, Riker RR, Skrobik Y. Alcohol withdrawal and delirium tremens in the critically ill: a systematic review and commentary. *Intensive Care Med* 2013;39:16–30.
- [23] Neto AS, Nassar Jr AP, Cardoso SO, Manetta JA, Pereira VG, Esposito DC, et al. Delirium screening in critically ill patients: a systematic review and meta-analysis. *Crit Care Med* 2012;40:1946–51.
- [24] Figueroa-Ramos MI, Arroyo-Novoa CM, Lee KA, Padilla G, Puntillo KA. Sleep and delirium in ICU patients: a review of mechanisms and manifestations. *Intensive Care Med* 2009;35:781–95.
- [25] Boettger S, Nuñez DG, Meyer R, Richter A, Fernandez SF, Rudiger A, et al. Delirium in the intensive care setting: a reevaluation of the validity of the CAM-ICU and ICDS versus the DSM-IV-TR in determining a diagnosis of delirium as part of the daily clinical routine. *Palliat Support Care* 2017:1–9.
- [26] Tomasi CD, Grandi C, Salluh J, Soares M, Giombelli VR, Cascaes S, et al. Comparison of CAM-ICU and ICDS for the detection of delirium in critically ill patients focusing on relevant clinical outcomes. *J Crit Care* 2012;27:212–7.
- [27] Numan T, van den Boogaard M, Kamper AM, Rood PJT, Peelen LM, Slooter AJC. Recognition of delirium in postoperative elderly patients: a multicenter study. *J Am Geriatr Soc* 2017 May 12.
- [28] Hayhurst CJ, Pandharipande PP, Hughes CG. Intensive care unit delirium: a review of diagnosis, prevention, and treatment. *Anesthesiology* 2016;125:1229–41.
- [29] Trogrlić Z, van der Jagt M, Bakker J, Balas MC, Ely EW, van der Voort PH, et al. A systematic review of implementation strategies for assessment, prevention, and management of ICU delirium and their effect on clinical outcomes. *Crit Care* 2015;19:157.
- [30] Higgins JPT, GSE. Cochrane handbook for systematic reviews of interventions, version 5.1.0 [updated March 2011]. 2011. The Cochrane Collaboration; 2011.
- [31] Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- [32] von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007;370:1453–7.
- [33] Barr J, Fraser GL, Puntillo K, Ely EW, Gelinas C, Dasta JF, et al. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Crit Care Med* 2013;41:263–306.
- [34] Moore L, Lavoie A, Bourgeois G, Lapointe J. Donabedian's structure-process-outcome quality of care model: validation in an integrated trauma system. *J Trauma Acute Care Surg* 2015;78:1168–75.
- [35] Donabedian A. The quality of care. How can it be assessed? *Jama* 1988;260:1743–8.
- [36] Foundation TR. R. 3.2.4 ed2016.

- [37] Bergeron N, Dubois MJ, Dumont M, Dial S, Skrobik Y. Intensive care delirium screening checklist: evaluation of a new screening tool. *Intensive Care Med* 2001;27:859–64.
- [38] Trzepacz PT, Mittal D, Torres R, Canary K, Norton J, Jimerson N. Validation of the delirium rating scale-revised-98. *J Neuropsychiatry Clin Neurosci* 2001;13:229–42.
- [39] Lutz A, Radtke FM, Franck M, Seeling M, Gaudreau JD, Kleinwachter R, et al. [The nursing delirium screening scale (NU-DESC)]. *Anesthesiologie, Intensivmedizin, Notfallmedizin* 43. *Schmerztherapie: AINS*; 2008. p. 98–102.
- [40] Association AP. DSM-IV-TR: diagnostic and statistical manual of mental disorders, Fourth Edition, text revision. American Psychiatric Association; 2000.
- [41] Ely EW, Margolin R, Francis J, May L, Truman B, Dittus R, et al. Evaluation of delirium in critically ill patients: validation of the confusion assessment method for the intensive care unit (CAM-ICU). *Crit Care Med* 2001;29:1370–9.
- [42] Smulter N, Lingehall HC, Gustafson Y, Olofsson B, Engstrom KG. Delirium after cardiac surgery: incidence and risk factors. *Interact Cardiovasc Thorac Surg* 2013;17:790–6.
- [43] Van Rompaey B, Schuurmans MJ, Shortridge-Baggett LM, Truijien S, Bossaert L. Risk factors for intensive care delirium: a systematic review. *Intensive Crit Care Nurs* 2008;24:98–107.
- [44] van den Boogaard M, Schoonhoven L, Maseda E, Plowright C, Jones C, Luetz A, et al. Recalibration of the delirium prediction model for ICU patients (PRE-DELIRIC): a multinational observational study. *Intensive Care Med* 2014;40:361–9.
- [45] Balas MC, Vasilevskis EE, Olsen KM, Schmid KK, Shostrom V, Cohen MZ. Effectiveness and safety of the awakening and breathing coordination, delirium monitoring/management, and early exercise/mobility bundle. *Crit Care Med* 2014;42.
- [46] Schweickert WD, Pohlman MC, Pohlman AS, Nigos C, Pawlik AJ, Esbrook CL, et al. Early physical and occupational therapy in mechanically ventilated, critically ill patients: a randomised controlled trial. *Lancet* 2009;373:1874–82.
- [47] Trogrlic Z, van der Jagt M, Bakker J, Balas MC, Ely EW, van der Voort PH, et al. A systematic review of implementation strategies for assessment, prevention, and management of ICU delirium and their effect on clinical outcomes. *Crit Care (London, England)* 2015;19:157.
- [48] Barr J, Pandharipande PP. The pain, agitation, and delirium care bundle: synergistic benefits of implementing the 2013 Pain, Agitation, and Delirium Guidelines in an integrated and interdisciplinary fashion. *Crit Care Med* 2013;41:S99–115.
- [49] Gusmao-Flores D, Salluh JI, Chalhub RA, Quarantini LC. The confusion assessment method for the intensive care unit (CAM-ICU) and intensive care delirium screening checklist (ICDSC) for the diagnosis of delirium: a systematic review and meta-analysis of clinical studies. *Crit Care (London, England)* 2012;16:R115.
- [50] Maldonado JR. Delirium in the acute care setting: characteristics, diagnosis and treatment. *Crit Care Clin* 2008;24. 657–722. vii.
- [51] Donabedian A. The quality of care: how can it be assessed? *J Am Med Assoc* 1988;260:1743–8.
- [52] Neta G, Glasgow RE, Carpenter CR, Grimshaw JM, Rabin BA, Fernandez ME, et al. A framework for enhancing the value of research for dissemination and implementation. *Am J Publ Health* 2015;105:49–57.
- [53] Sakr Y, Moreira CL, Rhodes A, Ferguson ND, Kleinpell R, Pickkers P, et al. The impact of hospital and ICU organizational factors on outcome in critically ill patients: results from the Extended Prevalence of Infection in Intensive Care study. *Crit Care Med* 2015;43:519–26.
- [54] Carmel S, Rowan K. Variation in intensive care unit outcomes: a search for the evidence on organizational factors. *Curr Opin Crit Care* 2001;7:284–96.
- [55] Chittawatanarat K, Chaiwat O, Morakul S, Pipanmekaporn T, Thawitsri T, Wacharasint P, et al. A multi-center Thai university-based surgical intensive care units study (THAI-SICU study): methodology and ICU characteristics. *Journal of the Medical Association of Thailand Chotmaihet thangphaet* 2014;97(Suppl 1):S45–54.
- [56] Morris R, Matthews A, Scott AP. Validity, reliability and utility of the Irish Nursing Minimum Data Set for General Nursing in investigating the effectiveness of nursing interventions in a general nursing setting: a repeated measures design. *Int J Nurs Stud* 2014;51:562–71.
- [57] Felton TW, Sander R, Al-Aloul M, Dark P, Bentley AM. Can a score derived from the Critical Care Minimum Data Set be used as a marker of organ dysfunction? – a pilot study. *BMC Res Notes* 2009;2:77.
- [58] Milani A, Mauri S, Gandini S, Magon G. Oncology nursing minimum data set (ONMDS): can we hypothesize a set of prevalent nursing sensitive outcomes (NSO) in cancer patients? *Ecancermedicallscience* 2013;7:345.
- [59] Pandharipande PP, Ely EW, Arora RC, Balas MC, Boustani MA, La Calle GH, et al. The intensive care delirium research agenda: a multinational, interprofessional perspective. *Intensive Care Med* 2017 Jun 13.
- [60] Ayllon Garrido N, Alvarez Gonzalez MJ, Gonzalez Garcia MP. Delirium in intensive care. Incidence and risk factors. *Enfermería Intensiva* 2007;18:138–43.
- [61] Bilge EU, Kaya M, Senel GO, Unver S. The incidence of delirium at the post-operative intensive care unit in adult patients. *Turk J Anaesthesiol Reanim* 2015;43:232–9.
- [62] Kanova M, Burda M, Povova J, Neiser J. Delirium in intensive care patients – a prospective study. [Czech]. *Ceská a Slov Neurol a Neurochir* 2015;78:662–7.
- [63] Lahariya S, Grover S, Bagga S, Sharma A. Delirium in patients admitted to a cardiac intensive care unit with cardiac emergencies in a developing country: incidence, prevalence, risk factor and outcome. *Gen Hosp Psychiatr* 2014;36:156–64.
- [64] Limpawattana P, Panitchote A, Tangvoraphonchai K, Suebsoh N, Emma W, Chanthonglang B, et al. Delirium in critical care: a study of incidence, prevalence, and associated factors in the tertiary care hospital of older Thai adults. *Aging Ment Health* 2016;20:74–80.
- [65] Norkiene I, Ringaitiene D, Kuzminkaite V, Sipylaitė J. Incidence and risk factors of early delirium after cardiac surgery. *BioMed Res Int* 2013;2013:323491.
- [66] Ouimet S, Kavanagh BP, Gottfried SB, Skrobik Y. Incidence, risk factors and consequences of ICU delirium. *Intensive Care Med* 2007;33:66–73.
- [67] Page VJ, Navarange S, Gama S, McAuley DF. Routine delirium monitoring in a UK critical care unit. *Crit Care (London, England)* 2009;13:R16.
- [68] Peterson JF, Pun BT, Dittus RS, Thomason JW, Jackson JC, Shintani AK, et al. Delirium and its motoric subtypes: a study of 614 critically ill patients. *J Am Geriatr Soc* 2006;54:479–84.
- [69] Roberts B, Rickard CM, Rajbhandari D, Turner G, Clarke J, Hill D, et al. Multi-centre study of delirium in ICU patients using a simple screening tool. *Aust Crit Care* 2005;18(6):8–9. 11–4 passim.
- [70] Shi CM, Wang DX, Chen KS, Gu XE. Incidence and risk factors of delirium in critically ill patients after non-cardiac surgery. *Chinese Med J* 2010;123:993–9.
- [71] Svenningsen H, Tonnesen E. Delirium incidents in three Danish intensive care units. *Nurs Crit Care* 2011;16:186–92.
- [72] van den Boogaard M, Schoonhoven L, van der Hoeven JG, van Achterberg T, Pickkers P. Incidence and short-term consequences of delirium in critically ill patients: a prospective observational cohort study. *Int J Nurs Stud* 2012;49:775–83.
- [73] Woien H, Balsliemke S, Stubhaug A. The incidence of delirium in Norwegian intensive care units; deep sedation makes assessment difficult. *Acta Anaesthesiol Scand* 2013;57:294–302.
- [74] Saklad MDM. Grading of patients for surgical procedures. *Anesthesiology* 1941;2:281–4.
- [75] Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Crit Care Med* 1985;13:818–29.
- [76] Nashef SA, Roques F, Sharples LD, Nilsson J, Smith C, Goldstone AR, et al. EuroSCORE II. *Eur J Cardio Thorac Surg* 2012;41:734–44. discussion 44–45.
- [77] Moreno RP, Metnitz PG, Almeida E, Jordan B, Bauer P, Campos RA, et al. SAPS 3—From evaluation of the patient to evaluation of the intensive care unit. Part 2: development of a prognostic model for hospital mortality at ICU admission. *Intensive Care Med* 2005;31:1345–55.