

Alzheimer's Disease—Failure of Drainage of Fluid from the Brain

An Expert Interview with Roxana Carare

Faculty of Medicine, Institute for Life Sciences, University of Southampton, Southampton General Hospital, Southampton, Hampshire, UK

DOI: <https://doi.org/10.17925/USN.2018.14.1.21>



Roxana Carare

Roxana Carare is a Professor in Clinical Neuroanatomy in the Faculty of Medicine, University of Southampton. Roxana graduated in General Medicine in 1994, completed her basic surgical training by 1996 and embarked on an academic career in Anatomy in 1998. She developed a strong international profile in research and is recognized for discovering the lymphatic drainage pathways of the brain that become clogged up with aggregated proteins, leading to Alzheimer's disease. She is a recipient of a Dementia Leaders Award from Alzheimer's Society UK, active member of the International Vascular Behavioral and Cognitive Disorders Society (VasCog), Cerebral Amyloid Angiopathy Society and Romanian Alzheimer's Society, serves on the Editorial Board of Neuropathology and Applied Neurobiology, and reviewer for specialist neuropathology and neurology journals, as well as major international funding agencies.

Keywords

Alzheimer's disease, amyloid-beta ($A\beta$), cerebral amyloid angiopathy, intramural periarterial drainage (IPAD), interstitial fluid (ISF)

Disclosure: Roxana Carare has nothing to declare in relation to this article.

Review Process: This is an expert interview and as such, has not undergone the journal's standard peer review process.

Authorship: The named author meets the International Committee of Medical Journal Editors (ICMJE) criteria for authorship of this manuscript, takes responsibility for the integrity of the work as a whole, and has given final approval for the version to be published.

Open Access: This article is published under the Creative Commons Attribution Noncommercial License, which permits any noncommercial use, distribution, adaptation, and reproduction provided the original author and source are given appropriate credit. © The Author 2018.

Received: January 9, 2018

Published Online: February 16, 2018

Citation: *US Neurology*. 2018;14(1):21–22

Corresponding Author: Roxana Carare, Faculty of Medicine, Institute for Life Sciences, University of Southampton, Southampton General Hospital, South Academic Block, MP806, Tremona Road, Southampton, Hampshire, SO16 6YD, UK. E: R.O.Carare@soton.ac.uk Facebook: @unisouthamptonmedicine

Support: No funding was received in the publication of this article.

The annual Alzheimer's Association International Conference (AAIC) took place in London, UK in July 2017. In an expert interview, Roxana Carare recaps on the topic of her presentation during the 'Emerging Concepts in Basic Science' session, clearance of interstitial fluid of the brain and pathogenesis of dementia.¹

Q. How is soluble amyloid-beta cleared from the brain in healthy individuals?

Amyloid-beta ($A\beta$) is produced in all brains, as it is key to the maintenance of brain health. Normally $A\beta$ is broken down by enzymes² and eliminated across the walls of the blood vessels into the blood,³ or is removed by intramural periarterial drainage (IPAD), described by the Carare group.⁴ Enzymes and transporters of $A\beta$ ⁵ into the blood fail with increasing age and with other risk factors for Alzheimer's disease, so the burden of removing $A\beta$ via IPAD from aging brains is increased. IPAD takes place along tiny channels (vascular basement membranes, composed of extracellular matrix) in the walls of arteries that supply blood to the brain.⁶


Q. How does this process change in Alzheimer's disease?

The direction of IPAD is opposite to that of blood flow and relies on the contraction of smooth muscle cells to function properly.⁷ With aging, high blood pressure and high cholesterol in midlife, diabetes and obesity, the arteries become stiffer and smooth muscle cells do not contract properly (Figure 1).^{8–10} This leads to a poor perfusion of the brain with blood, as well as reduced clearance of $A\beta$ from the brain.¹¹ A failure of efficient drainage of $A\beta$ and other proteins from the aging brain results in their deposition in the walls of blood vessels as sticky plaques, giving rise to cerebral amyloid angiopathy, a key feature of Alzheimer's disease (Figure 2).¹²

Q. What lifestyle factors might affect protein clearance?

Prevention or early treatment of metabolic diseases such as diabetes, high cholesterol, high blood pressure, low vitamin B and maintaining a healthy heart should also maintain healthy blood vessels in the brain, preventing Alzheimer's disease.^{13–15}

Figure 1: Clearance of A β from young versus old blood vessels




A β = beta-amyloid; ISF = interstitial fluid.

Q. What therapeutic strategies may help the clearance of A β from the brain?

There are already promising experimental studies demonstrating that compounds that increase the activity of the smooth muscle cells of blood vessels are able to reverse the features of Alzheimer's disease. Such compounds (for example cilostazol, a selective phosphodiesterase type 3 inhibitor in clinical trials in Japan and Edinburgh, UK) are able to slow the progression from mild cognitive impairment to full blown Alzheimer's disease.^{16,17} ■

Figure 2: Cerebral amyloid angiopathy—failure of elimination of A β along the walls of blood vessels



A: Schematic; B: Immuno-stained blood vessel where A β is red, smooth muscle actin cells are green, and collagen IV in the basement membrane is blue. A β = beta-amyloid.

- Carare RO, EC-01-01 Clearance of interstitial fluid of the brain and pathogenesis of dementia. *Alzheimers Dement*. 2017;13 Suppl. P180.
- Farris W, Schütz SG, Cirrito JR, et al. Loss of neprilysin function promotes amyloid plaque formation and causes cerebral amyloid angiopathy. *Am J Pathol*. 2007;171:241–51.
- Deane R, Wu Z, Sagare A, et al. LRP/amyloid beta-peptide interaction mediates differential brain efflux of Abeta isoforms. *Neuron*. 2004;43:333–44.
- Carare RO, Bernardes-Silva M, Newman TA, et al. Solutes, but not cells, drain from the brain parenchyma along basement membranes of capillaries and arteries: significance for cerebral amyloid angiopathy and neuroimmunology. *Neuropathol Appl Neurobiol*. 2008;34:131–44.
- Tarasoff-Conway JM, Carare RO, Osorio RS, et al. Clearance systems in the brain – implications for Alzheimer disease. *Nat Rev Neurol*. 2016;12:248.
- Morris AW, Sharp MM, Albigorothy NJ, et al. Vascular basement membranes as pathways for the passage of fluid into and out of the brain. *Acta Neuropathol*. 2016;131:725–36.
- Diem AK, MacGregor Sharp M, Gatherer M, et al. Arterial pulsations cannot drive intramural periaarterial drainage: significance for A β drainage. *Front Neurosci*. 2017;11:475.
- Hawkes CA, Gatherer M, Sharp MM, et al. Regional differences in the morphological and functional effects of aging on cerebral basement membranes and perivascular drainage of amyloid- β from the mouse brain. *Aging Cell*. 2013;12:224–36.
- Bueche CZ, Hawkes C, Garz C, et al. Hypertension drives parenchymal β -amyloid accumulation in the brain parenchyma. *Ann Clin Transl Neurol*. 2014;1:124–9.
- Hawkes CA, Gentleman SM, Nicoll JA, Carare RO. Prenatal high-fat diet alters the cerebrovasculature and clearance of β -amyloid in adult offspring. *J Pathol*. 2015;235:619–31.
- Weller RO, Hawkes CA, Carare RO, Hardy J. Does the difference between PART and Alzheimer's disease lie in the age-related changes in cerebral arteries that trigger the accumulation of A β and propagation of tau? *Acta Neuropathol*. 2015;129:763–6.
- Hawkes CA, Jayakody N, Johnston DA, et al. Failure of perivascular drainage of β -amyloid in cerebral amyloid angiopathy. *Brain Pathol*. 2014;24:396–403.
- Hainsworth AH, Yeo NE, Weekman EM, Wilcock DM. Homocysteine, hyperhomocysteinemia and vascular contributions to cognitive impairment and dementia (VCID). *Biochim Biophys Acta*. 2016;1862:1008–17.
- Mukaetova-Ladinska EB, Purshouse K, Andrade J, et al. Can healthy lifestyle modify risk factors for dementia? Findings from a pilot community-based survey in Chennai (India) and Newcastle (UK). *Neuroepidemiology*. 2012;39:163–70.
- Paillard-Borg S, Fratiglioni L, Xu WL, et al. An active lifestyle postpones dementia onset by more than one year in very old adults. *J Alzheimers Dis*. 2012;31:835–42.
- Maki T, Okamoto Y, Carare RO, et al. Phosphodiesterase III inhibitor promotes drainage of cerebrovascular β -amyloid. *Ann Clin Transl Neurol*. 2014;1:519–33.
- Saito S, Ihara M. New therapeutic approaches for Alzheimer's disease and cerebral amyloid angiopathy. *Front Aging Neurosci*. 2014;6:290.