

RELATIVE COMMUTANT PICTURES OF ROE ALGEBRAS

JÁN ŠPAKULA AND AARON TIKUISIS

ABSTRACT. Let X be a proper metric space, which has finite asymptotic dimension in the sense of Gromov (or more generally, straight finite decomposition complexity of Dranishnikov and Zarichnyi). New descriptions are provided of the Roe algebra of X : (i) it consists exactly of operators which essentially commute with diagonal operators coming from Higson functions (that is, functions on X whose oscillation tends to 0 at ∞), and (ii) it consists exactly of quasi-local operators, that is, ones which have finite ϵ -propagation (in the sense of Roe) for every $\epsilon > 0$. These descriptions hold both for the usual Roe algebra and for the uniform Roe algebra.

1. INTRODUCTION

The Roe algebra is a C^* -algebra constructed from a proper metric space, which encodes “coarse” or “large-scale” properties of the space (in the sense of Gromov). In typical applications, the space may be a complete, non-compact Riemannian manifold with bounded geometry, or finitely generated group with the word metric. The origins of this construction come from index theory, reflecting the insight that the Roe algebra is large enough to contain indices of many operators with which one wants to do index theory – such as geometric differential operators – yet small enough to have interesting and informative K -theory. It plays a central role in the coarse Baum–Connes conjecture, the study (and partial confirmation) of which has been a fruitful endeavor, leading to significant results concerning the Novikov conjecture and the scalar curvature of Riemannian manifolds [3, 12, 15, 21, 22, 24, 25, 29, 30, 31, 32]. It furthermore appears in work on the essential spectrum of Hamiltonian operators of quantum systems, Schrödinger operators, and various other operators, which are affiliated to the appropriate versions of Roe algebras [7, 8, 9, 19].

Roughly, the Roe algebra consists of bounded, locally compact operators on something like $L^2(X)$ (where X is the underlying space) which can be approximated by those with finite propagation. Here an operator a has “finite propagation” if it is localized near to the diagonal; one way of making this precise is, that there exists $R > 0$ such that for any $f, f' \in C_b(X)$ (acting on $L^2(X)$ as diagonal operators – by multiplication), if the supports of f and f' are separated by a distance

of at least R , then $faf' = 0$. Operators in the Roe algebra are required to be approximated in the *operator norm* by these finite propagation operators.

The Roe algebra is an interesting object to study from an operator algebraic perspective: that is, asking about its structure as an operator algebra, and how this structure relates to the proper metric space from which it is constructed. For example, Ozawa showed that exactness of a group can be characterized by amenability of the corresponding uniform Roe algebra ([18]). The question of how much of the large-scale structure of a space is remembered by the Roe algebra, was partially answered by JS and Rufus Willett: given two uniformly discrete proper metric spaces with Yu’s property A, their Roe algebras are $*$ -isomorphic if and only if the spaces are coarsely equivalent ([26]). In [28], Winter and Zacharias showed an interesting one-way connection between the asymptotic dimension of a metric space and the nuclear dimension of the corresponding uniform Roe algebra; the latter is a numerical invariant for amenable C^* -algebras which is crucial in recent results in the classification of amenable C^* -algebras. Their result is that the nuclear dimension of the Roe algebra is at most the asymptotic dimension of the underlying uniformly discrete proper metric space, and they asked the (still open) question of whether the reverse inequality also holds.

In this paper, we look at a fundamental question: exactly which operators *are* in the Roe algebra? In [22], Roe defined the concept of finite ϵ -propagation for an operator a on $L^2(X)$, as the following variant of finite propagation: a has finite ϵ -propagation if there exists $R > 0$ such that for any $f, f' \in C_b(X)$, if the supports of f and f' are separated by a distance of at least R , then $\|faf'\| \leq \epsilon\|f\| \cdot \|f'\|$. Operators with finite ϵ -propagation for all $\epsilon > 0$ have also been called *quasi-local operators* in the literature (originally from [21, Page 100]). It is a straightforward observation that, although limits of finite propagation operators need not have finite propagation, limits of finite ϵ -propagation operators have finite ϵ -propagation (that is, the set of quasi-local operators is norm-closed). Therefore, all operators in the Roe algebra are quasi-local.

The question we address is the converse: if an operator is quasi-local, is it in the Roe algebra, i.e., is it approximated by operators with finite propagation? We provide an affirmative answer in the situation that the space has finite asymptotic dimension (as predicted by Roe), and more generally under the hypothesis of straight finite decomposition complexity of Dranishnikov and Zarichnyi [4]. The latter is a weaker version of the “classical” finite decomposition complexity, as defined by Guentner, Tessera, and Yu [13, 14].

A motivation for asking whether quasi-local operators are in the Roe algebra, pointed out to the authors by Alexander Engel, is that whereas indices of genuine differential operators are known to be in

the Roe algebra, corresponding arguments only shows that indices of *pseudo*-differential operators (using their most natural representative) are quasi-local (see [5, Section 2]). Since the Roe algebra is better studied and understood than the C^* -algebra of all quasi-local operators, it is interesting and useful to know that a pseudo-differential operator belongs to it; indeed, our result answers [5, Question 6.4] under the assumption of straight finite decomposition complexity (this sort of assumption is anticipated in the question).

Additionally, we expose that the above question can be reformulated in terms of essential commutation with Higson functions, or in terms of relative commutants.

So far we have been a bit vague about what we mean by the Roe algebra (hiding behind the phrase “something like $L^2(X)$ ”). This is because in the literature there are two different versions of the Roe algebra, the “Roe algebra” and the “uniform Roe algebra”. Our results apply to both of these C^* -algebras, and indeed our main theorem is formulated in a way that encompasses both, as well as the “uniform algebra” $UC^*(X)$. The main result was proven by Lange and Rabinovich for the uniform Roe algebra of \mathbb{Z}^d in [17]. Engel proved a special case of the result, namely that for discrete groups G that are lattices in a Riemannian manifold with bounded geometry and polynomial volume growth, quasi-local operators that decay sufficiently quickly are in the Roe algebra ([6, Corollary 2.33]).¹

Let us now summarize the argument behind the main result: that quasi-local operators are in the Roe algebra (assuming straight finite decomposition complexity). Suppose for simplicity that X is a discrete space with asymptotic dimension at most 1 – for example a finitely generated free group. This case is much more restricted than finite decomposition complexity, but still difficult enough to allow us to convey the main ideas. Let a be a quasi-local operator. Asymptotic dimension at most 1 will allow us to decompose the space X into 2 pieces, $X^{(0)}$ and $X^{(1)}$, each piece being a disjoint union of sets that are far apart from each other and uniformly bounded in diameter. The characteristic functions $e^{(0)}, e^{(1)}$ of these pieces produce a partition of unity, and divides a into a sum of four pieces: $e^{(i)}ae^{(j)}$ over $i, j = 0, 1$. Each $e^{(i)}ae^{(i)}$ looks roughly like an infinite block matrix, indexed by the pieces from $X^{(i)}$. The hypothesis that a is quasi-local (finite ϵ -propagation) gives

¹In fact, Engel proved the result for quasi-local operators that decay sufficiently on *any* Riemannian manifold with bounded geometry and polynomial volume growth. For groups, polynomial growth implies virtual nilpotency ([11]), which in turn implies finite asymptotic dimension ([1, Corollary 68]). To our knowledge, it is not known whether polynomial volume growth implies finite asymptotic dimension (or even (straight) finite decomposition complexity) outside of the case of groups.

us a lot of control over the norm of the non-diagonal entries of this matrix, and a conditional expectation argument allows us to conclude that $e^{(i)}ae^{(i)}$ is not far away from its “restriction” to the diagonal (provided that the pieces in $X^{(i)}$ are sufficiently well separated), see Corollary 4.3. Since the pieces of the $X^{(i)}$ are uniformly bounded, the operator we get by expecting onto the diagonal has genuinely finite propagation. An algebraic trick allows us to view the asymmetric pieces $e^{(i)}ae^{(j)}$ as matrices in a similar way, so that we can likewise approximate each of them by finite propagation operators. In this way, we approximate a as a sum of four operators with finite propagation.

Outline. In Section 2 we introduce our general setup, with the Roe algebra, the uniform Roe algebra, and the uniform algebra as examples. We then state the main result, Theorem 2.8, in the language of our general setup. We give some background on asymptotic dimension and (straight) finite decomposition complexity in Section 2.1. The equivalence between quasi-locality and the relative commutant-type property is fairly straightforward, and laid out in Section 3. We use a more technical formulation of quasi-locality as a stepping stone towards proving that it implies being in the Roe algebra (assuming straight finite decomposition complexity), a proof that is carried out in Section 4. In Section 5, we prove that the relative commutant-type property is equivalent to essential commutation with Higson functions. The final section, Section 6, is concerned with the commutative (but non-separable) C^* -algebra $VL_\infty(X)$ that arises in our relative commutant-type property, looking at how well it determines X (up to coarse equivalence), and at its nuclear dimension (roughly, the covering dimension of its spectrum).

Acknowledgments. AT was supported by EPSRC EP/N00874X/1. JS was supported by Marie Curie FP7-PEOPLE-2013-CIG Coarse Analysis (631945). We would like to thank Alexander Engel, John Roe, Thomas Weighill, Stuart White, and Rufus Willett for comments and discussion relating to this piece.

2. DEFINITIONS AND THE MAIN RESULT

Let A be a C^* -algebra. We denote by A_1 the closed unit ball of A . For $a, b \in A$ and $\epsilon > 0$, we write $a \approx_\epsilon b$ to mean $\|a - b\| \leq \epsilon$. Define

$$A_\infty := l^\infty(\mathbb{N}, A) / \{(a_n)_{n=1}^\infty \in l^\infty(\mathbb{N}, A) : \lim_{n \rightarrow \infty} \|a_n\| = 0\},$$

which is a C^* -algebra.

We now set up a general situation to which our main result applies, encompassing both Roe algebras and uniform Roe algebras, as well as uniform algebras (see Example 2.5). Subsequently, we will state our main result in its full generality (Theorem 2.8)

Definition 2.1. Let X be a proper metric space. By an X -module, we mean a Hilbert space \mathcal{H} and an injective unital $*$ -homomorphism $C_b(X) \rightarrow \mathcal{B}(\mathcal{H})$, which is strictly continuous when viewing $C_b(X)$ and $\mathcal{B}(\mathcal{H})$ as multiplier algebras of $C_0(X)$ and $\mathcal{K}(\mathcal{H})$ respectively. We shall suppress the $*$ -homomorphism $C_b(X) \rightarrow \mathcal{B}(\mathcal{H})$, and treat $C_b(X)$ as a C^* -subalgebra of $\mathcal{B}(\mathcal{H})$.

For $R \geq 0$, an operator $a \in \mathcal{B}(\mathcal{H})$ has propagation at most R if for any $f, f' \in C_b(X)$, if the supports of f and f' are R -disjoint then $faf' = 0$. For $R \geq 0$ and $\epsilon > 0$, an operator $a \in \mathcal{B}(\mathcal{H})$ has ϵ -propagation at most R if for any $f, f' \in C_b(X)_1$, if the supports of f and f' are R -disjoint then $\|faf'\| < \epsilon$. An operator $a \in \mathcal{B}(\mathcal{H})$ is quasi-local if for every $\epsilon > 0$, it has finite ϵ -propagation.

Definition 2.2. Let X be a proper metric space and let \mathcal{H} be an X -module. Given an equicontinuous family $(e_j)_{j \in J}$ of positive contractions from $C_b(X)$ with pairwise disjoint supports, define the block cutdown map $\theta_{(e_j)_{j \in J}} : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ by

$$\theta_{(e_j)_{j \in J}}(a) := \sum_{j \in J} e_j a e_j$$

(using disjointness of the supports and the fact that the family is contractive, the right-hand sum converges in the strong operator topology).

Let $B \subseteq \mathcal{B}(\mathcal{H})$ be a C^* -subalgebra such that $C_b(X)BC_b(X) = B$. B is closed under block cutdowns if $\theta_{(e_j)_{j \in J}}(B) \subseteq B$ for every equicontinuous family $(e_j)_{j \in J}$ of positive contractions from $C_b(X)$ with pairwise disjoint supports.

For an equicontinuous family $(e_j)_{j \in J}$ of positive contractions from $C_b(X)$ with pairwise disjoint supports, the block cutdown map $\theta_{(e_j)_{j \in J}}$ defined above is evidently completely positive and contractive (c.p.c.). Note that multiplication by $C_b(X)$ commutes with block cutdowns:

$$f\theta_{(e_j)_{j \in J}}(a) = \theta_{(e_j)_{j \in J}}(fa) \quad \text{and} \quad \theta_{(e_j)_{j \in J}}(a)f = \theta_{(e_j)_{j \in J}}(af)$$

for $f \in C_b(X)$ and $a \in \mathcal{B}(\mathcal{H})$. Also note that

$$(2.1) \quad \|\theta_{(e_j)_{j \in J}}(a)\| = \sup_{j \in J} \|e_j a e_j\|.$$

Note that, if $(e_j)_{j \in J}$ is an equicontinuous family of positive contractions from $C_b(X)$ with uniformly bounded, pairwise disjoint supports, then $\theta_{(e_j)_{j \in J}}(a)$ has finite propagation, for every $a \in \mathcal{B}(\mathcal{H})$.

Definition 2.3. Let X be a proper metric space, \mathcal{H} an X -module, and let $B \subseteq \mathcal{B}(\mathcal{H})$ be a C^* -subalgebra such that $C_b(X)BC_b(X) = B$, and which is closed under block cutdowns. Define

- (i) $\text{Roe}(X, B) := \overline{\{b \in B : b \text{ has finite propagation}\}}^{\|\cdot\|}$, and
- (ii) $\mathcal{K}(X, B) := \overline{C_0(X)BC_0(X)}$.

If, in addition, we have

$$(2.2) \quad [C_0(X), B] \subseteq \mathcal{K}(X, B),$$

we shall call $\text{Roe}(X, B)$ a Roe-like algebra of X .

Remark 2.4. The condition (2.2) implies that $\mathcal{K}(X, B)$ is an ideal in $\text{Roe}(X, B)$. It is automatically satisfied in all the examples below, where in fact $C_0(X)B \subseteq \mathcal{K}(X, B)$ (and $\mathcal{K}(X, B)$ turns out to be the ideal of compact operators). Finally, it is not needed for the substantial part of this piece, so we shall explicitly refer to it when needed.

Example 2.5. Let X be a uniformly discrete proper metric space. Let \mathcal{H}' be an infinite dimensional, separable Hilbert space. Set $\mathcal{H}_u := l^2(X)$ and $\mathcal{H} := l^2(X, \mathcal{H}')$; $C_b(X)$ acts on both of these by pointwise multiplication, making them X -modules.

(i) With $B_u := \mathcal{B}(\mathcal{H}_u)$, we see that $C_b(X)B_uC_b(X) = B_u$, and B_u is closed under block cutdowns. In this case, $\text{Roe}(X, B_u) = C_u^*(X)$, the uniform Roe algebra, and $\mathcal{K}(X, B_u) = \mathcal{K}(\mathcal{H}_u)$. Since $C_0(X) \subseteq \mathcal{K}(\mathcal{H}_u)$, it follows that $C_0(X)B_u \subseteq \mathcal{K}(\mathcal{H}_u) = \mathcal{K}(X, B_u)$.

(ii) Set B equal to the set of all $b \in \mathcal{B}(\mathcal{H})$ which are locally compact, in the sense that for every $f \in C_0(X)$,

$$fb, bf \in \mathcal{K}(\mathcal{H}).$$

We see that $C_b(X)BC_b(X) = B$, and B is closed under block cutdowns. Then $\text{Roe}(X, B) = C^*(X)$, the Roe algebra, and $\mathcal{K}(X, B) = \mathcal{K}(\mathcal{H})$.

(iii) Assume that X has bounded geometry. Set B_0 equal to the closure of the set of all $b = (b_{x,y})_{x,y \in X} \in \mathcal{B}(\mathcal{H})$ for which the rank of $b_{x,y} \in \mathcal{B}(\mathcal{H}')$ is uniformly bounded. When $b = (b_{x,y})_{x,y \in X} \in \mathcal{B}(\mathcal{H})$ has entries with rank bounded by k , then so does any block cutdown map applied to b . Since each block cutdown map is continuous, it follows that B_0 is closed under block cutdowns. Continuity of multiplication ensures that

$$C_b(X)B_0C_b(X) = B_0.$$

When X has bounded geometry, then $\text{Roe}(X, B_0) = UC^*(X)$, the uniform algebra of X , defined as the closure of finite propagation operators $b = (b_{x,y})_{x,y \in X} \in \mathcal{B}(\mathcal{H})$ for which the rank of $b_{x,y}$ is uniformly bounded.

To see this, it is clear that $\text{Roe}(X, B_0)$ contains $UC^*(X)$. To show $\text{Roe}(X, B_0) \subseteq UC^*(X)$, it suffices to check that every finite propagation operator $a \in B_0$ is contained in $UC^*(X)$. For such a , say its propagation is less than $R > 0$. Set

$$K := \sup_{x \in X} |B_R(x)|,$$

which is finite due to the hypothesis of bounded geometry. Define $E_R : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ by $E_R((b_{x,y})_{x,y \in X}) := (c_{x,y})_{x,y \in X}$ where

$$c_{x,y} := \begin{cases} b_{x,y}, & d(x,y) < R; \\ 0, & d(x,y) \geq R. \end{cases}$$

Note that $\|E_R((b_{x,y})_{x,y \in X})\| \leq K \|(b_{x,y})_{x,y \in X}\|$ (this is a straightforward argument, see e.g., the proof of [28, Lemma 8.1]), so that in particular, E_K is continuous. Also note that $E_K(a) = a$. Since $a \in B_0$, it is a limit of a sequence of operators $b_n = (b_{x,y}^n)_{x,y \in X}$ such that for each n , there exists K_n bounding the rank of $b_{x,y}^n$ over all $x, y \in X$. Thus the same bound K_n applies to $E_K(b_n)$ so that $E_K(b_n) \in UC^*(X)$. By continuity of E_K , $a = \lim_{n \rightarrow \infty} E_K(b_n) \in UC^*(X)$.

In this example, we also have $\mathcal{K}(X, B_0) = \mathcal{K}(\mathcal{H})$, and since $B_0 \subseteq B$ (from (ii)), $C_0(X)B \subseteq \mathcal{K}(\mathcal{H}) = \mathcal{K}(X, B_0)$

(iv) Generalizing (ii), let X be any proper metric space and let \mathcal{H} be an adequate X -module in the sense of [22, Definition 3.4]. Recall that an operator $b \in \mathcal{B}(\mathcal{H})$ is *locally compact* if $C_0(X)b, bC_0(X) \subseteq \mathcal{K}(\mathcal{H})$. Set B equal to the set of all locally compact, bounded operators. One can easily see that $C_b(X)BC_b(X) = B$; it is also true that B is closed under block cutdowns.

To see this, let $b \in \mathcal{B}(\mathcal{H})$ be locally compact with $\|b\| \leq 1$, let $(e_j)_{j \in J}$ be an equicontinuous family of positive contractions in $C_b(X)$ with pairwise disjoint supports, and set $b' := \theta_{(e_j)_{j \in J}}(b)$, which we must prove is locally compact. As $\mathcal{K}(\mathcal{H})$ is closed, it suffices to show that for any $f \in C_c(X)$ with $\|f\| \leq 1$, $fb', b'f \in \mathcal{K}(\mathcal{H})$. Given $\epsilon > 0$, note that

$$b' \approx_{2\epsilon} \theta_{((e_j - \epsilon)_+)_{j \in J}}(b),$$

where $(e_j - \epsilon)_+ \in C_b(X)$ is given by $(e_j - \epsilon)_+(x) := \max\{e_j(x) - \epsilon, 0\}$. By equicontinuity and pairwise disjointness of the family (e_j) , we may choose δ such that if $d(x, y) < \delta$ and $j \neq j'$, then at most one of $e_j(x)$ or $e_{j'}(y)$ can be nonzero. Thus if $f \in C_c(X)$, then by compactness of its support, there are only finitely many j for which $f(e_j - \epsilon)_+ \neq 0$. Consequently,

$$\begin{aligned} fb' &\approx_{2\epsilon} f\theta_{((e_j - \epsilon)_+)_{j \in J}}(b) \\ &= f \sum_{j \in J} (e_j - \epsilon)_+ b(e_j - \epsilon)_+ \\ &= \sum_j f(e_j - \epsilon)_+ b(e_j - \epsilon)_+, \end{aligned}$$

and as this is a finite sum of elements of $\mathcal{K}(\mathcal{H})$, it is itself in $\mathcal{K}(\mathcal{H})$. As $\mathcal{K}(\mathcal{H})$ is closed and $\epsilon > 0$ is arbitrary, it follows that $fb' \in \mathcal{K}(\mathcal{H})$. Likewise, $b'f \in \mathcal{K}(\mathcal{H})$, establishing that b' is locally compact, and therefore that B is closed under block cutdowns.

In this example, we get $\text{Roe}(X, B) = C^*(X)$, the Roe algebra, and $\mathcal{K}(X, B) = \mathcal{K}(\mathcal{H}) = C_0(X)B$.

Definition 2.6. *Let X be a metric space. A bounded sequence $(f_n)_{n=1}^\infty$ from $C_b(X)$ is very Lipschitz if, for every $L > 0$, there exists n_0 such that f_n is L -Lipschitz for all $n \geq n_0$. Let $\text{VL}(X)$ denote the set of all very Lipschitz bounded sequences from $C_b(X)$. Define*

$$\text{VL}_\infty(X) := \text{VL}(X) / \{(f_n)_{n=1}^\infty \in \text{VL}(X) \mid \lim_{n \rightarrow \infty} \|f_n\| = 0\}.$$

$\text{VL}(X)$ is a C^* -subalgebra of $l^\infty(\mathbb{N}, C_b(X))$,² and therefore the quotient $\text{VL}_\infty(X)$ is a C^* -subalgebra of $(C_b(X))_\infty$.

E.g., if X is a finitely generated group G with the word metric, then $\text{VL}_\infty(X)$ can be identified with the fixed point algebra of $l^\infty(G)_\infty$ under the action of G induced by left translation on $l^\infty(G)$.

Recall the following definition from [23].

Definition 2.7. *Let X be a proper metric space. A function $g \in C_b(X)$ is a Higson function (also called a slowly oscillating function) if, for every $R > 0$ and $\epsilon > 0$, there exists a compact set $A \subseteq X$ such that for $x, y \in X \setminus A$, if $d(x, y) < R$ then $|g(x) - g(y)| < \epsilon$. The set of all Higson functions on X is denoted $C_h(X)$.*

E.g., if X is a finitely generated group G with the word metric, then $C_h(X) \subseteq l^\infty(X)$ is the preimage of the fixed point algebra of $l^\infty(G)/c_0(G)$ under the action of G induced by left translation on $l^\infty(G)$.

In the following, \mathcal{H} is an X -module, and we view both $\text{VL}_\infty(X)$ and $B \subseteq \mathcal{B}(\mathcal{H})$ as C^* -subalgebras of $\mathcal{B}(\mathcal{H})_\infty$, and consider the relative commutant

$$B \cap \text{VL}_\infty(X)'$$

It is easy to see that any finite propagation operator commutes with $\text{VL}_\infty(X)$, and by taking limits it follows that

$$\text{Roe}(X, B) \subseteq B \cap \text{VL}_\infty(X)'.$$

The main result is as follows. Recall that straight finite decomposition complexity, as introduced in [4], is a weakening of finite asymptotic dimension ([14, Theorem 4.1]). Both properties are defined in the following subsection.

²To check that the product of two very Lipschitz sequences is itself very Lipschitz, use the fact that if f, g are bounded functions, such that f is L -Lipschitz and g is L' -Lipschitz, then fg is $(\|f\|L' + \|g\|L)$ -Lipschitz.

Theorem 2.8. *Let X be a proper metric space, \mathcal{H} an X -module, and let $B \subseteq \mathcal{B}(\mathcal{H})$ be a C^* -subalgebra such that $C_b(X)BC_b(X) = B$, which is closed under block cutdowns, and such that (2.2) holds. For $b \in B$, the following are equivalent.*

- (i) $[b, f] = 0$ for all $f \in \text{VL}_\infty(X)$;
- (ii) b is quasi-local (it has finite ϵ -propagation for every $\epsilon > 0$);
- (iii) $[b, g] \in \mathcal{K}(X, B)$ (i.e., b essentially commutes with g) for all $g \in C_h(X)$.

If X has straight finite decomposition complexity, then these are also equivalent to

- (iv) $b \in \text{Roe}(X, B)$.

The equivalence of (i) and (ii) is fairly straightforward, and the equivalence of these conditions with (iii) seems to be known by workers in coarse geometry; we shall however provide a detailed proof for completeness. The implication (iv) \implies (i),(ii),(iii) is straightforward and holds in complete generality.

The implication (i) \implies (iv) was proven by Lange and Rabinovich for the uniform Roe algebra of \mathbb{Z}^d (i.e., the case $X = \mathbb{Z}^d$, $\mathcal{H} = l^2(X)$, and $B = \mathcal{B}(\mathcal{H})$) as in Example 2.5 (i) in [17] (see [19, Proposition 8] for a proof in English).

The result (ii) \implies (iv) was claimed by Roe in a remark on page 20 of [22] under a ‘‘finite dimensionality’’ assumption, but it was later found that his supposed proof was incomplete ([20]). The present paper is to the authors’ knowledge the first complete proof of a more general case (which is even more general than finite asymptotic dimension).

Question 2.9. *Is there a uniformly discrete countable metric space with bounded geometry, for which (i)-(iii) does not imply (iv) of Theorem 2.8?*

2.1. Coarse geometric notions. We collect some terminology from [13, 14, 4].

Definition 2.10. *Let X be a proper metric space, let $Z, Z' \subseteq X$, let \mathcal{X} and \mathcal{Y} be metric families (i.e. at most countable sets of subsets of X), and finally let $R \geq 0$.*

- We shall say that \mathcal{X} is uniformly bounded, if $\sup_{Y \in \mathcal{X}} \text{diam}(Y) < \infty$.
- We shall denote the metric neighbourhood of Z of radius R by $N_R(Z) := \{z \in X \mid d(z, Z) \leq R\}$. We further set

$$N_R(\mathcal{X}) := \{N_R(Y) : Y \in \mathcal{X}\}.$$

- The distance between Z and Z' is $d(Z, Z') := \inf\{d(z, z') : z \in Z, z' \in Z'\}$.

- A family $(Y_j)_{j \in J}$ is R -disjoint if $d(Y_j, Y_{j'}) > R$ for all $j \neq j'$; we write

$$\bigsqcup_{R\text{-disjoint}} Y_j$$

for the union of the Y_j to indicate that the family is R -disjoint.

- We say that Z R -decomposes over \mathcal{Y} , if we can decompose $Z = X_0 \cup X_1$ and

$$X_i = \bigsqcup_{R\text{-disjoint}} X_{ij}, \quad i = 0, 1,$$

such that $X_{ij} \in \mathcal{Y}$ for all i, j .

- We say that \mathcal{X} R -decomposes over \mathcal{Y} , denoted $\mathcal{X} \xrightarrow{R} \mathcal{Y}$, if every $Y \in \mathcal{X}$ R -decomposes over \mathcal{Y} .
- We say that X has asymptotic dimension at most n , if for every $r \geq 0$, we can decompose $X = X_0 \cup \dots \cup X_n$ and

$$X_i = \bigsqcup_{r\text{-disjoint}} X_{ij}, \quad i = 0, \dots, n,$$

such that the metric family $\{X_{ij} \mid i, j\}$ is uniformly bounded.

- We say that X has straight finite decomposition complexity, if for any sequence $0 \leq R_1 < R_2 < \dots$, there exists $m \in \mathbb{N}$ and metric families $\{X\} = \mathcal{X}_0, \mathcal{X}_1, \dots, \mathcal{X}_m$, such that $\mathcal{X}_{i-1} \xrightarrow{R_i} \mathcal{X}_i$ for $i = 1, \dots, m$, and the family \mathcal{X}_m is uniformly bounded.

The notion of straight finite decomposition complexity (sFDC) [4] is a priori weaker than the original notion of finite decomposition complexity of Guentner, Tessera and Yu [13, 14], see [4, Proposition 2.3]. The definition of finite decomposition complexity uses a certain “decomposition game”, which effectively means that the choices of R_i can depend on the previous decompositions $\mathcal{X}_1, \dots, \mathcal{X}_{i-1}$.

Already finite decomposition complexity is weaker than finite asymptotic dimension ([14, Theorem 4.1]).

3. PROOF OF (I) \Leftrightarrow (II)

To prove the main result, we begin with a technical-looking characterization of condition (ii).

Lemma 3.1. *Let X be a proper metric space, let \mathcal{H} be an X -module, and let $a \in \mathcal{B}(\mathcal{H})$. Then $\|[a, f]\| < \epsilon$ for every $f \in \text{VL}_\infty(X)_1$ if and only if there exists $L > 0$ such that $\|[a, f]\| < \epsilon$ whenever $f \in C_b(X)_1$ is L -Lipschitz.*

Remark 3.2. As we shall need to refer to the conclusion of the above lemma later, we shall fix the following notation. In the setup as in the above lemma, we write $a \in \text{Commut}(L, \epsilon)$ if $\|[a, f]\| < \epsilon$ whenever $f \in C_b(X)_1$ is L -Lipschitz.

Proof. The reverse implication is immediate from the definition of $\text{VL}(X)$. For the forward direction, we use a proof by contradiction. Suppose for a contradiction that, for every n there exists $f_n \in C_b(X)_1$ that is $(1/n)$ -Lipchitz and $\|[a, f_n]\| \geq \epsilon$.

Then evidently, $(f_n)_{n=1}^\infty \in \text{VL}(X)$ yet $\lim_\infty \|[a, f_n]\| \geq \epsilon$. This contradicts the hypothesis that $\|[a, \text{VL}_\infty(X)_1]\| < \epsilon$. \square

Proof of Theorem 2.8 (i) \Rightarrow (ii). Suppose that $[b, \text{VL}_\infty(X)_1] = 0$ and let $\epsilon > 0$. By Lemma 3.1, let $b \in \text{Commut}(L, \epsilon)$ (in the notation of Remark 3.2) for some $L > 0$.

We claim that b has ϵ -propagation at most L^{-1} . Certainly, suppose that $f, f' \in C_b(X)_1$ have L -disjoint supports. We may define $g \in C_b(X)$ such that $g|_{\text{supp}f} \equiv 1$, $g|_{\text{supp}f'} \equiv 0$ and g is L -Lipschitz. Hence, $\|[b, g]\| < \epsilon$. Consequently,

$$\|fbf'\| = \|fgbf'\| \leq \|[b, g]\| + \|fbg f'\| < \epsilon + 0,$$

as required. \square

Proof of Theorem 2.8 (ii) \Rightarrow (i). Suppose that b has finite ϵ -propagation for all $\epsilon > 0$. Assume that b is a contraction. We shall verify the condition in Lemma 3.1. Therefore, let $\epsilon > 0$ be given. Pick N such that $6/N < \epsilon/2$. By the hypothesis, let b have $(\epsilon/(2N^2))$ -propagation at most $R > 0$.

Let $f \in C_b(X)_1$ be $(2RN)^{-1}$ -Lipschitz. We claim that $\|[b, f]\| < \epsilon$. Surely, define sets

$$A_1 := f^{-1}\left(\left[0, \frac{1}{N}\right]\right), \quad A_i := f^{-1}\left(\left(\frac{i-1}{N}, \frac{i}{N}\right]\right), \quad i = 2, \dots, N.$$

These sets partition X and, for $|i - j| > 1$, A_i is $(2R)$ -disjoint from A_j . We may find a partition of unity $e_1, \dots, e_N \in C_b(X)$ such that e_i is supported in $N_{R/2}(A_i)$. It follows that the supports of e_i and e_j are R -disjoint when $|i - j| > 1$.

Thus,

$$(3.1) \quad \|e_i b e_j\| < \frac{\epsilon}{2N^2}.$$

Also,

$$(3.2) \quad f \approx_{1/N} \sum_{i=1}^N \frac{i}{N} e_i$$

and so

$$\begin{aligned}
\|[f, b]\| &\stackrel{(3.2)}{\leq} \frac{2}{N} + \left\| \left[\sum_{i=1}^N \frac{i}{N} e_i, b \right] \right\| \\
&= \frac{2}{N} + \left\| \left(\sum_{i=1}^N \frac{i}{N} e_i b \right) \left(\sum_{j=1}^N e_j \right) - \left(\sum_{i=1}^N e_i \right) \left(\sum_{j=1}^N \frac{j}{N} b e_j \right) \right\| \\
&= \frac{2}{N} + \left\| \sum_{i,j=1}^N \left(\frac{i}{N} - \frac{j}{N} \right) e_i b e_j \right\| \\
&\leq \frac{2}{N} + \sum_{|i-j|>1} \|e_i b e_j\| + \left\| \sum_{|i-j|\leq 1} \left(\frac{i}{N} - \frac{j}{N} \right) e_i b e_j \right\|.
\end{aligned}$$

The terms of the first sum are each dominated by $\frac{\epsilon}{2N^2}$ by (3.1), so this entire sum is less than $\epsilon/2$. The second sum can be broken into 4 sums with orthogonal terms (namely, note that when $i = j$, the terms vanish; what remains is $j = i + 1$ and $j = i - 1$, and we break each of these into even and odd parts). Each of the terms of the second sum has norm at most $1/N$; thus, we have

$$\|[f, b]\| < \frac{2}{N} + \frac{\epsilon}{2} + \frac{4}{N} < \epsilon,$$

as required. \square

4. PROOF OF (I) \Rightarrow (IV)

In this section, we prove that (i) \Rightarrow (iv) in Theorem 2.8. We begin by establishing a few general functional analytic facts.

Recall that the *strong* topology* on $\mathcal{B}(\mathcal{H})$ is the one in which a net (a_α) converges to $a \in \mathcal{B}(\mathcal{H})$ if and only if both $a_\alpha \rightarrow a$ and $a_\alpha^* \rightarrow a^*$ in the strong operator topology (i.e., $\|a_\alpha \xi - a \xi\| \rightarrow 0$ and $\|a_\alpha^* \xi - a^* \xi\| \rightarrow 0$ for every $\xi \in \mathcal{H}$). A *conditional expectation* from C^* -algebra A to a C^* -subalgebra B is a completely positive and contractive projection E from A to B satisfying

$$E(b_1 a b_2) = b_1 E(a) b_2$$

for all $b_1, b_2 \in B$ and $a \in A$.

Lemma 4.1. *Let \mathcal{H} be a Hilbert space and let G be a subgroup of the group of unitary operators, which is compact in the strong* topology. Then there is a unique conditional expectation $E_G : \mathcal{B}(\mathcal{H}) \rightarrow G'$ whose restriction to the unit ball is weak operator topology continuous. It satisfies*

$$(4.1) \quad \|E_G(a) - a\| \leq \sup_{u \in G} \|[a, u]\|, \quad a \in \mathcal{B}(\mathcal{H}).$$

Proof. Let μ_G be the normalized Haar measure on G (under the strong* topology). Fix $a \in \mathcal{B}(H)$, and consider the map $G \rightarrow \mathcal{B}(\mathcal{H})$ defined by $u \mapsto u^*au$. Then, with the strong* topology on the domain G and the weak operator topology on the range $\mathcal{B}(\mathcal{H})$, this map is continuous. We may therefore integrate, defining

$$E_G(a) := \text{WOT-}\int_G u^*au \, d\mu_G(u).$$

(Here, $\text{WOT-}\int_G \cdot d\mu_G$ indicates the Pettis integral, i.e., $E_G(a)$ is the unique operator satisfying

$$\langle E_G(a)\xi, \eta \rangle = \int_G \langle u^*au\xi, \eta \rangle,$$

for all $\xi, \eta \in \mathcal{H}$.) Using invariance of the Haar measure μ_G , one easily sees that $E_G(a)$ commutes with all of G .

We now check (4.1); for this, set $\gamma := \sup_{u \in G} \|[a, u]\|$. For $\eta, \xi \in \mathcal{H}$,

$$\begin{aligned} |\langle (E_G(a) - a)\eta, \xi \rangle| &= \left| \int_G \langle (u^*au - a)\eta, \xi \rangle \, d\mu_G(u) \right| \\ &\leq \int_G |\langle (u^*au - a)\eta, \xi \rangle| \, d\mu_G(u) \\ &\leq \int_G \|u^*au - a\| \|\eta\| \|\xi\| \, d\mu_G(u) \\ &= \int_G \|[u, a]\| \|\eta\| \|\xi\| \, d\mu_G(u) \\ &\leq \gamma \|\eta\| \|\xi\|. \end{aligned}$$

Therefore, (4.1) follows.

In particular, we conclude that if $a \in G'$ then $E_G(a) = a$. It is also straightforward to see that the function E_G is c.p.c., and therefore it is a conditional expectation.

On the unit ball of $\mathcal{B}(\mathcal{H})$, the integral defining E_G can be uniformly approximated in the weak operator topology by (finite) Riemann sums, which themselves are continuous in the weak operator topology. It follows that the restriction of E_G to the unit ball is continuous using the weak operator topology.

If $E : \mathcal{B}(\mathcal{H}) \rightarrow G'$ is another conditional expectation whose restriction to the unit ball is weak operator topology continuous, then for a

contraction $a \in \mathcal{B}(\mathcal{H})$,

$$\begin{aligned}
E_G(a) &= E(E_G(a)) && (E \text{ fixes } G') \\
&= E\left(\text{WOT-}\int_G u^* a u \, d\mu_G(u)\right) \\
&= \text{WOT-}\int_G E(u^* a u) \, d\mu_G(u) && (\text{WOT-continuity of } E|_{\mathcal{B}(\mathcal{H})_1}) \\
&= \text{WOT-}\int_G u^* E(a) u \, d\mu_G(u) && (E \text{ is a conditional expectation}) \\
&= E_G(E(a)) \\
&= E(a) && (E_G \text{ fixes } G').
\end{aligned}$$

Thus, $E = E_G$. □

Recall that an *atomic* abelian von Neumann algebra is a von Neumann algebra isomorphic to $l^\infty(X)$, for some set X . In the following, when $\mathcal{H} = l^2(X)$, then the conditional expectation $\mathcal{B}(l^2(X)) \rightarrow l^\infty(X)$ consists simply of taking an operator to its diagonal.

Corollary 4.2. *Let $D \subset \mathcal{B}(\mathcal{H})$ be an atomic abelian von Neumann algebra. Then there is a unique conditional expectation $E_D : \mathcal{B}(\mathcal{H}) \rightarrow D'$ whose restriction to the unit ball is weak operator topology continuous. It satisfies*

$$(4.2) \quad \|E_D(a) - a\| \leq \sup_{x \in D, \|x\| \leq 1} \|[a, x]\|, \quad a \in \mathcal{B}(\mathcal{H}).$$

Proof. Without loss of generality, D contains the identity operator. D is generated by a family of orthogonal projections $(p_j)_{j \in J}$, whose sum converges strongly to 1. Define

$$G := \left\{ \sum_{j \in J} (-1)^{\alpha_j} p_j : (\alpha_j)_{j \in J} \in (\mathbb{Z}/2)^J \right\}.$$

This is a strong* compact subgroup of the unitary group of D (it is homeomorphic to $(\mathbb{Z}/2)^J$ with the product topology), so that Lemma 4.1 applies to it. It is clear that it generates D as a von Neumann algebra, so that $G' = D'$. The conclusion follows from Lemma 4.1. □

Corollary 4.3. *Let X be a proper metric space, let \mathcal{H} an X -module, and let $a \in \mathcal{B}(\mathcal{H})$. Suppose $a \in \text{Commut}(L, \epsilon)$ for some $L, \epsilon > 0$ (in the notation of Remark 3.2). Let $(e_j)_{j \in J}$ be a family of positive contractions from $C_b(X)$ with $(2L^{-1})$ -disjoint supports, and define $e := \sum_{j \in J} e_j$. Then, with $\theta_{(e_j)_{j \in J}}$ from Definition 2.2,*

$$\|eae - \theta_{(e_j)_{j \in J}}(a)\| \leq \epsilon.$$

Proof. Set A_j equal to the support of e_j for each $j \in J$. We may find pairwise disjoint projections $p_j \in \mathcal{B}(\mathcal{H})$, for $j \in J$, such that p_j acts as a unit on e_j . Define D to be the von Neumann subalgebra generated by $\{p_j : j \in J\}$ (with unit $1_D = \sum_j p_j$), and let $E_D : \mathcal{B}(\mathcal{H}) \rightarrow D'$ be the unique conditional expectation provided by Corollary 4.2. Then one finds that for $x \in \mathcal{B}(\mathcal{H})$, $E_D(x) = \sum_{j \in J} p_j x p_j$ (converging in the strong operator topology), and therefore

$$E_D(eae) = \theta_{(e_j)_{j \in J}}(a).$$

Using $(2L^{-1})$ -disjointness of the family $(A_j)_{j \in J}$, for $f \in D_1$, there exists a function $\tilde{f} \in C_b(X)_1$ that is L -Lipschitz such that

$$f = 1_D \tilde{f}.$$

Therefore,

$$\begin{aligned} f(eae) &= \tilde{f}eae \\ &= e\tilde{f}ae \\ &\approx_\epsilon ea\tilde{f}e \\ &= (eae)f. \end{aligned}$$

Hence, by Corollary 4.2,

$$eae \approx_\epsilon E_D(eae) = \theta_{(e_j)_{j \in J}}(a),$$

as required. \square

Definition 4.4. Let X be a proper metric space, \mathcal{H} be an X -module, $a \in \mathcal{B}(\mathcal{H})$ and let \mathcal{X} be a metric family (of subsets of X). We say that a is block diagonal with respect to \mathcal{X} , if there exists an equicontinuous family $(e_j)_{j \in J}$ of positive contractions in $C_b(X)$ with pairwise disjoint supports, such that $a = \theta_{(e_j)_{j \in J}}(a)$, and the support of each e_j is contained in some set $Y_j \in \mathcal{X}$. Furthermore, in this case we shall denote $a_{Y_j} := e_j a e_j$ and call these operators blocks of a .

The next lemma sets up the ‘‘induction step’’ to be applied in the context of the decomposition game, in the proof of Theorem 2.8 (i) \Rightarrow (iv).

Lemma 4.5. Let X be a proper metric space, and let \mathcal{Y} be a metric family such that $\{X\} \xrightarrow{4L^{-1}+4} \mathcal{Y}$ for some $L > 0$. Let \mathcal{H} be an X -module, and let $a \in \mathcal{B}(\mathcal{H})$. Let $\epsilon > 0$ be such that $a \in \text{Commut}(L, \epsilon)$. Then we can write

$$a \approx_{8\epsilon} a_{00} + a_{01} + a_{10} + a_{11},$$

where each $a_{ii'}$ is of the form $\theta_{(f_k)_{k \in K}}(gag')$ for some contractions $g, g' \in C_b(X)$ and some family $(f_k)_{k \in K}$ of 1-Lipschitz positive contractions

in $C_b(X)$ with disjoint supports, such that the support of each f_k is contained in a set in $N_{L^{-1}+1}(\mathcal{Y})$.

Proof. By the decomposition assumption, we can write

$$X = X^{(0)} \cup X^{(1)}, \quad X^{(i)} = \bigsqcup_{\substack{j \in J \\ (4L^{-1}+4)\text{-disjoint}}} X_j^{(i)}, \quad i = 0, 1,$$

with $X_j^{(i)} \in \mathcal{Y}$ for each i, j . We may find a partition of unity consisting of 1-Lipschitz positive contractions $e_j^{(i)} \in C_b(X)$, for $i = 0, 1$ and $j \in J$, such that the support of $e_j^{(i)}$ is contained in $N_1(X_j^{(i)})$. It follows that for each i , the supports of $(e_j^{(i)})_{j \in J}$ are $(4L^{-1} + 2)$ -disjoint.

For each $i = 0, 1$, define $e^{(i)} := \sum_{j \in J} e_j^{(i)}$. Since

$$a = e^{(0)} a e^{(0)} + e^{(0)} a e^{(1)} + e^{(1)} a e^{(0)} + e^{(1)} a e^{(1)},$$

it suffices to find $a_{ii'} \approx e^{(i)} a e^{(i')}$ for each $i, i' \in \{0, 1\}$ with the required properties. (We will be precise about the degree of approximation – in short, it depends on whether i and i' are equal.)

For the case $i = i'$, Corollary 4.3 shows that

$$e^{(i)} a e^{(i)} \approx_{\epsilon} \theta_{(e_j^{(i)})_{j \in J}}(e^{(i)} a e^{(i)}) =: a_{ii}.$$

The latter operator is clearly block diagonal with respect to $N_1(\mathcal{Y})$ (hence also with respect to $N_{L^{-1}+1}(\mathcal{Y})$).

Turning now to the case $i \neq i'$, note that for fixed i , the family $(N_{L^{-1}+1}(X_j^{(i)}))_{j \in J}$ is $(2L^{-1} + 2)$ -disjoint. For each i, j , there exists $\hat{e}_j^{(i)} \in C_b(X)$ that is L -Lipschitz, that acts as the identity on $e_j^{(i)}$, and is supported on $N_{L^{-1}+1}(X_j^{(i)})$. For each i , define $\hat{e}^{(i)} := \sum_{j \in J} \hat{e}_j^{(i)}$. We have

$$\begin{aligned} e^{(i)} a e^{(i')} &= e^{(i)} \hat{e}^{(i)} a \hat{e}^{(i')} e^{(i')} \\ &\approx_{2\epsilon} e^{(i)} \hat{e}^{(i')} a \hat{e}^{(i)} e^{(i')}. \end{aligned}$$

For each $j, j' \in J$, there exists a 1-Lipschitz positive contraction $f_{j,j'} \in C_b(X)$ that is 1 on $N_{L^{-1}+1}(X_j^{(i)}) \cap N_{L^{-1}+1}(X_{j'}^{(i')})$, and is supported on the metric neighbourhood of this set of radius 1. In particular, the support of each $f_{j,j'}$ is contained in a set in $N_{L^{-1}+1}(\mathcal{Y})$, the family of supports of the family $(f_{j,j'})_{j,j' \in J}$ is $(2L^{-1})$ -disjoint, and $f := \sum_{j,j'} f_{j,j'}$ acts as an identity on (both sides of) $e^{(i)} \hat{e}^{(i')} a \hat{e}^{(i)} e^{(i')}$. Applying Corollary 4.3, we obtain

$$f e^{(i)} \hat{e}^{(i')} a \hat{e}^{(i)} e^{(i')} f \approx_{\epsilon} \theta_{(f_{j,j'})_{j,j' \in J}}(e^{(i)} \hat{e}^{(i')} a \hat{e}^{(i)} e^{(i')}).$$

Thus,

$$\begin{aligned} e^{(i)} a e^{(i')} &\approx_{2\epsilon} e^{(i)} \hat{e}^{(i')} a \hat{e}^{(i)} e^{(i')} \\ &= f e^{(i)} \hat{e}^{(i')} a \hat{e}^{(i)} e^{(i')} f \\ &\approx_{\epsilon} \theta_{(f_{j,j'})_{j,j' \in J}} \left(e^{(i)} \hat{e}^{(i')} a \hat{e}^{(i)} e^{(i')} \right) =: a_{ii'}. \end{aligned}$$

By construction, it is clear that $a_{ii'}$ is block diagonal with respect to $N_{L^{-1}+1}(\mathcal{Y})$.

Summarizing, we have $a \approx_{\epsilon+3\epsilon+3\epsilon+\epsilon} a_{00} + a_{10} + a_{01} + a_{11}$, and all the $a_{ii'}$ are of the right form. \square

We now strengthen the previous lemma by allowing an arbitrary metric family in place of $\{X\}$ and with a a correspondingly block diagonal operator.

Lemma 4.6. *Let X be a proper metric space, and let \mathcal{X} and \mathcal{Y} be metric families, such that $\mathcal{X} \xrightarrow{4L^{-1}+4} \mathcal{Y}$ for some $L > 0$. Let \mathcal{H} be an X -module, and let $a \in \mathcal{B}(\mathcal{H})$ be block diagonal with respect to \mathcal{X} . Let $\epsilon > 0$ be such that $a \in \text{Commut}(L, \epsilon)$. Then we can write*

$$(4.3) \quad a \approx_{8\epsilon} a_{00} + a_{01} + a_{10} + a_{11},$$

where each $a_{ii'}$ is of the form $\theta_{(f_k)_{k \in K}}(g a g')$ for some contractions $g, g' \in C_b(X)$ and some equicontinuous family $(f_k)_{k \in K}$ of positive contractions in $C_b(X)$ with disjoint supports, such that the support of each f_k is contained in a set in $N_{L^{-1}+1}(\mathcal{Y})$. In particular:

- (i) each $a_{ii'}$ is block diagonal with respect to $N_{L^{-1}+1}(\mathcal{Y})$,
- (ii) if $a \in \text{Commut}(L', \epsilon')$ for some $L', \epsilon' > 0$, then each $a_{ii'}$ is in $\text{Commut}(L', \epsilon')$ as well, and
- (iii) if $B \subseteq \mathcal{B}(\mathcal{H})$ is a C^* -subalgebra such that $C_b(X) B C_b(X) = B$ and B is closed under block cutdowns, and if a is in B , then each $a_{ii'}$ is in B as well.

Proof. Without loss of generality, both \mathcal{X} and \mathcal{Y} are closed under taking subsets. Start by letting (e_j) be an equicontinuous family of positive contractions in $C_b(X)$ with disjoint supports, such that $Y_j := \text{supp}(e_j) \in \mathcal{X}$. Applying Lemma 4.5 to each a_{Y_j} yields

$$a_{Y_j} \approx_{8\epsilon} a_{00}^j + a_{01}^j + a_{10}^j + a_{11}^j,$$

satisfying the conclusions of that lemma. Set

$$a_{ii'} := \sum_j a_{ii'}^j.$$

Then (4.3) follows from (2.1)

To see that each $a_{ii'}$ has the right form, fix i and i' . For each j , there exist contractions $g_j, g'_j \in C_b(X)$ and a family $(f_{j,k})_{k \in K_j}$ of 1-Lipschitz

positive contractions in $C_b(Y_j)$ with disjoint supports, such that

$$a_{ii'}^j = \theta_{(f_{j,k})_{k \in K_j}}(g_j a g_j').$$

and the support of each $f_{j,k}$ is of the form $Y \cap Z$ for some $Z \in N_{L^{-1}+1}(\mathcal{Y})$.

Then observe that

$$a_{ii'} = \theta_{(e_j^{1/4} f_{j,k})_{j \in J, k \in K_j}} \left(\left(\sum_j e_j^{1/4} g_j \right) a \left(\sum_j e_j^{1/4} g_j \right) \right),$$

where the family appearing in this block-cutdown formula, namely

$$(e_j^{1/4} f_{j,k})_{j \in J, k \in K_j}$$

is equicontinuous and contained in $N_{L^{-1}+1}(\mathcal{Y})$.

(i)-(ii) are immediate consequences of the form that $a_{ii'}$ takes. Since multiplication by $C_b(X)$ preserves block structure, and using (2.1), (iii) can also be seen to be a consequence of the form that $a_{ii'}$ takes. \square

We have seen in the previous lemma that we will need to work with “thickened” metric families, so we record the following straightforward observation.

Lemma 4.7. *Let \mathcal{X} and \mathcal{Y} be metric families, $R, S \geq 0$. Assume that $\mathcal{X} \xrightarrow{R} \mathcal{Y}$ and that $R - 2S \geq 0$. Then $N_S(\mathcal{X}) \xrightarrow{R-2S} N_S(\mathcal{Y})$.*

Proof of Theorem 2.8 (i) \Rightarrow (iv). Recall that we are given an operator b on an X -module \mathcal{H} satisfying $[b, f] = 0$ for all $f \in \text{VL}_\infty(X)$. Given $\epsilon > 0$, our task is to produce a finite propagation operator in B which is ϵ -far from b . Lemma 3.1 provides us with L_n for every

$$\epsilon_n := \epsilon / (2 \cdot 8^n),$$

such that $b \in \text{Commut}(L_n, \epsilon_n)$. Let

$$R_n := 4(L_n^{-1} + 1) + 2(L_{n-1}^{-1} + 1) + \cdots + 2(L_1^{-1} + 1).$$

As X has straight finite decomposition complexity (see Definition 2.10), we obtain metric families $\{X\} = \mathcal{X}_0, \mathcal{X}_1, \dots, \mathcal{X}_m$, such that $\mathcal{X}_{n-1} \xrightarrow{R_n} \mathcal{X}_n$ for $n \in \{1, \dots, m\}$ and \mathcal{X}_m is uniformly bounded. Note that Lemma 4.7 gives us that

$$N_{(L_n^{-1}+1)+\cdots+(L_1^{-1}+1)}(\mathcal{X}_{n-1}) \xrightarrow{4(L_n^{-1}+1)} N_{(L_n^{-1}+1)+\cdots+(L_1^{-1}+1)}(\mathcal{X}_n).$$

Thus, we can inductively apply Lemma 4.6, with L_n, ϵ_n , the operators obtained in the previous iteration, and metric families from the above display. After m steps, we will have approximated the operator b by an operator b' which is a sum of finitely many (4^m to be precise) operators in B which are block diagonal with respect to

$$N_{(L_m^{-1}+1)+\cdots+(L_1^{-1}+1)}(\mathcal{X}_m).$$

Since \mathcal{X}_m is uniformly bounded, so is the above family, and therefore operators which are block diagonal with respect to it have finite propagation; consequently, $b' \in \text{Roe}(B, X)$. Tracing through the estimates given by Lemma 4.6, we compute that the distance from b to b' is at most

$$8\epsilon_1 + 4(8\epsilon_2 + 4(8\epsilon_3 + 4(\dots))) = \epsilon \left(\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots \right) = \epsilon.$$

This finishes the proof. \square

Remark 4.8. When the asymptotic dimension of X is at most $d < \infty$, the induction component of the above proof can be removed: one can use the idea of Lemma 4.5 with a decomposition of X into $d + 1$ (instead of 2) uniformly bounded, $(4L^{-1} + 4)$ -disjoint families, and correspondingly approximate a be a sum of $(d + 1)^2$ block diagonal operators.

5. HIGSON FUNCTIONS

To prepare to prove (i) \Leftrightarrow (iii) of Theorem 2.8, we begin by considering a special class of Higson functions which are more closely related to our definition of $\text{VL}_\infty(X)$.

Definition 5.1. *Let X be a proper metric space. A function $g \in C_b(X)$ is a Lipschitz–Higson function if, for every $L > 0$, there exists a compact set $A \subseteq X$ such that $g|_{X \setminus A}$ is L -Lipschitz. The set of all Lipschitz–Higson functions on X is denoted $C_{lh}(X)$.*

Fix a proper metric space X and a point $x_0 \in X$. For $R > 0$, define $e_R \in C_0(X)$ by

$$(5.1) \quad e_R(x) := \max\{0, 1 - d(x, B_R(x_0))/R\}.$$

Observe that e_R is R^{-1} -Lipschitz, is 1 on $\bar{B}_R(x_0)$, and vanishes outside of $B_{2R}(x_0)$.

Lemma 5.2. $C_h(X) = C_{lh}(X) + C_0(X)$.

Proof. The inclusion \supseteq is straightforward. To go the other direction, let $g \in C_h(X)$. We shall produce $f \in C_{lh}(X)$ such that $f - g \in C_0(X)$. Without loss of generality, g is a positive contraction. Fix a point $x_0 \in X$.

Recursively define $R_0 := 0$ and $R_n \geq \max\{2(n+1), 2R_{n-1}\}$ such that if $x, y \in X \setminus B_{R_n}(x_0)$ and $d(x, y) < n + 1$ then $|g(x) - g(y)| < \frac{1}{2(n+1)}$. Using e_R from (5.1), set

$$g_1 := e_{R_1}g, \quad g_n := (e_{R_n} - e_{R_{n-1}})g, \quad n \geq 2.$$

Note that, for $n \geq 2$, g_n is such that if $d(x, y) < n$ then $|g_n(x) - g_n(y)| < \frac{1}{n}$, and that

$$(5.2) \quad g = \sum_{n=1}^{\infty} g_n,$$

converging pointwise (as at each point, at most two terms of the sum are nonzero).

Define $f_1 := g_1$.

Fix $n \geq 2$; we shall define a function f_n which approximates g_n , but is more Lipschitz. Define

$$A_i := g_n^{-1} \left[\frac{i}{n}, 1 \right], \quad i = 1, \dots, n,$$

and define $c_i \in C_b(X)$ by

$$c_i(x) := \max \left\{ 1 - \frac{d(x, A_i)}{n}, 0 \right\}, \quad i = 1, \dots, n,$$

which is $(1/n)$ -Lipschitz. Set

$$f_n := \frac{1}{n} \sum_{i=1}^n c_i,$$

which is also $(1/n)$ -Lipschitz, as it is an average of such. Moreover, $\|f_n - g_n\| \leq \frac{1}{n}$, and the support of g_n is contained in the support of f_n .

Set $f := \sum_{n=1}^{\infty} f_n$; as in (5.2), at each point, at most two terms of the sum are nonzero. Using this fact, one sees that $\sum_{n=n_0}^{\infty} f_n$ is $(2n_0)^{-1}$ -Lipschitz, for all $n_0 \geq 2$. Moreover, f agrees with this tail sum outside of $B_{2R_{n_0-1}}$, which proves that $f \in C_{lh}(X)$.

Similarly, since $f - g$ agrees with the tail $\sum_{n=n_0}^{\infty} (f_n - g_n)$ outside of $B_{2R_{n_0-1}}$, and this tail has norm at most $\frac{2}{n}$, it follows that $f - g \in C_0(X)$. \square

Remark 5.3. We note in passing that, due to the previous lemma, the Higson corona νX (defined as the compact Hausdorff space satisfying $C(\nu X) \cong C_h(X)/C_0(X)$) satisfies

$$C(\nu X) \cong C_{lh}(X)/C_0(X).$$

Now we set out two constructions to be used, producing a Lipschitz–Higson function from a very Lipschitz sequence and vice versa. Neither construction is canonical: both depend on a number of choices.

Let $(f_k)_{k=1}^{\infty} \in \text{VL}(X)$, let $(f_{k_i})_{i=1}^{\infty}$ be a subsequence, and let $(R_i)_{i=0}^{\infty} \subset (0, \infty)$ be a sequence such that $R_{i+1} \geq 6R_i$ for each i . From these, and using e_R from (5.1), define

$$(5.3) \quad g_{x_0, (f_{k_i})_{i=1}^{\infty}, (R_i)_{i=0}^{\infty}} := \sum_{i=1}^{\infty} f_{k_i} (e_{R_i} - e_{3R_{i-1}}) \in C_b(X).$$

Note that the functions in the summation are pairwise disjoint, so we can treat the series as converging pointwise. It is straightforward to see that $\|g_{x_0, (f_{k_i})_{i=1}^\infty, (R_i)_{i=0}^\infty}\| \leq \|(f_k)_{k=1}^\infty\|$.

Lemma 5.4. *Fix a proper metric space X and a point $x_0 \in X$. For $(f_k)_{k=1}^\infty \in \text{VL}(X)$, a subsequence $(f_{k_i})_{i=1}^\infty$, and a sequence $(R_i)_{i=0}^\infty \subset (0, \infty)$ such that $R_{i+1} \geq 6R_i$ for each i , let $g_{x_0, (f_{k_i})_{i=1}^\infty, (R_i)_{i=0}^\infty}$ be as defined in (5.3). Then*

$$g_{x_0, (f_{k_i})_{i=1}^\infty, (R_i)_{i=0}^\infty} \in C_{lh}(X).$$

Proof. Without loss of generality, we may assume $\|f_k\| \leq 1$ for all k . For ease of notation, we set

$$g := g_{x_0, (f_{k_i})_{i=1}^\infty, (R_i)_{i=0}^\infty}.$$

As in the definition of a Lipschitz–Higson function, let $L > 0$ be given. Pick i_0 such that $3R_{i_0-1} > 2/L$ and such that f_i is $(L/2)$ -Lipschitz for all $i \geq i_0$. We will be done when we show that the restriction of g to $X \setminus B_{R_{i_0}}(x_0)$ is L -Lipschitz.

For $i \geq i_0$, $e_{R_i} - e_{3R_{i-1}}$ is $(L/2)$ -Lipschitz, so that the product

$$h_i := f_{k_i}(e_{R_i} - e_{3R_{i-1}})$$

is L -Lipschitz. For $x, y \in X \setminus B_{R_{i_0}}(x_0)$, note that, by the definition of e_R and the condition $R_{i+1} \geq 6R_i$, that at least one of the following conditions holds.

- (i) $d(x, y) \geq R_{i_0}$, or
- (ii) There exists i such that $g(x) = h_i(x)$ and $g(y) = h_i(y)$.

In the first case,

$$|g(x) - g(y)| \leq 2\|g\| = 2 \leq 3LR_{i_0-1} \leq 3LR_{i_0} \leq Ld(x, y).$$

In the second case, since h_i is L -Lipschitz, it follows that $|g(x) - g(y)| \leq Ld(x, y)$. \square

Next, let $g \in C_h(X)$ be given, along with a sequence $(R_k) \in (0, \infty)$ such that $\lim_{k \rightarrow \infty} R_k = \infty$. From this data, define (using e_R from (5.1))

$$(5.4) \quad F_{x_0, g, (R_k)_{k=1}^\infty} := (f_k)_{k=1}^\infty \in l^\infty(\mathbb{N}, l^\infty(X)) \quad \text{where} \quad f_k := (1 - e_{R_k})g.$$

Lemma 5.5. *Fix a proper metric space X and a point $x_0 \in X$. For a Higson function $g \in C_{lh}(X)$ and a sequence $(R_k) \in (0, \infty)$ such that $\lim_{k \rightarrow \infty} R_k = \infty$, let $F_{g, (R_k)_{k=1}^\infty}$ be as defined in (5.4). Then*

$$F_{g, (R_k)_{k=1}^\infty} \in \text{VL}(X).$$

Proof. Without loss of generality, we may assume $\|g\| \leq 1$. As in the definition of $\text{VL}(X)$, let $L > 0$ be given. Pick $M \geq 2/L$ such that g is $(L/2)$ -Lipschitz on $X \setminus B_M(x_0)$. Pick k_0 such that $R_k \geq M$ for all $k \geq k_0$. For $k \geq k_0$, using the fact that $(1 - e_{R_k})$ is $(L/2)$ -Lipschitz

and vanishes on $B_M(x_0)$, it is easy to see that $f_k = g(1 - e_{R_k})$ is L -Lipschitz. \square

Proof of Theorem 2.8 (i) \Rightarrow (iii). By Lemma 5.2, (iii) is equivalent to $[g, b] \in \mathcal{K}(X, B)$ for all $g \in C_{lh}(X)$, which is the statement we will prove assuming (i). Assume that $[b, g] \notin \mathcal{K}(X, B)$ for some $g \in C_{lh}(X)$. Set $\epsilon := \|[b, g] + \mathcal{K}(X, B)\|$. Fix a point $x_0 \in X$, let $(R_k)_{k=1}^\infty \subset (0, \infty)$ be any sequence such that $\lim_{i \rightarrow \infty} R_i = \infty$, and define $(f_k)_{k=1}^\infty := F_{x_0, g, (R_k)_{k=1}^\infty}$ as in (5.4), that is,

$$f_k := (1 - e_{R_k})g.$$

By Lemma 5.5, $(f_k)_{k=1}^\infty \in \text{VL}(X)$. Since $e_{R_k} \in C_0(X)$ for each k , using the condition (2.2) we obtain

$$[f_k, b] = [g, b] - [e_{R_k}g, b] \in [g, b] + \mathcal{K}(X, B)$$

and therefore by the definition of ϵ ,

$$\|[f_k, b]\| \geq \epsilon.$$

Consequently, $\lim_{k \rightarrow \infty} \|[f_k, b]\| \geq \epsilon$, so that b does not commute with the image of $(f_k)_{k=1}^\infty$ in $\text{VL}_\infty(X)$. \square

Before we embark on the proof of Theorem 2.8 (iii) \implies (i), note that as a consequence of the Spectral Theorem, we may extend the X -module structure on $\mathcal{B}(\mathcal{H})$ from bounded continuous functions to bounded Borel functions on X . This is convenient in the following proof, since it allows us to easily “cut up” operators on \mathcal{H} using characteristic functions of Borel sets in X . (Of course, we cannot assume the algebra B is closed under multiplication by these bounded Borel functions.) We opt for this approach for the sake of readability, although it is possible to modify the proof to only use continuous functions for the price of more approximations.

Proof of Theorem 2.8 (iii) \Rightarrow (i). Since each of $\text{VL}_\infty(X)$, $C_h(X)$, and $\mathcal{K}(X, B)$ is $*$ -closed, it suffices to prove the theorem in the case that b is self-adjoint. We henceforth assume that $b = b^*$. Fix a point $x_0 \in X$, and to shorten notation in this proof, set

$$B_R := B_R(x_0).$$

For each $R > 0$, we will use χ_{B_R} to denote the support projection of a function whose cozero set is B_R .

Assume that $[b, f] \neq 0$ for some $f \in \text{VL}_\infty(X)$. Then in fact, $[b, f] \neq 0$ for some $f = (f_k)_{k=1}^\infty$ for which each f_k is a self-adjoint contraction; we fix this sequence. Let $0 < \epsilon < \|[b, f]\|$.

Consider now two cases.

Case 1. There exists $R_0 > 0$ such that for all $S > 0$ there are infinitely many k for which

$$\|\chi_{B_{R_0}}[b, f_k](1 - \chi_{B_S})\| > \frac{\epsilon}{5}.$$

Roughly, for this case, we will construct some g (of the form (5.3)) such that the block-column of $[b, g]$ corresponding to B_{R_0} doesn't converge to 0 at ∞ .

Note that as $k \rightarrow \infty$, $f_k|_{B_{R_0}}$ tends towards being constant; so without loss of generality, we may assume that $f_k|_{B_{R_0}}$ is constant. Adding a scalar to each f_k , we arrive at another sequence $(\hat{f}_k)_{n=1}^\infty$ with the same properties as $(f_k)_{n=1}^\infty$ (that is, self-adjoint and satisfying the Case 1 condition), such that $\hat{f}_k|_{B_{R_0}} \equiv 0$. From this it follows that $\chi_{B_{R_0}}[b, \hat{f}_k] = \chi_{B_{R_0}} b \hat{f}_k$ for all n , so that we have: for all $S > 0$ there exist infinitely many k such that

$$\|\chi_{B_{R_0}} b \hat{f}_k(1 - \chi_{B_S})\| > \frac{\epsilon}{5}.$$

Using R_0 as above and $k_0 := 0$, recursively choose $k_1 < k_2 < \dots$ and $R_1 \geq 6R_0$, $R_2 \geq 6R_1$, \dots as follows. Having chosen R_{i-1} , pick $k_i > k_{i-1}$ such that

$$\|\chi_{B_{R_0}} a \hat{f}_{k_i}(1 - \chi_{B_{6R_{i-1}}})\| > \frac{\epsilon}{5} \quad \text{and} \quad \|\hat{f}_{k_i} \chi_{B_{6R_{i-1}}}\| \leq \frac{\epsilon}{10\|a\|}.$$

(The second inequality can be arranged as \hat{f}_k converge to 0 on any given bounded subset of X .) Then, since $(e_R)_{R=1}^\infty$ converges strongly to 1, pick $R_i \geq 6R_{i-1}$ such that

$$\|\chi_{B_{R_0}} b \hat{f}_{k_i}(1 - \chi_{B_{6R_{i-1}}})e_{R_i}\| > \frac{\epsilon}{5}.$$

Since $e_{R_i} - e_{3R_{i-1}}$ differs from $(1 - \chi_{B_{6R_{i-1}}})e_{R_i}$ only on $B_{6R_{i-1}}$, it follows that

$$\|\chi_{B_{R_0}} b \hat{f}_{k_i}(e_{R_i} - e_{3R_{i-1}})\| > \frac{\epsilon}{10}.$$

Using these recursive choices, define $g := g_{x_0, (\hat{f}_{k_i})_{i=1}^\infty, (R_i)_{i=0}^\infty} \in C_{lh}(X) \subseteq C_h(X)$ by Lemma 5.4. If $[b, g] \in \mathcal{K}(X, B)$, then $\|[b, g](1 - \chi_{B_S})\| \rightarrow 0$ as $S \rightarrow \infty$. However, given $S \geq R_0$, there exists i such that $3R_{i-1} > S$. Then

$$\begin{aligned} \|[b, g](1 - \chi_{B_S})\| &\geq \|\chi_{B_{R_0}}[b, g](1 - \chi_{B_S})(\chi_{B_{2R_i}} - \chi_{B_{3R_{i-1}}})\| \\ &= \|\chi_{B_{R_0}} b \hat{f}_{k_i}(e_{R_i} - e_{3R_{i-1}})\| \\ &> \frac{\epsilon}{10}, \end{aligned}$$

which is a contradiction. This concludes the proof in Case 1.

Case 2. For every $R > 0$, there exists $S > 0$ such that, for all but finitely many $k \in \mathbb{N}$,

$$\|\chi_{B_R}[b, f_k](1 - \chi_{B_S})\| \leq \frac{\epsilon}{5}.$$

Without loss of generality, we may assume that $S > R$.

Roughly, for this case, we will construct some g (of the form (5.3)) such that the certain blocks on the diagonal of $[b, g]$ don't converge to 0 at ∞ .

In preparation for this, suppose we are given $R > 0$ and $K \in \mathbb{N}$. Let S be given by the Case 2 property. Then there exists $k \geq K$ such that

$$\|\chi_{B_R}[b, f_k](1 - \chi_{B_S})\| \leq \frac{\epsilon}{5},$$


and in addition, $\|[b, f_k]\| > \epsilon$ and $f_k|_{B_S}$ is $\frac{\epsilon}{5}$ -approximately constant. From the latter property, it follows that there is a scalar γ such that $f_k|_{B_S} \approx_{\epsilon/10} \gamma$, so that

$$(5.5) \quad \|\chi_{B_S}[b, f_k]\chi_{B_S}\| \leq 2 \cdot \frac{\epsilon}{10} \|b\| \leq \frac{\epsilon}{5}.$$

Since b and f_k are self-adjoint,

$$(5.6) \quad \|(1 - \chi_{B_S})[b, f_k]\chi_{B_R}\| = \|\chi_{B_R}[b, f_k](1 - \chi_{B_S})\| \leq \frac{\epsilon}{5}.$$

We now cut up the operator $T = [b, f_k]$ as follows:



That is, we use the equality

$$\begin{aligned} T &= (1 - \chi_{B_R})T(1 - \chi_{B_R}) + \chi_{B_R}T\chi_{B_S} + \chi_{B_R}T(1 - \chi_{B_S}) \\ &\quad + (\chi_{B_S} - \chi_{B_R})T\chi_{B_R} + (1 - \chi_{B_S})T\chi_{B_R} \end{aligned}$$

and the reverse triangle inequality to deduce

$$\begin{aligned}
& \|(1 - \chi_{B_R})[b, f_k](1 - \chi_{B_R})\| \\
& \geq \| [b, f_k] \| - (\| \chi_{B_R}[b, f_k]\chi_{B_S} \| + \| \chi_{B_R}[b, f_k](1 - \chi_{B_S}) \| \\
& \quad + \| (\chi_{B_S} - \chi_{B_R})[b, f_k]\chi_{B_R} \| + \| (1 - \chi_{B_S})[b, f_k]\chi_{B_R} \|) \\
& \stackrel{(5.5), (5.6)}{\geq} \| [b, f_k] \| - 4 \cdot \frac{\epsilon}{5} \\
& > \epsilon - \frac{4\epsilon}{5} \\
(5.7) \quad & = \frac{\epsilon}{5}.
\end{aligned}$$

In summary, we have shown that for every $R > 0$ and $K \in \mathbb{N}$, there exists $k \geq K$ such that

$$(5.8) \quad \|(1 - \chi_{B_R})[b, f_k](1 - \chi_{B_R})\| > \frac{\epsilon}{5}.$$

Now, start with $R_0 := 1$ and $k_0 := 0$, and (as in Case 1) we will choose $k_1 < k_2 < \dots$ and $R_1 \geq 6R_0, R_2 \geq 6R_1, \dots$ recursively. Given R_{i-1} , pick $k > k_{i-1}$ satisfying (5.8) for $R := 6R_{i-1}$, and set k_i equal to this k . That is, $k_i > k_{i-1}$ satisfies

$$\|(1 - \chi_{B_{6R_{i-1}}})[b, f_{k_i}](1 - \chi_{B_{6R_{i-1}}})\| > \frac{\epsilon}{5}.$$

Then, since $(\chi_{B_R})_{R=1}^\infty$ converges strongly to 1, there exists $R_i \geq 6R_{i-1}$ such that

$$\|\chi_{B_{R_i}}(1 - \chi_{B_{6R_{i-1}}})[b, f_{k_i}](1 - \chi_{B_{6R_{i-1}}})\chi_{B_{R_i}}\| > \frac{\epsilon}{5}.$$

Note that $\chi_{B_{R_i}}(1 - \chi_{B_{6R_{i-1}}}) = \chi_{B_{R_i}} - \chi_{B_{6R_{i-1}}}$.

Using these recursive choices, define $g := g_{x_0, (f_{k_i})_{i=1}^\infty, (R_i)_{i=0}^\infty} \in C_{lh}(X) \subseteq C_h(X)$ by Lemma 5.4. If $[b, g] \in \mathcal{K}(X, B)$, then $\|[b, g](1 - \chi_{B_S})\| \rightarrow 0$ as $S \rightarrow \infty$. However, given $S > 0$, there exists i such that $6R_{i-1} > S$. Then

$$\begin{aligned}
\|[b, g](1 - \chi_{B_S})\| & \geq \| (\chi_{B_{R_i}} - \chi_{B_{6R_{i-1}}})[b, g](\chi_{B_{R_i}} - \chi_{B_{6R_{i-1}}}) \| \\
& = \| (\chi_{B_{R_i}} - \chi_{B_{6R_{i-1}}})[b, f_{k_i}](\chi_{B_{R_i}} - \chi_{B_{6R_{i-1}}}) \| \\
& > \frac{\epsilon}{5},
\end{aligned}$$

again a contradiction. This concludes the proof. \square

6. MORE ABOUT $VL_\infty(X)$

6.1. To what extent does $VL_\infty(X)$ determine X ?

Definition 6.1. *Let X and Y be metric spaces. We say that a (not necessarily continuous) function $\phi : X \rightarrow Y$ is:*

- bornologous, if for every $R \geq 0$ there exists $S \geq 0$, such that for all $x, y \in X$, $d(x, y) \leq R$ implies $d(\phi(x), \phi(y)) \leq S$;
- cobounded, if $f^{-1}(y)$ is bounded for every $y \in Y$;
- coarse, if it is both cobounded and bornologous;
- a coarse equivalence, if it is bornologous, and there exists a bornologous $\psi : Y \rightarrow X$, such that both $\psi \circ \phi$ and $\phi \circ \psi$ are uniformly close to the identity maps (i.e. $\sup_{x \in X} d(\psi \circ \phi(x), x) < \infty$ and $\sup_{y \in Y} d(\phi \circ \psi(y), y) < \infty$). Note that in this case both ϕ and ψ are automatically coarse;
- locally Lipschitz, if there exist $\delta > 0$ and $T \geq 0$, such that $d(x, y) \leq \delta$ implies $d(f(x), f(y)) \leq Td(x, y)$, $x, y \in X$.
- a Lip-coarse equivalence, if it is a coarse equivalence, it is locally Lipschitz, and in the definition of coarse equivalence, ψ can be chosen to be locally Lipschitz as well.

Note that traditionally, coarse geometry does not concern itself with a local behaviour. However, as our main tool in this piece are Lipschitz functions, we will insist that the maps involved are locally Lipschitz. On the other hand, in the key setting in which the metric spaces involved are uniformly discrete, this requirement is automatic, and thus can be ignored.

Proposition 6.2. *Let X be a proper metric space. $\text{VL}_\infty(X)$ is a Lip-coarse invariant for X . More precisely, if X and Y are proper metric spaces which are Lip-coarsely equivalent via a (locally Lipschitz) map $\phi : X \rightarrow Y$, then composition by ϕ induces a *-isomorphism $\text{VL}_\infty(Y) \rightarrow \text{VL}_\infty(X)$.*

Proof. Suppose that X, Y are Lip-coarsely equivalent, so that there are locally Lipschitz coarse maps $\phi : X \rightarrow Y$ and $\psi : Y \rightarrow X$ such that the graphs of $\phi \circ \psi$ and $\psi \circ \phi$ are uniformly bounded. Denote by $\delta > 0$ and $T \geq 0$ the constants of local Lipschitzness of ϕ . As locally Lipschitz maps are continuous, ϕ and ψ induce maps $\phi_* : l^\infty(\mathbb{N}, C_b(Y)) \rightarrow l^\infty(\mathbb{N}, C_b(X))$ and $\psi_* : l^\infty(\mathbb{N}, C_b(X)) \rightarrow l^\infty(\mathbb{N}, C_b(Y))$.

First we show that $\phi_*(\text{VL}(Y)) \subseteq \text{VL}(X)$. Surely, let $(f_n)_{n=1}^\infty \in \text{VL}(Y)$ with $\|(f_n)_{n=1}^\infty\| \leq 1$. Let $L > 0$. Since ϕ is a coarse map, there exists $S > 0$ such that if $x, y \in X$ satisfy $d(x, y) \leq 2/L$ then $d(\phi(x), \phi(y)) < S$. Without loss of generality, we can assume that $S > \delta T$. Since $(f_n)_{n=1}^\infty \in \text{VL}(Y)$, there exists n_0 such that f_n is $(L\delta/S)$ -Lipschitz for all $n \geq n_0$. For $n \geq n_0$, let us show that $f_n \circ \phi$ is L -Lipschitz. Let $x, y \in X$, there are three cases:

- If $d(x, y) > 2/L$ then $|f_n(x) - f_n(y)| \leq 2\|f_n\| < Ld(x, y)$.
- If $\delta \leq d(x, y) \leq 2/L$, then $d(\phi(x), \phi(y)) < S$. Since f_n is $(L\delta/S)$ -Lipschitz,

$$\|f_n \circ \phi(x) - f_n \circ \phi(y)\| \leq L\delta \leq Ld(x, y).$$

- If $d(x, y) < \delta$, then

$$\|f_n(\phi(x)) - f_n(\phi(y))\| \leq \frac{L\delta}{S}d(\phi(x), \phi(y)) \leq \frac{L\delta T}{S}d(x, y) < Ld(x, y).$$

Thus, ϕ_* induces a map $\theta_\phi : \text{VL}_\infty(Y) \rightarrow \text{VL}_\infty(X)$; likewise, ψ_* induces a map $\theta_\psi : \text{VL}_\infty(X) \rightarrow \text{VL}_\infty(Y)$. Let us show that these maps are inverses. By the symmetry of their definition, it suffices to show that $\theta_\psi \circ \theta_\phi = \text{id}_{\text{VL}_\infty(Y)}$.

Let $(f_n)_{n=1}^\infty \in \text{VL}_\infty(Y)$. Let $\Gamma(\phi \circ \psi) \subset Y \times Y$ denote the graph of $\phi \circ \psi$. Since this is uniformly bounded, given $\epsilon > 0$, we may find n_0 such that, for $n \geq n_0$ and $(x, y) \in \Gamma(\phi \circ \psi)$,

$$|f_n(x) - f_n(y)| \leq \epsilon.$$

In other words, for $n \geq n_0$, $|f_n(x) - f_n(\phi(\psi(x)))| \leq \epsilon$, and thus,

$$\|f_n - \psi_* \circ \phi_*(f_n)\| \leq \epsilon.$$

Consequently, $\|(f_n)_{n=1}^\infty - \theta_\psi \circ \theta_\phi((f_n)_{n=1}^\infty)\|_{\text{VL}_\infty(Y)} \leq \epsilon$. Since ϵ is arbitrary, $\theta_\psi \circ \theta_\phi((f_n)_{n=1}^\infty) = (f_n)_{n=1}^\infty$ in $\text{VL}_\infty(Y)$. \square

Lemma 6.3. *Let X be a metric space, and let $E \subseteq X \times X$. The following are equivalent:*

- (i) E is uniformly bounded.
- (ii) For every $(f_n)_{n=1}^\infty \in \text{VL}(X)$ and $\epsilon > 0$, there exists n_0 such that, if $n \geq n_0$ and $(x, y) \in E$ then $|f_n(x) - f_n(y)| < \epsilon$.
- (iii) For every $(f_n)_{n=1}^\infty \in \text{VL}(X)$, there exists n_1 such that, if $(x, y) \in E$ then $|f_{n_1}(x) - f_{n_1}(y)| < 1$.

Proof. (i) \Rightarrow (ii) is by the definition of $\text{VL}(X)$. (ii) \Rightarrow (iii) is trivial.

For (iii) \Rightarrow (i), let $E \subseteq X \times X$ be a set that isn't uniformly bounded, and let us show that there exists $(f_n)_{n=1}^\infty \in \text{VL}(X)$ such that, for all n there exists $(x_n, y_n)_{n=1}^\infty \in E$ such that $|f_n(x) - f_n(y)| = 1$.

For each n , there exists $(x_n, y_n) \in E$ such that $d(x, y) > n$. Thus there exists a $(1/n)$ -Lipschitz function $f_n : X \rightarrow \mathbb{R}$ such that $f_n(x_n) = 0$, and $f_n(y_n) = 1$. It follows that $(f_n)_{n=1}^\infty \in \text{VL}(X)$, showing that (iii) doesn't hold. \square

Proposition 6.4. *Let X, Y be metric spaces. Let $\phi : Y \rightarrow X$ be a function.*

- (i) *If $\phi_*(\text{VL}(X)) \subseteq \text{VL}(Y)$ then ϕ is a bornologous map;*
- (ii) *If, moreover, the induced map $\theta_\phi : \text{VL}_\infty(X) \rightarrow \text{VL}_\infty(Y)$ is an isomorphism, then ϕ is a coarse equivalence.*

Proof. (i): Let $R > 0$. We must show that $E := \{(\phi(x), \phi(y)) : x, y \in X, d(x, y) < R\}$ is uniformly bounded. To this end, we will verify Lemma 6.3 (iii) for this set. Therefore, let $(f_n)_{n=1}^\infty \in \text{VL}(X)$.

Since $\phi_*((f_n)_{n=1}^\infty) \in \text{VL}(Y)$, there exists n_1 such that $f_{n_1} \circ \phi$ is R^{-1} -Lipschitz. Thus for $(\phi(x), \phi(y)) \in E$, i.e., $d(x, y) < R$,

$$|f_{n_1}(\phi(x)) - f_{n_1}(\phi(y))| = |f_{n_1} \circ \phi(x) - f_{n_1} \circ \phi(y)| < 1,$$

as required.

(ii): We must show two things: (a) for every $R > 0$, the set $\{(x, y) \in Y \times Y : d(\phi(x), \phi(y)) < R\}$ is uniformly bounded, and (b) there exists $R > 0$ such that, for all $x \in X$, there exists $y \in Y$ such that $d(x, \phi(y)) < R$.

(a): Let $R \geq 0$ be given. We will verify Lemma 6.3 (iii) for $E := \{(x, y) \in X \times X : d(\phi(x), \phi(y)) < R\}$. Therefore, let $(f_n)_{n=1}^\infty \in \text{VL}(X)$.

Since θ_ϕ is surjective, there exists $(g_n)_{n=1}^\infty \in \text{VL}(Y)$ such that $\lim_\infty \|f_n - g_n \circ \phi\| = 0$. By Lemma 6.3 (i) \Rightarrow (ii), there exists n_0 such that, if $n \geq n_0$ then g_n is $(3R)^{-1}$ -Lipschitz. Pick $n \geq n_0$ such that $\|f_n - g_n \circ \phi\| < \frac{1}{3}$.

Now, let $(x, y) \in E$, i.e., $d(\phi(x), \phi(y)) < R$. Then

$$f_n(x) \approx_{1/3} g_n(\phi(x)) \approx_{1/3} g_n(\phi(y)) \approx_{1/3} f_n(y),$$

as required.

(b): Proof by contradiction. Suppose for a contradiction that, for every n there exists $x_n \in X$ such that for all $y \in Y$,

$$(x_n, \phi(y)) \geq n.$$

Thus, there exists a $(1/n)$ -Lipschitz function $f_n : X \rightarrow [0, 1]$ such that $f_n(x_n) = 1$ and $f_n(\phi(y)) = 0$ for all $y \in Y$. Putting these together, we obtain $(f_n)_{n=1}^\infty \in \text{VL}(X)$ and $\|f_n\| = 1$ (since $f_n(x_n) = 1$), so that $\|(f_n)_{n=1}^\infty\| = 1$ in $\text{VL}_\infty(X)$. However, since $f_n(\phi(y)) = 0$ for all $y \in Y$, it follows that $\theta_\phi((f_n)_{n=1}^\infty) = 0$, which contradicts injectivity of θ_ϕ . \square

In other words, when $\text{VL}_\infty(X) \cong \text{VL}_\infty(Y)$, and the isomorphism comes from a map between Y and X , it follows that X and Y are coarsely equivalent. Here is the more interesting question:

Question 6.5. *Let X, Y be uniformly discrete metric spaces. If $\text{VL}_\infty(X) \cong \text{VL}_\infty(Y)$, must X and Y be coarsely equivalent?*

6.2. The nuclear dimension of $\text{VL}_\infty(X)$. $\text{VL}(X)$ and $\text{VL}_\infty(X)$ are commutative unital C^* -algebras, and therefore by Gelfand's Theorem, each are algebras of continuous functions on a compact Hausdorff space, namely the Gelfand spectrum of the respective algebras. As these C^* -algebras are nonseparable, their spectra are nonmetrizable. Here we show a relationship between the asymptotic dimension of X and the covering dimension (suitably interpreted) of these spectra. In fact we use the nuclear dimension of the algebras, which (for the spectra) corresponds to a version of covering dimension which is slightly modified (in this nonseparable case) from the original definition. The modification

is as follows. Let Y be a locally compact Hausdorff space; call an open set $U \subseteq Y$ a *preimage-open* set if it is the preimage of an open subset of \mathbb{R} under a continuous function $Y \rightarrow \mathbb{R}$. Then the nuclear dimension of $C_0(Y)$ is equal to the smallest number d such that every finite cover of Y consisting of preimage-open sets has a $(d+1)$ -colourable refinement consisting of preimage-open sets (see the proof of [16, Proposition 3.3], and [28, Proposition 2.4]; this fact is alluded to in the discussion before [28, Proposition 2.4]). (In the second countable situation, or more generally when Y is a normal space, all open sets are preimage-open, which is why nuclear dimension coincides with the usual definition of covering dimension in this case.)

Definition 6.6. ([10, §1.E]) *Let X be a metric space. Then the asymptotic dimension of X is at most $d \in \mathbb{N}$, written $\text{asdim}(X) \leq d$, if for every $R > 0$, there exists a cover of X of the form $(U_j^{(i)})_{i=0,\dots,d; j \in J}$, such that for each $i = 0, \dots, d$, the family $(U_j^{(i)})_{j \in J}$ is R -disjoint and uniformly bounded.*

Proposition 6.7. *Let X be a metric space. $\dim_{\text{nuc}} \text{VL}(X) \leq \text{asdim}(X)$ and $\dim_{\text{nuc}} \text{VL}_\infty(X) \leq \text{asdim}(X)$.*

Proof. As the nuclear dimension decreases when passing to quotients, it suffices to prove the first statement.

Set $d := \text{asdim}(X)$. Let $\mathcal{F} \subset \text{VL}(X)$ be a finite set and let $\epsilon > 0$ be given.

Using the definition of asymptotic dimension in a fairly straightforward way, for each $n \in \mathbb{N}$, we may find an infinite partition of unity $(e_j^{(i)}(n))_{j \in J(n); i=0,\dots,d}$, such that:

- (i) for each i , $(e_j^{(i)}(n))_{j \in J(n)}$ are pairwise orthogonal,
- (ii) each $e_j^{(i)}$ is $(1/n)$ -Lipschitz, and
- (iii) there is a uniform bound, $S(n)$, on the diameters of the supports of $e_j^{(i)}(n)$ (allowed to depend only on n).

Let us also pick a point $x_j^{(i)}(n)$ inside the support of $e_j^{(i)}(n)$, for each i, j , and n . For $n = 0$, define $J(0) := X$, $e_j^{(0)}(0)(x) := \delta_{j,x}$, $e_j^{(i)}(0) := 0$ for $i > 0$, $S(0) := 0$, and $x_j^{(i)}(0) = j \in X$.

Define a $*$ -homomorphism $\psi_n = (\psi_n^{(0)}, \dots, \psi_n^{(d)}) : C_b(X) \rightarrow \bigoplus_{i=0}^d l^\infty(J(n))$ coordinatewise by evaluation at $x_j^{(i)}(n)$. For $i = 0, \dots, n$, define a $*$ -homomorphism $\phi_n^{(i)} : l^\infty(J(n)) \rightarrow C_b(X)$ by

$$\phi_n^{(i)}(\lambda) = \sum_{j \in J(n)} \lambda(j) e_j^{(i)}(n),$$

with the sum converging pointwise, since in fact at each point $x \in X$, at most one summand is nonzero (by condition (i)). Note that if $f \in$

$C_b(X)$ is $(\epsilon/S(n))$ -Lipschitz then

$$(6.1) \quad \sum_{i=0}^d \phi_n^{(i)} \circ \psi_n^{(i)}(f) \approx_\epsilon f.$$

Let

$$\mathcal{F} = \{(f_{i,k})_{k=1}^\infty : i = 1, \dots, m\}.$$

For each k , we may find some $n_k \geq 0$ such that $f_{i,k}$ has $(\epsilon/S(n_k))$ -Lipschitz for all $i = 1, \dots, m$. Since $\mathcal{F} \subseteq \text{VL}(X)$, we can pick these n_k such that they converge to ∞ .

We now define a $*$ -homomorphism

$$\Psi := (\psi_{n_k})_{k=1}^\infty : l^\infty(\mathbb{N}, l^\infty(X)) \rightarrow \prod_k \bigoplus_{i=0}^d l^\infty(J(n_k)) \cong \bigoplus_{i=0}^d \prod_k l^\infty(J(n_k));$$

we may write $\Psi = (\Psi^{(0)}, \dots, \Psi^{(d)})$. For $i = 0, \dots, d$, define a $*$ -homomorphism

$$\Phi^{(i)} := (\phi_{n_k}^{(i)})_{k=1}^\infty : \prod_k l^\infty(J(n_k)) \rightarrow l^\infty(\mathbb{N}, l^\infty(X)).$$

Since $n_k \rightarrow \infty$, we see that the image of $\Phi^{(i)}$ is in fact contained in $\text{VL}(X)$. Moreover, by (6.1) and our choice of n_k , we find that

$$\sum_{i=0}^d \Phi^{(i)} \circ \Psi^{(i)}(f) \approx_\epsilon f$$

for $f \in \mathcal{F}$. Since $\prod_k l^\infty(J(n_k))$ has nuclear dimension zero, this is sufficient to prove that $\text{VL}(X)$ has nuclear dimension at most d . \square

We have an argument to get inequalities in the other direction, under the hypothesis that X has finite asymptotic dimension. For this, we begin with the following lemma.

Lemma 6.8. *Let X be a set and let $\eta > 0$. Let $f_1, \dots, f_m, e_1, \dots, e_n : X \rightarrow [0, \infty)$, and $\lambda_{i,j} \in [0, \infty)$ for $i = 1, \dots, n$, $j = 1, \dots, m$. Suppose that $f_j \approx_\eta \sum_{i=1}^n \lambda_{i,j} e_i$ for each j and $\sum_{j=1}^m \lambda_{i,j} = 1$ for each i . Then for each i , there exists $j(i)$ such that*

$$\{x : e_i(x) > m\eta\} \subseteq \{x : f_{j(i)}(x) > 0\}.$$

Proof. Fix i . Since $\sum_{j=1}^m \lambda_{i,j} = 1$, there exists some $j = j(i)$ such that $\lambda_{i,j} \geq 1/m$. For $x \in X$ such that $f_j(x) = 0$, it follows that

$$\begin{aligned} \frac{1}{m} e_i(x) &\leq \lambda_{i,j} e_i(x) \\ &\leq \sum_{i'=1}^n \lambda_{i',j} e_{i'}(x) \\ &\leq f_j(x) + \eta = \eta. \end{aligned}$$

This shows that $e_i(x) \leq m\eta$, as required. \square

Theorem 6.9. *If X has finite asymptotic dimension, then*

$$\text{asdim}(X) = \dim_{\text{nuc}} \text{VL}(X) = \dim_{\text{nuc}} \text{VL}_\infty(X).$$

Proof. Set $d := \dim_{\text{nuc}} \text{VL}_\infty(X)$. By Proposition 6.7, it suffices to show that $\text{asdim}(X) \leq d$. Let $R > 0$ be given, and we will partition X into $(d + 1)$ uniformly bounded, R -disjoint families.

By hypothesis, let $\text{asdim}(X) \leq m - 1$. Then from this, there exists a partition of unity $g_1, \dots, g_m \in \text{VL}_\infty(X)$, such that $g_j = (g_{j,l})_{l=1}^\infty$ where for each j, l , the support of $g_{j,l}$ decomposes as an l -disjoint, uniformly bounded family of subsets of X .

Set

$$\eta := \frac{1}{3(d+1)m}.$$

The only nonzero order zero maps from a matrix algebra into a commutative algebra occur when the matrix algebra is one-dimensional (this follows from [27, Proposition 3.2 (a)]). Hence, $\dim_{\text{nuc}} \text{VL}(X) \leq d$ implies that there exists $s \in \mathbb{N}$, a c.p.c. map $\psi = (\psi^{(0)}, \dots, \psi^{(d)}) : \text{VL}_\infty(X) \rightarrow \bigoplus_{i=0}^d \mathbb{C}^{\oplus s}$, and c.p.c. order zero maps $\phi^{(i)} : \mathbb{C}^s \rightarrow \text{VL}_\infty(X)$ such that

$$g_j \approx_\eta \sum_{i=0}^m \phi^{(i)} \circ \psi^{(i)}(g_j) \quad \text{and} \quad 1_{\text{VL}_\infty(X)} \approx_{1/2} \sum_{i=0}^m \phi^{(i)} \circ \psi^{(i)}(1).$$

By rescaling, we may assume that $\psi^{(i)}(1_{\text{VL}_\infty(X)}) = (1, \dots, 1)$. Write $\psi^{(i)}(g_j) = (\lambda_{(i,1),j}, \dots, \lambda_{(i,s),j}) \in [0, \infty)^s$ (since ψ is positive). By linearity, looking at the i' component of $\psi^{(i)}(1)$, we have

$$\sum_{j=1}^m \lambda_{(i,i'),j} = 1.$$

Each map $\phi^{(i)}$ lifts to a c.p.c. order zero map $(\phi_l^{(i)})_{l=1}^\infty : \mathbb{C}^s \rightarrow \text{VL}(X)$, by [16, Remark 2.4]. For all but finitely many l , we have

$$g_{j,l} \approx_\eta \sum_{i=0}^d \phi_l^{(i)}(\lambda_{(i,1),j}, \dots, \lambda_{(i,s),j}) \quad \text{and} \quad 1_{C_b(X)} \approx_{1/2} \sum_{i=0}^d \phi_l^{(i)}(1, \dots, 1). \quad (6.2)$$

Fix $l \geq R$ for which (6.2) holds, and such that the image of each minimal projection in \mathbb{C}^s under $\phi_l^{(i)}$ is $(m\eta/L)$ -Lipschitz. Write $e_{(i,1)}, \dots, e_{(i,s)}$ for these images under $\phi_l^{(i)}$ of the minimal projections in \mathbb{C}^s , and write $f_j := g_{j,l}$; thus (6.2) becomes

$$f_j \approx_\eta \sum_{i=0}^d \sum_{i'=1}^s \lambda_{(i,i'),j} e_{(i,i')} \quad \text{and} \quad 1 \approx_{1/2} \sum_{i=0}^d \sum_{i'=1}^s e_{(i,i')}. \quad (6.3)$$

We now apply Lemma 6.8 with (i, i') , $i = 0, \dots, d$, $i' = 1, \dots, s$ in place of the index $i = 1, \dots, n$. This tells us that for each $i = 0, \dots, d$ and $i' = 1, \dots, s$, there exists some $j(i, i')$ such that

$$B_{i,i'} := \{x \in X : e_{(i,i')} > m\eta\} \subseteq \{x \in X : f_{j(i,i')}(x) > 0\}.$$

Since the support of $f_{j(i,i')}$ ($= g_{j(i,i'),l}$) decomposes as a union of an l -disjoint uniformly bounded family, and $l \geq R$, we can partition $B_{i,i'}$ into an R -disjoint, uniformly bounded family, say

$$B_{i,i'} = \prod_{t \in T} A_{i',t}^{(i)}$$

We now consider the family $(A_{i',t}^{(i)})_{i'=1,\dots,s, t \in T}$. This family is a finite union of uniformly bounded families, whence it is uniformly bounded. Let us check that it is R -disjoint. Since for fixed i' we already have R -disjointness of $(A_{i',t}^{(i)})_{t \in T}$, we need to show that for $i'_1 \neq i'_2$ and $t_1, t_2 \in T$, the minimal distance between $A_{i'_1,t_1}^{(i)}$ and $A_{i'_2,t_2}^{(i)}$ is at least R . In other words, we need to show that the minimal distance between B_{i,i'_1} and B_{i,i'_2} is at least R .

We have that e_{i,i'_1} and e_{i,i'_2} are orthogonal, so if $x \in B_{i,i'_1}$ then since $e_{i,i'_1}(x) \neq 0$, it must be the case that $e_{i,i'_2}(x) = 0$. Consider now $y \in B_{i,i'_2}$, so that $e_{i,i'_2}(y) > m\eta$. Since e_{i,i'_2} is $(m\eta/R)$ -Lipschitz, it follows that $d(x, y) \geq R$, as required.

Finally let us show that

$$(A_{i',t}^{(i)})_{i=0,\dots,d, i'=1,\dots,s, t \in T}$$

is a cover of X , i.e., that $X = \bigcup_{i,i'} B_{i,i'}$. For $x \in X$, from the second part of (6.3), we have

$$1/2 < \sum_{i=0}^d \sum_{i'=1}^s e_{(i,i')}(x).$$

At most $d+1$ terms in this sum are nonzero, due to the pairwise orthogonality withing each family $(e_{(i,i')})_{i'=1}^s$. Therefore, there exists some i, i' such that $e_{(i,i')}(x) > \frac{1}{2(d+1)} = m\eta$. Thus, $x \in B_{i,i'}$ (by definition) as required. \square

We have the following consequence.

Corollary 6.10. *Suppose X is a metric space, and for all $m \in \mathbb{N}$, X contains a subspace Y_m such that $\text{asdim}(Y_m) \in [m, \infty)$. Then*

$$\dim_{\text{nuc}} \text{VL}(X) = \dim_{\text{nuc}} \text{VL}_{\infty}(X) = \infty.$$

Proof. It is not too hard to see that restriction to Y_m produces a surjective *-homomorphism $\text{VL}_{\infty}(X) \rightarrow \text{VL}_{\infty}(Y_m)$ (for surjectivity, the key

point is that an L -Lipschitz function on a closed subset of Y_m extends to an L -Lipschitz function on X). Hence we have

$$\dim_{\text{nuc}} \text{VL}(X) \geq \dim_{\text{nuc}} \text{VL}_{\infty}(Y_m) \geq m,$$

using [28, Proposition 2.3(iv)] for the first inequality and Theorem 6.9 for the second. \square

In [2, Theorem 7.2], it was shown that the asymptotic dimension of X is equal to the covering dimension of the Higson corona νX , *likewise provided that* $\text{asdim}(X) < \infty$.

Question 6.11. *Is $\dim_{\text{nuc}} \text{VL}(X) = \text{asdim}(X)$ always? Is $\dim_{\text{nuc}} \text{VL}(X) = \dim(\nu X)$ always?*

REFERENCES

- [1] G. Bell and A. Dranishnikov. Asymptotic dimension. *Topology Appl.*, 155(12):1265–1296, 2008.
- [2] A. N. Dranishnikov. Asymptotic topology. *Uspekhi Mat. Nauk*, 55(6(336)):71–116, 2000.
- [3] Alexander Dranishnikov. On Gromov’s positive scalar curvature conjecture for duality groups. *J. Topol. Anal.*, 6(3):397–419, 2014.
- [4] Alexander Dranishnikov and Michael Zarichnyi. Asymptotic dimension, decomposition complexity, and Haver’s property C. *Topology Appl.*, 169:99–107, 2014.
- [5] Alexander Engel. Index theory of uniform pseudodifferential operators. arXiv preprint 1502.00494v1.
- [6] Alexander Engel. Rough index theory on spaces of polynomial growth and contractibility. arXiv preprint 1505.03988v3.
- [7] Vladimir Georgescu. On the structure of the essential spectrum of elliptic operators on metric spaces. *J. Funct. Anal.*, 260(6):1734–1765, 2011.
- [8] Vladimir Georgescu. On the essential spectrum of the operators in certain crossed products, 2017. arXiv preprint 1705.00379.
- [9] Vladimir Georgescu and Andrei Iftimovici. Localizations at infinity and essential spectrum of quantum Hamiltonians. I. General theory. *Rev. Math. Phys.*, 18(4):417–483, 2006.
- [10] M. Gromov. Asymptotic invariants of infinite groups. In *Geometric group theory, Vol. 2 (Sussex, 1991)*, volume 182 of *London Math. Soc. Lecture Note Ser.*, pages 1–295. Cambridge Univ. Press, Cambridge, 1993.
- [11] Mikhail Gromov. Groups of polynomial growth and expanding maps. *Inst. Hautes Études Sci. Publ. Math.*, (53):53–73, 1981.
- [12] Erik Guentner and Jerome Kaminker. Exactness and the Novikov conjecture. *Topology*, 41(2):411–418, 2002.
- [13] Erik Guentner, Romain Tessera, and Guoliang Yu. A notion of geometric complexity and its application to topological rigidity. *Invent. Math.*, 189(2):315–357, 2012.
- [14] Erik Guentner, Romain Tessera, and Guoliang Yu. Discrete groups with finite decomposition complexity. *Groups Geom. Dyn.*, 7(2):377–402, 2013.
- [15] Gennadi Kasparov and Guoliang Yu. The coarse geometric Novikov conjecture and uniform convexity. *Adv. Math.*, 206(1):1–56, 2006.
- [16] Eberhard Kirchberg and Wilhelm Winter. Covering dimension and quasidiagonality. *Internat. J. Math.*, 15(1):63–85, 2004.

- [17] B. V. Lange and V. S. Rabinovich. Noethericity of multidimensional discrete convolution operators. *Mat. Zametki*, 37(3):407–421, 462, 1985.
- [18] Narutaka Ozawa. Amenable actions and exactness for discrete groups. *C. R. Acad. Sci. Paris Sér. I Math.*, 330(8):691–695, 2000.
- [19] V. S. Rabinovich, S. Roch, and B. Silbermann. Fredholm theory and finite section method for band-dominated operators. *Integral Equations Operator Theory*, 30(4):452–495, 1998. Dedicated to the memory of Mark Grigorievich Krein (1907–1989).
- [20] John Roe. Personal communication.
- [21] John Roe. An index theorem on open manifolds. I. *J. Differential Geom.*, 27(1):87–113, 1988.
- [22] John Roe. *Index theory, coarse geometry, and topology of manifolds*, volume 90 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1996.
- [23] John Roe. *Lectures on coarse geometry*, volume 31 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2003.
- [24] Thomas Schick. The topology of positive scalar curvature. In *Proceedings of the International Congress of Mathematicians Seoul 2014*, volume II, pages 1285–1308. Kyung Moon SA Co. Ltd., 2014.
- [25] G. Skandalis, J. L. Tu, and G. Yu. The coarse Baum-Connes conjecture and groupoids. *Topology*, 41(4):807–834, 2002.
- [26] Ján Špakula and Rufus Willett. On rigidity of Roe algebras. *Adv. Math.*, 249:289–310, 2013.
- [27] Wilhelm Winter. Covering dimension for nuclear C^* -algebras. *J. Funct. Anal.*, 199(2):535–556, 2003.
- [28] Wilhelm Winter and Joachim Zacharias. The nuclear dimension of C^* -algebras. *Adv. Math.*, 224(2):461–498, 2010.
- [29] Nick Wright. C_0 coarse geometry and scalar curvature. *J. Funct. Anal.*, 197(2):469–488, 2003.
- [30] Guoliang Yu. Zero-in-the-spectrum conjecture, positive scalar curvature and asymptotic dimension. *Invent. Math.*, 127(1):99–126, 1997.
- [31] Guoliang Yu. The Novikov conjecture for groups with finite asymptotic dimension. *Ann. of Math. (2)*, 147(2):325–355, 1998.
- [32] Guoliang Yu. The coarse Baum-Connes conjecture for spaces which admit a uniform embedding into Hilbert space. *Invent. Math.*, 139(1):201–240, 2000.