

Received Date : 25-Apr-2018

Accepted Date : 30-Apr-2018

Article type : Review Article

Biological Hydropersulfides and Related Polysulfides: A New Concept and Perspective in Redox Biology

Jon M. Fukuto^{1*}, Louis J. Ignarro^{2*}, Peter Nagy³, David A. Wink⁴, Christopher G. Kevil⁵, Martin Feelisch⁶, Miriam M. Cortese-Krott⁷, Christopher L. Bianco⁷, Yoshito Kumagai⁸, Adrian J. Hobbs⁹, Joseph Lin¹⁰, Tomoaki Ida¹¹, and Takaaki Akaike^{11*}

¹Department of Chemistry, Sonoma State University, Rohnert Park, CA 94928, ²Department of Molecular and Medical Pharmacology, UCLA School of Medicine, Center for the Health Sciences, Los Angeles, CA 90095, ³Department of Molecular Immunology and Toxicology, National Institute of Oncology, Budapest, Hungary, ⁴Tumor Biology Section, Radiation Biology Branch, National Cancer Institute, Bethesda, Maryland 20892, ⁵Department of Pathology, Louisiana State University Health Sciences Center, Shreveport, LA, ⁶NIHR Southampton Biomedical Research Center, University Hospital Southampton NHS Foundation Trust, Southampton, United Kingdom, ⁷Cardiovascular Research Laboratory, Department of Cardiology, Pneumology and Angiology, Medical Faculty, Heinrich Heine University, Dusseldorf, Germany, ⁸Environmental Biology Section, Faculty of Medicine, University of Tsukuba, Tsukuba, Ibaraki 305-8575, Japan, ⁹William Harvey Research Institute, Bart & London School of Medicine, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, UK, ¹⁰Department of Biology, Sonoma State University, Rohnert Park, CA 94928, ¹¹Department of Environmental Medicine and Molecular Toxicology, Tohoku University Graduate School of Medicine, Sendai 980-8575, Japan.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/1873-3468.13090

This article is protected by copyright. All rights reserved.

*To whom correspondence can be addressed: JMF, fukuto@sonoma.edu; LJI: Lignarro@gmail.com; TA, takaike@med.tohoku.ac.jp.

Keywords: hydropersulfides, cysteine persulfides, translational polysulfidation

Abstract

The chemical biology of thiols (RSH, e.g., cysteine and cysteine containing proteins/peptides) has been a topic of extreme interest for many decades due to their reported roles in protein structure/folding, redox signaling, metal ligation, cellular protection and enzymology. While many of the studies on thiol/sulfur biochemistry have focused on thiols, relatively ignored have been hydropersulfides (RSSH) and higher order polysulfur species (RSS_nH , RSS_nR , $n > 1$). Recent and provocative work has alluded to the prevalence and likely physiological importance of RSSH and related RSS_nH . RSSH of cysteine (Cys-SSH) has been found to be prevalent in mammalian systems along with Cys-SSH-containing proteins. The RSSH functionality has not been examined to the extent of other biologically relevant sulfur derivatives (e.g., sulfenic acids, disulfides, etc.), whose roles in cell signaling are strongly indicated. The recent finding of Cys-SSH biosynthesis and translational incorporation into proteins is an unequivocal indication of its fundamental importance and necessitates a more profound look into the physiology of RSSH. In this Review, we discuss the currently reported chemical biology of RSSH (and related species) as a prelude to discussing their possible physiological roles.

Introduction

The chemical biology of thiols (RSH, e.g. cysteine and cysteine containing proteins/peptides) has been a topic of extreme interest for many decades due to their reported roles in protein structure/folding, redox signaling, metal ligation, cellular protection and enzymology [1-5]. That is, oxidation of thiols to covalently linked disulfides (RSSR) is a critical aspect of protein folding and structure, modification (typically oxidative) of critical thiols can result in alteration of protein function/activity, thiol sulfur atoms are used in the binding of metals in numerous metalloproteins, thiol peptides scavenge otherwise toxic electrophiles, and thiolates (RS^-) serve as potent nucleophiles important in the catalytic activity of numerous

enzymes. Moreover, oxidized thiol species such as thiyl radicals ($RS\cdot$), sulfenic acids ($RSOH$) and sulfinic acids ($RS(O)OH$) also possess important chemical properties that Nature has exploited for various purposes. For example, $RS\cdot$ is used as an oxidant in the biosynthesis of DNA [6] and $RSOH/RS(O)OH$ bind specific metals in metalloproteins, conferring activity [7]. Thus, the diverse chemistry of thiols and derived species are fundamental to various aspects of cell biochemistry, function and signaling.

In the recent past, most studies of thiol/sulfur biochemistry have focused primarily on RSH/RS^- , $RS\cdot$, $RSSR$, $RSOH$, nitrosothiols ($RSNO$), $RS(O)OH$ and sulfonic acids ($RS(O)_2OH$). Relatively ignored have been hydropersulfides ($RSSH$) and higher order polysulfur species (i.e. RSS_nH , RSS_nR , $n > 1$, note: the term “hydropersulfide” and the abbreviation “ $RSSH$ ” will be used to denote both the protonated and deprotonated species unless otherwise specified). Indeed, until very recently, the discussion of the chemical biology of hydropersulfides and polysulfides has been primarily relegated to studies of alliums and other food sources rich in these species [8] and to scant and specific examples of proteins with presumed hydropersulfides (of unknown function) [9-11]. Also, hydropersulfides have been implicated in the biochemistry of several specific enzymes such as mercaptopyruvate sulfur transferase (MST) [12], $NIFS$ (a bacterial enzyme involved in metallocluster biosynthesis [13]) and rhodanese [14], among others [15], some of which with no established physiological function. In spite of these examples, the idea that hydropersulfides are fundamental signaling/biochemical intermediates with important physiological functions (akin to previously mentioned sulfur species such as sulfenic acids, disulfides or S-nitrosothiols) has not been generally considered. To be sure, important papers by Toohey, Wood and others, some of them published more than 25 years ago [16-18], indicated the possible general importance of hydropersulfides in biological systems. In spite of these somewhat prophetic reports, the idea that hydropersulfides (and related/derived polysulfides) are important and general biological mediators/effectors has remained unappreciated. However, there has been a resurgence of this idea due to recent and substantive reports of their biological prevalence and likely importance. In particular, a new and watershed report demonstrates translational cysteine hydropersulfide incorporation into proteins (*vide infra*), consistent with the idea of fundamentally important physiological roles for hydropersulfides. This finding, along with other recent work on the unique chemical and biological properties of hydropersulfides and related polysulfides, serve as the impetus for providing a perspective in redox biology.

Accepted Article

It is the tenet of this article that hydropersulfides and related polysulfides represent a greatly underappreciated and fundamentally important aspect of sulfur-based biochemistry/signaling that is involved in a diverse array of physiological and/or biochemical functions. Moreover, future elucidation of these effects and functions will lead to a more profound and complete understanding of the general utility of sulfur chemistry in biological/mammalian systems. Herein are discussed some of the recent reports indicating that hydropersulfides and polysulfides can play a role in mammalian biology, as well as the possible future directions of this field.

The Generation and Prevalence of Hydropersulfides and Polysulfides in

Eukaryotes: The past and apparent lack of general interest in the biology of per- and polysulfides likely stemmed from the fact that their natural existence was only indirectly alluded to (based on non-specific assays) and limited to only a select number of enzymes/proteins such as xanthine oxidase [10] and aldehyde oxidase [11]. Until recently, the ubiquitous and highly prevalent nature of polysulfur species in mammalian systems was unknown. Recent and significant interest in the biological actions and signaling of hydrogen sulfide (H_2S) [19] spurred interest in H_2S -related and/or derived species as possible biological mediators/effectors. One of the most important early studies regarding H_2S pharmacology and physiology was by Blackstone and coworkers [20] who reported that pharmacological administration of H_2S induced a reversible hibernative state in mice. This study revealed that low levels of H_2S can have dramatic physiological effects and is not just a toxic species (which is observed at higher levels). An important aspect of the presence of H_2S in biological systems is the possibility that RSSH and higher order polysulfides can be formed [21,22].

One of the earliest and seminal studies drawing attention to the possible prevalence of polysulfur species (specifically hydropersulfides) in proteins was by Mustafa and coworkers [23] who utilized a modified biotin-switch assay to detect protein hydropersulfides and suggested that 10-25% of numerous liver proteins contained protein hydropersulfides (this assay, as conducted, may not have been specific for hydropersulfides [2,24]). These investigators also reported that protein “sulphydration”¹ led to enhanced activity for glyceraldehyde 3-phosphate dehydrogenase (although brought into question [25,26]) and promoted actin polymerization. Subsequent reports generally support the idea of the ubiquitous and highly prevalent nature of hydropersulfides and polysulfides in biological (i.e. eukaryotic) systems. For example, Ida and coworkers [27] utilized electrophilic trapping

Accepted Article

followed by LC-MS analysis to find that small molecule hydroperulfides such as cysteine hydroperulfide (Cys-SSH) and glutathione hydroperulfide (GSSH) as well as higher order (additional sulfur atoms, e.g. RSSSH) hydroperulfides were present in mammalian cells, tissue and plasma. Indeed, levels of GSSH were found to be greater than 50 micromolar in mouse heart and liver and greater than 150 micromolar in mouse brain. Levels of Cys-SSH, although significantly lower, were found to be in the low micromolar range in mouse heart, liver and brain. Using a tag-switch-tag method [27], Ida and coworkers also detected numerous protein hydroperulfides, confirming the results of Mustafa and coworkers that the existence of protein hydroperulfide is widespread. Assuming the value of GSSH levels in mouse brain to be accurate (i.e. >150 micromolar), GSSH may be the most prevalent cysteine-containing sulfur species behind only glutathione (GSH) itself (which is typically between 1 and 10 mM in cells) in certain tissues.

The Nagy lab also developed a hydroperulfide-specific assay, referred to as the protein hydroperulfide detection protocol (ProPerDP method), amenable to whole cells and tissues and confirmed the existence and prevalence of protein hydroperulfides in cells at levels in a steady-state range of approximately 1-12 micrograms/milligram protein [28]. An advantage of this method compared to others developed previously is that it utilized an alkylation step prior to cellular lysis thus precluding cell lysis-mediated artifactual oxidation/persulfidation. Further work by Longen and coworkers [29], also using a mass spectrometry-based assay (entitled quantitative PerSulfide Site Identification or qPerS-SID), found that cells treated with H₂S resulted in the generation of numerous protein hydroperulfides (with significant identification overlap with the above-mentioned work by Ida and coworkers [27]). They also found that H₂S-mediated protein hydroperulfide formation appeared to exhibit specificity among numerous potential "sulfhydration" sites. That is, among numerous thiols on a single protein, only certain thiols were subject to hydroperulfide formation. Thus, there appears to be little doubt of the biological prevalence of hydroperulfide formation in peptides and proteins. This contention is supported by the finding by the Akaike group [30] of translational incorporation of cysteine hydroperulfide into proteins. Indeed, it is this major discovery that provides much of the impetus for writing this perspective, which will hopefully draw attention to this provocative discovery and its relevance to the field of sulfur biochemistry and physiology.

Up until 2017, hydropersulfide formation in peptides and proteins was thought to result primarily from a posttranslational modification of existing cysteine residues in proteins. That is, reaction of an oxidized protein cysteine (possibly a disulfide or sulfenic acid, among others) could react with H₂S giving the corresponding hydropersulfide (Reactions 1 and 2).



These reactions have been studied extensively [21,22,31,32] and they represent possible routes of endogenous hydropersulfide formation. It needs to be mentioned, however, that formation of hydropersulfides via Reaction 1 has been questioned on the basis of the low reducing power of H₂S, its relatively low concentration and the high concentrations of GSH [33]. Regardless, since H₂S is biosynthesized via a variety of enzymes (*vide infra*) and the presence of oxidized sulfurs (i.e. disulfides, sulfenic acids) in biological systems is well established, generation of hydropersulfides via Reactions 1 or 2 is clearly possible. Indeed, hydropersulfides are likely involved in numerous equilibria with other sulfur species and, although at low steady state levels, are biologically accessible via these equilibria [34]. However, it remains to be determined whether hydropersulfide formation via these reactions (or others) is significant and serves a biological function. That is, does Nature generate hydropersulfides as a response or reaction to some physiological state, cellular stress or biochemical requirement? Do hydropersulfides serve a biological function and are, therefore, generated in a regulated manner?

These questions have recently been answered by the finding that protein hydropersulfides are incorporated during translation via the actions of cysteinyl tRNA synthetases (CARS) [30] (Figure 1). The Akaike group has provided strong evidence that CARS are capable of converting Cys-SH to Cys-SSH via a pyridoxal phosphate-dependent process using a second Cys-SH as the sulfur atom donor (independent of ATP and tRNA). The CARS-synthesized Cys-SSH can then form a tRNA-bound Cys-SSH adduct (also via CARS catalysis) resulting in the incorporation of Cys-SSH into proteins thus generating a protein containing a hydropersulfide functionality. This process is reminiscent of, although not mechanistically

analogous to, the incorporation of selenocysteine (SeCys) into selenoproteins [35,36]. Figure 2 schematically illustrates and compares the mechanisms of Cys-SSH and SeCys generation via translational machinery. A fundamental difference between the incorporation of Cys-SSH and SeCys into proteins is that SeCys is biosynthesized from a tRNA-bound serine while Cys-SSH is premade from Cys-SH by CARS prior to association with the tRNA, which is also catalyzed by CARS. Regardless, both SeCys and Cys-SSH are biosynthesized by translational “machinery” prior to incorporation into proteins. The fact that hydropersulfides are both biosynthesized and attached to tRNA by CARS and subsequently translationally incorporated into proteins indicates they have a potential biological action and are not merely an artifact of H₂S biosynthesis.

Considering the analogy between SeCys and Cys-SSH, it is worthwhile to further compare these two amino acids. SeCys is considered to be Nature’s 21st amino acid since it is biosynthesized pre-translationally (i.e., is not a post-translational modification), specifically encoded (using UGA, which is normally the stop codon) and incorporated into proteins to serve a specific function. Analogously, Cys-SSH may represent Nature’s 22nd amino acid (in eukaryotes) for most of the same reasons (pyrrolysine is considered to be the 22nd amino acid in Archaea [37]). However, the case for Cys-SSH as the 22nd amino acid in eukaryotes is not yet as strong as it is for SeCys since the nature of the codon, the regulation of the process and the mechanism(s) that allows for specific incorporation are not yet established. Interestingly, based on their chemical properties, SeCys and Cys-SSH can be considered to be functionally similar. Comparing these amino acids to Cys-SH, they are both superior nucleophiles, more potent reductants, and more acidic (i.e. exist primarily in the anionic form). In the case of SeCys, all of these properties have been reported to be important with respect to its biological functions [38]. For example, the high nucleophilicity and reducing capabilities of SeCys are critical to the activities of the redox proteins glutathione peroxidase and thioredoxin reductase. The same would be expected for the biological utility of protein hydropersulfides. However, an important distinction between SeCys and Cys-SSH is the fact that Cys-SSH can be easily converted to Cys-SH, thus generating a “normal” cysteine protein (i.e. Reaction 3 or Reactions 4 and 5), whereas SeCys is essentially a permanent functional group on selenoproteins. Thus, Cys-SSH can be viewed as a potentially fleeting SeCys-like functionality allowing proteins containing hydropersulfides to change their chemical properties depending on, for example, the cellular redox environment.



Previous to the discovery of Cys-SSH formation via CARS, the biosynthesis of Cys-SSH was known to occur primarily via the actions of two enzymes, cystathionine beta-synthase (CBS) and cystathionine gamma-lyase (CSE). These enzymes appear to be responsible for at least a portion of the Cys-SSH found in cells since, for example, CSE knock out cells were significantly lower in intracellular hydropersulfides (and polysulfides) and CSE over-expressing cells possessed high levels [27]. Previous work by Sen and coworkers using an improved maleimide-based assay also showed that CSE-delete mice lacked the ability to form hydropersulfides on the p65 subunit of NF- κ B [39], affecting activity and Mustafa et al. [23] also reported that CSE expression led to hydropersulfide formation in proteins such as GAPDH, β -tubulin and actin. The idea that CSE and/or CBS are generally responsible for the Cys-SSH found in cells has been questioned since the levels of enzymatic substrates and the reported kinetics of the enzymes would predict that H₂S is the major product and not Cys-SSH. That is, both CSE and CBS are somewhat promiscuous enzymes capable of catalyzing numerous reactions and based on the kinetic constants for these reactions and the approximate steady-state levels of all the possible enzymatic substrates and reactive species, it has been argued that Cys-SSH is merely a fleeting intermediate. It should be noted, however, that this treatment may oversimplify the system and does not, for example, account for other factors such as Cys-SS-Cys uptake and redox regulation of the enzymes. Of significant importance, however, is the recent finding by Akaike and coworkers who have also demonstrated that CARS-mediated Cys-SSH biosynthesis can be responsible for much of the intracellular Cys-SSH [30]. In fact, this group has proposed that, in some cases, biosynthesis of Cys-SSH by CARS may be responsible for the majority of the Cys-SSH found in cells and tissues *in vivo*. Regardless of how hydropersulfides are made, it is clear they are prevalent (*vide supra*) and likely to possess important biological function.

Thus far, the discussion has focused almost exclusively on RSSH species. It is also noteworthy that polysulfides ($\text{RSS}_n\text{R}'$, $n > 1$, $\text{R}' = \text{hydrogen or alkyl}$) are prevalent in biological systems. Indeed, Ida and coworkers [27] have found numerous small molecule polysulfides (e.g. glutathione trisulfide (GSSSG), cysteine trisulfide (aka thiocystine, Cys-SSS-Cys), glutathione tetrasulfide (GSSSSG), cysteine tetrasulfide (Cys-SSSS-Cys), etc.) in human and mouse cells, tissue or plasma. Moreover, the kinetic analysis of the enzymes CSE and CBS mentioned above also predicts that polysulfides are likely products. The mechanism(s) for the generation of these species in biological systems has not been determined and is likely to be due to a complex series of reactions involving, initially, hydropersulfides and possibly inorganic polysulfide species (e.g. $\text{HS}_2^-/\text{S}_2^{2-}$, $\text{HS}_3^-/\text{S}_3^{2-}$, etc.).

The Chemical Biology of Hydropersulfides and Polysulfides: The biological actions of hydropersulfides and polysulfides will undoubtedly be due to their unique and distinct chemistry. Thus, in order to speculate as to the biological functions of hydropersulfides and polysulfides, it is essential to first have a discussion regarding the chemistry of hydropersulfides and polysulfides and compare these chemical properties to other biologically relevant sulfur species. As mentioned above, hydropersulfides are superior to thiols as nucleophiles and reducing agents (Figure 3). Although not yet established, it can also be predicted that the metal ligation properties are distinct. It has been proposed that RSSH is equivalent to a chemically hyperactivated RSH which led to the idea that “anything a thiol can do, a hydropersulfide can do better” [25]. Although this can be why hydropersulfides are made in some cases (to enhance the activity of thiol proteins), it is at least safe to say that hydropersulfides possess distinct chemistry from that of simple thiols and that this may be important to the biological utility of hydropersulfides. One important distinction between thiols and hydropersulfides is the fact that thiols can be nucleophilic while hydropersulfides can be either nucleophilic or electrophilic depending on the protonation state. That is, a persulfide anion (RSS^-) is extremely nucleophilic while the protonated RSSH species is electrophilic (akin to an oxidized sulfur species like a disulfide or sulfenic acid). Thus, hydropersulfides have the properties of both thiols and disulfides. It is also noteworthy that RSSH is oxidized relative to RSH (RSSH is at the same oxidation state as RSSR) and yet RSSH/ RSS^- is a superior reductant compared to RSH/ RS^- . It may seem counter-intuitive that the oxidation of a molecule results in the generation of a superior reductant, but this is exactly the case with the conversion of a thiol to a hydropersulfide. Indeed, this situation led to the idea that RSSH generation can be a biological response to cellular oxidative stress since an oxidized species generated under this stress is actually a superior reductant capable of possibly protecting against potentially deleterious oxidation events (due to its reducing capabilities, *vide infra*).

Thiyl radicals ($RS\cdot$) are used biologically as potent oxidants capable of, for example, breaking a C-H bond via hydrogen atom abstraction [6] and thus initiating radical chemistry. It has been postulated that $RS\cdot$ species may be involved in fundamental cellular redox signaling as well [40]. The fact that $RS\cdot$ is a fairly potent one-electron oxidant is an indication of the relatively poor ability of RSH to serve as a reductant. On the other hand, perthiyl radicals ($RSS\cdot$) are relatively weak oxidants and are readily formed from one-electron oxidation of RSSH/ $RSS\cdot$ [41,42]. RSSH/ $RSS\cdot$ is a good reductant due to the stability of the oxidized $RSS\cdot$ species. The stability of $RSS\cdot$ is akin to the stabilization of analogous nitroxide systems ($R_2NO\cdot$). In both cases the stability of the unpaired electron is likely due to an overlap of the orbital containing the unpaired electron with the adjacent orbital containing a lone pair of electrons [43]. The lack of reactivity of $RSS\cdot$ is evidenced by the fact that it does not readily react with O_2 and NO, both established scavengers of other radical species [41,44]. Although it has been proposed that the one-electron redox properties of RSSH/ $RSS\cdot$ may have biological function (i.e. can serve as a “redox gate” [41]), this has yet to be shown.

Thus far, only the chemical biology of hydropersulfides has been discussed. As briefly mentioned above, polysulfides are also prevalent (albeit at lower levels than, for example, Cys-SSH or GSSH). Like disulfides, polysulfides are electrophilic and thus readily react with nucleophiles. As “soft” electrophiles, these sulfur-based species react primarily with “soft” nucleophiles such as thiols, phosphines and cyanide [45,46]. The reaction of a nucleophile with a polysulfide presumably occurs via direct displacement of a sulfur containing leaving group (Figure 4). For example, considering the reaction of an electrophilic disulfide with an appropriate nucleophile (e.g. a thiolate), the site of nucleophilic attack is primarily governed by the quality of the corresponding leaving group. That is, the most electrophilic sulfur, which is the site of nucleophilic attack, is the one that is attached to the best leaving group (i.e. the thiolate with a conjugate acid having the lowest pK_a) [47]. In the case of higher order polysulfides, the same trend is expected. Importantly, the pK_a of an RSSH species is typically 1-2 pK_a units lower than the corresponding RSH species [42] (and possibly much lower [32]), indicating that reactions with an $RSS\cdot$ leaving group should predominate over a reaction with an $RS\cdot$ leaving group. Thus, an RSS_nR ($n > 1$) species should be more electrophilic than the corresponding disulfide. It is also important to note that increasing the length of the polysulfur chain should also increase the inherent nucleophilicity of the sulfur atoms (due to an extended alpha-effect [19,48]). Thus, an RSSSR species should be more nucleophilic compared to a simple disulfide. Although this is chemically expected and justifiable, quantitation of this phenomenon has not been reported. Regardless, the chemical biology of higher order polysulfides can be

distinct from disulfides and these differences may be important to their biology (although this has not been specifically addressed thus far).

Physiological Implications and Future Work: Hydropersulfides are biosynthesized, incorporated into proteins and present as small molecules/peptides. Numerous descriptive studies have appeared in the literature reporting that the presence of hydropersulfides in proteins can elicit a functional change. Despite this, the overriding question remains: What is it that hydropersulfides specifically do? What is unique about hydropersulfides that prompted Nature to adopt them and evolve numerous synthetic processes to generate them (both as small molecule species and in proteins)? Is there something unique and novel about the hydropersulfide functional group that distinguishes it from, for example, thiols and/or selenols and are these unique properties a necessary aspect of fundamental biological/physiological signaling? At present, there are no clear or definitive answers to these questions, although some speculative proposals have been forwarded based on what is currently known regarding the distinct and novel chemistry of these species (*vide infra*). For example, the enhanced reductive capabilities and nucleophilicity of hydropersulfides compared to thiols may predict that their formation could be the result of cellular exposure to oxidative and/or electrophilic stress. Interestingly, proteins involved in Cys-SSH biosynthesis can be induced under oxidative/electrophilic stress, consistent with the idea that their synthesis can be a response to these stresses [34]. In this regard, it is important to reiterate that a hydropersulfide is an oxidized thiol and yet is a superior reductant compared to a thiol. That is, the oxidation of a thiol to a hydropersulfide, which could occur under oxidizing cellular conditions, generates a superior reductant. The seemingly incongruous and counterintuitive situation that a superior reductant is made via oxidation may be an important factor to the biological utility of hydropersulfides. To be sure, this is not a completely novel chemical phenomenon. For example, oxidation of an amine can generate a hydroxylamine, which is a superior one-electron reductant compared to the corresponding amine. The generation of a hydropersulfide in the face of oxidative/electrophilic stress may serve to counteract the deleterious effects associated with these stresses, akin to the way glutathione/glutathione transferase scavenges electrophiles, and antioxidants such as ascorbate or the tocopherols limit oxidative radical damage. Importantly, Paul and Snyder proposed that hydropersulfide formation in proteins may possibly protect the protein cysteine thiol from oxidation under oxidative stress [49]. However, the chemical nature of this protection was not discussed and it should be understood that conversion of a thiol to a hydropersulfide does represent an oxidation. More recently, it has been proposed that conversion of a protein thiol to a hydropersulfide results in a species

that when oxidized further is easily reversible (since oxidation of the sacrificial sulfane sulfur would be expected) under cellular reducing conditions [25,50]. Thus, irreversible oxidative modifications of protein hydropersulfides would not occur as they would for thiol functions in proteins. Importantly, selenoproteins function similarly in that they are also resistant to overoxidation to irreversibly modified species [38,51].

Clearly, other aspects of hydropersulfide chemistry may be important to their biological utility. Although not unequivocally demonstrated, the formation of a hydropersulfide at a protein active site cysteine could confer increased nucleophilic reactivity, possibly increasing enzyme activity. Thus, thiol proteins that may have important function in an oxidative and/or electrophilic stress environment may have increased activity if converted to a hydropersulfide. Also, as alluded to previously, the more physiologically accessible redox couple for $\text{RSS}^-/\text{RSS}\cdot$ compared to $\text{RS}^-/\text{RS}\cdot$ may allow the oxidation of a protein cysteine to the corresponding hydropersulfide to serve as an electron transfer “redox gate” [41]. That is, electron transfer through an $\text{RS}^-/\text{RS}\cdot$ couple is prohibitive since $\text{RS}\cdot$ is too oxidizing and RS^- is too poor a reductant – however, oxidation of the protein thiol to a hydropersulfide could allow electrons to flow since $\text{RSS}\cdot$ is a reasonably good reductant and RSS^- is not nearly as oxidizing as $\text{RS}\cdot$. Moreover, $\text{RSS}\cdot$ has been shown to be relatively unreactive towards the paramagnetic signaling species NO and O_2 , allowing the $\text{RSS}^-/\text{RSS}\cdot$ couple to be usable in aerobic and NO -rich environments [41,44]. It should be stressed that this idea is speculative at this time and there are no documented examples of this phenomenon. This is presented only because the chemical properties would predict such a scenario.

Clearly, there is a great deal of work to be done to fully elucidate hydropersulfide physiology and the associated mechanisms. The recent work of the Akaike group demonstrating translational incorporation of cysteine hydropersulfides into proteins reveals the importance of hydropersulfides in fundamental biochemistry and provides ample impetus for the further elucidation of hydropersulfide chemistry and biology. Some of the more pressing questions regarding biological hydropersulfides are the following:

1. How is hydropersulfide incorporation into proteins regulated? Protein hydropersulfides are translationally incorporated via the actions of CARS. What regulates this activity (i.e. what are the factors that determine whether CARS incorporates Cys-SH or Cys-SSH

into proteins)? Also, protein hydroperoxides can conceivably be incorporated into proteins via sulfur exchange chemistry (i.e. protein cysteines reacting with hydroperoxides or oxidized protein cysteines reacting with H₂S). With this in mind, it will be important to determine which process, translational incorporation or post-translational sulfur exchange chemistry, is the most relevant and is the relevance variable depending on cellular conditions? It will also be important to determine the specific roles (direct or indirect) of the transsulfuration enzymes such as 3-mercaptopyruvate sulfurtransferase (3MST), which is another protein capable of synthesizing hydroperoxides, in hydroperoxide production.

The temporal aspects of hydroperoxides in proteins represents an important issue in delineating the signaling and function of this modification. Thus, it is essential to determine how and when hydroperoxides are generated as well as determine how and when they are removed. In this regard, Doka and coworkers [28] have reported that intracellular protein hydroperoxide levels are highly dependent on the presence and activity of thioredoxin (Trx), an important enzyme involved in the reduction of oxidized thiol proteins. Thus, it will be important to consider the expression, activity and regulation of Trx (and possibly other reductases) when speculating on the function and signaling associated with protein hydroperoxides.

2. What determines which protein cysteines end up as hydroperoxides? It appears that only certain cysteine residues in proteins are “sulfurated” (see [33] for a discussion of this term) during translation, and possibly during sulfur exchange processes. Since amino acid incorporation into proteins is governed by codons, it will be important to discern how CARS interprets codons as a mechanism for specific hydroperoxide incorporation into proteins. With respect to post-translational sulfuration, it is appropriate to mention studies of the enzyme rhodanese, a protein with a catalytic cysteine that is known to be sulfurated. Rhodanese is capable of transferring a sulfane sulfur from a donor molecule to cyanide ion (-CN) to make thiocyanate (-SCN) via an intermediary catalytic cysteine hydroperoxide. Numerous proteins have been identified to possess a “rhodanese homology domain” associated with their catalytic cysteine residues, implicating these residues as potential sites of sulfuration [14,18,52]. For example, the phosphatase Cdc25, involved in cell-cycle regulation, possesses a catalytic rhodanese homology domain [53], possibly implicating persulfide formation as a regulatory mechanism in this system.

3. What cellular conditions are conducive to hydropersulfide formation? It has been hypothesized that oxidative and/or electrophilic stress can lead to increases in hydropersulfide generation. This idea is largely based on the fact that hydropersulfides are superior to thiols as reductants and nucleophiles; chemical attributes that may be important in combating oxidative and electrophilic stressors. In partial support of this idea, it has been shown that increased expression of CSE in A549 cells (a lung epithelial cell line) led to increased intracellular hydropersulfides and increased resistance to H₂O₂ toxicity [27]. Although this result is consistent with the idea that hydropersulfides are protective against oxidative/electrophilic stress, clearly further work must be performed to fully address this hypothesis.

4. Why are there multiple biosynthetic pathways for hydropersulfides? The recent discovery of CARS-mediated Cys-SSH generation begs the question of why there are multiple and biochemically distinct processes capable of making the same molecules (i.e. CSE/CBS, CARS, etc.). Clearly, some rationale can include compartmentalization of Cys-SSH for disparate biochemical purposes, differential regulation for accommodating varying cellular stresses/signals or simple redundancy to ensure the availability of a crucial biological effector. Answering this fundamental question will greatly assist in determining the physiological significance of hydropersulfides. Regardless, the fact that there are multiple processes to generate Cys-SSH is consistent with the idea that hydropersulfides are important biological species with varied and essential functions.

5. How does hydropersulfide incorporation into a protein affect protein activity/function? As mentioned above, numerous proteins have been reported to possess hydropersulfide residues, at least part of the time. However, it is not clear how hydropersulfides affect the activity and function of all of these proteins. Based on the current literature, it appears that hydropersulfide formation mostly confers a loss or decrease in enzyme activity. For example, the cysteine-hydrolase papain [22] and the cysteine dehydrogenase GAPDH [26] are inhibited when sulfurated. Sulfuration of erythrocyte enzymes via exposure to a hydropersulfide generating system led to the inhibition of numerous enzymes [54]. Similarly, liver enzymes treated with CSE/cystine (a system that generates Cys-SSH) were also found to be either inhibited or unaffected [55]. However, hydropersulfide formation in the protein parkin has been reported to enhance its catalytic activity, possibly leading to a protection against the development of Parkinson's disease [56]. Further studies are clearly warranted to investigate the possible alteration of protein activity as well as determination of possible functional changes.

6. What are the chemical properties that are important to the biological utility of hydropersulfides? As discussed previously, hydropersulfides should possess all of the chemical properties of a thiol, albeit enhanced. Although this is clear, it is not known which, if any, of these chemical properties are important to hydropersulfide function in biological systems. The enhanced chemistry may not yet be realized at the active sites of proteins. It should also be noted that the chemistry of a hydropersulfide function is, to a certain extent, partly disulfide and partly selenol. Since selenols and disulfides are already established biological effectors, it will be important to determine how and why the functions of hydropersulfides are distinct from the functions of these species. Also, the one-electron redox chemistry of hydropersulfides and thiols are distinct, with $\text{RSS}\cdot/\text{RSSH}$ being superior reductants compared to $\text{RS}\cdot/\text{RSH}$ and $\text{RSS}\cdot$ being much poorer oxidants compared to $\text{RS}\cdot$. Nature has found a use for $\text{RS}\cdot$ as an oxidant in numerous enzymatic systems [6]. Since $\text{RS}\cdot$ is a much stronger oxidant than $\text{RSS}\cdot$, the use of $\text{RSS}\cdot$ as an oxidant in analogous systems is not expected. The question remains, however, whether Nature has also found a use for $\text{RSS}\cdot$ or the $\text{RSS}\cdot/\text{RSS}\cdot$ redox couple. The stability of $\text{RSS}\cdot$ indicates that $\text{RSS}\cdot/\text{RSSH}$ can conceivably act as a one-electron reductant similar to the antioxidants ascorbate and tocopherols. Indeed, the antioxidant properties of RSSH have already been described [42] and it is intriguing to speculate that endogenously generated hydropersulfides also function in this manner. Interestingly, numerous enzymes proposed to possess hydropersulfides also have redox activity. For example, xanthine oxidase, an enzyme that catalyzes the oxidation of (hypo)xanthine with subsequent reduction of O_2 to superoxide ($\text{O}_2\cdot^-$), was among the first to be proposed to contain a hydropersulfide [10]. Regardless, it remains to be determined whether the $\text{RSS}\cdot/\text{RSS}\cdot$ couple has biological utility.

7. What are the biological roles and functions of the higher order polysulfide species? Although this article has focused mostly on RSSH species, it has been mentioned that polysulfides (RSS_nR , $n>1$, R = hydrogen or alkyl) are also prevalent and related to hydropersulfides. Indeed, the generation of these higher order polysulfides is a result of H_2S and hydropersulfide formation since all of these species can be related via a series of equilibria. In fact, it is going to be difficult to distinguish among the possible biological activities of all these species since they are related to each other via equilibrium reactions and, therefore, can all be present simultaneously in a biological milieu. Regardless, it is entirely possible, if not probable, that polysulfides are not merely storage forms of H_2S or RSSH species, but rather possess important biological activities and properties on their own. Discerning these activities/properties will be important in the full elucidation of the chemical biology and physiology of the hydropersulfides.

8. Are hydropersulfides involved in metal/metalloprotein biology? Thus far, much of the focus on hydropersulfide biochemistry has primarily involved interactions with other sulfur species (*vide supra*). Due to the presumed novel ligation and redox properties of hydropersulfides, metals and metalloproteins are also likely sites of biological reactivity. Previous studies have demonstrated that the RSSH species can reduce ferric heme proteins such as cytochrome c under conditions where thiols or H₂S do not [22] (and, as mentioned above, the biological relevance/utility of the one-electron redox properties of hydropersulfides are worthy of further investigation). It also seems possible, if not likely, that hydropersulfides will interact with biologically non-redox active metal ions such as Zn²⁺ as well. As a borderline “soft” ion, Zn²⁺ can coordinate a wide array of biological ligands containing S, N and O atoms. For example, Zn-finger proteins bind Zn²⁺ via histidine nitrogens and cysteine thiols [57]. The regulation of Zn²⁺ “re-distribution” (i.e. movement of Zn²⁺ from storage sites to protein binding sites) is an important field due to the prevalence and of Zn-containing proteins in signaling and fundamental biochemistry [58]. Indeed, one mechanism of mobilization of Zn²⁺ from sites of sequestration is thought to be via redox chemistry at Zn-binding thiolates, coupling redox signaling with Zn-dependent signaling/biochemistry. It is intriguing to speculate that cysteine persulfide anions can also be involved in Zn²⁺ coordination, allowing a different redox couple (compared to thiolate) to affect Zn²⁺ binding, release and/or mobilization. Moreover, it is highly likely that the coordination chemistry of a persulfide anion versus a thiolate will be significantly different, thus altering the affinity of proteins for Zn²⁺ and possibly resulting in a chemically distinct pool of Zn²⁺ (e.g. where the redox chemistry of the Zn²⁺ ligands allows for release under different redox conditions).

One aspect of the interaction of hydropersulfides with metals in a biological system that has been examined is the effect of persulfides on electrophilic heavy metal toxicity [59]. Abiko and coworkers [60] found that detoxification of methyl mercury can occur via interaction with H₂S, hydropersulfides and polysulfides. Similarly, Cd²⁺ toxicity can also be abated by the presence of hydropersulfides [61]. Importantly, Zn²⁺ and Cd²⁺ are chemically and biochemically similar in that they both readily bind thiol proteins and indeed a facet of Cd²⁺ toxicity is thought to involve interference with Zn²⁺ homeostasis [62]. The fact that Cd²⁺ and Zn²⁺ are chemically similar and persulfides alter the pathobiology of Cd²⁺ is consistent with the idea that endogenous persulfides can also be involved in Zn²⁺ regulation.

9. Are hydropersulfides involved in the etiology of disease or involved in redox signaling? One of the most important aspects of hydropersulfide/polysulfide chemical biology is with respect to their possible roles in disease. As alluded to above, hydropersulfides possess the proper chemical properties to protect cells from certain chemical stresses, although this remains to be established. Moreover, hydropersulfides possess chemical properties that are distinct from thiols (and other derived species), allowing them to perform biochemistry not available to simple thiols. Considering all of this, these species may be involved in many diseases, disorders or developmental pathways whereby “oxidative” or electrophilic stress and/or signaling have been implicated and/or where thiol modification processes are proposed. Moreover, if hydropersulfides are fundamental to cellular protection, they may also be important aspects of, for example, cancer cell viability, neuronal cell degeneration, regulation of apoptosis and/or redox-mediated developmental pathways. These are highly speculative ideas, but it is clear that the impact of hydropersulfides on disease states, disorders, developmental signaling or cellular protection is potentially significant.

The points listed above likely represent only the “tip-of-the-iceberg” with regard to pressing questions in this field. Moreover, details of the physiological implications of hydropersulfides and polysulfides remain to be addressed once some of the above questions are answered. It appears likely that the current ideas regarding the biological signaling associated with thiol sulfur modification will need to be revisited and redefined in the light of the recent findings pertaining to the prevalence of hydropersulfides and polysulfides. Moreover, hydropersulfide/polysulfide-derived species that are analogous to established thiol-derived species will also be essential to include in any conversation involving sulfur species. For example, thiol oxidation to the corresponding sulfenic acid is a well-known and established process under certain biological conditions [63]. It will now be equally important to consider the analogous reaction with hydropersulfides under those same conditions. That is, oxidation of a hydropersulfide to the corresponding perthiosulfenic acid (RSSOH) is an equally likely, if not more likely, fate for hydropersulfides [64]. It is unnecessary to list all of the biologically relevant thiol modification reactions but yet important to indicate that any type of thiol modification reaction (many of which have been associated with thiol-based signaling) can occur and are likely with hydropersulfides, possibly with important signaling/biochemical consequences. Regardless, the potential importance and impact of hydropersulfides and polysulfides on the fields of fundamental biochemistry and (patho)physiology, among many others, appears significant and burgeoning and, in our opinion, it would be unwise not to consider these novel species as important molecules in the chemical biology of nature.

Footnote

¹ The term “sulfhydration” was used to denote the addition/insertion of a sulfane sulfur (basically atomic sulfur) to/into a molecule. That is, the conversion of RSH to RSSH was referred to as “sulfhydration” of RSH. This term may not be the most accurate description of the chemistry since “hydration” describes reactions that are simply represented by the addition of water. For example, the addition of water to an olefin to generate the corresponding alcohol is a hydration reaction. The term “sulfhydration” may have been derived from the term “sulfhydryl”, which has been used to describe an -SH group (note that “sulf” denotes a sulfur atom and “hydryl” denotes a hydrogen and is distinct from “hydration”). Thus, it may be tempting to invoke the term “sulfhydrylation”. However, this terminology is also not an accurate description of the chemistry since conversion of RSH to RSSH is not the addition of an -SH (sulfhydryl) group but, as described above, the addition of a sulfur atom. Therefore, as Toohey discusses and recommends [33], “sulfhydration” should be replaced by “sulfuration” for these reactions since the conversion of RSH to RSSH is equivalent to the addition of a sulfur atom.

References

- [1] Go Y.M. and Jones D.P. (2013). Thiol/disulfide redox states in signaling and sensing. *Crit Rev Biochem Mol Bio.* 48, 173-191.
- [2] Paulsen, C.E. and Carroll, K.S. (2013). Cysteine-mediated redox signaling: Chemistry, Biology and Tools for Discovery. *Chem. Rev.* 113, 4633-4679.
- [3] Winterbourn, C.C. and Hampton, M.B. (2008). Thiol chemistry and specificity in redox signaling. *Free Radic. Biol. Med.* 45, 549-561.
- [4] Reddie, K.G. and Carroll, K.S. (2008). Expanding the functional diversity of proteins through cysteine oxidation. *Curr. Opin. Chem. Biol.* 12, 746-754.
- [5] Jacob, C., Giles, G.I., Giles, N.M., and Sies, H. (2003). Sulfur and Selenium: The role of oxidation state in protein structure and function. *Angew Chem Int Ed.* 42, 4742-4758.
- [6] Stubbe, J. and van der Donk, W.A. (1998). Protein radicals in enzyme catalysis. *Chem Rev.* 98, 705-762.
- [7] Bonnet, D., Stevens, J.M., de Sousa, R.A., Sari, M.A., Mansuy, D., and Artaud, I. (2001). New inhibitors of iron-containing nitrile hydratases. *J Biochem (Tokyo).* 130, 227-233.

- [8] Munchberg, U., Anwar, A., Mecklenburg, S., and Jacob, C. (2007). Polysulfides as biologically active ingredients of garlic. *Org Biomol Chem.* 5, 1505-1518.
- [9] Calabrese, L., Federici, G., Bannister W.H., Bannister, J.V., Rotilio, G., and Finazzi-Agro, A. (1975). Labile sulfur in human superoxide dismutase. *Eur J Biochem.* 56, 305-309.
- [10] Massey, V. and Edmondson, D. (1970). On the mechanism of inactivation of xanthine oxidase by cyanide. *J Biol Chem.* 245, 6595-6598.
- [11] Branzoli, U. and Massey, V. (1974). Evidence for an active site persulfide residue in rabbit liver aldehyde oxidase. *J Biol Chem.* 249, 4346-4349.
- [12] Nagahara, N., Nagano, M., Ito, T., and Suzuki, H. (2016). Redox regulation of mammalian 3-mercaptopyruvate sulfurtransferase. *Meth Enzymol.* 554, 229-254.
- [13] Zheng, L., White, R.H., Cash, V.L., and Dean, D.R. (1994). Mechanism for the desulfurization of L-cysteine catalyzed by the *nofS* gene product. *Biochemistry.* 33, 4714-4720.
- [14] Cipollone, R., Ascenzi, P., and Visca, P. (2007). Common themes and variations in the rhodanese superfamily. *IUBMB Life.* 59, 51-59.
- [15] Mueller, E.G. (2006). Trafficking in persulfides: Delivering sulfur in biosynthetic pathways. *Nature Chem Biol.* 2, 185-194.
- [16] Toohey, J.I. (1989). Sulphane sulphur in biological systems: a possible regulatory role. *Biochem J.* 264, 625-632.
- [17] Wood, J.L. (1987). Sulfane sulfur. *Meth Enzymol.* 143, 25-29.
- [18] Toohey, J.I. (2011). Sulfur signaling: Is the agent sulfide or sulfane. *Anal Biochem.* 412, 1-7.
- [19] Nagy, P. (2015). Mechanistic chemical perspective of hydrogen sulfide signaling. *Meth Enzymol.* 554, 3-29.
- [20] Blackstone, E., Morrison, M., and Roth, M.B. (2005). H₂S induces a suspended animation-like state in mice. *Science.* 308, 518.
- [21] Cavallini, D., Federici, G., and Barboni, E. (1970). Interactions of proteins with sulfide. *Eur. J. Biochem.* 14, 169-174.
- [22] Francoleon, N.E., Carrington, S.J., and Fukuto, J.M. (2011). The reaction of H₂S with oxidized thiols: Generation of persulfides and implications to H₂S biology. *Arch. Biochem. Biophys.* 516, 146-153.

- [23] Mustafa, A.K., Gadalla, M.M., Sen, N., Kim, S., Mu, W., Gazi, S.K., Barrow, R.K., Yang, G., Wang, R., and Snyder, S.H. (2009). H₂S Signals through protein S-sulfhydration. *Sci Sign.* 2, 1-8.
- [24] Pan, J. and Carroll, K.S. (2013). Persulfide reactivity in the detection of protein S-sulfhydration. *ACS Chem. Biol.* 8, 1110-1116.
- [25] Ono, K., Akaike, T., Sawa, T., Kumagai, Y., Wink, D.A., Tantillo, D.J., Hobbs, A.J., Nagy, P., Xian, M., Lin, J. and Fukuto, J.M. (2014). The redox chemistry and chemical biology of H₂S, hydropersulfides and derived species: Implications of their possible biological activity and utility. *Free Radic. Biol. Med.* 77, 82-94.
- [26] Jarosz, A.P., Wei, W., Gauld, J.W., Auld, J., Ozcan, F., Aslan, M., and Mutus, B. (2015). Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) is inactivated by S-sulfuration in vitro. *Free Rad Biol Med.* 89, 512-521.
- [27] Ida, T., Sawa, T., Ihara, H., Tsuchiya, Y., Watanabe, Y., Kumagai, Y., Suematsu, M., Motohashi, H., Fujii, S., Matsunaga, T., Yamamoto, M., Ono, K., Devarie-Baez, N.O., Xian, M., Fukuto, J.M. and Akaike, T. (2014). Reactive cysteine persulfides and S-polythiolation regulate oxidative stress and redox signaling. *Proc. Natl., Acad., Sci., USA.* 111, 7606-7611.
- [28] Doka, E., Pader, I., Biro, A., Johansson, K., Cheng, Q., Ballago, K., Prigge, J.R., Pastor-Flores, D., Dick, T.P., Schmidt, E.E., Arner, E.S.J., and Nagy, P. (2016). A novel persulfide detection method reveals protein persulfide- and polysulfide-reducing function of thioredoxin and glutathione systems. *Sci Adv.* 2, e1500968.
- [29] Longen, S., Richter, F., Kohler, Y., Wittig, I., Beck, K.F., and Pfeilschifter, J. (2016). Quantitative persulfide site identification (qPerS-SID) reveals protein targets of H₂S releasing donors in mammalian cells. *Scientific Reports.* 6, 29808.
- [30] Akaike, T., Ida, T., Wei, F.W., Nishida, M., Kumagai, Y., Alam, M., Ihara, H., Sawa, T., Matsunaga, T., Kasamatsu, S., Nishimura, A., Morita, M., Tomizawa, K., Nishimura, A., Watanabe, S., Inaba, K., Shima, H., Tanuma, N., Jung, M., Fujii, S., Watanabe, Y., Ohmuraya, M., Nagy, P., Feelisch, M., Fukuto, J.M. and Motohashi, H. (2017). Cysteinyl-tRNA synthase governs cysteine polysulfidation and mitochondrial bioenergetics. *Nature Comm.* 8, 1177, 1-15.
- [31] Vasas, A., Doka, E., Fabian, I., and Nagy, P. (2015). Kinetic and thermodynamic studies on the disulfide-bond reducing potential of hydrogen sulfide. *Nitric Oxide.* 46, 93-101.

- [32] Cuevasanta, E., Lange, M., Bonanata, J., Coitino, E.L., Ferrer-Sueta, G., Filipovic, M.R., and Alvarez, B. (2015). Reaction of hydrogen sulfide with disulfide and sulfenic acid to form the strongly nucleophilic persulfide. *J. Biol. Chem.* 290, 26866-26880.
- [33] Toohey, J.I. (2012). The conversion of H₂S to sulfane sulfur. *Nature Rev Mol Cell Biol.* 13, 803.
- [34] Alvarez, L., Bianco, C. L., Toscano, J. P., Lin, J., Akaike, T., and Fukuto, J.M. (2017). The chemical biology of hydropersulfides and related species: Possible roles in cellular protection and redox signaling. *Antiox. Redox Sign.* 27, 622-633.
- [35] Turanov, A.A., Xu, X.M., Carlson, B.A., Yoo, M.H., Gladyshev, V.N., and Hatfield, D.L. (2011). Biosynthesis of selenocysteine, the 21st amino acid in the genetic code, and a novel pathway for cysteine biosynthesis. *Adv Nutr.* 2, 122-128.
- [36] Lu, J. and Holmgren, A. (2009). Selenoproteins. *J Biol Chem.* 284, 723-727.
- [37] Srinivasan, G., James, C.M., and Krzycki, J.A. (2002). Pyrrolysine encoded by UAG in Archaea: Charging of a UAG-decoding specialized tRNA. *Science.* 296, 1459-1462.
- [38] Reich, H.J. and Hondal, R.J. (2016). Why Nature chose selenium. *ACS Chem Biol.* 11, 821-841.
- [39] Sen, N., Paul, B.D., Gadalla, M.M., Mustafa, A.K., Sen, T., Kim, S., and Snyder, S.H. (2012). Hydrogen sulfide-linked sulfhydration of NF- κ B mediates its antiapoptotic actions. *Mol. Cell.* 45, 13-24.
- [40] Winterbourn, C.C. (2015). Are free radicals involved in thiol-based redox signaling? *Free Radic Biol Med.* 80, 164-170.
- [41] Bianco, C.L., Chavez, T.A., Sosa, V., Nguyen, Q.N.N., Tantillo, D.J., Ichimura, A.S., Toscano, J. P., and Fukuto, J.M. (2016). The chemical biology of the persulfide (RSSH)/perthiyl (RSS \cdot) redox couple and possible role in biological redox signaling. *Free Rad. Biol. Med.* 101, 20-31.
- [42] Everett, S.A. and Wardman, P. (1995). Perthiols as antioxidants: Radical-scavenging and prooxidative mechanisms. *Meth Enzymol.* 251, 55-69.
- [43] Mahoney, L.R., Mendenhall, G.D., and Ingold, K.U. (1973). Calorimetric and equilibrium studies on some stable nitroxide and iminoxyl radicals. Approximate O-H bond dissociation energies in hydroxylamines and oximes. *J Am Chem Soc.* 95, 8610-8614.

- [44] Chauvin, J.R., Griesser, M., and Pratt, D.A. (2017). Hydropersulfides: H-Atom transfer agents par excellence. *J. Am. Chem. Soc.* 139, 6484-6493.
- [45] Saund, S.S., Sosa, V., Henriquez, S., Nguyen, Q.N.N., Bianco, C.L., Soeda, S., Millikin, R., White, C., Le, H., Tantillo, D.J., Kumagai, Y., Akaike, T., Lin, J., and Fukuto, J.M. (2015). The chemical biology of hydropersulfides (RSSH): Chemical stability, reactivity and redox roles. *Arch. Biochem. Biophys.* 588, 15-24.
- [46] Pearson, R.G. (1987). Recent advances in the concept of hard and soft acids and bases. *J. Chem. Educ.* 64, 561-567.
- [47] Hiskey, R.G. and Carrol, F.I. (1961). Chemistry of aliphatic disulfides. II. Cyanide cleavage of unsymmetrical disulfides. *J. Am. Chem. Soc.* 83, 4644-4647.
- [48] Fina, N.J. and Edwards, J.O. (1973). The Alpha Effect. A Review. *Int J Chem Kin.* V, 1-23.
- [49] Paul, B.D., and Snyder, S.H. (2012). H₂S Signaling through protein sulfhydration and beyond. *Nature Rev., Mol Cell Biol.* 13, 499-506.
- [50] Millikin, R., Bianco, C.L., White, C., Saund, S.S., Henriquez, S., Sosa, V., Akaike, T., Kumagai, Y., Soeda, S., Toscano, J.P., Lin, J. and Fukuto, J.M. (2016). The chemical biology of protein hydropersulfides: Studies of a possible protective function of biological hydropersulfide generation. *Free Rad. Biol. Med.* 97, 136-147.
- [51] Snider, G.W., Ruggles, E., Kahn, N., and Hondal, R.J. (2013). Selenocysteine confers resistance to inactivation by oxidation in thioredoxin reductase: Comparison of selenium and sulfur enzymes. *Biochemistry*, 52, 5472-5481.
- [52] Bordo, D. and Bork, P. (2002). The rhodanese/Cdc25 phosphatase superfamily. *EMBO Reports*. 3, 741-746.
- [53] Hofmann, K., Bucher, P., and Kajava, A.V. (1998). A model of Cdc25 phosphatase catalytic domain and Cdk-interaction surface based on the presence of a rhodanese homology domain. *J Mol Biol.* 282, 195-208.
- [54] Valentine, W.N., Toohey, J.I., Paglia, D.E., Nakatani, M., and Brockway, R.A. (1987). Modification of erythrocyte enzyme activities by persulfides and methanethiol: Possible regulatory role. *Proc. Natl. Acad. Sci., USA.* 84, 1394-1398.

- [55] Ogasawara, Y., Suzuki, T., Ishii, K., and Tanabe, S. (1997). Modification of liver cytosol enzyme activities promoted in vitro by reduced sulfur species generated from cystine with γ -cystathionase. *Biochim Biophys Acta*. 1334, 33-43.
- [56] Vandiver, M.S., Paul, B.D., Xu, R., Karuppagounder, S., Rao, F., Snowman, A.M., Ko, H.S., Lee, Y.I., Dawson, V.L., Dawson, T.M., Sen, N., and Snyder, S.H. (2013). Sulfhydration mediates neuroprotective actions of parkin. *Nat. Commun.* 4, 1626.
- [57] Laity, J.H., Lee, B.M., and Wright, P.E. (2001). Zinc finger proteins: New insights into structural and functional diversity. *Curr Opin Struct Biol.* 11, 39-46.
- [58] Krezel, A., Hao, Q., and Maret, W. (2007). The zinc/thiolate redox biochemistry of metallothionein and the control of zinc ion fluctuations in cell signaling. *Arch Biochem Biophys.* 463, 188-200.
- [59] Kumagai, Y. and Abiko, Y. (2017). Environmental electrophiles: protein adducts, modulation of redox signaling and interaction with persulfides/polysulfides. *Chem Res Toxicol.* 30, 203-219.
- [60] Abiko, Y., Yoshida, E., Ishii, I., Fukuto, J.M., Akaike, T., and Kumagai, Y. (2015). Involvement of reactive persulfides in biological dimethylmercury sulfide formation. *Chem Res Toxicol.* 28, 1301-1306.
- [61] Shinkai, Y., Masuda A., Akiyama, M., Xian M., and Kumagai, Y. (2017). Cadmium-mediated activation of the HSP90/HSF1 pathway regulated by reactive persulfides/polysulfides. *Toxicol Sci.* 156, 412-421.
- [62] Martelli, A., Rousselet, E., Dycke, C., Bouron, A., and Moulis, J.M. (2006). Cadmium toxicity in animal cells by interference with essential metals. *Biochimie.* 88, 1807-1814.
- [63] Conte, M.L. and Carroll, K.S. (2013). The redox biochemistry of protein sulfenylation and sulfinylation. *J. Biol. Chem.* 288, 26480-26488.
- [64] Heppner, D.E., Hristova, M., Ida, T., Mijuskovic, A., Dustin, C.M., Bogdandi, V., Fukuto, J.M., Dick, T.P., Nagy, P., Li, J., Akaike, T., van der Vliet, A. (2018). Cysteine perthiosulfenic acid (Cys-SSOH): A novel intermediate in thiol based redox signaling? *Redox Biol.* 14, 379-385.

Figure legends

Figure 1 A schematic drawing for the co-translational formation of CysSSH and protein polysulfidation as catalyzed by CARSs. Cysteine persulfide (CysSSH) formation and protein polysulfidation are mediated by the novel CysSSH-producing enzyme CARS. CBS and CSE are involved in the cysteine biogenesis in canonical transsulfuration pathway. Cystine transport by cystine-glutamate transporter system mediated by xCT is also sustaining the intracellular cysteine. Cysteine biosynthesis and cystine uptake are known to be up-regulated by Nrf2 (NF-E2-related factor 2), and the persulfides and polysulfides formed are reductively metabolized by glutathione reductase/thioredoxin (GR/Trx) system.

Figure 2 Schematics of the biosynthetic pathways for A) Cys-SSH and B) SeCys. Cysteine tRNA synthetase (CARS); Serine tRNA synthetase (SARS); Phosphoserine-tRNA kinase (PSTK); Selenophosphate synthetase 2 (SEPHS2); Sep (O-phosphoserine) tRNA; SeCys (Selenocysteine) tRNA synthetase (SEPSECS).

Figure 3 Comparative properties of RSS[•]/RSSH versus RS[•]/RSH.

Figure 4 Reaction of a nucleophile with a polysulfur species.

Figure 1




Figure 2




Figure 3




Figure 4

