

Stochastic modelling and projection of mortality improvements using a hybrid parametric/semi-parametric age-period-cohort model *

Erengul Dodd^{1,2,4}, Jonathan J. Forster^{1,4,5}, Jakub Bijak^{1,3,4}, and Peter W. F. Smith^{1,3,4}

¹Southampton Statistical Sciences Research Institute, University of Southampton, Southampton, SO17 1BJ, UK

²Mathematical Sciences, University of Southampton, Southampton, SO17 1BJ, UK

³Social Sciences, University of Southampton, Southampton, SO17 1BJ, UK

⁴ESRC Centre for Population Change, University of Southampton, Southampton, SO17 1BJ, UK

⁵Department of Statistics, University of Warwick, Coventry, CV4 7AL, UK

Abstract

We propose a comprehensive and coherent approach for mortality projection using a maximum likelihood method which benefits from full use of the substantial data available on mortality rates, their improvement rates, and the associated variability. Under this approach we fit a negative binomial distribution to overcome one of the several limitations of existing approaches such as insufficiently robust mortality projections as a result of employing a model (e.g. Poisson) which provides a poor fit to the data. We also impose smoothness in parameter series which vary over age, cohort, and time in an integrated way. Generalised Additive Models (GAMs), being a flexible class of semi-parametric statistical models, allow us to differentially smooth components, such as cohorts, more heavily in areas of sparse data for the component concerned. While GAMs can provide a reasonable fit for the ages where there is adequate data, estimation and extrapolation of mortality rates using a GAM at higher ages is problematic due to high variation in crude rates. At these ages, parametric models can give a more robust fit, enabling a borrowing of strength across age groups. Our projection methodology assumes a smooth transition between a GAM at lower ages and a fully parametric model at higher ages.

Keywords: Age-period-cohort model, generalised additive model, overdispersed data, projection, expert opinion.

1 Introduction

Recent mortality improvements in most countries have led to higher life expectancies. Since this has significant social policy implications, for example in such areas as pensions and healthcare, modelling and projecting mortality rates becomes imperative. For example, in the insurance industry, the risk of higher than expected annuity payments, or the so-called ‘longevity risk’ needs to be quantified for solvency requirements. The mortality projections are not only important for

*The majority of this work was supported through a research contract (‘Review of Mortality Projections’) between the UK Office for National Statistics (ONS) and the University of Southampton. This included helpful discussions with ONS colleagues and members of the Mortality Assumptions Expert Users Group, convened by ONS. Additional support was provided by the ESRC Centre for Population Change - Phase II (ES/K007394/1). All the views presented in this paper are of the authors only.

pensions or healthcare but also in allocating resources and government planning, for example, for housing, education and labour market. This requires probabilistic models for mortality rates. In the past decade, a vast literature on probabilistic mortality models has been developed. However, very few of them are suitable for the entire age range. In this paper, instead of focusing only on older ages, we present a mortality projection methodology for the entire population.

Probabilistic mortality projection models can be broadly summarised under four categories: generalised bilinear, generalised linear, semi-parametric and generalised additive models. The original Lee-Carter model (Lee and Carter, 1992) is the pioneering method in this area. This model is an example of a bilinear model and has two factors, i.e. age and period, to model and forecast mortality rates. The other bilinear models include extensions of the Lee-Carter model, for example three-factor models that encompass the cohort effect or Poisson error structure instead of an implied normality assumption (see e.g. Renshaw and Haberman (2006); Brouhns et al. (2002)). Whilst these models can provide a satisfactory fit, they also have some undesirable features. In particular their parameter estimates can be sensitive to the range of years used for fitting, they are challenging to estimate efficiently.

Alternatively, linear (rather than bilinear) models with age and period as factors were investigated by Renshaw and Haberman (2003). Different linear structures were developed and compared by Cairns et al. (2009, 2011a). Many studies show that coherent mortality forecasts can be obtained for different populations using these models (e.g. Cairns et al. (2011b); Li and Hardy (2011); Börger and Aleksic (2014)). Currie et al. (2004) proposed modelling mortality as a smooth function in two dimensions (age and time) using P-spline methodology, although such an approach can be difficult to incorporate into a projection since it ignores the majority of the observed historical mortality experience and is too sensitive to mortality in the base year, even after smoothing (Li et al., 2010). Besides, this approach does not allow for coherent projections for different populations (Börger and Aleksic, 2014). Approaches which fully account for uncertainty include Cairns et al. (2006, 2011b), Bennett et al. (2015) and Hilton et al. (2019).

In particular, Hilton et al. (2019) provide a Bayesian approach to producing mortality projections based on the use of generalised additive models (GAMs) for the majority of the age range, but with an inclusion of a parametric model at older ages where the data are sparse. Their approach allows for smooth functions of age and cohort effects and provides estimates of mortality at young ages as well as extreme ages. Bayesian models incorporate multiple sources of uncertainty and expert opinion in a natural way. However, implementing a full Bayesian approach might be expensive, especially if computing marginal likelihoods requires high dimensional integrals and posterior distributions are analytically intractable.

In this paper we present the maximum likelihood approach to the methodology presented in Hilton et al. (2019). Additionally, we incorporate expert opinion in our projections. We model the mortality improvements instead of mortality rates allowing for overdispersion. This is because we believe mortality improvements can be modelled by a stable process which is required to be projected forward based on the past experience. Projection of mortality improvement rates is advocated by Plat (2011), Haberman and Renshaw (2012, 2013) and more recently by Börger and Aleksic (2014). In the United Kingdom (UK), the Continuous Mortality Investigation (CMI) introduced Age-Period-Cohort Improvement (APCI) model as a new mortality projection method (CMI, 2016). However the CMI does not use the APCI model as a stochastic model to project future mortality rates. The CMI uses the APCI model to simply obtain the initial mortality improvements (separated by age, period and cohort related improvements) for projections. Richards et al. (2019) implemented the APCI model as a fully stochastic model. They compared this model to the Age-Period, Age-Period-Cohort and Lee-Carter models and found that the APCI model fits the data better than these other models considered in their paper.

In CMI (2016), the CMI recognises that the Office for National Statistics (ONS) dataset show

considerable overdispersion, relative to a Poisson error distribution. In the existence of overdispersion, modelling the observed number of deaths under a single parameter distribution such as a Poisson distribution (where the variance is restricted to be equal to the mean) will lead to underestimation of uncertainty. To allow for overdispersion, we use a more flexible negative binomial distribution in modelling. We also impose smoothness in parameter series which vary over age, cohort, and time in an integrated way. GAMs, being a flexible class of semi-parametric statistical models, allow us to differentially smooth components, such as cohorts, more heavily in areas of sparse data for the component concerned.

While GAMs can provide a reasonable fit for the ages where there is adequate data, estimation and extrapolation of mortality rates using a GAM at higher ages is problematic due to high variation in crude rates. At these ages, parametric models can give a more robust fit, enabling a borrowing of strength across age groups. Our projection methodology is based on a smooth transition between a GAM at lower ages and a fully parametric model at higher ages. We model infant rates separately and propose a new method to model and predict them. Since spline-based methods are used widely in the literature (especially for the entire age range), as discussed above, we compare our results to the P-spline model where relevant.

The rest of the paper is organised as follows: In Section 2, we introduce the data and our methodology for modelling the mortality improvements, and how a smooth transition from the smoothing spline to an old-age model is attained. In Section 3, we present our estimates for mortality rates and investigate the robustness of the proposed methodology. In Section 4 we present our mortality projections and provide comparisons with the UK national population projections. Our conclusions are in Section 5.

2 The data and the model

We use the UK population data between 1961 and 2013 obtained from the Human Mortality Database (Human Mortality Database, 2019). The data include the mid-year exposures and number of deaths for each year of age for males and females.

Here we propose a model that contains terms which specifically account for variation of mortality differences over time and between different ages and cohorts. Let m_{xt} denote central mortality rates at age x in year t , then we consider as the initial model specification

$$\log \frac{m_{xt}}{m_{xt-1}} = \alpha_x + \kappa_t + \gamma_{t-x} \quad (1)$$

where α_x can be interpreted as a baseline annual mortality improvement at age x , κ_t as the level of mortality improvement in year t and γ_{t-x} represents cohort differences in mortality improvement since cohorts are indexed by year of birth ($t - x$).

Model (1) is an age-period-cohort model for log-mortality differences (mortality logratios). Here we represent mortality improvements as logratios, rather than as relative differences, where the model (1) would be expressed as

$$\frac{m_{xt} - m_{xt-1}}{m_{xt-1}} = \alpha_x + \kappa_t + \gamma_{t-x}.$$

For all but large mortality rates, differences between $\log \frac{m_{xt}}{m_{xt-1}}$ and $\frac{m_{xt} - m_{xt-1}}{m_{xt-1}}$ are negligible. This model is similar in structure to models proposed by Renshaw and Haberman (2003) and the non-spatial component of Bennett et al. (2015).

Note that in terms of mortality rates, model (1) can be expressed as

$$\log m_{xt} = \mu_x + \alpha_x t + \kappa_t + \gamma_{t-x} \quad (2)$$

where there is a straightforward correspondence between the κ_t and γ_{t-x} parameters of models (1) and (2). Due to the linear relationship between age, period and cohort components of the model, constraints are required in order to identify these effects. The identifiability constraints we use for model (1) are

$$\sum \kappa_t = 0, \quad (3)$$

$$\sum \gamma_{t-x} = 0, \quad (4)$$

$$\sum (t-x)\gamma_{t-x} = 0. \quad (5)$$

That is, the period effect is constrained to sum to zero. Similarly, the cohort effect is constrained so that the sum of effects is zero, and display zero growth over the whole range of cohorts.

In Section 3 we present parameter estimates for model (1), together with μ_x . This model, being simply a generalised linear model, is efficient to fit. Furthermore its parameter estimates seem to be robust to the time window used to fit the models. Börger and Aleksic (2014) advocate the use of this model for projecting mortality, and we also find that it has the required properties of adequately and robustly fitting the observed data.

In the literature it is very common to estimate the model parameters based on the Poisson log-likelihood. However, under the Poisson model the variance is restricted to be equal to the mean, an assumption which is implausible for a large inhomogeneous population. A more flexible model would be a negative binomial model where the log-likelihood is

$$l(\boldsymbol{\theta}, a) = \sum_{x,t} a \log \left(\frac{a}{E_{xt} m_{xt}(\boldsymbol{\theta}) + a} \right) + \sum_{x,t} d_{xt} \log \left(\frac{E_{xt} m_{xt}(\boldsymbol{\theta})}{E_{xt} m_{xt}(\boldsymbol{\theta}) + a} \right) + \sum_{x,t} \log \Gamma(a+d_{xt}) - n \log \Gamma(a).$$

Here $\boldsymbol{\theta}$ represents the model parameters ($\alpha_x, \kappa_t, \gamma_{t-x}$), d_{xt} is the observed death count, E_{xt} is the central exposure to risk at age x in year t , a is the dispersion parameter such that the variance is $E_{xt} m_{xt}(\boldsymbol{\theta}) + (E_{xt} m_{xt}(\boldsymbol{\theta}))^2/a$ and n is the number of positive values of E_{xt} .

One disadvantage of model (1) is that the maximum likelihood estimates of some of the model parameters do not vary smoothly (Figure 1). However, this can be easily overcome by adopting an estimation method which penalises roughness in the series of estimates for model (1) (e.g. penalised likelihood or Bayesian). One possible way of obtaining smoother estimates is to modify (2) to yield a generalised additive model of the form:

$$m_{xt} = \exp (s_\mu(x) + s_\alpha(x) t + \kappa_t + s_\gamma(t-x)), \quad (6)$$

In (6), s_μ , s_α and s_γ denote arbitrary smooth functions, which can be estimated by balancing goodness-of-fit to the observed data with smoothness of the corresponding function (Wood, 2006). This can be fitted by using standard gam packages in R (e.g. using gam function in `mgcv` package).

For the highest ages x , for which observed mortality experience is sparse, we recommend that the baseline mortality μ_x and the age-specific mortality differences α_x are estimated by using parametric models, for example log-linear model or logistic model, with parameters estimated from the mortality data for the older ages. The resulting log-linear model, with $\mu_x = \mu + \mu_X x$ and $\alpha_x = \alpha + \alpha_X x$ has the form

$$m_{xt} = \exp (\mu + \mu_X x + (\alpha + \alpha_X x) t + \kappa_t + s_\gamma(t-x)), \quad \text{for } x \geq x_0 \quad (7)$$

and logistic model has the form

$$m_{xt} = \frac{\beta \exp (\mu + \mu_X x + (\alpha + \alpha_X x) t + \kappa_t + s_\gamma(t-x))}{1 + \exp (\mu + \mu_X x + (\alpha + \alpha_X x) t)}, \quad \text{for } x \geq x_0 \quad (8)$$

where x_0 is an optimal age to make the transition from smooth to linear model, κ_t and $s_\gamma(t-x)$ are the estimates obtained from fitting (6) to the main body of data ($0 < x < x_0$). In (8), κ_t and $s_\gamma(t-x)$ can be considered as an unstandard ‘offset’ as they only appear in the numerator and therefore can be fitted using a general optimisation function (e.g. `optim` or `nlm`) in R .

The log-linear model has therefore the following estimates of the baseline mortality

$$\mu_x = \begin{cases} s_\mu(x) & x < x_0 \\ \mu + \mu_X x & x \geq x_0 \end{cases}$$

and mortality improvement

$$\alpha_x = \begin{cases} s_\alpha(x) & x < x_0 \\ \alpha + \alpha_X x & x \geq x_0 \end{cases}$$

for both males and females. For the logistic model these estimates are, respectively,

$$\mu_x = \begin{cases} s_\mu(x) & x < x_0 \\ \log\left(\frac{\beta \exp(\mu + \mu_X x)}{1 + \exp(\mu + \mu_X x)}\right) & x \geq x_0 \end{cases}$$

and

$$\alpha_x = \begin{cases} s_\alpha(x) & x < x_0 \\ \log\left(\frac{\beta \exp(\mu + \mu_X x + \alpha + \alpha_x x)}{1 + \exp(\mu + \mu_X x + \alpha + \alpha_x x)}\right) - \log\left(\frac{\beta \exp(\mu + \mu_X x)}{1 + \exp(\mu + \mu_X x)}\right) & x \geq x_0 \end{cases}$$

also for both males and females. In this paper we only present the results using the logistic model for older ages.

We treat infant (age 0) mortality separately. Here, we exclude the period effect κ_t , and fit the model

$$m_{0t} = \exp(\mu_0 + \alpha_0 t + s_\gamma(t)) \quad (9)$$

where $s_\gamma(t)$ is the estimate of the cohort effect for $x = 0$ obtained from fitting (6) to the main body of data ($0 < x < x_0$). We have investigated the dependence of infant mortality rates on both the time and the cohort effects, estimated from the rest of the data. It transpired that infant mortality has a unique pattern of period variation and therefore the time effect was a very weak predictor for the infant rates. On the other hand, the infant mortality exhibits strong dependence on the cohort effect estimated from the rest of the data (see Figure 9).

3 Estimation of mortality rates

Dodd et al. (2018) suggest that for England and Wales mortality data, an optimal age (x_0) at which to make the transition from smooth to logistic model is 93 for males and 91 for females, based on 2010-2012 mortality data. We assume that these thresholds are fixed over time. This is a strong assumption and the transition age x_0 at each year might be included in the model as an unknown parameter. However, the added complexity required for different threshold ages may not be justified since our preliminary investigation shows negligible effect on mortality projections.

Figure 1 presents the maximum likelihood estimates of the parameters of model (1) under the Poisson distribution (black solid line) and negative binomial distribution (red solid line) for males aged between 1 and 92 years, using data for the period 1961-2013.

For model (1) to be considered as a credible alternative to the commonly used P-spline methodology for smoothing observed mortality rates, the fit of the model to the observed data should not be significantly worse. With regard to an assessment of model fit, Figure 2 presents the




Figure 1: Maximum likelihood estimates of the parameters of model (1) under the Poisson model (black line) and the negative binomial model (red line), data for males, 1961-2013.




Figure 2: Comparison of residuals, data for males, 1961-2013: the P-spline approach (left panel) and model (1) assuming the Poisson distribution (right panel). For each year and age group the residual is categorised according to its absolute value and plotted with a corresponding colour ranging from green (small residuals) through to red (large residuals).

square of Pearson residuals from the P-spline method, and from model (1) assuming the Poisson distribution.

It can be observed from Figure 2 that model (1) fits the data at least as well as the P-spline method. Indeed, by conventional goodness-of-fit measures (residual deviance), model (1) fits significantly better than the P-spline model, even allowing for its increased complexity in terms of the number of degrees of freedom required for parameter estimation. Model (1) seems to do a better job of estimating mortality in the age range 15-20 (at the start of the ‘accident hump’ related to external causes of death during early adulthood, especially for men). Both models have difficulty fitting the 1919 cohort (see Cairns et al., 2014), but arguably this cohort is of limited significance for population projection. Both models, however, fail to fit when assessed by conventional goodness-of-fit measures. Evidence for this is the large number of Pearson residuals with absolute value greater than 3. On the other hand, estimates which allow for overdispersion, either model (1), fitted by maximising a negative binomial likelihood, or a P-spline fitted by quasi-likelihood produce residuals within a much more acceptable range (see Figure 3). Therefore, we use the negative binomial model to estimate the model parameters for the rest of our analyses.




Figure 3: Comparison of residuals, data for males, 1961-2013: the P-spline approach allowing for overdispersion (left panel) and model (1) assuming the negative binomial distribution (right panel). For each year and age group the residual is categorised according to its absolute value and plotted with a corresponding colour ranging from green (small residuals) through to red (large residuals).

One advantage of the P-spline approach is that it provides estimates of mortality rates that vary smoothly over age and time, as illustrated in Figure 4, which accounts for overdispersion through the maximum Poisson quasi-likelihood estimation. Significant cohort effects, and cohorts with large annual mortality improvements can be also seen in this figure. Under this P-spline method, the projections will only depend on the most recent years and they would be largely insensitive to historical data.

For model (1), the maximum likelihood estimates of some of the model parameters, illustrated in Figure 1, do not vary smoothly and as a consequence the estimated mortality rates, presented in Figure 5, are also more irregular than would be desirable.




Figure 4: Heatmap of the fitted mortality improvements for the P-spline model allowing for overdispersion through a penalised quasi-likelihood method, data for males, 1961-2013. For each year and age group the estimated mortality improvement is categorised according to its absolute value and plotted with a corresponding colour ranging from green (large decrease) through to red (large increase).

To obtain smoother estimates we use model (6). Figure 6 displays the estimates for the resulting smooth model (6), superimposed over the corresponding estimates for model (2) on a mortality improvement scale. Not surprisingly, the estimates for model (6) are much more regular and have the desired smoothness, and the fitted mortality rates, displayed in Figure 7, are also smoother. There is an increase in residual deviance, but this is compensated by a corresponding decrease in the effective complexity of the model. Note also that the vertical strips correspond to the year




Figure 5: Heatmap of the fitted mortality improvements for model (1), data for males, 1961-2013. For each year and age group the estimated mortality improvement is categorised according to its absolute value and plotted with a corresponding colour ranging from green (large decrease) through to red (large increase).

effect, which we do not smooth.




Figure 6: Estimates of the parameters of model (6) (red lines), superimposed over the corresponding estimates for model (2) (black lines) on a mortality improvement scale, data for males, 1961-2013.

Figure 8 presents the combined estimates of the parameters of models (9), (6) and (8). Under the logistic model (8) mortality rates flatten off, converging to a limiting rate β as x tends to infinity. We estimate β as 1.48 for males and 0.99 for females – the values which we set as constant over time.

The observed and fitted infant mortality using model (9) are displayed in Figure 9. If the cohort effect is not considered in the model, the estimates of infant mortality rates would be a straight line (on the log scale). By borrowing information on cohort effect from the rest of the data by age, we can identify the cohort improvements in infant mortality rates, e.g. around 1990.

Finally, we investigate the robustness of the proposed methodology by exploring the sensitivity of the estimated mortality rates in a later year (2011) to changes in the data used to estimate the model. Two different approaches are taken. In the first, we compare the estimates of 2011 mortality rates and 2011-12 mortality improvements for model (6) fitted for ages $1 \leq x < x_0$




Figure 7: Heatmap of the fitted mortality improvements for model (6), data for males, 1961-2013. For each year and age group the estimated mortality improvement is categorised according to its absolute value and plotted with a corresponding colour ranging from green (large decrease) through to red (large increase).




Figure 8: Estimates of the parameters of models (9), (6) and (8) (blue, black and red lines, respectively), data for 1961-2013, for males (upper panels; $x_0 = 93$) and females (lower panels; $x_0 = 91$).




Figure 9: Estimates of infant mortality rates using model (9) (red lines), compared with observed rates (black lines), data for 1961-2013. (Left panel) males, (right panel) females.

by using data from 1961-2013, with the equivalent estimates fitted for 1971-2013 and 1981-2013; see Figure 10. Then, we compare the estimates of 2011 mortality rates and 2011-12 mortality obtained by using data from 1961-2011 with the equivalent estimates fitted to the 1961-2012 and 1961-2013 series; see Figure 11. In Figures 10, 11 and 12 we also provide a comparison with the P-spline approach.

Both in Figures 10 and 11 there is a big difference between our proposed model and the P-spline method in terms of the fit for 2011. Under the P-spline method there is a quite dramatic mortality improvement at around age 20, which we are not picking up in our model. This is because under the P-spline model the big mortality improvement at late teens in current years are projected forward as an age specific improvement, without taking the historical data into account (see Figure 4). Since our model takes the whole period into account instead of only the current years, we do not see such a dramatic improvement in mortality around age 20. This is the largest difference between the future mortality projections produced by the two models.




Figure 10: Estimated 2011 mortality rates (left panel), and 2011-12 mortality improvements (right panel) for males, using model (6) and different historical fitting periods.

In the proposed projection methodology, existing cohort components are included in projections, but the period effects (and any as yet unobserved cohorts) are projected as zero, which is consistent with the model. Hence, the mortality improvements for 2011-12 in Figures 10 and 11 ignore any estimated period effect for 2012, even where the 2012 data have been used in fitting. Our actual proposal would be to project forward from the final year of observed data (2013) in which case the base mortality rates and mortality improvements are presented in Figure 12.

Note that the scale of Figure 12 is different than the scale of Figure 10 and therefore the difference between different historical fitting periods is more obvious. The 1961-2013 or 1971-2013 datasets broadly show similar patterns. However, if we ignore the data from 1961 to 1980, we move to a regime where there are much bigger mortality improvements at age around 20. As a result, not surprisingly, when we lose almost 40% of the data we see some sensitivity in mortality improvements.




Figure 11: Estimated 2011 mortality rates (left panel), and 2011-12 mortality improvements (right panel) for males, using model (6) and different recent fitting periods.




Figure 12: Estimated 2013 mortality rates (left panel), and 2013-14 mortality improvements (right panel) for males, using model (6) and different historical fitting periods.

4 Projection of mortality rates

Based on the parameter estimates of a model such as (6), providing point projections over any future time horizon is straightforward. Such a projection only requires extrapolation of the time effects κ_t for future years t , and the cohort effects γ_{t-x} for future birth cohorts. Because the time effects κ_t are constrained to zero in the final observed year and the estimated κ_t series approximates a random walk with zero-mean increments (see Figure 6), then it is reasonable to forecast κ_t to be zero for future t . Uncertainty about these forecasts is incorporated by assuming normally distributed increments with variance, σ_κ^2 , estimated from the κ_t series. A similar argument suggests that it is reasonable to also set cohort effects for future cohorts to zero. We do not include uncertainty about future cohorts, as these cohorts are likely to have a negligible effect on population mortality over the forecast horizon.

The confidence intervals can be calculated using the standard deviation obtained from the covariance matrix of the estimated model parameters, the variance of the observed period effect and the additional variance for the expert opinion, if used. More precisely for the main model (6) we have

$$\text{Var}(\log \hat{m}_{xt}) = \text{Var}(s_\mu(x) + s_\alpha(x)t + s_\gamma(t-x)) + \sigma_\kappa^2, \quad (10)$$

where the first term on the right hand side is the variance of a linear function of GAM parameters which can be computed using standard output from GAM fitting in R. For the old-age model, applying the delta method (see e.g. Schervish (1995)), we have

$$\text{Var}(\log \hat{m}_{xt}) = \Delta_{xt}^T V \Delta_{xt} + \sigma_\kappa^2 \quad (11)$$

where V is the covariance matrix of the model parameters $(\beta, \mu, \mu_X, \alpha, \alpha_X)$ and

$$\Delta_{xt} = \begin{pmatrix} 1 \\ \delta_{xt} \\ x\delta_{xt} \\ t\delta_{xt} \\ xt\delta_{xt} \end{pmatrix}$$

and

$$\delta_{xt} = 1 - \frac{\exp(\mu + \mu_X x + (\alpha + \alpha_X x)t)}{1 + \exp(\mu + \mu_X x + (\alpha + \alpha_X x)t)}.$$

Where expert opinion is incorporated in the forecasts, it is also incorporated in the uncertainty with the expert uncertainty and parameter uncertainty being weighted correspondingly.

We present our 2025 and 2055 projections for males and females in Figure 13.

In Figures 14 and 15 we compare the projections for 2025 and 2055 to the respective values from the 2014-based national population projections of the ONS, which use the past and projected data from the period and cohort life tables, in a range of variants: principal, high, and low (ONS, 2015a,b). The projection methodology of the ONS is based on a P-spline model, and the technical details can be found in ONS (2016).

We understand that the discontinuity of the ONS rates after age around 110 is due to merging the estimated q_x for different constituent countries of the UK. There is some discrepancy between ONS projections and our projections, especially around the accident hump. As mentioned before this is because of the current high improvement rates that is projected forward under the P-spline method. For both males and females, from age around 60 onwards the projections are consistent with each other.

It has to be noted that the ONS projections are moderated by experts, whereas our proposed model is completely data driven. We believe that the use of expert knowledge is a useful tool




Figure 13: Mortality projections for 2025 (left panel) and 2055 (right panel) for males and females.




Figure 14: Comparison of 2025 mortality projections for males (left panel) and females (right panel) with ONS projections.




Figure 15: Comparison of 2055 mortality projections for males with ONS projections.

for moderating predictions provided by the model. A slightly modified version of the proposed approach can allow us to make full use of all available sources of information, including expert opinion.

The ONS projections assume a convergence in annual mortality improvement to a constant value over a fixed time horizon (currently 25 years) across all ages, for every cohort born after 1960. For older cohorts, the convergence is imposed on the series of cohort mortality improvements. Our modification remains in the spirit of the approach proposed earlier, with the age-specific mortality improvements (α_x) converging to a common, expert-specified value and the cohort effects converging to zero over a 25 year time horizon. We incorporate the convergence using the same weight function given in (12):

$$w_h = \begin{cases} 1 - 3 \left(\frac{h}{25}\right)^2 + 2 \left(\frac{h}{25}\right)^3 & \text{for } 0 \leq h \leq 25 \\ 0 & \text{for } h \geq 25 \end{cases} \quad (12)$$

where h represents the projection period. So, for example, the age-specific mortality improvements incorporating the experts in period h , $\hat{\alpha}_{x,h}^e$, becomes

$$\hat{\alpha}_{x,h}^e = \begin{cases} \alpha^e(1 - w_h) + \hat{\alpha}_x w_h & \text{for } 0 \leq h \leq 25 \\ \alpha^e & \text{for } h \geq 25 \end{cases} \quad (13)$$

Here $\hat{\alpha}_x$ is the estimated year-on-year improvement by age effect using (6), (8) and (9). In line with the ONS mortality assumptions, we assume the value of the target expert-based mortality improvement is 1.2%, independent of age and sex, i.e. $\alpha^e = -1.2\%$.

When projecting the mortality rates, one final adjustment is made to fix any divergence between male and female rates at very old ages (see Figure 13). This is because we do not believe that the male and female mortality rates will start to diverge at very high ages following steady convergence up to this point. This also applies where the divergence occur after the rate functions cross. Therefore starting from the age where the difference between male and female mortality rates starts to increase (if that occurs) we keep the difference between the male and female mortality rates at a constant value and we obtain weighted mortality rates using

$$m_{xt}^m = m_{xt}^m w_{xh}^l + \left(m_{xt}^f + \min(m_{xt}^m - m_{xt}^f) \right) (1 - w_{xh}^l)$$

and

$$m_{xt}^f = \left(m_{xt}^m - \min(m_{xt}^m - m_{xt}^f) \right) w_{xh}^l + m_{xt}^f (1 - w_{xh}^l)$$

where

$$w_{xh}^l = \frac{l_{xh}^m}{l_{xh}^m + l_{xh}^f}$$

and l_{xh} represent the expected number of survivors to exact age x in year h from a birth population of size $l_0 = 100,000$ and the weights are used as a proxy for survivorship probabilities at age x and any future year for males and females.

The comparison of 2025 and 2055 projections after these adjustments can be seen in Figure 16.




Figure 16: Mortality projections for 2025 (left panel) and 2055 (right panel) for males and females.

In Figures 17 and 18, we compare our 2025 and 2055 mortality projections, this time incorporating the expert opinion and other adjustments we mentioned, with the 2014-based ONS projections. These adjustments, especially incorporating the expert opinion, visibly decreases the difference between two sets of projections. Also, restricting continuous year-on-year improvements by age enables us to avoid implausible long term projections at ages where we estimate high mortality improvements from the model.

Finally, in order to investigate the forecast performance of the proposed model, we present the 2014-2017 forecasts against the out of sample outcomes obtained from ONS (2018). Note that we used mortality data between 1961 and 2013 to estimate the model parameters. In Figure 19 we present the mortality projections for males under the proposed model, the realised mortality rates for a 4-year forecast horizon and the 2014-based mortality projections of the ONS for the same horizon.

Although we have not implemented any formal tests of how well these forecasts subsequently performed, for the majority of the ages, the mortality projections under the proposed model adheres acceptably well to the realised mortality rates.




Figure 17: Comparison of 2025 mortality projections for males (left panel) and females (right panel) with ONS projections.




Figure 18: Comparison of 2055 mortality projections for males with ONS projections.





Figure 19: Comparison of out of sample mortality rates (+) for males with the mortality projections under the proposed model (-) and ONS projections (-).

5 Conclusion

In this paper we have developed a method for estimating and projecting mortality rates, which takes advantage of the ease with which a wide range of smooth and parametric models can routinely be fitted. We model mortality improvement using generalised additive models for ages where we have reliable data and a parametric model at older ages where the data is sparse. Our methodology is based on a smooth transition between a GAM at lower ages and a fully parametric model at higher ages. To obtain the estimates we use maximum likelihood method. The approach described in this paper provides a computationally straightforward way of estimating and projecting mortality rates across the whole age range, including older ages where data are sparse or non-existent. Furthermore, our approach allows uncertainty and expert opinion to be coherently incorporated.

Under a fully Bayesian estimation method all different sources of uncertainty – in data series, model parameters, including the choice of the model cut-off x_0 , as well as expert judgement – would be treated jointly in a coherent, fully probabilistic manner. However, this would come at a considerable expense in terms of computing effort. In contrast, our method is very simple, and is therefore computationally very cheap and easy to implement, as it requires no Markov chain Monte Carlo sampling and is based on pre-existing R functions. For that reason, the approach proposed in this paper offers an appealing alternative for implementing a sophisticated and robust analytical method in actuarial practice.

References

- Bennett, J. E., Li, G., Foreman, K., Best, N., Kontis, V., Pearson, C., Hambly, P. and Ezzati, M. (2015) The future of life expectancy and life expectancy inequalities in England and Wales: Bayesian spatiotemporal forecasting. *The Lancet*, **386**, 163 – 170.
- Börger, M. and Aleksic, M.-C. (2014) Coherent projections of age, period, and cohort dependent mortality improvements. In *Living to 100 Symposium. Orlando, Fla. January*, vol. 8, 2014.
- Brouhns, N., Denuit, M. and Vermunt, J. K. (2002) A Poisson log-bilinear regression approach to the construction of projected lifetables. *Insurance: Mathematics and Economics*, **31**, 373–393.
- Cairns, A. J., Blake, D. and Dowd, K. (2006) A two-factor model for stochastic mortality with parameter uncertainty: Theory and calibration. *Journal of Risk and Insurance*, **73**, 687–718.
- Cairns, A. J., Blake, D., Dowd, K., Coughlan, G. D., Epstein, D. and Khalaf-Allah, M. (2011a) Mortality density forecasts: An analysis of six stochastic mortality models. *Insurance: Mathematics and Economics*, **48**, 355–367.
- Cairns, A. J., Blake, D., Dowd, K., Coughlan, G. D., Epstein, D., Ong, A. and Balevich, I. (2009) A quantitative comparison of stochastic mortality models using data from England and Wales and the United States. *North American Actuarial Journal*, **13**, 1–35.
- Cairns, A. J., Blake, D., Dowd, K., Coughlan, G. D. and Khalaf-Allah, M. (2011b) Bayesian stochastic mortality modelling for two populations. *Astin Bulletin*, **41**, 29–59.
- CMI (2016) CMI Mortality Projections Model consultation – technical paper. Institute and Faculty of Actuaries.
- Currie, I. D., Durban, M. and Eilers, P. H. (2004) Smoothing and forecasting mortality rates. *Statistical modelling*, **4**, 279–298.

- Dodd, E., Forster, J. J., Bijak, J. and Smith, P. W. (2018) Smoothing mortality data: the english life tables, 2010–2012. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, **181**, 717–735.
- Haberman, S. and Renshaw, A. (2012) Parametric mortality improvement rate modelling and projecting. *Insurance: Mathematics and Economics*, **50**, 309–333.
- (2013) Modelling and projecting mortality improvement rates using a cohort perspective. *Insurance: Mathematics and Economics*, **53**, 150–168.
- Hilton, J., Dodd, E., Forster, J. J. and Smith, P. W. F. (2019) Projecting UK mortality using Bayesian generalised additive models. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **68(1)**, 29–49.
- Human Mortality Database (2019) Human Mortality Database. URL<http://www.mortality.org/cgi-bin/hmd>.
- Lee, R. D. and Carter, L. R. (1992) Modeling and forecasting US mortality. *Journal of the American statistical association*, **87**, 659–671.
- Li, J. S.-H., Hardy, M. and Tan, K. S. (2010) Developing mortality improvement formulas: The canadian insured lives case study. *North American Actuarial Journal*, **14**, 381–399.
- Li, J. S.-H. and Hardy, M. R. (2011) Measuring basis risk in longevity hedges. *North American Actuarial Journal*, **15**, 177–200.
- ONS (2015a) National population projections: 2014-based statistical bulletin. URL<https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationprojections/bulletins/nationalpopulationprojections/2015-10-29>.
- (2015b) Past and projected data from the period and cohort life tables: 2014-based, UK, 1981 to 2064. URL<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/lifeexpectancies/bulletins/pastandprojecteddatafromtheperiodandcohortlifetables/2014baseduk1981to2064>.
- ONS (2016) National Population Projections: 2014-based Reference Volume, Series PP2. *Tech. rep.*, Office for National Statistics. URL<https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationprojections/compendium/nationalpopulationprojections/2014basedreferencevolumeseriespp2>.
- ONS (2018) Population estimates and deaths by single year of age for england and wales and the uk, 1961 to 2017. URL<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/lifeexpectancies/adhocs/009189populationestimatesanddeathsbysingleyearofageforenglandandwalesandtheuk1961to2017>.
- Plat, R. (2011) One-year value-at-risk for longevity and mortality. *Insurance: Mathematics and Economics*, **49**, 462–470.
- Renshaw, A. and Haberman, S. (2003) Lee–Carter mortality forecasting: A parallel generalized linear modelling approach for England and Wales mortality projections. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **52**, 119–137.
- Renshaw, A. E. and Haberman, S. (2006) A cohort-based extension to the Lee–Carter model for mortality reduction factors. *Insurance: Mathematics and Economics*, **38**, 556–570.

- Richards, S. J., Currie, I. D., Kleinow, T. and Ritchie, G. P. (2019) A stochastic implementation of the apci model for mortality projections. *British Actuarial Journal*, **24**.
- Schervish, M. J. (1995) *Theory of Statistics*. Springer.
- Wood, S. (2006) *Generalized additive models: an introduction with R*. CRC press.