

Past and future global transformation of terrestrial ecosystems under climate change

Nolan, Connor¹, Jonathan T. Overpeck^{2,1}, Judy R.M. Allen³, Patricia M. Anderson⁴, Julio L. Betancourt⁵, Heather A. Binney⁶, Simon Brewer⁷, Mark B. Bush⁸, Brian M. Chase⁹, Rachid Cheddadi⁹, Morteza Djamali¹⁰, John Dodson¹¹, Mary E. Edwards^{6,12}, William D. Gosling¹³, Simon Haberle¹⁴, Sara C. Hotchkiss¹⁵, Brian Huntley³, Sarah J. Ivory¹⁶, A. Peter Kershaw¹⁷, Soo-Hyun Kim¹⁵, Claudio Latorre¹⁸, Michelle Leydet¹⁰, Anne-Marie Lézine¹⁹, Kam-Biu Liu²⁰, Yao Liu²¹, A.V. Lozhkin²², Matt S. McGlone²³, Robert A. Marchant²⁴, Arata Momohara²⁵, Patricio I. Moreno²⁶, Stefanie Müller²⁷, Bette L. Otto-Bliesner²⁸, Caiming Shen²⁹, Janelle Stevenson³⁰, Hikaru Takahara³¹, Pavel E. Tarasov²⁷, John Tipton³², Annie Vincens³³, Chengyu Weng³⁴, Qinghai Xu³⁵, Zhuo Zheng³⁶, & Stephen T. Jackson^{37*,1}.

¹University of Arizona, Department of Geosciences, Tucson, AZ 85721, USA

²University of Michigan, School for Environment and Sustainability, Ann Arbor, MI 48109, USA

³University of Durham, Department of Biosciences, Durham DH1 3LE, UK

⁴University of Washington, Department of Earth and Space Sciences, Seattle, WA 98195, USA

⁵U.S. Geological Survey, National Research Program, Reston, VA 20192, USA

⁶University of Southampton, Geography and Environment, Southampton SO17 1BJ, UK

⁷University of Utah, Department of Geography, Salt Lake City, UT 84112, USA

⁸Florida Institute of Technology, Department of Biological Sciences, Melbourne, FL 32901, USA

⁹ Centre National de la Recherche Scientifique, UMR 5554, Institut des Sciences de l'Evolution de Montpellier, Université Montpellier, Bat.22, CC061, Place Eugène Bataillon, 34095 Montpellier, cedex5, France

¹⁰ Aix Marseille Université, Avignon Université, CNRS, IRD, Institut Méditerranéen de Biodiversité et d'Ecologie, 13545 Aix-en Provence Cedex 04, France

¹¹University of New South Wales, Palaeontology, Geobiology and Earth Archives Research Centre (PANGAEA), Sydney NSW 2052, Australia and State Key Laboratory of Loess and Quaternary Geology, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an, 71002, Shaanxi, China

¹²College of Natural Sciences and Mathematics, University of Alaska Fairbanks, Fairbanks, AK 99775, USA

¹³University of Amsterdam, Institute for Biodiversity and Ecosystem Dynamics, P.O. Box 94248, 1090 GE Amsterdam, The Netherlands and The Open University, Department of Environment, Earth & Ecosystems, Walton Hall, Milton Keynes, MK7 6AA, UK

¹⁴Australian National University, Department of Archaeology and Natural History, Canberra, Australia

¹⁵University of Wisconsin, Department of Botany, Madison, WI 53706, USA

¹⁶Pennsylvania State University, Department of Geosciences, State College, PA, 16802, USA

¹⁷Monash University, School of Earth, Atmosphere, and Environment, Melbourne VIC 3800, Australia

¹⁸Pontificia Universidad Católica de Chile, Departamento de Ecología, Institute of Ecology and Biodiversity (IEB), Santiago, Chile

¹⁹Sorbonne Université, CNRS-IRD-MNHN, LOCEAN/IPSL Laboratory, 4 Place Jussieu, 75005 Paris, France

²⁰Louisiana State University, Department of Oceanography and Coastal Sciences, Baton Rouge, LA 70803, USA

²¹Northern Arizona University, School of Informatics, Computing, and Cyber Systems, Flagstaff, AZ 86011, USA

²²North-East Interdisciplinary Scientific Research Institute, Far East Branch Russian Academy of Sciences, Magadan 685000, Russia

- ²³Landcare Research, Lincoln 7640, New Zealand
- ²⁴University of York, York Institute for Tropical Ecosystems, Environment Department, York YO10 5NG, UK
- ²⁵Chiba University, Graduate School of Horticulture, Matsudo-shi Chiba 271-8510 Japan
- ²⁶Universidad de Chile, Departamento de Ciencias Ecológicas, IEB and (CR)2, Santiago, Chile
- ²⁷Freie Universität Berlin, Institute of Geological Sciences, D-12249 Berlin, Germany
- ²⁸National Center for Atmospheric Research, Climate and Global Dynamics Laboratory, Boulder, CO 80307, USA
- ²⁹Yunnan Normal University, Key Laboratory of Plateau Lake Ecology and Global Change, Kunming, Yunnan 650092, China
- ³⁰Australian National University, School of Culture, History, and Language, Canberra, Australia
- ³¹Kyoto Prefectural University, Graduate School of Life and Environmental Sciences, Kyoto, 606-8522, Japan
- ³²University of Arkansas, Department of Mathematical Sciences, Fayetteville, AR 72701, USA
- ³³Centre Européen de Recherche et d'Enseignement des Géosciences de l'Environnement (CEREGE), 13545 Aix-en-Provence cedex 4, France
- ³⁴Tongji University, School of Ocean and Earth Science, Shanghai, China
- ³⁵Hebei Normal University, Institute of Nihewan Archaeology and College of Resource and Environmental Sciences, Shijiazhuang 050024, China
- ³⁶Sun Yat-Sen University, School of Earth Science and Engineering, Guangdong Provincial Key Lab of Geodynamics and Geohazards, Guangzhou 510275, China
- ³⁷U.S. Geological Survey, Southwest Climate Adaptation Science Center, Tucson, AZ 85721, USA*Corresponding author at stjackson@usgs.gov

Abstract. Impacts of global climate change on terrestrial ecosystems are imperfectly constrained by ecosystem models and direct observations. Pervasive ecosystem transformations occurred in response to warming and associated climatic changes during the last glacial-to-interglacial transition, which was comparable in magnitude to warming projected for the next century under high-emission scenarios. We reviewed 594 published paleoecological records to examine compositional and structural changes in terrestrial vegetation since the last glacial period, and to project the magnitude of ecosystem transformations under alternative future emission scenarios. Our results indicate that terrestrial ecosystems are highly sensitive to temperature change, and suggest that without major reductions in greenhouse-gas emissions to the atmosphere, terrestrial ecosystems worldwide are at risk of major transformation, with accompanying disruption of ecosystem services and impacts on biodiversity.

Terrestrial-ecosystem function is governed largely by composition and physical structure of vegetation (1-3), and climate-change impacts on vegetation can potentially cause disruption of ecosystem services and loss of biodiversity (4,5). It is critical to assess the likely extent of ecosystem transformation as global greenhouse-gas (GHG) emissions increase (6), and to understand the full potential magnitude of impacts should current GHG emission rates continue unabated.

Ecosystem transformation generally involves replacement of dominant plant species or functional types by others, whether recruited locally or migrating from afar. Observations from around the globe indicate that current climate change may already be driving substantial changes in vegetation composition and structure (3). Ecosystem change is accelerated by mass mortality of incumbent dominants (7,8), and widespread dieback events and other large disturbances are already underway in many forests and woodlands (9-11), with further mortality events predicted under increasing temperatures and drought (3,9,10,12). Replacement of pre-disturbance dominants by other species and growth-forms has been widely documented (8,13,14). In addition, evidence is accumulating for geographic range-shifts in individual species, and climate change is interacting with invasive species, fire regimes, land-use, and CO₂ increase to drive vegetation changes in many regions (15,16).

Beyond observations of recent and ongoing change, models indicate ecosystem transformation under climate projections for the 21st Century. These include dynamic global vegetation models (3,17), species-distribution models (18), and comparison of the multivariate climate distance between biomes with that between modern and future climates (19). However, capacity for assessing the magnitude of ecosystem transformation under future climate scenarios is limited by the difficulty of evaluating model performance against empirical records, particularly when projected climate states are novel (19,20).

Paleoecological records of past ecological responses to climate change provide an independent means for gauging the sensitivity of ecosystems to climate change. High-precision time-series studies indicate that local and regional ecosystems can shift rapidly, within years to decades, under abrupt climate change (21-23), but sites with such detailed chronologies are scarce. Here we pursue an approach based on a global network of published, radiocarbon-dated paleoecological records of terrestrial vegetation composition and structure since the Last Glacial Maximum (LGM), ca. 21,000 yr BP (24). Most postglacial warming happened 16,000 -10,000 yr BP, although it commenced earlier in parts of the Southern Hemisphere (25,26). Global warming between the LGM and the early Holocene (10,000 yr BP) was on the order of 4-7° C, with more warming over land than oceans (26,27). These estimates are roughly comparable to the magnitude of warming the Earth is projected to undergo in the next 100- 150 years if GHG emissions are not reduced significantly (28). The magnitude of change in vegetation composition and structure since the last glacial period provides an index of the magnitude of ecosystem change that might be expected under warming of similar magnitude in the coming century (29). Although the rate of projected future global warming is at least an order of magnitude greater than that of the last glacial-to-interglacial transition (26), a glacial-to-modern comparison provides a conservative estimate of the extent of ecological transformation to which the planet will be committed under future climate scenarios.

We reviewed and evaluated paleoecological (pollen and macrofossil) records from 594 sites worldwide (fig. S1), all drawn from peer-reviewed published literature, to determine the magnitude of post-glacial vegetation change. We adopted an expert-judgment approach, in which paleoecologists with relevant regional experience compiled published records (table S1), reviewed the data, diagrams, and accompanying papers, and inferred the composition and

structure of the glacial-age and Holocene vegetation at each site (24). For the purposes of our analyses, we defined the last glacial period (LGP) as the interval between 21,000 and 14,000 yr BP. Although postglacial warming was underway in many regions by 16,000 yr BP (25), continental ice-sheets were still extensive 14,000 yr BP, and some climate regimes remained essentially ‘glacial’ in nature, particularly in the Northern Hemisphere (30). Extending the LGP window to 14,000 yr BP provides a larger array of records for the assessment, both in glaciated and unglaciated terrains, and renders our analysis more conservative (i.e., climatic and vegetation contrasts with the Holocene are likely to decrease between 21,000 and 14,000 yr BP).

For each record, experts were asked to classify the magnitude of compositional change and structural change since the LGP as large, moderate, or low, and to provide detailed justification for their judgment (24) (table S2). This placed all the diverse records into a common framework for comparison. For sites that experienced moderate to large ecological change, experts were also asked to assess the role of climate change (large, moderate, or none) in driving the observed vegetation change. For each of these four judgments, experts were asked to state their level of confidence as high, medium, or low. In assessing the role of climate change, experts were asked to focus specifically on whether climate change since the LGP was sufficient to drive the observed changes, acknowledging that other factors (e.g., human activity, post-glacial CO₂ increase, megafaunal dynamics) may have also played important roles. For sites with a long history of human land-use), experts used Holocene records predating widespread land clearance as a benchmark for comparison with the LGP records.

Our results indicate that the magnitude of past glacial-to-interglacial warming was sufficient at most locations across the globe to drive changes in vegetation composition that were moderate (26% of sites) to large (72%), as well as moderate (28%) to large (67%) structural changes (Fig. 1, table S3). These changes were particularly evident at mid- to high latitudes in the Northern Hemisphere, as well as in southern South America, tropical and temperate southern Africa, the Indo-Pacific region, Australia, Oceania, and New Zealand (Fig 1A). Compositional change at most sites in the Neotropics was moderate to large, but three sites showed little or no compositional change, all with medium to high confidence (fig. S2). Shifts in vegetation structure were also moderate to large at mid-to high-latitude sites, although a few sites showed low change (Fig. 1B). The Neotropics had nine sites with little or no structural change (Fig. 1B),

all with high-confidence assessments (fig. S2). These sites have been occupied by tropical forest ecosystems since the LGM, although most have undergone moderate to large compositional change (31,32). For nearly all sites that experienced moderate or large ecological change, climate change since the LGM was judged as sufficient to explain the observed changes with high confidence (table S4). Atmospheric CO₂ concentrations also increased from 190 to 280 ppm during the deglaciation, interacting with and in some cases modulating ecological responses to climate change. However, CO₂ changes alone cannot account for postglacial vegetational changes (Supplementary Text).

Independent of the expert-judgment process, we used the estimated anomaly in mean annual temperature between the LGM and the present (pre-industrial) as a proxy for the overall magnitude of climate change since the LGM (24). LGM temperature estimates were derived using an assimilated proxy-data/model integration (27). Low-change sites were largely concentrated in regions where the estimated temperature anomaly was relatively small (Fig. 1). To explore this relationship further, we plotted the frequency distribution of the difference between estimated LGM and present-day mean-annual temperatures for individual sites in each of the three ecological-response categories. Nearly all sites with low compositional change between the LGM and today are associated with small projected temperature anomalies (median 2.4°C), whereas sites with moderate to high compositional change have larger temperature anomalies (Fig. 2A). Results for structural changes are similar, although a greater number of sites with low structural change include larger temperature anomalies (Fig. 2B). This difference is not surprising, because compositional change in vegetation can occur without an accompanying change in vegetation structure (Fig. 1). Europe and eastern North America experienced unusually large temperature changes since the LGM, owing to depressed temperatures near the large ice sheets, and these regions show significant compositional and structural changes since the LGM. However, results from other parts of the globe indicate that widespread ecosystem changes were driven by much smaller temperature changes (fig. S3). We repeated our analysis using the TraCE-21ka model simulations (33,34), which yield lower magnitude of LGM-to-Holocene climate change (35); despite the potential conservative bias, results for compositional and structural change (fig. S4) were similar to those in Fig. 2. Temperature differences between LGM and the present were significantly greater for sites with

large ecological change than those with low to moderate change, using both paleoclimate estimates (27, 33) (table S2).

We also used our database of ecological change since the LGM to assess the global distribution of the probability of large compositional and structural change given GHG emission scenarios (Representative Concentrations Pathways (RCP) 2.6, 4.5, 6.0 and 8.5, each as simulated by the CCSM4 coupled climate model) (24,36). The range of LGM-to-present temperature changes (Fig. 2) overlaps with the range of temperature changes projected for the coming century under these scenarios (Fig. 3A; fig. S5). We quantified the relationship between temperature and ecological change using a logistic spline regression with ordered categories (37). We fit models for compositional and structural change using the temperature change since the LGM as the independent predictor variable. In both models, LGM-to-modern temperature change is a significant predictor of ecosystem change ($p < 0.001$). We then used these models to predict the risk of large change for the future range of projected global temperature changes (Fig. 3B) and to map the probability of large change under RCP 2.6 and RCP 8.5 (Fig. 3C-3F) at the end of the 21st century (see fig. S6 for RCP 4.5 and RCP 6.0). Under RCP 2.6 the probability of large compositional change is less than 45% over most of the globe (Fig. 3C) and the probability of large structural change is generally less than 30% (Fig. 3E). In contrast, under the business-as-usual emissions scenario, RCP 8.5, the probability of large compositional change and large structural change are both greater than 60% (Fig. 3D, 3F). Analyses using the TraCE-21ka model yielded similar patterns (fig. S7).

Our study uses a single variable, mean annual temperature, as a metric for the broader array of climatic changes that can drive vegetation change, and it compares vegetation and climate states separated by 10,000-20,000 years. Future climate change, like that in the past, will be multivariate, involving shifts in seasonal temperatures, seasonal precipitation, climate extremes, and variability regimes. As mean annual temperature increases, other ecologically important variables will change, often in complex or counterintuitive ways (20,38,39), and ecological responses will often be episodic or non-linear (8, 13-15). Although the temperature increases since the LGM provide crude analogs for ongoing and future climate changes – for example, boundary conditions and forcings are different now (26,40,41) – our results nevertheless provide concrete evidence that vegetation composition and structure are sensitive to

changes in mean annual temperature of the magnitudes forecast for the coming century, and that vegetation transformations will become increasingly extensive as temperatures increase. Under the RCP8.5 scenario, the rate of warming will be on the order of 65 times greater than the average warming during the last deglaciation (26). Furthermore, the warming between the LGP and the Holocene occurred within the range of previous glacial/interglacial temperatures, while projected future changes will exceed those experienced over the past 2 million years (26). Although many ecological processes (e.g., species migration, colonization, succession) will likely lag climate changes, ecosystem transformations will probably be accelerated by disturbance and mortality events, land use, and invasive species (7-15).

We therefore conclude that terrestrial vegetation over the entire planet is at substantial risk of major compositional and structural change in the absence of dramatically reduced GHG emissions. Much of this change could occur during the 21st century, especially where vegetation disturbance is accelerated or amplified by human impacts (7). Many emerging ecosystems will be novel in composition, structure, and function (42), and many will be ephemeral under sustained climate change; equilibrium states may not be attained until the 22nd Century or beyond. Compositional transformation will affect biodiversity, via disintegration and reorganization of communities, replacement of dominant or keystone species, pass-through effects on higher trophic levels, and ripple effects on species interactions (16,43). Structural transformation will have particularly large consequences for ecosystem services (4), including achievement of nature-based development solutions under the United Nations' Sustainable Development Goals (<https://sustainabledevelopment.un.org/sdgs>). Structural changes will also influence biodiversity, driving alterations in habitats and resources for species at higher trophic levels. Compositional and structural changes may also induce potentially large changes to carbon sources and sinks, as well as to atmospheric moisture recycling and other climate feedbacks. Our results suggest that impacts on planetary-scale biodiversity, ecological functioning, and ecosystem services increase substantially with increasing GHG emissions, particularly if warming exceeds that projected by the RCP2.6 emission scenario (i.e., 1.5°C).

1. F.S. Chapin III, P.A. Matson, H.A. Mooney. *Principles of Terrestrial Ecosystem Ecology*. (Springer, New York, NY, 2002).
2. S. Díaz et al., The plant traits that drive ecosystems: Evidence from three continents. *Journal of Vegetation Science* **15**, 295-304 (2004).
3. J. Settele *et al.*, “Terrestrial and inland water systems” in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, C.B. Field *et al.*, Eds. (Cambridge Univ. Press, 2014), pp 271-359.
4. Millennium Ecosystem Assessment. *Ecosystems and Human Well-Being: Synthesis*. (Island Press, 2005).
5. B.J. Cardinale *et al.*, Biodiversity loss and its impact on humanity. *Nature* **486**, 59-67 (2012).
6. *United Nations Framework Convention on Climate Change (UNFCCC)* (1992).
7. J.T. Overpeck, D. Rind, R. Goldberg, Climate-induced changes in forest disturbance and vegetation. *Nature* **343**, 51-53 (1990).
8. C.D. Allen, D.D. Breshears. Drought-induced shift of a forest-woodland ecotone: rapid landscape response to climate variation. *Proceedings of the National Academy of Sciences* **95**, 14839-14842 (1998).
9. C.D. Allen, D. D. Breshears, N.G. McDowell, On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* **6**, 1-55 (2015).
10. W.R.L. Anderegg, J.M. Kane, L.D.L. Anderegg, Consequences of widespread tree mortality triggered by drought and temperature stress. *Nature Climate Change* **3**, 30–36 (2013).

11. G.F. Midgley, W.J. Bond, Future of African terrestrial biodiversity and ecosystems under anthropogenic climate change. *Nature Climate Change* **5**, 823-829 (2015).
12. G.P. Asner *et al.*, Progressive forest canopy water loss during the 2012-2015 California drought. *Proceedings of the National Academy of Sciences* **113**, E249-E255 (2016).
13. J.F. Johnstone *et al.*, Changing disturbance regimes, ecological memory, and forest resilience. *Frontiers in Ecology and the Environment* **14**, 369-378 (2016).
14. C.H. Guiterman, E.Q. Margolis, C.D. Allen, D.A. Falk, T.W. Swetnam, Long-term persistence and fire resilience of oak shrubfields in dry conifer forests of northern New Mexico. *Ecosystems* DOI: 10.1007/s10021-017-0192-2 (2017).
15. C.I. Millar, N.L. Stephenson, Temperate forest health in an era of emerging megadisturbance. *Science* **349**, 823-826 (2015).
16. C. Parmesan, M.E. Hanley, Plants and climate change: complexities and surprises. *Annals of Botany* **116**, 849-864 (2015).
17. I.C. Prentice *et al.*, "Dynamic global vegetation modeling: quantifying terrestrial ecosystem responses to large-scale environmental change" in *Terrestrial Ecosystems in a Changing World*, J.G. Canadell, D.E. Pataki, L.F. Pitelka, Eds. (Springer, Heidelberg, 2007), pp. 175-192.
18. G.E. Rehfeldt, N.L. Crookston, M.V. Warwell, J.S. Evans, Empirical analyses of plant-climate relationships for the western United States. *International Journal of Plant Sciences* **167**, 1123-1150 (2006).
19. J.W. Williams, S.T. Jackson, J.E. Kutzbach, Projected distributions of novel and disappearing climates by 2100 AD. *Proceedings of the National Academy of Sciences* **104**, 5738-5742 (2007).

20. S.T. Jackson, J.W. Williams, Modern analogs in Quaternary paleoecology: here today, gone yesterday, gone tomorrow? *Annual Review of Earth and Planetary Sciences* **32**, 495-537 (2004).
21. J.R.M. Allen, W.A. Watts, B. Huntley, Weichselian palynostratigraphy, palaeovegetation and palaeoenvironment; the record from Lago Grande di Monticchio, southern Italy. *Quaternary International* **73/74**, 91-110 (2000).
22. J.W. Williams, D.M. Post, L.C. Cwynar, A.F. Lotter, A.J. Levesque, Rapid and widespread vegetation responses to past climate change in the North Atlantic. *Geology* **30**, 971-974 (2002).
23. A. Correa-Metrio *et al.*, Rapid climate change and no-analog vegetation in lowland Central America during the last 86,000 years. *Quaternary Science Reviews* **38**, 63-75 (2012).
24. Materials and methods are provided in the Supplementary Materials.
25. J.D. Shakun, A.E. Carlson, A global perspective on Last Glacial Maximum to Holocene climate change. *Quaternary Science Reviews* **29**, 1801-1816 (2010).
26. V. Masson-Delmotte *et al.*, “Information from Paleoclimate Archives”, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, T.F. Stocker *et al.*, Eds. (Cambridge Univ. Press, 2013). pp. 383-464.
27. J.D. Annan, J. C. Hargreaves, A new global reconstruction of temperature changes at the Last Glacial Maximum. *Climate of the Past* **9**, 367-376 (2013).
28. M. Collins *et al.*, 2013: “Long-term climate change: Projections, commitments and irreversibility” in *Climate Change 2013: The Physical Science Basis. Contribution of*

- Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, T.F. Stocker *et al.*, Eds. Cambridge University Press, 2013), pp. 1029-1136.
29. J. Guiot, W. Cramer. Climate change: the 2015 Paris Agreement thresholds and Mediterranean basin ecosystems. *Science* **354**, 465-468 (2016).
 30. P.U. Clark *et al.*, The last glacial maximum. *Science* **325**, 710-714 (2009).
 31. P.A. Colinvaux, P.E. De Oliveira, J.E. Moreno, M.C. Miller, M.B. Bush. A long pollen record from lowland Amazonia: Forest and cooling in glacial times. *Science* **274**, 85-88 (1996).
 32. M.B. Bush, P.E. De Oliveira, P.A. Colinvaux, M.C. Miller, E. Moreno. Amazonian paleoecological histories: one hill, three watersheds. *Palaeogeography Palaeoclimatology Palaeoecology* **214**, 359-393 (2004).
 33. Z. Liu *et al.*, Transient simulation of last deglaciation with a new mechanism for Bølling-Allerød warming. *Science* **325**, 310-314 (2009).
 34. F. He, “Simulating transient climate evolution of the last deglaciation with CSM3,” dissertation, University of Wisconsin, Madison, WI (2011)
 35. Z. Liu *et al.*, The Holocene temperature conundrum. *PNAS* **111**, E3501-E3505 (2014).
 36. G.A. Meehl *et al.*, Climate system response to external forcings and climate change projections in CCSM4. *Journal of Climate* **25**, 3661-3683 (2012).
 37. S.N. Wood, N. Pya, B. Saefken, Smoothing parameter and model selection for general smooth models. *Journal of the American Statistical Association*. <http://arxiv.org/abs/1511.03864> (2016)

38. S.T. Jackson, J.T. Overpeck, Responses of plant populations and communities to environmental changes of the Late Quaternary. *Paleobiology* **26** (Supplement), 194-220 (2000).
39. S.T. Jackson, J.L. Betancourt, R.K. Booth, S.T. Gray, Ecology and the ratchet of events: climate variability, niche dimensions, and species distributions. *Proceedings of the National Academy of Sciences* 106: 19685–19692 (2009).
40. D.L. Hartmann *et al.*, “Observations: Atmosphere and surface”, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, T.F. Stocker *et al.*, Eds. (Cambridge Univ. Press, 2013). pp.159-254. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
41. P.U. Clark *et al.*, 2016. Consequences of twenty-first-century policy for multi-millennial climate and sea-level change. *Nature Climate Change* **6**, 360-369 (2016).
42. R.J. Hobbs, E.S. Higgs, C. Hall, editors. *Novel Ecosystems: When and How Do We Intervene in the New Ecological World Order?* (Wiley-Blackwell, Chichester, 2013).
43. C. Bellard, C. Bertelsmeier, P. Leadley, W. Thuiller, F. Courchamp, Impacts of climate change on the future of biodiversity. *Ecology Letters* **15**, 365-377 (2012).
44. K. Fægri, J. Iversen. *Text-Book of Pollen Analysis. Third Revised Edition.* (Munksgaard, Copenhagen, 1975).
45. T. Webb III, J.H. McAndrews J.H. Corresponding patterns of contemporary pollen and vegetation in central North America. *Geological Society of America Memoir* **145**, 267-299 (1976).

46. T. Webb III, R.A. Lasenki, J.C. Bernabo. Sensing vegetational patterns with pollen data: choosing the data. *Ecology* **59**, 1151-1163 (1978).
47. H.J.B. Birks, H.H. Birks. *Quaternary Palaeoecology*. (University Park Press, Baltimore, MD, 1980).
48. I.C. Prentice. Pollen representation, source area, and basin size: toward a unified theory of pollen analysis. *Quaternary Research* **23**, 76-86 (1985).
49. B.H. Huntley & T. Webb, III. *Vegetation History*. (Kluwer, Dordrecht, The Netherlands, 1988).
50. S.T. Jackson, S.T. Pollen and spores in Quaternary lake sediments as sensors of vegetation composition: theoretical models and empirical evidence. *Sedimentation of Organic Particles*, A. Traverse, Ed. (Cambridge University Press, 1994), pp. 253-286.
51. S.T. Jackson, J.T. Overpeck, T. Webb III, S.E. Keatts, K.H. Anderson. 1997. Mapped plant macrofossil and pollen records of Late Quaternary vegetation change in eastern North America. *Quaternary Science Reviews* **16**, 1-70 (1997).
52. H.A. Binney *et al.* The distribution of late-Quaternary woody taxa in northern Eurasia: evidence from a new macrofossil database. *Quaternary Science Reviews* **28**, 2445-2464 (2009).
53. S. Sugita. Theory of quantitative reconstruction of vegetation I: pollen from large sites REVEALS regional vegetation composition. *Holocene* **17**, 229-241 (2007).
54. S. Sugita. Theory of quantitative reconstruction of vegetation II: All you need is LOVE. *Holocene* **17**, 243-257 (2007).

55. A. Dawson *et al.* Quantifying pollen-vegetation relationships to reconstruct forests using 19th-century forest composition and pollen data. *Quaternary Science Reviews* 137:156-175 (2016).
56. S. Brewer, S.T. Jackson, J.W. Williams. Paleoecoinformatics: applying geohistorical data to ecological questions. *Trends in Ecology and Evolution* **27**, 104-112 (2012).
57. J.W. Williams *et al.* 2018. The Neotoma Paleoecology Database: a multi-proxy, international community-curated data resource. *Quaternary Research* **89**,156-177 (2018).
58. S.T. Jackson. Representation of flora and vegetation in Quaternary fossil assemblages: known and unknown knowns and unknowns. *Quaternary Science Reviews* **49**, 1-15 (2012).
59. H.J.B. Birks, H.H. Birks, B. Ammann. The fourth dimension of vegetation. *Science* **354**, 412-413 (2016).
60. K.J. Edwards, R.M. Fyfe, S.T. Jackson. The first 100 years of pollen analysis. *Nature Plants* **3**, 17001 (2017). <http://www.nature.com/articles/nplants20171>.
61. H.H. Birks., “Plant macrofossils”, in *Tracking Environmental Change Using Lake Sediments. Developments in Paleoenvironmental Research, Vol 3*, J.P. Smol, H.J.B. Birks, W.M. Last, R.S. Bradley, K. Alverson, Eds. (Springer, Dordrecht, Netherlands, 2002), pp. 49-74.
62. H.H. Birks, “Plant macrofossil introduction”, in *Encyclopedia of Quaternary Science, Second Edition*, S.A. Elias, C.J. Mock, Eds. (Elsevier, Amsterdam, 2013), pp. 593-612.
63. S.T. Jackson, R.K. Booth, “Validation of pollen studies”, in *Encyclopaedia of Quaternary Sciences*, S.A. Elias, Ed. (Elsevier, Amsterdam, 2007), pp. 2413-2422.
64. S.T. Jackson *et al.*, Inferring local to regional changes in forest composition from Holocene macrofossils and pollen of a small lake in central Upper Michigan, USA. *Quaternary Science Reviews* **98**, 60-73 (2014).

65. J.L. Betancourt, T.R. Van Devender, P.S. Martin, Eds. *Packrat Middens: The Last 40,000 Years of Biotic Change*. (University of Arizona Press, Tucson, AZ, 1990).
66. Lyford, M.E., S.T. Jackson, S.T. Gray & R.J. Eddy. 2004. Validating the use of woodrat (*Neotoma*) middens for documenting natural invasions. *Journal of Biogeography* 31:333-342.
67. M.R. Lesser, S.T. Jackson. Reliability of macrofossils in woodrat (*Neotoma*) middens for detecting low-density tree populations. *Paleobiology* **37**, 603-615 (2011).
68. R.S. Bradley. *Paleoclimatology: Reconstructing Climates of the Quaternary, Third Edition*. (Academic Press, Oxford, UK, 2014).
69. H.E. Wright, Jr. *et al.*, Eds. *Global Climates Since the Last Glacial Maximum*. (University of Minnesota Press, Minneapolis, 1993).
70. J. Kutzbach *et al.* Climate and biome simulations for the past 21,000 years. *Quaternary Science Reviews*, **17**, 473-506 (1998).
71. MARGO Project Members, Constraints on the magnitude and patterns of ocean cooling at the Last Glacial Maximum. *Nature Geoscience* **2**, 127-132 (2009).
72. J.D. Shakun *et al.*, Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation. *Nature* **484**, 49-54 (2012).
73. Bartlein, P. J., *et al.*, Pollen-based continental climate reconstructions at 6 and 21 ka: a global synthesis. *Climate Dynamics* **37**, 775-802 (2011).
74. Stocker, Thomas F., *et al.*, "Technical summary." *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. T.F. Stocker *et al.*, Eds. (Cambridge Univ.

Press, 2013). pp.33-115. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

75. S.N. Wood, Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society (B)* **73**, 3-36 (2011).
76. W.J. Bond, G.F. Midgley. A proposed CO₂-controlled mechanism of woody plant invasion in grasslands and savannas. *Global Change Biology* **6**, 865-869 (2000).
77. F.J. Bragg *et al.* Stable isotope and modelling evidence for CO₂ as a driver of glacial–interglacial vegetation shifts in southern Africa. *Biogeosciences* **10**, 2001-2010 (2013).
78. G.D. Farquhar, J.R. Ehleringer, K.T. Hubick. Carbon isotope discrimination and photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology* **40**, 503-537 (1989).
79. G.D. Farquhar, K.T. Hubick, A.G. Condon, R.A. Richards. “Carbon isotope fractionation and plant water-use efficiency”, in *Stable Isotopes in Ecological Research*, P.W. Rundel, J.R. Ehleringer, K.A. Nagy, Eds., (Springer New York, NY, 1989), pp. 21-40.
80. H.W. Polley. Implications of rising atmospheric carbon dioxide concentration for rangelands. *Journal of Range Management* **50**, 562-577 (1997).
81. J.R. Petit *et al.* Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica. *Nature* **399**, 429-436 (1999).
82. E. Monnin *et al.* Atmospheric CO₂ concentrations over the last glacial termination. *Science* **291**, 112-114 (2001).

83. T.R. van Devender, J.L. Betancourt, M. Wimberly. Biogeographic implications of a packrat midden sequence from the Sacramento Mountains, south-central New Mexico. *Quaternary Research* **22**, 344-360 (1984).
84. Y. Garcin, A. Vincens, D. Williamson, J. Guiot, Wet phases in tropical southern Africa during the last glacial period. *Geophysical Research Letters* **33**, L07703 (2006.)
85. C.A. Holmgren *et al.* Evidence against a Pleistocene desert refugium in the Lower Colorado River Basin. *Journal of Biogeography* **41**, 1769-1780 (2014).
86. S. Lim, B.M. Chase, M. Chevalier, P.J. Reimer, 50,000 years of vegetation and climate change in the southern Namib Desert, Pella, South Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* **451**, 197-209 (2016).
87. B.M. Chase, M. Chevalier, A. Boom, A.S. Carr, The dynamic relationship between temperate and tropical circulation systems across South Africa since the last glacial maximum. *Quaternary Science Reviews* **174**, 54—62 (2017).
88. S.I. Higgins, S. Scheiter. Atmospheric CO₂ forces abrupt vegetation shifts locally, but not globally. *Nature* **488**, 209 (2012).
89. T.M. Shanahan *et al.* CO₂ and fire influence tropical ecosystem stability in response to climate change. *Scientific Reports* **6**, 29587 (2016).
90. K. Izumi, A.M. Lézine, Pollen-based biome reconstructions over the past 18,000 years and atmospheric CO₂ impacts on vegetation in equatorial mountains of Africa. *Quaternary Science Reviews* **152**, 93-103.

91. B. Bereiter *et al.*, Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before present. *Geophysical Research Letters* **42**, 542-549 (2015).
92. A. Staal, S.C. Dekker, M. Hirota, E.H. van Nes, Synergistic effects of drought and deforestation on the resilience of the south-eastern Amazon rainforest. *Ecological Complexity* **22**, 65-75 (2015).
93. D.H. Urrego, B.A. Niccum, C.F. La Drew, M.R. Silman, M.B. Bush, Fire and drought as drivers of early Holocene tree line changes in the Peruvian Andes. *Journal of Quaternary Science* **26**, 28-36 (2011).
94. B.G.Valencia, D.H. Urrego, M.R. Silman, M.B. Bush, From ice age to modern: a record of landscape change in an Andean cloud forest. *Journal of Biogeography* **37**, 1637-1647 (2010).

ACKNOWLEDGMENTS

The paper benefitted from thoughtful comments of S.T. Gray and three anonymous reviewers.

Funding: This research was supported by the National Science Foundation (DEB-1241851, AGS-1243125, EAR-1304083) and by the Department of the Interior Southwest Climate Adaptation Science Center. Research in northeast Siberia was funded by Russian Academy of Sciences, FEB (15-I-2-067) and the Russian Foundation for Fundamental Research (15-05-06420). **Author Contributions:** STJ, CN, and JTO designed project; all authors collected data; CN analyzed data with advice from JTO, STJ, SB, and JT; STJ, CN, and JTO wrote paper with text contributions in Supplementary Materials from BMC, MBB, MEE, JLB, BH, YL, and SI, and further contributions from all authors. **Competing Interests:** The authors declare no competing interests. **Data and Materials:** All data are available in the main paper or the supplementary materials.




Fig. 1. Vegetation differences between the last glacial period and the present. Each square represents an individual paleoecological site. Color density indicates the magnitude of estimated vegetation change since the last glacial period (21,000- 14,000 yr BP). Background shading denotes the estimated temperature anomaly between the last glacial maximum (LGM) 21,000 years ago and today, based on assimilated proxy-data and model estimates (27). **(A)** Composition. **(B)** Structure.




Fig. 2. Estimated temperature differences for different categories of vegetation response.

Box plots of the estimated mean-annual temperature differences between the last glacial maximum and today in each of the three vegetation-change categories (low, moderate, large), for (A) composition and (B) structure. Low vegetational changes are associated largely with relatively small temperature anomalies, while moderate and large changes are associated with larger post-LGM temperature differences, indicating that magnitude of temperature change plays an important role in the magnitude of vegetation change. The glacial temperature anomalies are based on (27). Analyses using the TraCE-21ka simulation show similar patterns (fig. S4).




Fig. 3. Estimated vegetation change under future-climate scenarios. (A) Box plots of the estimated mean-annual temperature differences between today and future-climate simulations for individual sites (using the nearest grid-point). Most sites show relatively small temperature change under the low-emissions scenario (RCP2.6), with substantially higher change under the high-emission scenario (RCP8.5). (B) Probability of large change in vegetation composition (orange) and structure (green) as a function of temperature change. (C-F) Estimated probability of large compositional and structural change by the end of the 21st century (average of 2081-2100) under RCP 2.6 (C,E) and RCP 8.5 (D,F). Probabilities (B-F) are estimated from a logistic-spline

regression model fit using LGM-to-modern temperature change as a predictor variable and observed LGM-to-modern vegetation changes (large *versus* not-large) as the response variable. Future temperature increases are calculated as an average of 2081-2100 under the model scenarios, minus an average of 1985-2005 from the CCSM4 historical simulation. Analyses using the TraCE-21ka simulation show similar patterns (fig. S7).

SUPPLEMENTARY MATERIALS

Materials and Methods

Supplementary Text

Figs. S1 to S8

Tables S1 to S4

Instructions for Assessors

Captions for Additional Data Table S1

Additional Data Table S1

References (44-94)



Supplementary Materials for

Past and future global transformation of terrestrial ecosystems under climate change

Nolan, Connor¹, Jonathan T. Overpeck^{2,1}, Judy R.M. Allen³, Patricia M. Anderson⁴, Julio L. Betancourt⁵, Heather A. Binney⁶, Simon Brewer⁷, Mark B. Bush⁸, Brian M. Chase⁹, Rachid Cheddadi⁹, Morteza Djamali¹⁰, John Dodson¹¹, Mary E. Edwards^{6,12}, William D. Gosling¹³, Simon Haberle¹⁴, Sara C. Hotchkiss¹⁵, Brian Huntley³, Sarah J. Ivory¹⁶, A. Peter Kershaw¹⁷, Soo-Hyun Kim¹⁵, Claudio Latorre¹⁸, Michelle Leydet¹⁰, Anne-Marie Lézine¹⁹, Kam-Biu Liu²⁰, Yao Liu²¹, A.V. Lozhkin²², Matt S. McGlone²³, Robert A. Marchant²⁴, Arata Momohara²⁵, Patricio I. Moreno²⁶, Stefanie Müller²⁷, Bette L. Otto-Bliesner²⁸, Caiming Shen²⁹, Janelle Stevenson³⁰, Hikaru Takahara³¹, Pavel E. Tarasov²⁷, John Tipton³², Annie Vincens³³, Chengyu Weng³⁴, Qinghai Xu³⁵, Zhuo Zheng³⁶, & Stephen T. Jackson^{37*,1}.

correspondence to: stjackson@usgs.gov

This PDF file includes:

Materials and Methods

Supplementary Text

Figs. S1 to S8

Tables S1 to S4

Instructions for assessors

Captions for supplementary data table S1

Other Supplementary Materials for this manuscript includes the following:

Additional Data Table S1

Materials and Methods

Paleoecological Inference and Expert Judgment

Sources of paleoecological inference concerning terrestrial ecosystem properties are diverse, and vary in level of detail, quality, and geographic coverage. For this study, we used pollen and plant macrofossil assemblages, which are widely applied, robust proxies that provide direct evidence for composition and structure of terrestrial vegetation. Our use of fossil pollen, plant macrofossils, or both, collectively provides broad geographic coverage within and among the vegetated continents and islands. Here, we discuss the characteristics of the paleoecological data we use, and justify our application of a transparent process of expert judgment that builds on more than a century of community experience in paleoecological inference and causal attribution. We describe specific procedures and protocols in the following section.

Pollen and plant macrofossil data derive from a wide variety of depositional settings, ranging from sediments of lakes and wetlands to land-surfaces buried under glacial and fluvial deposits to rodent middens stashed in caves. Pollen data can usually be expressed in a single numerical currency (percentages of a terrestrial taxon sum), but pollen percentages are influenced by site-specific processes (dependent on basin size, depositional environment, local vegetation, and regional topography, among others), and by differential pollen productivity and dispersal among taxa (44-50). Pollen accumulation rates and plant macrofossils, when available, can help inform and constrain interpretation of pollen percentages. Plant macrofossil data themselves lack a single universal currency beyond 'presence', so numerical comparisons among assemblages or sites are difficult and rare (51,52). No formal algorithm currently allows incorporation of information from macrofossils, pollen accumulation rates, or depositional context into interpretation of pollen percentages, and many macrofossil assemblages, particularly from buried land surfaces and rodent middens, lack associated pollen data. Numerical methods and formal tools have been developed to support vegetational inferences from paleoecological data, particularly pollen percentages (20,48,49,53-55). However, most of these tools are best suited for relatively uniform data and site types, and most require calibration or reference data sets from comparable modern sites. Such reference data sets do not exist in many regions. Furthermore, although the development of global databases for paleoecological inference is rapidly advancing (56,57), raw data for many sites and some entire regions remain unavailable, further hampering global application of numerical analyses.

The sole universal approach for inferring past vegetation composition and structure from these diverse data types remains the informed judgment of experts with relevant training and experience.

Such expert judgment involves examining the available data with a particular question in mind, and weighing the evidence in context of broader knowledge, experience, and relevant literature to arrive at a conclusion. Although often aided by formal numerical analyses, expert judgment is also informed by site-specific knowledge (basin morphometry, depositional environment, relevant taphonomic processes), region-specific knowledge (flora, vegetation, topography, comparative modern pollen/vegetation datasets), taxon-specific knowledge (taphonomic and assemblage-representation properties), and general knowledge of the vegetation-sensing properties of fossil assemblages, including potential biases and distortions (50,58).

Attainment of expertise in paleoecology involves immersion in understanding the processes that intervene between source vegetation and fossil assemblages, in mastering the literature of paleoecological interpretation, and in identifying the idiosyncracies of individual sites and regions. Interpretation of fossil pollen percentages is based on a century of intensive study (20,49,59,60), and paleoecologists draw on a rich literature of theoretical and empirical studies to guide their interpretations (44-50,53-55,58). Interpretation of plant macrofossil assemblages from lakes, peatlands, and other 'wet' sites is also founded on theoretical and empirical understanding of the physical processes by which these assemblages are formed and what they do and do not record (51,52,61-64). Plant macrofossils in rodent (*Neotoma* and other genera) middens represent a particular case in which the behavior and foraging preferences of the animals need to be considered (65-67), along with other factors (58).

In the current study, scientists with experience in the various regions and data sources reviewed each record and accompanying literature, and provided their best judgments concerning the degree of vegetational change between LGP and today for each site and their confidence in those judgments (see below). To ensure transparency and consistency, each expert also provided explicit justification for each assessment they made, noting the specific features of the LGP pollen or macrofossil assemblages, contrasting them with modern assemblages or modern vegetation, and identifying potential sources of uncertainty (e.g., taxonomic smoothing, depositional or dispersal processes). Judgments of the type used in this study are thus the same as those incorporated into the discussions and conclusions of virtually every paper in the paleoecological literature. Those judgments are often supported by numerical or other analytical tools, but they ultimately draw on skills and knowledge built on a century of paleoecological inquiry. Such informed judgment is at the heart of scientific inference, and applying it in our study allows us to maximize global coverage, using multiple data sources from diverse depositional settings, drawing on expertise from every region of the globe for which suitable pollen or macrofossil data exist. The judgments convert the diverse array of sites and data into a single, uniform currency.

Compilation of Last Glacial Period (LGP) paleoecological records

Teams of regional experts (Table S1) compiled 594 published paleoecological records (pollen, plant macrofossil, or both) from all vegetated continents and oceanic islands with data within the period 21,000-14,000 yr BP (Figure S1, Additional Data table S1), which we operationally define as the Last Glacial Period (LGP). The sole criterion for inclusion was that a site have pollen or macrofossil data (or both) reliably dating to the period between 21,000 and 14,000 yr BP.

Assessment of ecological change

Regional experts compared LGP records from each site with ‘modern’ vegetation or palaeoecological data. They examined the data or diagrams directly, and where necessary or relevant, consulted original publications, using text discussions and multivariate analyses (clustering, ordination, zonation) as information sources to infer vegetation composition and structure. Magnitude of vegetational change in terms of composition and structure from LGP to ‘modern’ was classified as large, moderate, or low for each site based on the definitions of those terms in Table S2. This classification summarizes the paleoecological inferences and judgments described above. For each assessment, the assessors provided a justification that describes the data and inferences underlying the assessed change (“CompJustification” and “StructureJustification” in Additional Data Table S1).

For most sites (504 sites, 85% of total), the ‘modern’ benchmark for assessing change was vegetational or paleoecological (pollen, macrofossil) data from the past two centuries. In some cases, human land-use or lack of suitable paleoecological data required an earlier benchmark. The late Holocene (past 2000-3000 yr) was used for 57 sites (10% of total), and 33 sites (5% of total) used paleoecological data from the early or mid-Holocene. Typically, benchmark data from the same paleoecological site were used in the comparison, but experts also employed their knowledge of modern or natural vegetation of the respective region. If paleoecological data from a site did not extend into the Holocene, experts often used Holocene paleoecological data from other, comparable sites in the same region. No relationship was observed between the assessment of ecological change and the temporal benchmark used; each of the benchmarks used (modern, late Holocene, early-mid-Holocene) yielded a mix of low, moderate, and large changes in vegetation composition and structure.

The degree of confidence in the assessments of ecological change was rated as high, medium, or low. All assessments were based on the definitions in Table S2 and the instructions in the Instructions for Assessors (see Other Supplementary Materials). Authors provided succinct but informative justifications for each individual assessment of compositional and structural change, sufficient to allow a paleoecologically literate reader to understand the empirical basis for their assignment. A complete list of sites with accompanying metadata, assessments, and justifications, is provided in Additional Data table S1.

Attribution of Vegetational Change

Climate change is among many mechanisms potentially capable of driving vegetational change. For each site where ecological changes were assessed as moderate or large, the role of climate in driving those changes was also assessed (large, moderate, or none). For this decision, expert judgment drew from multiple considerations, including understanding of modern ecosystems and communities and their controls, independent paleoclimatic evidence from multiple proxies (paleontological, geochemical, lithological) deriving from multiple sources (ice-cores, marine sediments, lake sediments, paleosols, glacial deposits, fossil groundwater, boreholes) (68) and climate-model simulations (69,70). The inclusion of ‘none’ as a choice, although rarely used, was to ensure that experts considered alternatives to climate as a primary driver (including succession, neutral assembly, ecological drift, human activities, soil development, biogeochemical changes, migration, and evolution). The degree of confidence in assessments of the role of climate was rated as high, medium, or low.

Estimates of temperature change since LGP

Site-level changes in mean annual temperature since LGM were estimated based on two different datasets: the model/proxy integration of Annan and Hargreaves (2013) (27) and the TraCE-21ka transient climate model run (33). For each dataset the estimate of temperature change for the single closest grid box to each site was used. The distribution of temperature changes is shown in box plots, in subsets according to degree of assessed ecological change (figure 2, figure S4).

The main paleoclimate dataset used is the last glacial maximum (LGM) temperature anomaly reconstruction from Annan and Hargreaves (2013) (27). This reconstruction uses fully coupled GCMs from the PMIP3/CMIP5 set of models run with LGM boundary conditions and proxy data in a multiple linear regression framework to obtain an estimate of the surface air-temperature anomaly from LGM to pre-industrial on a $2.5^\circ \times 2.5^\circ$ grid. The proxy data include marine sea-surface temperature reconstructions from Multiproxy Approach to Reconstructing Global Oceans (71), the multiproxy deglaciation synthesis (72), and pollen-derived LGM temperatures from selected continental sites (73).

Estimated LGM-to-preindustrial temperature change based solely on the models is $3.1\text{--}5.9^\circ\text{C}$ (27). With the data constraints, the final resulting LGM cooling estimate relative to preindustrial temperatures is $4.0 \pm 0.8^\circ\text{C}$. Application of the proxy data in this framework reduced uncertainty by arbitrating disagreements between the nine individual models and by improving spatial skill (27). The influence of proxy data on the mean estimates of temperature change between LGM and the preindustrial period was very small (27).

As an independent check on the temperature estimates and analyses, we also used the TraCE-21ka (simulation of the transient climate evolution over the last 21,000 years) (33,34), a 21,000 year run of the National Center for Atmospheric Research’s Community Climate System Model version 3 (CCSM3) with time-evolving forcing including atmospheric CO_2 concentrations, insolation, and land ice sheets. This is the only fully-coupled, moderately high-resolution transient simulation of the LGM to Holocene climate evolution. We estimated the LGM to late Holocene temperature change from TraCE by subtracting the average temperature for 0-6ka from the average temperature from 14-21ka. These large windows were chosen with

the intention of smoothing out noise in the transient simulation and getting a coarse estimate of temperature change at each grid point. The estimate of LGM-to-late-Holocene temperature change from TraCE at the grid point nearest to each paleoecological site is plotted in figure S4, an expanded version of Figure 2.

The boxplots suggest a relationship between the degree of ecosystem change and temperature change. We tested for statistical significance of this relationship using the Wilcoxon rank-sum test. The temperature change experienced by sites with Large ecological change was statistically significantly different from that at sites with Moderate and Low ecological change (table S3). The temperature differences between sites with Moderate versus Low ecological change were not statistically significant. Results were similar for the paleo-temperature estimates from the Annan and Hargreaves reconstructions and from TraCE (Table S3).

Projections of future temperature change

For each LGP paleoecological site we also obtained estimates for the projected future mean annual temperature change under various emissions scenarios. We used the difference between the average temperature for the last 20 years (2081-2100) of the future simulations under four different representative concentration pathway (RCP) scenarios and the average temperature for the last 20 years of the historical simulations (1985-2005) at the nearest grid point using simulations of the Community Climate System Model version 4 (CCSM4) run for CMIP5 and included in IPCC AR5 (36,74). All four RCPs – RCP 8.5, 6.0, 4.5, and 2.6 (also known as RCP 3.0-PD) – were used in our analysis.

These RCPs represent net radiative forcing of 8.5, 6.0, 4.5, and 2.6 W/m² at the end of the 21st century (36). RCP 8.5 is a “business-as-usual” scenario that ends up with ~1370 ppm CO₂ equivalent and global average temperature increase of approximately 5°C. RCP 6.0 is a stabilization scenario where radiative forcing is stabilized shortly after 2100. The CO₂ equivalent at 2100 is 936 ppm and the resulting global average temperature increase is 3-4°C. The RCP 4.5 and 2.6 scenarios represent aggressive emissions cutting. RCP 4.5 results in 650 ppm CO₂ equivalent and a global average temperature increase of 2.4°C by 2100. This scenario represents aggressive emission stabilization without overshoot. The most aggressive mitigation pathway is RCP 2.6. This scenario results in 490 ppm CO₂ equivalent and a global average temperature increase of 1.5°C at 2100. The emissions in this scenario peak at a radiative forcing of 3.0 W/m² in the mid-21st century and decline to 2.6 W/m² by 2100 (36).

The distributions of projected temperature change at our LGP paleoecological sites are shown as box plots in main text figure 3A and figure S5.

Estimating risk of future ecological change

We used our database of LGM-to-modern ecological change and associated temperature change to project the risk of future ecosystem transformation under different emission scenarios (see above). We developed a regression model between temperature change and ecological change using a logistic spline regression, also known as a generalized additive model (GAM). These models were fit in R using

the mgcv package (37,75). We fit one model for compositional change and one for structural change. For both models the independent variable was the temperature change from the LGM to present at the closest grid box to each site from (27) (Fig. 3) or TraCE-21ka (fig. S7). Ecological change served as the dependent variable. We used the ordered categorical family of GAMs where Small/No change is coded as 1, Moderate change is coded as 2, and Large change is coded as 3. The smooth terms of both models are highly significant ($p < 0.001$).

We used these regression models to predict the probability of large, moderate, and low compositional and structural change for the full range of LGM-to-present temperature changes (0-30°C) and for the full range of future projected temperature changes (0-12°C) (fig. S7). The results in figure S7 are from the same GAM plotted on different x-axes. We also mapped these probabilities spatially for all of the representative concentration pathways (Figure 3, fig. S6).

Supplementary Text

Confidence by region

In fig. S2 we plot the confidence (low, medium, high) for assessment of compositional and structural change for the entire data set and for each individual region. Confidence for large compositional and structural change was high for nearly all regions; confidence was lower for sites assessed as having moderate and (especially) low change. The lower confidence associated with moderate to low change derived largely from poor taxonomic resolution in pollen assemblages, and from uncertainties associated with long-distance pollen transport, which may respectively mask vegetational changes and confound interpretations (44-50). These pollen considerations also resulted in moderate confidence for many high-latitude Northern Hemisphere pollen sites (Northern Eurasia, North America). (In unforested vegetation (particularly tundra, but also some grassland and steppe) low pollen productivity (low NPP, prevalence of insect-pollinated plants with poor pollen dispersal) renders pollen assemblages susceptible to high relative influx of pollen (particularly tree pollen) from distant sources. Vegetation in such regions is also often dominated by grasses (Poaceae), sedges (Cyperaceae), composites (Asteraceae), chenopods (Chenopodiaceae), and other taxa that cannot be differentiated palynologically below the family level.)

Many North American sites were assessed as having medium confidence for structural change (in contrast to high confidence for compositional change); most of these were rodent-midden sites from the western United States. Middens primarily sample floristic composition, not relative or absolute abundance, and are thus more reliable indicators of vegetation composition than structure (58,66).

Temperature change by region

In fig. S3 we plot the frequency distributions of the estimated mean-annual temperature differences between the last glacial maximum and today for each region of the globe (27), in each of the three LGM-to-modern vegetation-change categories (low, moderate, large) for composition and structure. Although some regions (North America, Europe, New Zealand, Japan) have very high sampling density (figs. S1, S2), they do not dominate the patterns observed in Figure 2. Furthermore, although regions in proximity to continental ice sheets (North America, Europe) have many sites with extreme temperature anomalies ($>10^{\circ}\text{C}$) between LGM and today, large vegetation changes are associated at many sites (typically further from ice sheets) with smaller temperature anomalies. Observed vegetational changes at sites near ice sheets may have occurred with much smaller temperature differences.

Comparing estimates of LGM-to-modern temperature change

In fig. S4 we plot the frequency distributions of the mean-annual temperature change from LGM to present vs. vegetation change using two different products for the temperature change: Annan and Hargreaves 2013 (27) and TraCE-21ka (33), as described above.

The Annan and Hargreaves estimate may overestimate the temperature change required to drive the observed ecological changes, because not all the paleoecological records extend to LGM. Our temperature-change estimates using TraCE (LGP to late Holocene) provide an independent check, with a bias opposite that of Annan and Hargreaves 2013. TraCE may underestimate the temperature change necessary to drive the observed vegetation change because of its cold Holocene bias (35) and because of the broad time ranges used to estimate the temperature change (particularly including the post-LGM warming trend observed in many regions between 21,000 and 14,000 yr BP). TraCE is based solely on climate model forcings and physics, and thus provides independent validation of our inferences, free of any possible bias from pollen and other paleoclimate proxy data. (As noted above, such bias is small even for the Annan and Hargreaves 2013 dataset.)

The range of temperature estimates and the general pattern in which greater vegetation change is associated with higher temperature change are consistent between the two LGM temperature products (fig. S4; table S3)). We repeated the GAM modeling of ecological change as a function of estimated temperature change using the TraCE temperature-change estimates. Results were consistent with those obtained using the Annan & Hargreaves 2013 estimates (fig. S7).

Role of climate and confidence

For all cases in which compositional or structural change were judged to be moderate or large, assessors were asked to judge the role of climate in driving the observed changes, as well as their confidence in that judgment (table S4). Climate was assessed with high confidence to have played a large role in the observed changes in composition and structure for 85% of the sites, and was judged as having a large role with medium or high confidence for 88% of sites. Climate was assessed to have played a low role in driving ecological change at less than 1% of sites (6 out of 582 for composition and 1 out of 564 for structure).

Role of Carbon Dioxide in Forcing Vegetational Changes

Some studies suggest that atmospheric CO₂ concentrations may be a primary driver of vegetation change, particularly in savanna and grassland regions (76), even overriding changes in climate (77). This hypothesis is based on advantages conferred by increased CO₂ concentrations on C₃ plants relative to C₄ plants in terms of quantum yield, wherein reduced transpiration raises soil moisture and water-use efficiency (78-80), increases woody-plant resprouting rates (76), or both.

Global temperature and atmospheric CO₂ concentrations are strongly linked (81); both increase significantly across the last glacial-interglacial transition (82), and both are predicted to rise in the future (74). Our conclusion that substantial changes in vegetation are likely to occur in the future does not require that the relative importance of these two controls be disentangled. The spatial scale of our study does, however, allow for some degree of insight regarding the role of CO₂ versus climate in driving vegetation changes observed in the past.

At the global scale, atmospheric CO₂ levels are essentially homogeneous; when CO₂ rises in one part of the globe, it rises everywhere else proportionally. In contrast to CO₂, climate change is much more heterogeneous; increasing global temperature may manifest itself climatically in any number of ways across the globe (40), and climatic changes since the LGM have been spatially heterogeneous (26). Although temperatures have increased globally since the LGM - and as a result the hydrological cycle has been invigorated, resulting in wetter conditions in many regions - the degree of change varies significantly between regions, and in some parts of the world the LGM was in fact more humid than the Holocene (83-87). Thus, although certain ecosystems may be particularly sensitive to CO₂ forcing (88), or may exhibit threshold responses to changing CO₂ levels (89), if CO₂ was a dominant global driver of postglacial vegetation change, it would be expected to override, or strongly modify, climatic influences, and a widespread deglacial trend towards more arboreal taxa would be observed.

Paleoecological records indicate, however, that several regions show a clear decline in arboreal taxa during the deglacial period, in spite of increasing CO₂, but consistent with increasing drought stress

as a result of reduced precipitation and/or increasing temperatures (fig. S8). In other regions, where a deglacial increase in arboreal taxa is recorded, the variability observed is much more tightly coupled to changes in hydroclimate than CO₂ (fig. S8; see also (90)). These disparities do not mean that CO₂ is unimportant in vegetation change, but suggest instead that CO₂ changes may modulate vegetational responses to climate change, rather than being the sole or dominant driver.






Fig. S1. Distribution of paleoecological sites used in the assessment. Site dots colored by region.









Fig. S2

Confidence (high, medium, low) of individual assessments for each category of vegetation change (low, moderate, large), across the entire data set (Global) and individually for each region.






Fig. S3

Regional decomposition of Figure 2. Box plots show the estimated mean-annual temperature differences between the last glacial maximum and today in each of the three vegetation-change categories (low, moderate, large), for composition and structure.




Fig. S4

Comparison of the estimated temperature change obtained from the Annan and Hargreaves 2013 data (27) and the TraCE simulations (33), in each of the vegetation-change categories. The Annan and Hargreaves 2013 plots are as in Figure 2; the corresponding TraCE-21ka estimates are plotted for comparison. The TraCE plots show smaller temperature differences than the Annan and Hargreaves 2013 plots, but show a similar pattern of increasing magnitude of vegetational change with increasing temperature change.




Fig. S5

Box plots of the estimated mean-annual temperature differences between today and future-climate simulations for individual sites (using the nearest grid-point). RCP2.6 and RCP8.5 are as in Fig. 3a; this figure includes RCP4.5 and RCP6.0 for comparison. Projected temperature changes generally increase with higher radiative forcing.




Fig. S6

Estimated probability of large compositional and structural change by the end of the 21st century (average of 2081-2100) under the four representative concentration pathway (RCP) scenarios: RCP2.6, RCP4.5, RCP6.0, and RCP 8.5. Results for RCP2.6 and RCP8.5 are as in Fig. 3c; RCP4.5 and RCP6.0 are added for comparison. Global risk of large vegetational change increases with magnitude of radiative forcing.




Fig. S7

Probability of large change in vegetation composition and structure as a function of temperature change, based on the generalized additive models (GAMs) using two sets of temperature-difference estimates as the independent variable: Annan and Hargreaves 2013 LGM to pre-industrial temperature change (27) (as in Fig. 3b), and the TraCE-21ka LGM to late Holocene temperature change (33). Although the shape of the function differs between the two data sets, both show sensitivity of vegetation change to temperature change. Upper figure shows the model within the range of projected future temperature change for the 21st century. Lower figure shows results of the same models with an expanded x-axis to show the entire range of

temperatures used in developing the model. The excursion for ‘Structure-TraCE’ response between 10 and 18 is driven by several sites in Europe within this temperature range that were assessed as having moderate compositional change.




Fig. S8

Comparison of atmospheric CO₂ concentrations as measured from Antarctic ice cores (a) (91) with a selection of records that highlight the relative importance of CO₂ and climatic factors as drivers of vegetation change. At Lake Bosumtwi, Ghana, $\delta^{13}\text{C}$ data (woody C₃ plants versus C₄ grasses) indicate an increase in forest cover since the Last Glacial Maximum (LGM) (b, c) (89). Although this is consistent with the broad changes in atmospheric CO₂, vegetation change is much more tightly linked to changes in hydroclimate, as indicated by δD data (rainfall amount/intensity) from the same site. While the late Holocene persistence of forest at the site may be enabled by high CO₂ levels (89), it may also, or alternatively, be related to a mid-Holocene establishment of a stable forest state that is naturally resistant to perturbation (92). At Pella, on the southern margin of the Namib Desert, fossil pollen data indicate that arboreal vegetation was dominant during the LGM, contrary to predictions based on CO₂ forcing, and grasses became an important feature of the vegetation only during the CO₂-rich Holocene (d, e) (86). A similar pattern is observed in the North American southwest, with macrofossil data from the Lower Colorado River Basin revealing that significant C₄ grass cover only became established during the Holocene, with the decline of extra-local arboreal taxa (f, g) (85). In South America, records from the Andes indicate no significant change in forest cover at higher elevations sites such as Consuelo, and vegetation change at lower elevations is driven by climatic drivers, with no post-glacial increase in forest cover that can be clearly linked to changes in CO₂ (h) (93,94).

Table S1.

Team members responsible for compilation and assessment of specific regions.

REGION	TEAM MEMBERS
North America	Jackson, Nolan, Betancourt, Overpeck
Neotropics	Bush, Gosling
Subtropical South America	Latorre
Southern South America	Moreno
Europe	Huntley, Allen
Northern Eurasia	Tarasov, Muller, Binney
Japan	Takahara, Momohara
Beringia	Edwards, Lozhkin, Anderson, Binney
China	Y. Liu, Xu, Zheng, Shen, K.B. Liu, Weng
Middle East	Djamali, Cheddadi, Brewer
North Africa	Cheddadi, Djamali, Brewer
East and Tropical Africa	Ivory, Lezine, Marchant, Vincens
South Africa and Madagascar	Chase
Indo Pacific and Oceania	Stevenson, Hotchkiss, Kim
Australia	Kershaw, Haberle, Dodson

New Zealand	McGlone
-------------	---------

Table S2.

Working definitions used in assessments of paleoecological records.

Term	Definition
Composition	The species composition of the dominant vegetation in the record (Note: Although pollen records do not support species level discrimination, species turnover can be inferred from genus- and family-level changes)
Physiognomy/Structure	The physical structure of the dominant vegetation (e.g. coniferous forest, grassland, steppe, deciduous woodland, etc)
Large change	Complete or nearly complete replacement of dominant species or growth forms. (Note: does not imply complete biogeographic turnover; former dominants may remain in region, but not as dominants. Conversely, current dominants may be present at LGM , but not as dominants)
Moderate change	Partial replacement of dominant species or growth forms
Small change	Little or no replacement; dominant species and growth forms of today are unchanged at LGM
High confidence	Few or no substantial uncertainties, high quality data, inferences clear and unequivocal
Medium confidence	Some substantial uncertainties related to questions of data quality or clarity of inference
Low confidence	Significant uncertainties in the quality or interpretation of the data

Table S3.

Summary of the results of the Wilcoxon rank sum test for statistically significant difference in temperature change from LGM to modern between different levels of ecological change. The table shows the test statistic, W , and the p -values for different pairs of ecological change groups using two different estimates of the temperature change from LGM to modern.

Ecological Change Group 1	Ecological Change Group 2	Climate Variable	W	P-Value
Large Composition	Moderate Composition	Annan and Hargreaves	20926	3.228e-12
Large Composition	Moderate Composition	<u>TraCE</u>	22642	1.786e-9
Large Composition	Low Composition	Annan and Hargreaves	999.5	0.0003344
Large Composition	Low Composition	<u>TraCE</u>	1260.5	0.002809
Moderate Composition	Low Composition	Annan and Hargreaves	776.5	0.2982
Moderate Composition	Low Composition	<u>TraCE</u>	763.5	0.2629

Ecological Change Group 1	Ecological Change Group 2	Climate Variable	W	P-Value
Large Structure	Moderate Structure	Annan and Hargreaves	21602	4.432e-11
Large Structure	Moderate Structure	<u>TraCE</u>	24164	2.722e-7
Large Structure	Low Structure	Annan and Hargreaves	3243	3.361e-5
Large Structure	Low Structure	<u>TraCE</u>	3674.5	0.0004939
Moderate Structure	Low Structure	Annan and Hargreaves	2295	0.4374
Moderate Structure	Low Structure	<u>TraCE</u>	2341.5	0.5381

Table S4.

Summary of assessments, ordered by confidence, of the role of climate in driving observed moderate or large compositional change (top) or structural change (bottom). Columns represent the assessed role of climate in driving the observed ecological changes. Rows are the confidence assigned to those judgments by the assessors. The numbers are the number of sites in each combination of role of climate and confidence. The role of climate was not assessed for sites for which the ecological change was assessed as low (12 sites for composition and 30 sites for structure).

Composition	Large Role of Climate	Moderate	Low	N/A
High Confidence	511	11	0	0
Medium	40	13	1	0
Low	3	1	2	0
N/A	0	0	0	12
Total	555	25	3	12

Structure	Large Role of Climate	Moderate	Low	N/A
High Confidence	499	8	0	0
Medium	22	26	0	0
Low	5	3	1	0
N/A	0	0	0	30
Total	526	37	1	30

Instructions for Assessors

The following document was sent to all assessment team members, along with templates for data compilation and submission.

Assessment:

- Based on expert evaluation of all available and reliable paleoecological records from individual sites, in all regions of the planet.
- Our primary focus is on climate-driven vegetational change. Vegetation has, of course, been altered greatly by human activities in many regions. Thus, an important subsidiary question is whether substantial vegetational changes happened (or would have happened) in the absence of human activities.
- Assessments in this initial analysis will be based on “expert judgment” by paleoecologists who have experience in collection, analysis, and interpretation of the relevant data.
- Assessment will be qualitative, based on judgment of degree of contrast between glacial-age vegetation states and modern or Holocene vegetation states
- Assessments will be categorical, typically involving three classes (see below)
- Assessments will require judgment of confidence (also three classes)

Central Operational Questions (to be asked for each individual record):

1. What has been the magnitude of vegetational change, *in terms of vegetational composition*, since the last glacial period? *Choose from three outcomes:*
 - a. Large
 - b. Moderate
 - c. None
2. What is the level of confidence for this assessment? *Choose from three outcomes:*
 - a. High confidence
 - b. Medium confidence
 - c. Low confidence
3. If the answer to Question 1 was “Large” or “Moderate”, what role did climate change since the last glacial period play in driving change of this magnitude? *Choose from three outcomes:*
 - a. Large
 - b. Moderate
 - c. None
4. What is the level of confidence for this assessment? *Choose from three outcomes:*
 - a. High confidence
 - b. Medium confidence
 - c. Low confidence

5. What has been the magnitude of vegetational change, *in terms of vegetation physiognomy*, since the last glacial period? *Choose from three outcomes:*
 - a. Large
 - b. Moderate
 - c. None
6. What is the level of confidence for this assessment? *Choose from three outcomes:*
 - a. High confidence
 - b. Medium confidence
 - c. Low confidence
7. If the answer to Question 5 was “Large” or “Moderate”, what role did climate change since the last glacial period play in driving change of this magnitude? *Choose from three outcomes:*
 - a. Large
 - b. Moderate
 - c. None
8. What is the level of confidence for this assessment? *Choose from three outcomes:*
 - a. High confidence
 - b. Medium confidence
 - c. Low confidence

N.B. concerning Questions 3 and 7: Assessments should focus on judging whether climate change since the last glacial period was *sufficient* to drive vegetation change of “large” magnitude. Climate change as an explanation of high-magnitude vegetational change is not mutually exclusive of other processes (e.g., human activity, post-glacial CO₂ increase, megafaunal decline, etc.). In some regions with a strong human signature (e.g., Europe), judgment can be facilitated by examining the magnitude of change that is recorded before extensive human land-clearance. Thus, comparison of glacial-age with early Holocene assemblages may be useful in many cases.

N.B. concerning coverage: In some regions, some sites with temporal coverage to >14 ka may occur on land surfaces formerly occupied by continental or montane ice-sheets. *These sites should be included in the analysis, as long as they have records extending more than ~1000 years after site inception.* Thus, for example, a lake on glaciated terrain with basal pollen assemblages dating to 17 ka might be included, whereas a lake with basal sediments dating to 14.5 ka should be excluded. The purpose of the 1000-year rule is to ensure that sufficient time elapses after deglaciation to allow primary successional processes to play out.

Provisional Operational Definitions:

Vegetation terms:

Composition: The species composition of the dominant vegetation in the record. Although most pollen records do not support species-level discrimination, species turnover can of course be inferred from genus- and family-level changes. Macrofossils can also support such inferences, particularly among dominant types.

Physiognomy: The physical structure of the dominant vegetation (e.g., coniferous forest, grassland, steppe, tundra, deciduous woodland, etc.).

Magnitude of vegetation change:

Large: Complete or nearly complete replacement of dominant species or growth-forms. (This does not imply complete *biogeographic* turnover; former dominants may remain in the region, but not as dominants. Conversely, current dominants may have occurred at the locale during LGM, but not as dominants.)

Moderate: Partial replacement of dominant species or growth-forms

None: Little or no replacement; dominant species or growth-forms today are unchanged

Role of climate change:

Large: Climate change since the last glacial period is by itself sufficient to explain the magnitude of observed vegetation change. (Other factors may well have played a role, but in the absence of those factors, a similar magnitude of vegetation change would have been observed.)

Moderate: Climate change since the last glacial period explains at least part of the observed magnitude of vegetation change. (Other factors, in addition to climate change, are necessary to explain the magnitude of vegetation change).

None: Climate change is entirely insufficient to explain the magnitude of vegetation change observed since the last glacial period. Other factors are wholly responsible for the magnitude of change.

Level of confidence:

High: Few or no substantial uncertainties; high-quality data, inferences clear and unequivocal

Medium: Some substantial uncertainties arising from questions of data quality or clarity of inferences from the data

Low: Significant uncertainties in quality or interpretation of data

Additional Data table S1 (separate file)

List of paleoecological sites used in analysis with data and metadata:

Region:	Region in fig. S1, S2, S3, and Table 1.
Site:	Site name
Sitelats:	Latitude
Sitelons:	Longitude
CompChange:	Compositional change since LGP
CompConf:	Confidence of compositional change assessment
CompClim:	Role of climate change since LGP in driving compositional change
CompClimConf:	Confidence of climate-change role for composition
StructureChange:	Structural change since LGP
StructureConf:	Confidence of structural change assessment
StructureClim:	Role of climate change since LGP in driving structure change
StructureClimConf:	Confidence of climate-change role for structure
CompJustification:	Description of the ecological change the underlies the composition change assessment
StructureJustification:	Description of ecological change that underlies the structure change assessment
Sitetemps_ah:	Temperature difference since LGM (Annan&Hargreaves2013)
Sitetemps_trace1:	Temperature change since LGP (TraCE21ka)
Sitetemps_ccsm4rcp85:	Projected temperature change (RCP8.5)
Sitetemps_ccsm4rcp26:	Projected temperature change (RCP2.6)
Sitetemps_ccsm4rcp45:	Projected temperature change (RCP4.5)
Sitetemps_ccsm4rcp60:	Projected temperature change (RCP6.0)
ReferenceDOI:	Digital Object Identifier (DOI) for the site's reference
Reference:	Citation for paleoecological site