

## FORUM ORIGINAL RESEARCH COMMUNICATION

**Urinary excretion of sulfur metabolites****and risk of cardiovascular events and all-cause mortality in the  
general population**

Joost C. van den Born<sup>1\*</sup>, Anne-Roos S. Frenay<sup>1\*</sup>, Anne M. Koning<sup>1,2</sup>, Matthias Bachtler<sup>3</sup>, Ineke J. Riphagen<sup>4,5</sup>, Isidor Minović<sup>4</sup>, Martin Feelisch<sup>6</sup>, Marinda M. Dekker<sup>1</sup>, Marian L.C. Bulthuis<sup>1</sup>, Ron. T. Gansevoort<sup>4</sup>, Jan-Luuk Hillebrands<sup>1</sup>, Andreas Pasch<sup>3</sup>, Stephan J.L. Bakker<sup>4,5#</sup>, Harry van Goor<sup>1#</sup>

\*These authors contributed equally #These authors share last authorship

<sup>1</sup>Department of Pathology and Medical Biology, Division of Pathology, University Medical Center Groningen (UMCG), University of Groningen (RUG), Groningen, the Netherlands

<sup>2</sup>Department of Surgery, UMCG, RUG, Groningen, the Netherlands

<sup>3</sup>Department of Clinical Research, University of Bern, Bern, Switzerland

<sup>4</sup>Department of Internal Medicine, Division of Nephrology, UMCG, RUG, the Netherlands.

<sup>5</sup>Top Institute Food and Nutrition, Wageningen, the Netherlands

<sup>6</sup>Clinical and Experimental Sciences, Faculty of Medicine, and NIHR Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, United Kingdom

Running head: Urine sulfur metabolites in the general population

Word count: 4530

Number of references: 54

Number of grey scale illustrations: 2

Number of color illustrations: 2 (online only)

Keywords:

epidemiology, cardiovascular events, mortality, sulfate, thiosulfate, hydrogen sulfide

**Corresponding author**

H. van Goor, PhD

Department of Pathology & Medical Biology

University Medical Center Groningen

Hanzeplein 1, 9713, GZ, HP EA10, Groningen, the Netherlands

Phone: +31 50 3619529

Fax: +31 50 3619107

Email: [h.van.goor@umcg.nl](mailto:h.van.goor@umcg.nl)

## Abstract

Thiosulfate and sulfate are metabolites of hydrogen sulfide (H<sub>2</sub>S), a gaseous signaling molecule with cardiovascular protective properties. Urinary thiosulfate and sulfate excretion are associated with favorable disease outcome in high-risk patient groups. We investigated the relationship between urinary excretion of sulfur metabolites and risk of cardiovascular (CV) events and all-cause mortality in the general population. Subjects (n=6839) of the Prevention of Renal and Vascular End-stage Disease (PREVEND) study were followed prospectively. At baseline, 24-h urinary excretion of thiosulfate and sulfate were determined. Median urinary thiosulfate and sulfate excretion were 1.27 (ICR 0.89-2.37)  $\mu$ mol/24 h and 15.7 (IQR 12.0-20.3) mmol/24 h, respectively. Neither thiosulfate, nor sulfate excretion showed an independent association with risk of CV events. Sulfate, but not thiosulfate, was inversely associated with risk of all-cause mortality, independent of potential confounders (hazard ratio 0.73 (95% confidence interval 0.63-0.84),  $P < 0.001$ ). This association appeared most pronounced for normolipidemic subjects ( $P_{\text{interaction}} = 0.019$ ). The strong association between sulfate excretion and mortality in the general population emphasizes the (patho)physiological importance of sulfate or its precursor H<sub>2</sub>S. We hypothesize that urinary sulfate excretion, which is inversely associated with all-cause mortality in the general population, holds clinical relevance as a beneficial modulator in health and disease.

## Introduction

In the general population, cardiovascular disease (CVD) is the leading cause of death worldwide(28). Despite evolving efforts to control known risk factors(39), the incidence of CVD continues to increase, warranting the exploration of novel pathways for cardiovascular (CV) risk reduction.

This study focuses on the urinary excretion of the sulfur metabolites thiosulfate ( $S_2O_3^{2-}$ ) and sulfate ( $SO_4^{2-}$ ). These metabolites arise from the oxidation of sulfur containing amino acids (SAAs) in the transsulfuration pathway(15). One of the intermediates in this process is hydrogen sulfide ( $H_2S$ ), a gaseous signaling molecule with protective properties(41), including the potential to counteract CVD(30). Several preclinical studies have shown  $H_2S$  to be protective in CV disease. In mice, deficiency of cystathionine  $\gamma$ -lyase (CSE) - one of the  $H_2S$  producing enzymes - and consequent decreased systemic  $H_2S$  levels are accompanied by hyperhomocysteinemia, hypertension and impaired vasorelaxation(53). Exogenous administration of  $H_2S$  or  $H_2S$  donors offers protection in models of atherosclerosis(3, 23, 30), cardiac injury(36, 37, 42), renal disease(17, 37), and stroke(35, 50).

Sulfate itself is a dietary constituent and involved in various (patho)physiological processes(6, 22). Through sulfate conjugation or sulfation, it is responsible for the biotransformation and detoxification of many endogenous and exogenous substances(6, 8, 22). Sulfate may thereby directly contain disease development. Conversely, sulfated toxic intermediates, including indoxyl sulfate and p-cresyl sulfate, are implicated in carcinogenesis, and heart failure (2, 10, 49). As sulfation is generally considered to increase hydrophilicity and thereby to promote renal elimination of its targets(5), variations in urinary sulfate excretion may reflect the need for sulfate-mediated detoxification. The way in which sulfate conjugation is related to the toxicity of these specific compounds is yet unknown.

Previous studies of high-risk populations have shown that urinary excretion of thiosulfate and sulfate are associated with a favorable CV risk profile and survival in renal transplant recipients(45) and preservation of renal function in patients with diabetes(1, 46). Our

group has shown both urinary excretion and clearance of sulfate to be associated with a decreased rehospitalization rate and increased patient survival in chronic heart failure (CHF)(18). The apparent link to CV (patho)physiology led us to hypothesize that urinary excretion of sulfur metabolites is inversely associated with CV events and mortality in the general population. The aim of the study was to determine the association of urinary thiosulfate and sulfate excretion with risk of CV events and all-cause mortality in the general population.

### *Cohort characteristics*

Participants had a median 24-h urinary thiosulfate excretion of 1.27 (0.89-2.37)  $\mu\text{mol}/24\text{ h}$  and a median 24-h urinary sulfate excretion of 15.7 (12.0-20.3)  $\text{mmol}/24\text{ h}$  and a. Baseline characteristics are presented in Table 1, overall and categorized by gender-stratified quintiles of sulfate excretion. The mean age was  $53.4 \pm 12.1$  years and 50% of the subjects ( $n=3420$ ) were female. 24-h urinary urea excretion, a rough estimate of dietary protein intake, rises with every quintile of urinary sulfate excretion. Subjects in the highest quintile were younger and had a larger body surface area (BSA). Systolic blood pressure and heart rate were lower in this group. Subjects in the lowest quintile of urinary sulfate excretion more often had a history of CV events, which was accompanied by the highest use of anti-hypertensive and lipid lowering treatment. This quintile contained the most current smokers and its subjects had the lowest estimated glomerular filtration rate (eGFR).

### *Factors associated with urinary excretion of thiosulfate and sulfate*

Univariable and multivariable linear regression analyses identified gender, BSA, diabetes and current smoking as potential confounders for 24-h urinary thiosulfate excretion (Table 2). The same analyses identified gender, age, BSA, history of CVD, current smoking and consumption of more than 1 alcoholic beverage per day as potential confounders for daily urinary sulfate excretion (Table 3).

24-h urinary thiosulfate excretion (Table 4) is positively associated with systolic and diastolic blood pressure, anti-diabetic treatment, glucose, eGFR and 24-h urinary albumin and sulfate excretion, and inversely associated with high-density lipoprotein (HDL). 24-h urinary sulfate excretion is positively associated with diastolic blood pressure, glucose,

eGFR and 24-h urinary albumin and thiosulfate excretion, and inversely associated with heart rate, anti-hypertensive and lipid lowering treatment, HDL and high sensitivity C-reactive protein (hs-CRP).

#### *Urinary excretion of thiosulfate and sulfate and risk of cardiovascular events*

In the follow-up (8.2 (7.7-8.8) years), 504 CV events were recorded. 239 subjects (47.4%) had ischemic heart disease of whom 36 (15.1%) died. A cerebrovascular event occurred in 111 subjects (22%), of whom 4 (3.6%) died.

Crude Cox proportional hazards analyses showed a significant inverse association of 24-h urinary sulfate, but not thiosulfate excretion with risk of CV events (Table 5, model 1 for 24-h urinary sulfate excretion, hazard ratio (HR) per doubling 0.87 (0.76-0.99),  $P=0.039$ ). This association lost its statistical significance by adjustment for potential confounding factors (gender, age, BSA, history of cardiovascular disease, diabetes, current smoking and consumption of >1 alcoholic beverage per day; Table 5, model 2 for 24-h urinary sulfate excretion, HR per doubling 0.89 (0.77-1.03),  $P=0.107$ ).

#### *Urinary excretion of thiosulfate and sulfate and risk of all-cause mortality*

During follow-up a total of 445 subjects (6.5%) died. Of these, 120 (27.0%) died of a cardiovascular cause.

Most deaths occurred in the lowest quintiles of 24-h urinary thiosulfate and sulfate excretion (109 (8.0%), log rank test,  $P=0.036$  and 151 (11%), log rank test,  $P<0.001$ , respectively). The Kaplan-Meier plots (Figure 1 and 2) demonstrate the differential distribution of quintiles of thiosulfate and sulfate excretion among survivors and non-survivors ( $\chi^2$  test,  $P=0.011$ , Figure 1 and  $\chi^2$  test,  $P<0.001$ , Figure 2, respectively), with overrepresentation of the lowest quintile, particularly of sulfate excretion, in the non-survivors population.

Despite the significant difference in the survival distributions for quintiles of thiosulfate excretion determined by the log-rank test, Cox proportional hazards analysis showed no association between 24-h urinary thiosulfate excretion and all-cause mortality (Table 6, model 1 for 24-h urinary thiosulfate excretion, HR per doubling 0.93 (0.86-1.01),  $P=0.100$ ).

For 24-h urinary sulfate excretion Cox proportional hazards analysis did show a significant inverse association with all-cause mortality (Table 6, model 1 for 24-h urinary sulfate excretion, HR per doubling 0.65 (0.57-0.74),  $P < 0.001$ ), which remained significant after adjustment for potential confounders (Table 6, model 2 for 24-h urinary sulfate excretion, HR per doubling 0.73 (0.63-0.84),  $P < 0.001$ ), and further adjustment for 24-h urinary urea excretion (HR per doubling 0.66 (0.54-0.80),  $P < 0.001$ ). Crude Cox regression also showed an association of urinary sulfate excretion with death of a CV cause (HR per doubling 0.64 (0.51-0.82),  $P < 0.001$ ). However, after correction for potential confounders, this association was no longer significant (HR per doubling 0.77 (0.58-1.02),  $P = 0.070$ ).

Restricted cubic splines showed no significant deviances from linear associations with all-cause mortality for either 24-h urinary thiosulfate or sulfate excretion (Figure 3,  $P_{\text{nonlinearity}} = 0.08$  and 0.10, respectively).

Stratified analyses of the association between 24-h urinary sulfate excretion and all-cause mortality showed consistent hazard ratios across various subgroups, except for those stratified by cholesterol (normo- vs. dyslipidemia) (Figure 4,  $P_{\text{interaction}} = 0.019$ ). Stratification by renal function ( $\text{eGFR} \geq 60$  vs.  $< 60$  ml/min) and the presence of diabetes (no diabetes vs. diabetes) resulted in borderline significant differences between groups (Figure 4,  $P_{\text{interaction}} = 0.057$  and  $P_{\text{interaction}} = 0.054$ , respectively). The corresponding hazard ratios were lower for subjects with normolipidemia, those with impaired renal function ( $\text{eGFR} < 60$  ml/min) and those without diabetes.

### *Causal path analysis*

In causal path analyses, following adjustment for potential confounders, the association of 24-h urinary sulfate excretion and all-cause mortality was further adjusted for hemodynamic parameters (systolic and diastolic blood pressure, heart rate and anti-hypertensive treatment), lipid profile (total cholesterol, HDL, triglycerides and lipid lowering treatment), hs-CRP and eGFR. None of these adjustments affected the statistical significance of the association between 24-urinary sulfate excretion and all-cause mortality (Table 7, model 2-5, all  $P < 0.001$ ).

## Discussion

In this large cohort of individuals from the general population 24-h urinary sulfate excretion was found to be inversely associated with risk of CV events and all-cause mortality. While the association with all-cause mortality remained significant after adjustment for potential confounding factors, the association with CV events did not. This may create the impression that sulfate excretion is not directly connected to CVD and related mortality. However, studies on renal transplant recipients(45) and CHF patients(18) have shown urinary sulfate excretion to be associated with a beneficial CV risk profile and patient survival. It should also be noted that the definition of a CV event is limited to acute incidents, whereas CV causes of death also include chronic forms of CVD, such as CHF. Nevertheless, the association of sulfate excretion with death of a CV cause, found in our study, also lost its statistical significance on correction for potential confounders. As of yet, this unexpected finding is without explanation.

24-h urinary thiosulfate excretion, in turn, was not found to be associated with either risk of CV events or all-cause mortality, although Kaplan Meier analysis did show significantly different survival distributions for quintiles of thiosulfate excretion. This is in contrast with previous findings in the before mentioned high-risk population of renal transplant recipients, which, apart from sulfate, also link thiosulfate to a favorable CV risk profile and patient survival(45). The discrepancy may indicate that thiosulfate excretion is triggered in disease conditions, precluding determination of an association with baseline values in our cohort of predominantly healthy individuals from the general population. This hypothesis is substantiated by the increase in urinary thiosulfate excretion found in renal transplant recipients compared to healthy controls(45). Furthermore, whereas sulfate is an end-product, thiosulfate is an intermediate metabolite and therefore less stable(15). In fact, several studies have demonstrated reconversion of thiosulfate into H<sub>2</sub>S(16, 24, 27, 48). Possibly, its dynamic nature provides another explanation for the absence of a robust relationship between urinary thiosulfate excretion and CV events or mortality, especially in a heterogeneous group of subjects from the general population. Besides, thiosulfate treatment has recently been shown to offer protection in experimental models of



hypertensive cardiac(37) and renal(38) disease, as well as diabetes(25). In the latter, this has been related to the activation of thiosulfate sulfurtransferase(25).

Several factors may underlie the inverse association between 24-h urinary sulfate excretion and all-cause mortality. For one thing, dietary intake of sulfate and the SAAs methionine and cysteine is known to augment urinary sulfate excretion(21, 22, 33). In the present study this is confirmed by the strong association with 24-h urinary urea excretion, which roughly reflects dietary protein intake. Diets high in protein or specific SAAs have been associated with decreased risk of CVD and mortality(19, 29, 31, 32, 47). However, results have been inconsistent(19, 29, 31, 32, 44, 47). Here, adjustment for 24-h urinary urea excretion on top of potential confounders had no effect on the statistical significance of the association between urinary sulfate excretion and all-cause mortality. This is in line with previous studies of renal transplant recipients and patients with diabetes mellitus type 1 nephropathy in which the association of sulfate excretion with all-cause mortality or renal disease progression was shown to be independent of dietary protein intake(1, 45).

Perhaps the degree of enzymatic production of sulfate from SAAs is a stronger determinant. This process links urinary sulfate excretion to H<sub>2</sub>S, which is an intermediate in the transsulfuration pathway(15). Also, H<sub>2</sub>S from other sources - including the reduction of bound sulfur and the reduction of sulfate by bacteria in the gut - contributes to the overall sulfate production(12, 34). Thus, urinary sulfate excretion may, at least in part, reflect endogenous production of H<sub>2</sub>S, which is of particular interest as this gaseous signaling molecule is known to be involved in CV (patho)physiology(30). In experimental models, exogenous administration of H<sub>2</sub>S or H<sub>2</sub>S donors has been shown to protect against atherosclerosis(3, 23, 30), cardiac injury(36, 37, 42), renal disease(17, 37), and stroke(35, 50). Clinically, reduced plasma levels of H<sub>2</sub>S-related metabolites have been associated with obesity and diabetes mellitus type 2(51). Research has shown H<sub>2</sub>S to be involved in blood pressure regulation(54), lipid metabolism(13, 52), inflammation(40) and renal function(17). However, in causal path analysis, adjustment for associated variables did not identify any of these factors as the underlying mechanism for the association between urinary sulfate excretion and all-cause mortality.

In our study, subjects from the highest quintile of sulfate excretion did show the highest eGFR. This is in accordance with results from previous studies on the relationship of sulfate (excretion) and renal function(1, 9, 45, 46). As eGFR itself is strongly associated with a reduced risk of CVD and all-cause mortality(4) one may assume this explains the inverse association of urinary sulfate excretion with all-cause mortality found in our study. However, as mentioned above, adjustment of this association for eGFR on top of potential confounders had no effect on its statistical significance. Besides, under steady-state conditions the urinary excretion of any metabolite is not determined by renal function, but rather represents its metabolic turnover which has to be matched to avoid systemic accumulation. Subgroup analysis revealed a lower HR for subjects with an eGFR below 60 ml/min compared to those with normal renal function. Although the interaction was only borderline significant, the observed trend substantiates the notion that urinary sulfate excretion is not a reflection of renal function. Interestingly, in CHF patients both urinary excretion and clearance, but not the plasma concentration of sulfate have been found to be associated with favorable disease outcome(18). Collectively, these findings suggest that the renal handling of sulfate is of importance. Unfortunately, in the present study sulfate clearance could not be assessed.

Subgroup analysis also uncovered a lower HR for subjects with normal cholesterol levels compared to those with dyslipidemia. The interaction between cholesterol levels and the association of sulfate excretion with all-cause mortality is likely connected to the process of steroid sulfation. Sulfate conjugation increases hydrophilicity and thereby promotes urinary excretion of steroids, but also modulates cholesterol function(6). Possibly, an increased need for sulfation explains the relatively small advantage of sulfate excretion in subjects with dyslipidemia.

Considering the before mentioned evidence from previous studies of high-risk populations, the trend towards an interaction between the presence of diabetes and the association of sulfate excretion with all-cause mortality is surprising. Unexpectedly, the association appeared to be most pronounced for subjects without diabetes, the meaning of which remains to be elucidated.

This study has several limitations. Firstly, it was carried out in a single center and includes almost only Caucasian subjects. Consequently, our results may not be representative for other populations. Also, our data are observational and therefore do not allow causality of the relationship between urinary sulfate excretion and all-cause mortality to be established. Furthermore, 24-h urinary urea excretion provides only a rough estimate of dietary protein intake and is therefore inferior to actual dietary information, which unfortunately was not available. Strengths of our study include the large number of subjects and their extensive characterization, the use of 24-h urine samples and the long duration of follow-up.

In conclusion, 24-h urinary sulfate, but not thiosulfate excretion is inversely associated with all-cause mortality in a large, general population based cohort, in particular in subjects with normolipidemia. Whether functional properties of sulfate itself or rather its relationship with H<sub>2</sub>S underlies this association remains to be elucidated. Accordingly, further research is warranted to unravel the role of sulfate and its urinary excretion in physiology and disease.

## Innovation

Urinary sulfate excretion and health benefit go hand in hand, not only in high-risk patient groups, but also in the general population. The strong inverse association between urinary excretion of sulfate and all-cause mortality in a large, general population based cohort emphasizes the (patho)physiological importance either of sulfate itself or of its precursor hydrogen sulfide. Accordingly, sulfate excretion holds promise as a marker of physiological disturbance and may even serve as a target for nutritional intervention to promote healthy aging.

## Materials and methods

### *Study population*

Data on subjects of the Prevention of Renal and Vascular End-stage Disease (PREVEND) study were used for this analysis. The PREVEND study was designed to prospectively investigate the natural course of albuminuria and its relation to renal and cardiovascular

disease in a large cohort drawn from the general population, and has previously been described in detail(14). In brief, from 1997 to 1998 a total of 85421 inhabitants of Groningen, the Netherlands were approached with a short questionnaire (regarding demographics, medication use, and pregnancy) and a vial to collect an early morning urine sample. Pregnancy and type-1 diabetes were exclusion criteria. In 40856 responding subjects urinary albumin concentration was determined. Subsequently, 6000 participants with a urinary albumin concentration  $\geq 10$  mg/l and 2592 randomly selected subjects with a urinary albumin concentration  $< 10$  mg/l enrolled in the study. In total, the PREVEND cohort consists of 8592 individuals. The study was approved by the medical ethics committee of the University of Groningen and conducted in accordance with the Declaration of Helsinki. Written informed consent was obtained from all participants. For the present analysis, urine samples from the second survey (in 2001 to 2003) were selected and this time-point was considered baseline. Due to unavailability of urine samples or data on urine volume 16 of 6855 potential subjects had to be excluded. Of the remaining 6839 participants - aged 32 to 80 years - 967 (14.2%) had albuminuria, whereas 5854 (85.8%) did not. For 18 participants (0.3%) this data was missing.

### *Data collection*

All subjects of the PREVEND study visited the outpatient research unit twice for baseline investigation. At the first visit, participants filled out a questionnaire on demographics, general health, CVD history, medication, smoking habits and alcohol consumption. Height, weight and waist circumference were measured. In addition, a fasting blood sample was collected and stored at  $-80^{\circ}\text{C}$  until analysis. At the second visit, blood pressure was measured in supine position every minute for 8 minutes with an automatic Dinamap XL Model 9300 series device (Johnson-Johnson Medical, Tampa, FL), and the mean of the last 2 recordings was listed. Additionally, participants collected 24-h urine after oral and written instructions. Urine samples were stored at  $-20^{\circ}\text{C}$  until analysis.

### *Laboratory measurements*

Serum total cholesterol, serum glucose and serum and urinary creatinine were measured by Kodak Ektachem dry chemistry (Eastman Kodak, Rochester, NY). High-density

lipoprotein (HDL) cholesterol was determined with a homogenous method (direct HDL, Aeroset TM System, Abbott Laboratories, Abbott Park, IL). Triglycerides were measured enzymatically.

Urinary albumin concentrations and high sensitivity C-reactive protein (hs-CRP) were determined by nephelometry (Dade Behring Diagnostics, Marburg, Germany). Urinary urea was measured with a MEGA clinical chemistry analyzer (Merck, Darmstadt, Germany) by a photometric test with the urease-GIDH method. Urinary sulfate was determined by ion exchange chromatography (type 861; Metrohm, Herisau, Switzerland), using a Metrosep A Supp 4 - 250/4.0 column. Intra- and inter-assay variations were 2.0% and 4.3%, respectively. Urinary thiosulfate was measured by reverse-phase HPLC as previously described(26, 45). In brief, 25 µl of urine was derivatized with 5 µl of 46 mM monobromobimane, 25 µl of acetonitrile and 25 µl of 160 mM HEPES/16 mM EDTA pH 8 buffer (Invitrogen, Carlsbad, CA) for 30 min. Derivatization was stopped by addition of 50 µl of 65 mM methanosulfonic acid (Fluka, Buchs, Switzerland), and proteins were removed by re-centrifugation. Intra- and inter-assay variations were 8.6% and 9.3%, respectively.

#### *Definitions and Primary Outcome Definition*

Body surface area (BSA) was defined following the Dubois & Dubois formula:  $0.007184 \times (\text{Height}^{0.725} \times \text{Weight}^{0.425})$ (7). Diabetes mellitus was defined according to the guidelines of the American Diabetes Association as a fasting glucose level  $\geq 7.0$  mmol/l and/or the use of anti-diabetic medication(43). Estimated glomerular filtration rates (eGFR) were calculated following the Chronic Kidney Disease Epidemiology collaboration equation (CKD-EPI) formula(20). Urinary concentrations of creatinine, albumin, thiosulfate, and urea were multiplied by the 24-h urinary volume to determine 24-h urinary excretion. Information on death, cardiovascular disease and hospitalization for cardiovascular disease was obtained from the participant's questionnaire and the Dutch national registry of all hospital discharge diagnoses (Prismant). Data were coded according to the International Statistical Classification of Diseases (ICD-10) and the International Classification of Health Interventions(11). All-cause mortality was set as the primary endpoint and cardiovascular events were set as the secondary endpoint of this study.

### Statistical analyses

Statistical analyses were performed with the Statistical Package for Social Sciences (SPSS Statistics, IBM Corporation, Armonk, NY) version 22.0, STATA/SE software (Release 13; StataCorp, College Station, TX) and R version 3.0.1 (Vienna, Austria). Graphs were drawn in GraphPad Prism (version 5.0, GraphPad Software, La Jolla, California, USA).

Distributions of variables were visualized with histograms and Q-Q plots. Normally distributed continuous data are presented as mean  $\pm$  standard deviation (SD). Skewed data are presented as median (interquartile range IQR) and were normalized prior to analyses by logarithmic transformation (triglycerides, glucose, hs-CRP, and 24-h urinary albumin, sulfate, and thiosulfate excretion). 24-h urinary thiosulfate and sulfate excretion values were log-transformed according to the base of two, to allow interpretation of hazard ratios (HRs) per doubling. Nominal data are presented as  $n$  (%).

Differences in baseline characteristics between groups were determined using One-way ANOVA for normally distributed continuous data, the Kruskal-Wallis test for skewed continuous data, and the Chi-square test for nominal data. Chi-square tests were also used to compare the distributions of quintiles of 24-h urinary thiosulfate and sulfate excretion among survivors and non-survivors.

To compare the survival distributions for quintiles of 24-h urinary thiosulfate and sulfate excretion Kaplan-Meier plots and log-rank tests were applied. Follow-up time was defined as the period from the date of urine collection until the date of death or the end of follow-up (on January 1, 2009).

Univariable linear regression analyses were performed for 24-h urinary thiosulfate and sulfate excretion and potential confounding factors, followed by multivariable analyses with backward selection. Subsequently, the association between 24-h urinary thiosulfate and sulfate excretion and potential intermediary factors was determined using univariable linear regression analyses.

As several subjects showed missing values for one or more baseline variables and bias can be introduced when subjects with missing values are excluded, multiple imputation

was used to obtain 5 imputed datasets. Using the imputed dataset, Cox proportional hazards analysis was applied to study the association between 24-h urinary thiosulfate and sulfate excretion and risk of CV events and all-cause and CV mortality. Follow-up time was defined as the period from the date of urine collection until the date of a CV event, death or the end of follow-up (on January 1, 2009). Participants were censored if they moved to an unknown destination or, in the case of CV events, if they died of a non-CV cause. HRs are reported with 95% confidence intervals (95% CI). Cox regression analysis with restricted cubic splines with 3 knots was used to test for potential non-linearity of the associations of 24-h urinary thiosulfate and sulfate excretion with all-cause mortality. Finally, Cox proportional hazards analysis with adjustment for potential confounders was applied to assess HRs for the association of 24-h urinary sulfate excretion with all-cause mortality across various subgroups.

### Acknowledgements

The PREVEND study was financially supported by several grants from the Dutch Kidney Foundation. JCvdB received financial support by grant 2015-07 from the J.K. de Cock Foundation and the Groningen University Institute for Drug Exploration (GUIDE). HvG is supported by a grant from the Dutch Kidney foundation (P13-114). SJLB, IJR and IM received financial support from grant CH-001 and CH-003 of the Top Institute Food and Nutrition. MF is supported by the National Institute for Health Research through the NIHR Southampton Biomedical Research Centre. Funding agencies had no role in the design of the study, the collection, analysis, and interpretation of the data, or the decision to submit the manuscript for publication.

The authors thank Beatrix Blanchard for her excellent technical support with the thiosulfate measurements at University Hospital Bern, Switzerland.

### Author Disclosure Statement

No competing financial interests exist.

## References

1. Andrésdóttir G, Bakker SJL, Hansen HP, Parving H-H, Rossing P. 2013. Urinary sulphate excretion and progression of diabetic nephropathy in type 1 diabetes. *Diabet. Med. J. Br. Diabet. Assoc.* 30(5):563–66
2. Cao X-S, Chen J, Zou J-Z, Zhong Y-H, Teng J, et al. 2015. Association of indoxyl sulfate with heart failure among patients on hemodialysis. *Clin. J. Am. Soc. Nephrol. CJASN.* 10(1):111–19
3. Cheung SH, Kwok WK, To KF, Lau JYW. 2014. Anti-atherogenic effect of hydrogen sulfide by over-expression of cystathionine gamma-lyase (cse) gene. *PloS One.* 9(11):e113038
4. Chronic Kidney Disease Prognosis Consortium, Matsushita K, van der Velde M, Astor BC, Woodward M, et al. 2010. Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality in general population cohorts: a collaborative meta-analysis. *Lancet Lond. Engl.* 375(9731):2073–81
5. Coughtrie MW, Bamforth KJ, Sharp S, Jones AL, Borthwick EB, et al. 1994. Sulfation of endogenous compounds and xenobiotics—interactions and function in health and disease. *Chem. Biol. Interact.* 92(1–3):247–56
6. Dawson PA. 2013. Role of sulphate in development. *Reprod. Camb. Engl.* 146(3):R81–89
7. Du Bois D, Du Bois EF. 1989. A formula to estimate the approximate surface area if height and weight be known. 1916. *Nutr. Burbank Los Angel. Cty. Calif.* 5(5):303–311–313
8. Florin T, Neale G, Gibson GR, Christl SU, Cummings JH. 1991. Metabolism of dietary sulphate: absorption and excretion in humans. *Gut.* 32(7):766–73
9. Freeman RM, Richards CJ. 1979. Studies on sulfate in end-stage renal disease. *Kidney Int.* 15(2):167–75
10. Glatt H, Engelke CE, Pabel U, Teubner W, Jones AL, et al. 2000. Sulfotransferases: genetics and role in toxicology. *Toxicol. Lett.* 112–113:341–48
11. *ICD-10: International statistical classification of diseases and related health problems.* 2011. Geneva: World Health Organization



12. Ishigami M, Hiraki K, Umemura K, Ogasawara Y, Ishii K, Kimura H. 2009. A source of hydrogen sulfide and a mechanism of its release in the brain. *Antioxid. Redox Signal.* 11(2):205–14
13. Jain SK, Micinski D, Lieblong BJ, Stapleton T. 2012. Relationship between hydrogen sulfide levels and hdl-cholesterol, adiponectin, and potassium levels in the blood of healthy subjects. *Atherosclerosis.* 225(1):242–45
14. Joosten MM, Gansevoort RT, Mukamal KJ, Lambers Heerspink HJ, Geleijnse JM, et al. 2014. Sodium excretion and risk of developing coronary heart disease. *Circulation.* 129(10):1121–28
15. Kabil O, Banerjee R. 2014. Enzymology of h<sub>2</sub>s biogenesis, decay and signaling. *Antioxid. Redox Signal.* 20(5):770–82
16. Koj A, Frendo J, Janik Z. 1967. [35s]thiosulphate oxidation by rat liver mitochondria in the presence of glutathione. *Biochem. J.* 103(3):791–95
17. Koning AM, Frenay A-RS, Leuvenink HGD, van Goor H. 2015. Hydrogen sulfide in renal physiology, disease and transplantation - the smell of renal protection. *Nitric Oxide.* 2015 Apr 30;46:37-49
18. Koning AM, Meijers W, Minović I, Post A, Feelisch M, et al. 2016. The fate of sulfate in chronic heart failure. *Am. J. Physiol. Heart Circ. Physiol.* ajpheart.00645.2016.
19. Larsson SC, Håkansson N, Wolk A. 2015. Dietary cysteine and other amino acids and stroke incidence in women. *Stroke J. Cereb. Circ.* 46(4):922–26
20. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF, et al. 2009. A new equation to estimate glomerular filtration rate. *Ann. Intern. Med.* 150(9):604–12
21. Magee EA, Curno R, Edmond LM, Cummings JH. 2004. Contribution of dietary protein and inorganic sulfur to urinary sulfate: toward a biomarker of inorganic sulfur intake. *Am. J. Clin. Nutr.* 80(1):137–42
22. Markovich D. 2001. Physiological roles and regulation of mammalian sulfate transporters. *Physiol. Rev.* 81(4):1499–1533
23. Meng G, Ma Y, Xie L, Ferro A, Ji Y. 2015. Emerging role of hydrogen sulfide in hypertension and related cardiovascular diseases. *Br. J. Pharmacol.* 172(23):5501–11

24. Mikami Y, Shibuya N, Kimura Y, Nagahara N, Ogasawara Y, Kimura H. 2011. Thioredoxin and dihydrolipoic acid are required for 3-mercaptopyruvate sulfurtransferase to produce hydrogen sulfide. *Biochem. J.* 439(3):479–85
25. Morton NM, Beltram J, Carter RN, Michailidou Z, Gorjanc G, et al. 2016. Genetic identification of thiosulfate sulfurtransferase as an adipocyte-expressed antidiabetic target in mice selected for leanness. *Nat Med.* 2016 Jul;22(7):771-9.
26. Newton GL, Dorian R, Fahey RC. 1981. Analysis of biological thiols: derivatization with monobromobimane and separation by reverse-phase high-performance liquid chromatography. *Anal. Biochem.* 114(2):383–87
27. Olson KR, DeLeon ER, Gao Y, Hurley K, Sadauskas V, et al. 2013. Thiosulfate: a readily accessible source of hydrogen sulfide in oxygen sensing. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 305(6):R592-603
28. Organisation mondiale de la santé. 2014. *Global status report on noncommunicable diseases 2014: attaining the nine global noncommunicable diseases targets; a shared responsibility*. Geneva: World Health Organization
29. Pedersen AN, Kondrup J, Børsheim E. 2013. Health effects of protein intake in healthy adults: a systematic literature review. *Food Nutr. Res.* 2013 Jul 30;57
30. Polhemus DJ, Lefer DJ. 2014. Emergence of hydrogen sulfide as an endogenous gaseous signaling molecule in cardiovascular disease. *Circ. Res.* 114(4):730–37
31. Robin S, Maupoil V, Groubatch F, Laurant P, Jacqueson A, Berthelot A. 2003. Effect of a methionine-supplemented diet on the blood pressure of wistar-kyoto and spontaneously hypertensive rats. *Br. J. Nutr.* 89(4):539–48
32. Robin S, Maupoil V, Laurant P, Jacqueson A, Berthelot A. 2004. Effect of a methionine-supplemented diet on the blood pressure of sprague-dawley and deoxycorticosterone acetate-salt hypertensive rats. *Br. J. Nutr.* 91(6):857–65
33. Sabry ZI, Shadarevian SB, Cowan JW, Campbell JA. 1965. Relationship of dietary intake of sulphur amino-acids to urinary excretion of inorganic sulphate in man. *Nature.* 206(987):931–33
34. Shen X, Carlström M, Borniquel S, Jädert C, Kevil CG, Lundberg JO. 2013. Microbial regulation of host hydrogen sulfide bioavailability and metabolism. *Free Radic. Biol. Med.* 60:195–200

35. Shi H-Q, Zhang Y, Cheng M-H, Fan B-S, Tian J-S, et al. 2016. Sodium sulfide, a hydrogen sulfide-releasing molecule, attenuates acute cerebral ischemia in rats. *CNS Neurosci. Ther.* 22(7):625–32
36. Snijder PM, de Boer RA, Bos EM, van den Born JC, Ruifrok W-PT, et al. 2013. Gaseous hydrogen sulfide protects against myocardial ischemia-reperfusion injury in mice partially independent from hypometabolism. *PloS One.* 8(5):e63291
37. Snijder PM, Frenay AS, de Boer RA, Pasch A, Hillebrands J, et al. 2014. Exogenous administration of thiosulfate, a donor of hydrogen sulfide, attenuates angiotensin ii-induced hypertensive heart disease in rats. *Br J Pharmacol.* 2015 Mar;172(6):1494-504.
38. Snijder PM, Frenay A-RS, Koning AM, Bachtler M, Pasch A, et al. 2014. Sodium thiosulfate attenuates angiotensin ii-induced hypertension, proteinuria and renal damage. *Nitric Oxide Biol. Chem. Off. J. Nitric Oxide Soc.* 42:87–98.
39. SPRINT Research Group, Wright JT, Williamson JD, Whelton PK, Snyder JK, et al. 2015. A randomized trial of intensive versus standard blood-pressure control. *N. Engl. J. Med.* 373(22):2103–16
40. Szabó C. 2007. Hydrogen sulphide and its therapeutic potential. *Nat. Rev. Drug Discov.* 6(11):917–35
41. Szabo C. 2010. Medicinal chemistry and therapeutic applications of the gasotransmitters NO, CO, and H<sub>2</sub>S and their prodrugs. In *Burger's Medicinal Chemistry and Drug Discovery*, pp. 265–368. Hoboken, NJ, USA: John Wiley & Sons, Inc. Seventh Edition ed.
42. Szabó G, Veres G, Radovits T, Gero D, Módis K, et al. 2011. Cardioprotective effects of hydrogen sulfide. *Nitric Oxide Biol. Chem.* 25(2):201–10
43. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. 1997. Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care.* 20(7):1183–97
44. Tuttle KR, Milton JE, Packard DP, Shuler LA, Short RA. 2012. Dietary amino acids and blood pressure: a cohort study of patients with cardiovascular disease. *Am. J. Kidney Dis. Off. J. Natl. Kidney Found.* 59(6):803–9

45. van den Berg E, Pasch A, Westendorp WH, Navis G, Brink EJ, et al. 2014. Urinary sulfur metabolites associate with a favorable cardiovascular risk profile and survival benefit in renal transplant recipients. *J. Am. Soc. Nephrol. JASN*. 2014 Jun;25(6):1303-12
46. van den Born JC, Frenay A-RS, Bakker SJL, Pasch A, Hillebrands J-L, et al. 2016. High urinary sulfate concentration is associated with reduced risk of renal disease progression in type 2 diabetes. *Nitric Oxide Biol. Chem. Off. J. Nitric Oxide Soc.* 55–56:18–24
47. Vasdev S, Singal P, Gill V. 2009. The antihypertensive effect of cysteine. *Int. J. Angiol. Off. Publ. Int. Coll. Angiol. Inc.* 18(1):7–21
48. Villarejo M, Westley J. 1963. Mechanism of rhodanese catalysis of thiosulfate-lipoate oxidation-reduction. *J. Biol. Chem.* 238:4016–20
49. Wang C-H, Cheng M-L, Liu M-H, Shiao M-S, Hsu K-H, et al. 2016. Increased p-cresyl sulfate level is independently associated with poor outcomes in patients with heart failure. *Heart Vessels*. 31(7):1100–1108
50. Wang J-F, Li Y, Song J-N, Pang H-G. 2014. Role of hydrogen sulfide in secondary neuronal injury. *Neurochem. Int.* 64:37–47
51. Whiteman M, Gooding KM, Whatmore JL, Ball CI, Mawson D, et al. 2010. Adiposity is a major determinant of plasma levels of the novel vasodilator hydrogen sulphide. *Diabetologia*. 53(8):1722–26
52. Wu D, Zheng N, Qi K, Cheng H, Sun Z, et al. 2015. Exogenous hydrogen sulfide mitigates the fatty liver in obese mice through improving lipid metabolism and antioxidant potential. *Med. Gas Res.* 5(1):1
53. Yang G, Wu L, Jiang B, Yang W, Qi J, et al. 2008. H<sub>2</sub>S as a physiologic vasorelaxant: hypertension in mice with deletion of cystathionine gamma-lyase. *Science*. 322(5901):587–90
54. Zhao W, Zhang J, Lu Y, Wang R. 2001. The vasorelaxant effect of h<sub>2</sub>s as a novel endogenous gaseous k(atp) channel opener. *EMBO J.* 20(21):6008–16

## Abbreviations

BMI; body mass index

BSA; body surface area

CI; confidence interval

CKD-EPI; Chronic Kidney Disease Epidemiology collaboration equation

CSE; cystathionine  $\gamma$ -lyase

CV; cardiovascular

CVD; cardiovascular disease

DPB; diastolic blood pressure

EDTA; Ethylenediaminetetraacetic acid

eGFR; estimated glomerular filtration rate

HDL; high-density lipoprotein

HEPES; 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

HR; hazard ratio

hs-CRP; high sensitivity C-reactive protein

H<sub>2</sub>S; hydrogen sulfide

ICD-10; International Statistical Classification of Diseases

IQR; interquartile range

PREVEND; Prevention of Renal and Vascular End-stage Disease

SBP; systolic blood pressure

SD; standard deviation

## Tables

**Table 1. Baseline characteristics by gender-stratified quintiles of 24-h urinary sulfate excretion**

	Overall (n=6839)	1 <sup>st</sup> quintile (n=1367)	2 <sup>nd</sup> quintile (n=1368)	3 <sup>rd</sup> quintile (n=1368)	4 <sup>th</sup> quintile (n=1368)	5 <sup>th</sup> quintile (n=1368)	P- value
24-h urinary sulfate, mmol/24 h *	15.7 [12.0-20.3]	♀: < 10.0 ♂: < 12.8	♀: 10.0-12.8 ♂: 12.8-16.2	♀: 12.8-15.3 ♂: 16.2-19.7	♀: 15.4-18.6 ♂: 19.7-23.9	♀: > 18.6 ♂: > 23.9	
<b>Demographics</b>							
Female, n (%)	3420 (50)	684 (50)	684 (50)	684 (50)	684 (50)	684 (50)	<b>1</b>
Age, years	53.4 ± 12.1	55.7 ± 12.9	54.8 ± 12.6	53.5 ± 12.0	52.1 ± 11.6	50.9 ± 10.6	<b>&lt;0.001</b>
BSA, m <sup>2</sup>	1.93 ± 0.20	1.88 ± 0.20	1.9 ± 0.19	1.9 ± 0.18	2.0 ± 0.19	2.0 ± 0.21	<b>&lt;0.001</b>
Systolic blood pressure, mmHg	126 ± 18.9	128 ± 20.5	127 ± 19.5	126 ± 18.6	125 ± 18.5	125 ± 17.2	<b>&lt;0.001</b>
Diastolic blood pressure, mmHg	73.4 ± 9.10	73.6 ± 9.1	73.8 ± 9.1	73.5 ± 9.4	73 ± 9.1	73.3 ± 8.8	0.259
Heart rate, bpm	68.4 ± 10.1	69.4 ± 11.1	68.9 ± 10.2	67.9 ± 10.0	68.1 ± 9.5	68.0 ± 9.9	<b>&lt;0.001</b>
History of	316	94 (7)	62 (5)	71 (5)	49 (4)	40 (3)	<b>&lt;0.001</b>

cardiovascular disease, <i>n</i> (%)	(4.8)							23 1
Diabetes, <i>n</i> (%)	440 (7)	92 (7)	84 (6)	74 (6)	89 (7)	101 (8)	0.264	
Smoking status, <i>n</i> (%)								<0.00 1
Never	1974 (29)	337 (25)	395 (29)	417 (31)	403 (30)	422 (31)		
Former	2903 (43)	460 (34)	576 (43)	594 (44)	614 (45)	659 (49)		
Current	1895 (28)	555 (41)	381 (28)	348 (26)	342 (25)	269 (20)		
Alcohol consumption, <i>n</i> (%)								<0.00 1
Never/hardly ever	1737 (26)	437 (32)	409 (30)	304 (22)	298 (22)	289 (21)		
Up to 1/day	3266 (48)	583 (43)	620 (46)	701 (52)	690 (51)	672 (50)		
> 1/day	1772 (26)	333 (25)	323 (24)	354 (26)	371 (27)	391 (29)		
Medication								
Anti-hypertensive treatment, <i>n</i> (%)	1408 (24)	334 (27)	319 (26)	276 (23)	236 (20)	243 (21)		<0.00 1
Lipid lowering treatment, <i>n</i> (%)	590 (10)	165 (14)	129 (11)	115 (10)	99 (8)	82 (7)		<0.00 1
Anti-diabetic treatment, <i>n</i> (%)	233 (4)	57 (5)	34 (3)	39 (3)	47 (4)	56 (5)	0.052	

### Laboratory measurements

Total cholesterol, mmol/l	5.4 ± 1.1	5.4 ± 1.1	5.5 ± 1.0	5.4 ± 1.0	5.5 ± 1.0	5.5 ± 1.1	0.106
HDL, mmol/l	1.2 ± 0.3	1.2 ± 0.3	1.2 ± 0.3	1.2 ± 0.3	1.3 ± 0.3	1.3 ± 0.3	0.230
Triglycerides, mmol/l*	1.1 [0.8-1.6]	1.2 [0.9-1.7]	1.1 [0.8-1.6]	1.1 [0.8-1.6]	1.1 [0.8-1.6]	1.1 [0.8-1.6]	<b>&lt;0.00</b> <b>1</b>
Glucose, mmol/l*	4.8 [4.4-5.3]	4.8 [4.4-5.4]	4.8 [4.4-5.3]	4.8 [4.4-5.3]	4.8 [4.4-5.4]	4.8 [4.5-5.4]	<b>0.010</b>
hs-CRP, mg/l*	1.4 [0.6-3.1]	1.5 [0.7-3.3]	1.4 [0.7-3.3]	1.3 [0.6-3.1]	1.3 [0.6-2.7]	1.4 [0.6-3.1]	<b>0.003</b>
eGFR, ml/min per 1.73 m <sup>2</sup>	85.8 ± 14.3	84.1 ± 14.9	84.3 ± 15.1	85.9 ± 14.2	87.0 ± 13.5	87.4 ± 13.5	<b>&lt;0.00</b> <b>1</b>
24-h urinary albumin, mg/24 h *	8.7 [6.0-16]	7.7 [5.1- 14.9]	7.9 [5.7- 15.1]	8.7 [6.1- 16.0]	9.1 [6.4- 16.3]	9.9 [6.9- 18.8]	<b>&lt;0.00</b> <b>1</b>
24- h urinary thiosulfate, μmol/24 h *	1.27 [0.89- 2.37]	1.04 [0.73- 1.56]	1.15 [0.83- 1.87]	1.29 [0.9-2.3]	1.44 [1.0-3.0]	1.76 [1.1-3.7]	<b>&lt;0.00</b> <b>1</b>
24-h urinary urea, mmol/24 h	367 ± 125	245 ± 75.6	315 ± 70.7	360 ± 78.2	411 ± 83.7	504 ± 127	<b>&lt;0.00</b> <b>1</b>

Normally distributed continuous data are presented as mean ± SD \*Skewed data are presented as median (IQR)

BSA; body surface area, BMI; body mass index, bpm; beats per min HDL; high-density lipoprotein, hs-CRP; high sensitivity C-reactive protein, eGFR; estimated glomerular filtration rate, SD; standard deviation, IQR; interquartile range



**Table 2. Univariable and multivariable linear regression analyses of 24-h urinary thiosulfate excretion and potential confounders**

	24-h urinary thiosulfate excretion *			
	Univariable regression		Multivariable regression	
	Coefficient	P-value	Coefficient	P-value
<b>Demographics</b>				
Female gender	-0.116	<b>&lt;0.001</b>	-0.043	<b>0.005</b>
Age	-0.019	0.107		
BSA	0.164	<b>&lt;0.001</b>	0.116	<b>&lt;0.001</b>
History of cardiovascular disease	0.024	0.056		
Diabetes	0.072	<b>&lt;0.001</b>	0.056	<b>&lt;0.001</b>
Current smoking	-0.172	<b>&lt;0.001</b>	-0.157	<b>&lt;0.001</b>
> 1 alcoholic consumption/day	0.008	0.51		
* Skewed data, normalized by logarithmic transformation				
BSA; body surface area				

**Table 3. Univariable and multivariable linear regression analyses of 24-h urinary sulfate excretion and potential confounders**

	24-h urinary sulfate excretion *			
	Univariable regression		Multivariable regression	
	Coefficient	P-value	Coefficient	P-value
<b>Demographics</b>				
Female gender	-0.266	<b>&lt;0.001</b>	-0.135	<b>&lt;0.001</b>
Age	-0.110	<b>&lt;0.001</b>	-0.126	<b>&lt;0.001</b>
BSA	0.339	<b>&lt;0.001</b>	0.243	<b>&lt;0.001</b>
History of cardiovascular disease	-0.033	<b>0.007</b>	-0.034	<b>0.004</b>
Diabetes	0.017	0.160		
Current smoking	-0.156	<b>&lt;0.001</b>	-0.148	<b>&lt;0.001</b>
> 1 alcoholic consumption/day	0.064	<b>&lt;0.001</b>	0.041	<b>&lt;0.001</b>

\* Skewed data, normalized by logarithmic transformation

BSA; body surface area

**Table 4. Univariable linear regression analyses of 24-h urinary thiosulfate and sulfate excretion and potential intermediary factors**

	24-h urinary thiosulfate excretion *		24-h urinary sulfate excretion *	
	Coefficient	P-value	Coefficient	P-value
<b>Demographics</b>				
Systolic blood pressure	0.027	<b>0.027</b>	-0.005	0.693
Diastolic blood pressure	0.046	<b>&lt;0.001</b>	0.054	<b>&lt;0.001</b>
Heart rate	-0.015	0.213	-0.084	<b>&lt;0.001</b>
<b>Medication</b>				
Anti-hypertensive treatment	0.015	0.236	-0.051	<b>&lt;0.001</b>
Lipid lowering treatment	-0.003	0.792	-0.052	<b>&lt;0.001</b>
Anti-diabetic treatment	0.058	<b>&lt;0.001</b>	0.012	0.358
<b>Laboratory measurements</b>				
Total cholesterol	-0.023	0.054	0.014	0.263
HDL	-0.044	<b>&lt;0.001</b>	-0.80	<b>&lt;0.001</b>
Triglycerides *	0.092	0.762	-0.004	0.774
Glucose *	0.070	<b>&lt;0.001</b>	0.064	<b>&lt;0.001</b>
hs-CRP *	-0.005	0.685	-0.72	<b>&lt;0.001</b>
eGFR	0.071	<b>&lt;0.001</b>	0.199	<b>&lt;0.001</b>
24-h urinary albumin *	0.042	<b>&lt;0.001</b>	0.124	<b>&lt;0.001</b>
24-h urinary thiosulfate *	-	-	0.289	<b>&lt;0.001</b>
24-h urinary sulfate *	0.289	<b>&lt;0.001</b>	-	-

\* Skewed data, normalized by logarithmic transformation

HDL; high-density lipoprotein, hs-CRP; high sensitivity C-reactive protein, eGFR; estimated glomerular filtration rate

**Table 5. Cox proportional hazard models of the association of 24-h urinary thiosulfate and sulfate excretion and potential confounders with cardiovascular events**

24-h urinary thiosulfate excretion (μmol/24 h)						
Gender-stratified quintiles						
	per doubling	♀: < 0.79	♀: 0.79-	♀: 1.07-	♀: 1.41-	♀: > 2.38
		♂: < 0.84	1.06	1.41	2.38	♂: > 3.31
			♂: 0.84-	♂: 1.15-	♂: 1.74-	♂: > 3.31
			1.15	1.74	3.31	
No. of cases	503	101	113	96	90	103
Person-years	50.452	10.079	10.188	10.066	10.067	10.051
Model 1	1.05 (0.97-	0.98 (0.74-	1.08 (0.83-	0.93 (0.70-	0.87 (0.66-	1.0
	1.13)	1.28)	1.41)	1.23)	1.16)	(reference)
	P=0.258	P=0.858	P=0.567	P=0.611	P=0.346	
Model 2	1.04 (0.96-	0.83 (0.63-	1.03 (0.78-	0.87 (0.66-	0.91 (0.69-	1.0
	1.12)	1.10)	1.34)	1.16)	1.21)	(reference)
	P=0.342	P=0.192	P=0.858	P=0.341	P=0.531	
24-h urinary sulfate excretion (mmol/24 h)						
Gender-stratified quintiles						
	per doubling	♀: < 10.0	♀: 10.0-	♀: 12.8-	♀: 15.4-	♀: > 18.6
		♂: < 12.8	12.8	15.3	18.6	♂: > 23.9
			♂: 12.8-	♂: 16.2-	♂: 19.7-	♂: > 23.9
			16.2	19.7	23.9	
No. of cases	504	131	102	102	82	87

30

Person- years	50.539	9357	10.120	10.155	10.506	10.400
Model 1	0.87 (0.76- 0.99) <b>P=0.039</b>	1.67 (1.27- 2.19) <b>P&lt;0.001</b>	1.20 (0.91- 1.60) P=0.203	1.20 (0.90- 1.60) P=0.212	0.93 (0.69- 1.26) P=0.649	1.0 (reference)
Model 2	0.89 (0.77- 1.03) P=0.107	1.04 (0.78- 1.39) P=0.781	0.82 (0.61- 1.10) P=0.193	0.92 (0.69- 1.23) P=0.557	0.79 (0.59- 1.07) P=0.132	1.0 (reference)

Model 1: crude, Model 2: model 1, adjusted for potential confounders (sex, age, BSA, history of cardiovascular disease, diabetes, current smoking, consumption of more than 1 alcoholic beverage per day)

HR; hazard ratio, CI; confidence interval, BSA; body surface area

**Table 6. Cox proportional hazard models of the association of 24-h urinary thiosulfate and sulfate excretion and potential confounders with all-cause mortality**

24-h urinary thiosulfate excretion (μmol/24 h)						
Gender-stratified quintiles						
		♀: < 0.79	♀: 0.79-	♀: 1.07-	♀: 1.41-	
	per	♂: < 0.84	1.06	1.41	2.38	♀: > 2.38
	doubling		♂: 0.84-	♂: 1.15-	♂: 1.74-	♂: > 3.31
			1.15	1.74	3.31	
No. of cases	445	109	102	84	68	82
Person-years	53.733	10.699	10.923	10.737	10.696	10.678
Model 1	0.93 (0.86-	1.30 (0.97-	1.19 (0.89-	1.01 (0.74-	0.83 (0.60-	1.0 (reference)
	1.01)	1.73)	1.59)	1.37)	1.14)	
	P=0.100	P=0.078	P=0.239	P=0.962	P=0.243	
Model 2	0.94 (0.86-	1.06 (0.79-	1.10 (0.82-	0.91 (0.67-	0.86 (0.62-	1.0 (reference)
	1.02)	1.42)	1.49)	1.24)	1.18)	
	P=0.143	P=0.708	P=0.502	P=0.545	P=0.350	
24-h urinary sulfate excretion (mmol/24 h)						
Gender-stratified quintiles						
		♀: < 10.0	♀: 10.0-	♀: 12.8-	♀: 15.4-	
	per	♂: < 12.8	12.8	15.3	18.6	♀: > 18.6
	doubling		♂: 12.8-	♂: 16.2-	♂: 19.7-	♂: > 23.9
			16.2	19.7	23.9	
No. of cases	445	151	90	81	69	54

32

Person- years	53.847	10.324	10.766	10.817	10.943	10.999
Model 1	0.65 (0.57- 0.74) <b>P&lt;0.001</b>	3.06 (2.25- 4.18) <b>P&lt;0.001</b>	1.72 (1.23- 2.41) <b>P=0.002</b>	1.54 (1.09- 2.17) <b>P=0.014</b>	1.29 (0.91- 1.84) P=0.159	1.0 (reference)
Model 2	0.73 (0.63- 0.84) <b>P&lt;0.001</b>	1.48 (1.07- 2.06) <b>P=0.019</b>	0.98 (0.69- 1.38) P=0.896	1.04 (0.73- 1.47) P=0.831	1.03 (0.72- 1.47) P=0.892	1.0 (reference)

Model 1: crude, Model 2: model 1, adjusted for potential confounders (sex, age, BSA, history of cardiovascular disease, diabetes, current smoking and consumption of more than 1 alcoholic beverage per day)

HR; hazard ratio, CI; confidence interval, BSA; body surface area



**Table 7. Cox proportional hazard models of the association of 24-h urinary sulfate excretion with all-cause mortality, causal path analysis**


Model	24-h urinary sulfate excretion <sup>†</sup>	
	HR (95% CI)	P-value
1	0.73 (0.63-0.84)	<b>&lt;0.001</b>
2	0.75 (0.65-0.86)	<b>&lt;0.001</b>
3	0.72 (0.63-0.83)	<b>&lt;0.001</b>
4	0.75 (0.65-0.86)	<b>&lt;0.001</b>
5	0.73 (0.63-0.84)	<b>&lt;0.001</b>

<sup>†</sup>HR per doubling \* Skewed data, normalized by logarithmic transformation

Model 1: adjusted for potential confounders (sex, age, BSA, history of cardiovascular disease, diabetes, current smoking, consumption of more than 1 alcoholic beverage per day), Model 2: model 1, additionally adjusted for hemodynamic parameters (SBP, DBP, heart rate and anti-hypertensive treatment), Model 3: model 1, additionally adjusted for lipid profile (total cholesterol, HDL and triglycerides, lipid lowering treatment), Model 4: model 1, additionally adjusted for hs-CRP, Model 5: model 1, additionally adjusted for eGFR


HR; hazard ratio, CI; confidence interval, BSA; body surface area, BMI; body mass index, SBP; systolic blood pressure, DPB; diastolic blood pressure, HDL; high-density lipoprotein, hs-CRP; high sensitivity C-reactive protein, eGFR; estimated glomerular filtration rate

Figure legends




**Figure 1. Survival distributions of urinary thiosulfate excretion quintiles**

**A.** Gender-stratified quintiles of 24-h urinary thiosulfate excretion are differentially distributed among survivors and non-survivors with overrepresentation of the lowest quintile in the non-survivors population ( $\chi^2$  test,  $P=0.011$ ). **B.** Kaplan-Meier plot with log-rank test for survival showing the highest mortality rate for the lowest quintile of thiosulfate excretion ( $P=0.034$ ). (To see this illustration in color the reader is referred to the web version of this article at [www.liebertonline.com/ars](http://www.liebertonline.com/ars))




**Figure 2. Survival distributions of urinary sulfate excretion quintiles**

**A.** Gender-stratified quintiles of 24-h urinary sulfate excretion are differentially distributed among survivors and non-survivors with overrepresentation of the lowest quintile in the non-survivors population ( $\chi^2$  test,  $P < 0.001$ ). **B.** Kaplan-Meier plot with log-rank test for survival showing the highest mortality rate for the lowest quintile of sulfate excretion ( $P < 0.001$ ). (To see this illustration in color the reader is referred to the web version of this article at [www.liebertonline.com/ars](http://www.liebertonline.com/ars))



**Figure 3. No deviances from linear associations with all-cause mortality for urinary thiosulfate and sulfate excretion**

Unadjusted associations estimated by Cox proportional hazards analyses based on restricted cubic splines show no deviances from linear associations for 24-h urinary thiosulfate and sulfate excretion ( $P_{\text{nonlinearity}}=0.08$  and  $0.10$ , respectively). Median 24-h urinary thiosulfate and sulfate excretion ( $1.27 \mu\text{mol}$  and  $15.7 \text{ mmol}/24 \text{ h}$ , respectively) are the reference standards. Grey area indicate 95% confidence intervals.



**Figure 4. Associations between urinary sulfate excretion and all-cause mortality in different subgroups**

Hazard ratios for all-cause mortality in various subgroups, consistently showing positive associations between 24-h urinary sulfate excretion and survival. Only cholesterol levels (normo- vs. dyslipidemia) show significant interaction ( $P_{\text{interaction}}=0.019$ ). Stratifications by renal function ( $\text{eGFR} \geq 60$  vs.  $< 60$  ml/min) and the presence of diabetes (no diabetes vs. diabetes) reveal borderline significant differences between groups ( $P_{\text{interaction}}=0.057$  and  $P_{\text{interaction}}=0.054$ , respectively). Corresponding hazard ratios are lower for subjects with normolipidemia, subjects with impaired renal function ( $\text{eGFR} < 60$  ml/min) and subjects without diabetes.

\*After adjustment for potential confounders (sex, age, BSA, history of cardiovascular disease, diabetes, current smoking and consumption of more than 1 alcoholic beverage per day)