

**Title:** Grafted human iPSC-derived neural progenitor cells express integrins and extend long-distance axons within the developing corticospinal tract.

**Authors:** Lindsey H Forbes<sup>1</sup> & Melissa R Andrews<sup>1, 2\*</sup>

<sup>1</sup> School of Medicine, University of St Andrews, St Andrews, United Kingdom

<sup>2</sup> Biological Sciences, University of Southampton, Southampton, United Kingdom

**\*Corresponding Author:** Melissa R Andrews; [m.r.andrews@soton.ac.uk](mailto:m.r.andrews@soton.ac.uk)

Melissa R Andrews

Biological Sciences

University of Southampton

3059, Life Sciences Bldg 85

Highfield Campus

Southampton, SO17 1BJ

United Kingdom

+44 (0)2380 592432

Number of words: 6065

Number of figures: 11

**Highlights**

- Increasing  $\alpha 9$  integrin expression in human iPSC-derived NPCs increases neurite outgrowth *in vitro*.
- Human iPSC-derived NPCs transplanted into the rat developing sensorimotor cortex project within and follow the endogenous CST.
- Exogenous expression of  $\alpha 9$  integrin is retained within the axonal compartment of human iPSC-derived NPCs up to 8 weeks following transplantation.

**Keywords**

$\alpha 9$  integrin, induced pluripotent stem cell, transplantation, neural progenitor cell, sensorimotor cortex.

**Abstract**

After spinal cord injury, regeneration of adult motor axons such as axons in the corticospinal tract (CST) is severely limited. Alongside the inhibitory lesion environment, most neuronal subtypes in the mature CNS are intrinsically unrepairable. With age, expression of growth-promoting proteins in neurons, such as integrins, declines. Integrin receptors allow communication between the extracellular matrix and cell cytoskeleton and their expression in axons facilitates growth and guidance throughout the ECM. The  $\alpha 9\beta 1$  integrin heterodimer binds to tenascin-C (TN-C), an extracellular matrix glycoprotein expressed during development and after injury. In the mature CST however, expression of the  $\alpha 9$  integrin subunit is downregulated, adding to the intrinsic inability of axons to regenerate. Our previous work has shown the  $\alpha 9$  integrin subunit is not trafficked within axons of mature CST or rubrospinal tracts. Thus, here we have utilized human induced pluripotent stem cell (iPSC)-derived neural progenitor cells (NPCs) to increase expression of  $\alpha 9$  integrin within the developing rat CST.

We demonstrate that hNPCs express endogenous levels of both  $\alpha 9$  and  $\beta 1$  integrin subunits as well as cortical neuron markers such as Ctip2 and Tbr1. In addition, lentivirus-mediated  $\alpha 9$  integrin overexpression in hNPCs resulted in increased neurite outgrowth in the presence of TN-C *in vitro*. Following transplantation into the sensorimotor cortex of newborn rats, both WT and  $\alpha 9$ -expressing hNPCs extend along the endogenous CST and retain expression of  $\alpha 9$  throughout the length of the axonal compartment for up to 8 weeks following transplantation. These data highlight the growth potential of transplanted human iPSCs which may be a future target for regenerative therapies after nervous system injury.

## Introduction

Following injury to the mature central nervous system (CNS), the repair capacity is limited due to a myriad of intrinsic and extrinsic factors within the lesion site. After spinal cord injury (SCI), a characteristic glial scar densely packed with chondroitin sulphate proteoglycans (CSPGs) forms preventing further damage but also preventing regrowth of damaged axons (Reviewed in Silver and Miller, 2004; Reviewed in Tran et al., 2018). In addition, other inhibitory extracellular proteins including the myelin-associated protein, Nogo-A, are present and actively prevent regrowth (Reviewed in Schwab and Strittmatter, 2014). Conversely, there is a downregulation of growth-promoting proteins, such as integrins (Bi et al., 2001; Condic, 2001; Franssen et al., 2015; Reviewed in Nieuwenhuis et al., 2018) and Trk receptors (Lu et al., 2001), within mature CNS axons resulting in a poor neuronal regenerative response. Integrin receptors, known mediators of cell-cell and cell-matrix interactions, are transmembrane receptors comprised of one  $\alpha$  and one  $\beta$  subunit. Within the developing CNS, integrins regulate cell cytoskeleton dynamics resulting in neurite outgrowth, axonal elongation and migration. Specifically, the  $\alpha 9$  subunit, which forms a functional heterodimer with the  $\beta 1$  subunit ( $\alpha 9\beta 1$ ) binds tenascin-C (TN-C), the predominant extracellular matrix (ECM) glycoprotein in the CNS. Although several integrins including  $\alpha 9$  are downregulated in the adult CNS, re-expression of integrins *in vitro* and *in vivo* rescues this inhibition resulting in increased axonal growth in the presence of inhibitory ECM proteins (Condic, 2001), including TN-C (Andrews et al., 2009; Cheah et al., 2016) which is secreted by reactive astrocytes. Recently, however, we have demonstrated that overexpressed integrin subunits (via viral vectors) are not transported within axons of the adult corticospinal or rubrospinal tract (CST and RST, respectively) *in vivo* (Andrews et al., 2016) presenting a challenge for gene therapy-mediated transmembrane receptor expression.

The field of regenerative medicine has also taken significant advantage of the recent discovery and development of induced pluripotent stem cell (iPSC) technology giving rise to infinite cell sources with high growth potential. Specifically, iPSCs and the numerous cell types which have successfully been derived from them, have great potential in the field of CNS regeneration whether through direct cell replacement and/or creation of a pro-regenerative environment (Lu et al., 2012; 2014; Nori et al., 2011; Tornero et al., 2013). In the current study, we use human iPSC-derived neural progenitor cells (hNPCs) as a vehicle to enhance  $\alpha 9$  integrin expression within the CST following transplantation into the developing sensorimotor cortex. We show iPSC-hNPCs express a basal level of  $\alpha 9$  integrin that can be augmented using lentiviral transduction. This overexpression leads to a significant increase in neurite outgrowth of cultured  $\alpha 9$ -hNPCs when grown on a TN-C substrate compared to controls. Following transplantation into the naïve sensorimotor cortex of neonatal rats, we demonstrate that both  $\alpha 9$ -hNPCs and wild type (WT) hNPCs survive for up to 8 weeks and extend axons within the CST reaching the pyramids within the medulla. Together these data highlight the ability of

human iPSC-derived NPCs to develop and integrate within the rodent CNS as well as increase integrin activity within the CST that may contribute to future repair of the injured CNS.

## Methods

### Culture of human iPSC-derived NPCs

Human iPSC-derived NPCs (Axol Bioscience) were cultured as per manufacturer's instructions with some modifications. Briefly, cells were cultured on 20µg/mL poly-L-ornithine (PLO) (Sigma) and 10µg/mL laminin (Sigma) at a density of  $5 \times 10^4/\text{cm}^2$ . Cells were maintained in Neural Maintenance Medium (Axol Bioscience) and passaged using StemPro Accutase (Gibco™). For ICC analysis, cells were cultured on acid-washed glass coverslips coated with PLO and laminin as above.

### Production of 2<sup>nd</sup> generation lentivirus

HEK293T producer cells were cultured in 10 cm plates (Nunc) and transfected with three second generation plasmids [5µg psPAX2, 2.1µg pMD2.G-VSV.G (Naldini et al., 1996) and either 10µg LV-PGK-α9-eYFP (Andrews et al., 2016) or 6.5µg LV-CMV-farnesylated GFP or fGFP (Andrews et al., 2009)] using TransIT®LT-1 transfection reagent (Mirus) at a ratio of 3:1 (reagent [µL]: DNA [µg]). The transfection mix was incubated with the producer cells for 24 hours before being replaced with fresh complete DMEM (10% FBS [Seralab] and 1% Pen/Strep [Gibco™] in DMEM [Gibco™]). Viral supernatant was collected from cells at 48 hours and 72 hours after initial transfection.

### Transduction of iPSC-derived hNPCs

72 hours after thawing, iPSC-derived hNPCs (Axol Bioscience) were transduced with either LV-α9-eYFP or LV-fGFP supernatant, containing 2µg/mL hexadimethrine bromide (polybrene, Sigma) for 4 hours at 37°C. Following incubation, viral media was removed and cells were washed with Neural Maintenance Media, followed by incubation with fresh Neural Maintenance Media. Cells were cultured for a further 4-5 days before collection for transplantation, ICC analysis, or neurite outgrowth assays.

### Immunocytochemistry

Cells were fixed using 4% paraformaldehyde (PFA) and coverslips were washed in triplicate with PBS before being incubated with blocking solution (10% goat serum in PBS). Coverslips were incubated with primary antibodies diluted in blocking solution overnight at 4°C. The following primary antibodies were used: rabbit anti-BDNF, rabbit anti-Ctip2, rabbit anti-DCX, rabbit anti-GFAP, rabbit anti-MBP, rabbit anti-Tbr1, rabbit anti-tenascin-C, rabbit anti-TrkB, rabbit anti-α9 integrin, mouse anti-β1 integrin, mouse anti-vinculin, mouse anti-βIII-tubulin and rabbit anti-βIII-tubulin (see Table 1 for concentrations). The following day, coverslips were incubated with secondary antibodies (at 1:1000 dilution; goat anti-mouse AlexaFluor 488 or 568, goat anti-rabbit AlexaFluor 488 or 568; Invitrogen), and/or phalloidin (AlexaFluor 647 phalloidin; 1:250; Invitrogen) for 2 hours. Cell nuclei were stained using DAPI (1:10,000; Thermo Scientific™) for 10 minutes before coverslips were mounted onto slides with FluorSave (Calbiochem) and imaged using a Leica D5500 epifluorescent microscope (Leica) with an attached DFC550 camera (Leica).

### Western blotting

Cell lysates were collected from hNPC cultures using 1X radioimmunoprecipitation assay (RIPA) buffer (containing 50nM Tris, 150mM NaCl, 1% NP-40 and 0.5% sodium

deoxycholate) containing 25X complete protease inhibitor (Roche). Protein sample concentrations were determined using a bicinchoninic acid (BCA) assay kit (Thermo Scientific<sup>TM</sup>) as per manufacturer's instructions. Protein samples containing 40µg protein diluted in water and NuPage loading dye were separated by electrophoresis on a 4-12% bis-tris pre-cast gel (Invitrogen). Protein samples were denatured at 90°C for 10 minutes and were run alongside a PageRuler Plus Prestained protein ladder (Thermo Fisher). The gel was run with 1X NuPage MOPS SDS buffer and transferred to a nitrocellulose membrane (GELifesciences) with 1X NuPage transfer buffer. The membranes were blocked in 5% milk powder and stained overnight at 4°C with the following primary antibodies: rabbit anti-α9 integrin, mouse anti-β1 integrin, rabbit anti-GFP, and mouse anti-β-actin (see Table 1 for concentrations). The corresponding HRP secondary antibodies (HRP-mouse and HRP-rabbit, Invitrogen) were used at a dilution of 1:15,000 before analysis using enhanced chemiluminescence (ECL, GE Healthcare) and imaging with a LAS-3000 Lite imaging system. Images were processed and analyzed using Image J.

### **Neurite outgrowth assays and analysis**

Neurite outgrowth assays were carried out on Permanox<sup>®</sup> 8-well chamberslides (Thermo Scientific<sup>TM</sup>). Wildtype, α9-hNPCs and GFP-hNPCs were cultured on 20µg/mL PLO and 10µg/mL chicken TN-C (n=4) or varying concentrations of human TN-C (Millipore): 1µg/mL (n=3), 5µg/mL (n=3) or 10µg/mL (n=3). Cells were plated at a density of 5x10<sup>4</sup>/well and incubated for 72 hours at 37°C. The cells were fixed using 4% PFA and stained using the primary antibodies anti-GFP and anti-βIII-tubulin. Analysis and tracing of neurite outgrowth was performed using Neuron J (Meijering et al., 2004; Schneider et al., 2012) and statistical analysis was performed using Microsoft Excel and Prism 5 GraphPad. A total of 30 neurite/axon lengths were measured for each condition. One replicate consisted of one vial of hNPCs cultured in one chamberslide (with 2 wells/condition). Data sets were analyzed using one-way analysis of variance (ANOVA) with Tukey post hoc.

### **Preparation of cells for transplantation**

For transplantation, WT hNPCs or α9-hNPCs were collected at day 8-9 in culture using StemPro Accutase, resuspended at a density of 1x10<sup>6</sup>/µL in Neural Maintenance Medium supplemented with Sure Boost (Axol Bioscience) and stored at RT until use.

### **Transplantation of WT and α9-eYFP-expressing hNPCs**

All experimental and surgical procedures were conducted in accordance with the UK Animals (Scientific Procedures) Act 1986. Newborn Sprague-Dawley rats received transplants of either WT hNPCs (n=26) or α9-eYFP-expressing hNPCs (n=17). Rat pups between the age of postnatal day 0-2 (P0-P2) were anesthetized on ice for 3-5 minutes. Pups were transferred to a surgical platform containing ice to maintain anesthesia. For surgery, each pup was secured using a custom-made 3D-printed rat pup head frame (kindly printed by Dr Robert Hammond, Univ of St Andrews). Using stereotaxic coordinates, each pup received either a unilateral or bilateral transplant in two specific sites with 1x10<sup>6</sup> cells per target site (1µL/injection) into layer 5 of the neonatal sensorimotor cortex. Coordinates that produced the most on-target injections were: AP 0.0 mm, ML 1.5/-1.5 mm, DV -0.4 mm. Cells were injected manually using a Hamilton syringe with a custom-made 30gauge stainless steel needle over the course of 1 minute with the needle left in place for a further 2 minutes to prevent backflow. Following surgery, pups were transferred to a heated cage to recover before being returned to their mother. On-target injections included pups where a noticeable scar was evident within the target site. All pups with off-target injections were excluded from the study.

## Tissue perfusion and sectioning

Tissue was collected at the following time points: 2 weeks (n=9); 4 weeks (n=7\*); 6 weeks (n=9); 7 weeks (n=10) and 8 weeks (n=8) (Table 2; \* indicates animals at 4 weeks from WT group only). Any rats with off-target injections are not included in Table 2. Rats were administered a terminal dose of sodium pentobarbital and transcardially perfused with 0.9% saline solution (pH 7.4) and 4% PFA (pH 7.4). The brain and spinal cord were removed from each animal, post-fixed in 4% PFA for 48-72 hours and cryoprotected in 30% sucrose solution for a further 48-72 hours. Tissue was cryo-sectioned in either the coronal or sagittal plane using a sliding microtome (Leica) at a thickness of 40µm. Sections were stored in 0.05% sodium azide in PBS at 4°C prior to processing for immunohistochemistry (IHC).

## Immunohistochemistry

Sections were incubated in blocking solution for 2 hours at RT and then with primary antibody overnight at 4°C. Primary antibodies used were as follows: rabbit anti-DCX, rabbit anti-GFAP, rabbit anti-GFP, mouse anti-HuNu, mouse anti-hNCAM, rabbit anti-TN-C and rabbit anti-Tbr1 (see Table 1 for full details). Tissue sections were incubated with the corresponding secondary antibodies (all at a dilution of 1:750; goat anti-mouse AlexaFluor 488 or 568, goat anti-rabbit AlexaFluor 488 or 568; Invitrogen) for 2 hours at RT. All nuclei were stained using DAPI (1:10,000; Thermo Scientific™). Tissue sections were then mounted onto 0.25% gelatin-coated slides and coverslipped using FluorSave (Calbiochem). Slides were imaged using a Leica D5500 epifluorescent microscope (Leica) with an attached DFC550 camera (Leica). Images were processed and collated using Image J and Microsoft Powerpoint.

## Results

### Human iPSC-derived NPCs express growth-promoting proteins *in vitro*

A primary goal of this study was to characterize iPSC-derived hNPCs *in vitro* by analyzing expression of neuronal and growth-promoting proteins as well as neurite outgrowth capacity on TN-C. To analyze endogenous integrin expression ( $\alpha 9$  and  $\beta 1$  subunits) within the hNPCs, both western blotting (WB) and immunocytochemistry (ICC) were used. WB data indicated expression of both  $\alpha 9$  (Fig. 1A) and  $\beta 1$  (Fig. 1B) integrin subunits in iPSC-derived hNPCs *in vitro* which was further confirmed by ICC (Fig. 1C-F). Interestingly, cultured hNPCs also express an endogenous level of TN-C (Fig. 1G, H), the ligand for the  $\alpha 9\beta 1$  heterodimer, a finding previously shown with PC12 cells (Andrews et al., 2009).

Alongside integrin expression, further ICC analysis revealed iPSC-derived hNPCs express the progenitor cell marker doublecortin (DCX) (Fig. 1I, J) and the growth-promoting proteins including brain derived neurotrophic factor (BDNF) (Fig. 1K, L) and its receptor TrkB (Fig. 1M, N). The hNPCs used within these experiments were pre-programmed to differentiate into cerebral cortical neurons (Shi et al., 2012b), however further cell types could be detected within the culture including a small proportion of GFAP-positive cells (approximately 5%) indicative of astrocytes (Fig. 1P). No cells were positive for myelin basic protein (MBP) following 21 days in culture (Fig. 1Q). The majority of the culture, however was positive for  $\beta$ III-tubulin (approximately 95%) (Fig. 1O) indicating a high proportion of cells had a neuronal phenotype. To confirm previous literature (Shi et al., 2012b) for this cell population, the hNPCs were analyzed for the expression of deep layer cortical neuron markers Ctip2 and Tbr1, with a high proportion (approximately 70%) of the cell population expressing these proteins within the cell nucleus (Fig. 1R-W).

## **Increasing $\alpha 9$ integrin expression in hNPCs results in increased neurite outgrowth in the presence of TN-C *in vitro***

In addition to demonstrating endogenous expression of the  $\alpha 9$  integrin subunit, we also overexpressed  $\alpha 9$  integrin (tagged to eYFP) within iPSC-derived hNPCs using lentiviral transduction, resulting in an approximate 90% transduction efficiency. As a control, expression of fGFP was used in addition to untransduced WT hNPCs. Following ICC analysis for anti-GFP, expression of both  $\alpha 9$ -eYFP and fGFP were observed within the cell bodies and neurites of the hNPCs (Fig. 2A-F), which was further confirmed by WB (Fig. 2G).

In order for cells to migrate and axons to extend as well as interact with the extracellular environment, integrins need to be enriched at nascent focal adhesion sites. This is known as integrin clustering, a process that prompts intracellular signaling which coincides with cytoskeletal reorganization. Following overexpression of  $\alpha 9$ -eYFP within hNPCs, the extending neurites displayed dense enrichment of  $\alpha 9$ -eYFP *in vitro* (Fig. 2H-J). Furthermore, localization of  $\alpha 9$  integrin-eYFP was observed in vinculin- and actin-rich focal adhesion sites (Supplemental Figure 1). In addition, we have observed a subtle enhancement of  $\beta 1$  integrin expression upon overexpression of  $\alpha 9$  integrin in hNPCs, evaluated by western blot (Supplemental Figure 2). This suggests that  $\alpha 9$  overexpression may lead to increased  $\alpha 9\beta 1$  heterodimer formation and/or upregulation of  $\beta 1$  integrin subunits.

The function of exogenously expressed  $\alpha 9$ -eYFP protein in hNPCs was analyzed using a neurite outgrowth assay in the presence of either chicken or human TN-C, at varying concentrations after 72 hours in culture. As previously shown, inducing  $\alpha 9$  integrin expression in PC12 cells stimulates neurite outgrowth in the presence of TN-C, an inhibitory growth substrate for cells lacking  $\alpha 9$  integrin (Andrews et al., 2009). On chicken TN-C (10 $\mu$ g/mL), a significant difference in neurite outgrowth was observed for LV- $\alpha 9$ -hNPCs compared to both fGFP-expressing hNPCs ( $P < 0.001$ ) and WT hNPCs ( $P < 0.001$ ) (Fig. 3A-C, M, Q). A small amount of neurite outgrowth was observed from the two control groups grown on TN-C (Fig. 3B, C, M), likely due to low levels of endogenous  $\alpha 9$  and  $\beta 1$  integrin expression (Fig. 1). To determine whether the hNPCs showed any specificity for human TN-C, neurite outgrowth was analyzed on varying concentrations of human TN-C (1 $\mu$ g/mL, 5 $\mu$ g/mL and 10 $\mu$ g/mL). Results from these assays show an increase in neurite outgrowth from the LV- $\alpha 9$ -hNPCs as compared to the other groups most significantly with 1 $\mu$ g/mL human TN-C (Fig. 3D-L, N-Q) further highlighting the ability of  $\alpha 9$  expression to increase outgrowth in the presence of TN-C. However, as the concentration of human TN-C was increased to 5 $\mu$ g/mL and 10 $\mu$ g/mL, the outgrowth from both the fGFP-hNPCs and WT hNPCs also increased, while the neurite outgrowth in the  $\alpha 9$ -hNPCs group did not increase further. Specifically, neurite outgrowth of fGFP-hNPCs grown on 1 $\mu$ g/mL human TN-C was found to be 48.8  $\mu$ m  $\pm$  1.1 ( $n = 3$ ), WT hNPCs 49.1  $\mu$ m  $\pm$  4.1 ( $n = 3$ ) and  $\alpha 9$ -eYFP-hNPCs 226.0  $\pm$  17.2 ( $n = 3$ ), whereas with 10 $\mu$ g/mL human TN-C, the outgrowth from both fGFP-hNPCs and WT hNPCs more than doubled to 108.1  $\mu$ m  $\pm$  22.5 and 103.0  $\mu$ m  $\pm$  25.9, respectively, whilst outgrowth from  $\alpha 9$ -hNPCs remained relatively constant at 220.4  $\mu$ m  $\pm$  31.6 (Fig. 3O-Q).

## **Human iPSC-derived NPCs transplanted into the developing rat sensorimotor cortex project within the intrinsic pyramidal tract**

We next assessed the ability of hNPCs to survive, express exogenous growth-promoting integrin receptors and extend axonal processes following transplantation into the rat neonatal sensorimotor cortex. Regeneration of the adult CST, one of the main pathways governing motor control, has been shown to have particularly limited regenerative capacity (Reviewed in Tuszynski and Steward, 2012). We recently demonstrated that axonal localization of growth-promoting integrins, including  $\alpha 9$ , is hindered within the mature CST compared to the developing CST and sensory pathways (Andrews et al., 2016).

In this study, pups aged between P0-P2 received grafts of either WT hNPCs or  $\alpha 9$  hNPCs into layer 5 of the sensorimotor cortex and tissue was analyzed at 2, 4, 6, 7 and 8 weeks post-transplantation. Injections that were off-target, specifically where no part of the hNPC bolus was observed within the target site, have been excluded from the study. This was determined by identifying the deep layers of the sensorimotor cortex which was immunopositive for Tbr1-expressing cells (Supplementary Fig. 3). Grafts were detected using human-specific antibodies for human nuclear antigen (HuNu), to identify the nuclei of the hNPCs (Fig. 4A, B), and human neural cell adhesion molecule (hNCAM) to identify the projections emanating from the graft site (Fig. 4C-H, J, J\*). In 20.9% of pups (n = 9 out of the total 43), most of which were from the later time points (6, 7 and 8 weeks), no surviving hNPCs were detected and instead an injection scar was present at the intended injection site, identified by an upregulation of GFAP staining (data not shown).

In the remaining 79.1% of pups (n=34), graft survival of iPSC-derived hNPCs into the sensorimotor cortex was observed from 2 weeks up to 8 weeks. Across all time points, the formation of the cell bolus varied from one which spread out (Fig. 4A) to a bolus that was more densely packed (Fig. 4B). In some grafts, rosette formation was observed, a common feature of neural stem cells (Fig. 5) (Shi et al., 2012a). In addition, there was an extensive number of hNCAM-positive axonal projections observed within regions of the intrinsic corticospinal pathway including the internal capsule, cerebral peduncles and areas of the brain stem, specifically in the pons and pyramids (Fig. 4). Furthermore, there was variation in thickness of hNCAM-positive projections. Some projections appeared very narrow and thin (Fig. 4C, D), whilst others appeared not as single fibres but as thicker bundles (Fig. 4G) suggesting fibres may have bundled together over time. In following the trajectory of the hNCAM-positive axonal projections at the 2 week time point, axonal projections were found within the pyramids, specifically within the PKC $\gamma$ -positive region used to identify the pyramidal tract (Fig. 4I) (Bradbury et al., 2002). More specifically, when analyzing WT hNPCs grafts, a total of 3.8% (n=1) of WT hNPCs showed projections terminated within the medullary pyramids. Similarly, a total of 3.8% (n=1) of grafts showed projections terminated within the pons. A further 23.1% (n=6) of grafts showed hNCAM-positive projections as far as the cerebral peduncles whilst 34.6% (n=9) showed projections reaching the internal capsule. In 23.1% (n=6) of grafts, there projections only found emanating within the localised region of the cell bolus within the cortex, whilst a further 11.5% (n=3) showed no NCAM-positive projections due to potential degeneration of the cell bolus. Interestingly, we did not find projections reaching the pyramids at later time points (4, 6, 7, or 8 weeks).

As this was a xenogeneic transplant, the host immune response was analyzed following grafting of hNPCs using an antibody against IBA-1, a marker for microglia, the main immune cell of the CNS and against GFAP, to identify astrocytes which can become reactive with an immune response. At early time points of 2 weeks (Fig. 6A-D), microglial activity was increased slightly compared to control tissue (contralateral hemisphere, Fig. 8I) for both WT and  $\alpha 9$ -eYFP hNPC grafts. Over time however, microglial activity further increased demonstrated with a clear increase in positive IBA-1 staining at the graft site (Fig. 6K-N). GFAP immunoreactivity showed a similar pattern of activity for astrocytes surrounding the graft site. At 2 weeks post-transplantation, GFAP expression was mildly increased (Fig. 6E-H) compared to control tissue (Fig. 6J) and by 8 weeks this was markedly increased (Fig. 6O-R and T) indicative of reactive astrocytes at the graft site. Furthermore, there was no indication that the small population of astrocytes (~5%) that were initially present in the hNPC cultures contributed to the GFAP-immunoreactivity with minimal to no overlap in staining between transplants and GFAP (Supplemental Fig. 4). The expression of the ECM protein TN-C within the tissue was also analyzed following transplantation of both WT hNPCs and  $\alpha 9$ -eYFP-

expressing hNPCs. Similar to the staining observed with GFAP and IBA-1, expression of TN-C was increased at the site of transplant across all time points (Fig. 6U-W).

To further characterize iPSC-hNPC transplants *in vivo*, expression of cortical neuron markers were assessed. We have shown cortical neuron markers were expressed *in vitro* prior to transplantation (Fig. 1). IHC analysis confirmed this expression was maintained following transplantation, with a large proportion of the cells retaining their cortical identity and expressing the deep-layer cortical neuron marker, Tbr1, from 2 weeks up until 8 weeks (the longest time point analyzed) (Fig. 7A, B). Furthermore, we also investigated whether these cells matured or remained immature over the 2 month time period of analysis, using the progenitor cell marker doublecortin (DCX). The hNPCs retained DCX expression as long as the cells were present suggesting that at least some of the cells remained in a progenitor cell state over a period of 2 months post-transplantation (Fig. 7C-H). When endogenous expression of rat DCX was analyzed within the cortex, expression was high at 2 weeks postnatally/post-transplant, yet declined thereafter (data not shown).

### **Exogenously expressed $\alpha 9$ integrin is retained in transplanted hNPCs and localized throughout the axonal compartment *in vivo***

We next investigated the extent to which axons from transplanted  $\alpha 9$ -expressing hNPCs could project within a developing neonatal system whilst maintaining integrin expression within axons. Analysis of  $\alpha 9$ -eYFP hNPC grafts identified that  $\alpha 9$ -eYFP was expressed throughout the projected axons in small punctate dots (Fig. 8) at each time point from 2 weeks to 8 weeks indicating axonal expression, transport and/or targeting of this exogenous protein.

Further analysis of  $\alpha 9$ -eYFP hNPC grafts confirmed that fibers projecting from these cells followed the intrinsic CST from the sensorimotor cortex through the internal capsule, cerebral peduncles and pons. Specifically, the transplanted  $\alpha 9$ -hNPCs reached the pyramids by 2 weeks post-transplantation in the best case, comparable to WT hNPCs (Fig. 9). Interestingly at later time points, similar to WT hNPCs, axons projecting from these cells did not project as far as they had at 2 weeks post-transplantation and in many cases  $\alpha 9$ -hNPCs did not project axons as far as WT hNPCs (Fig. 9). For example, at 6 weeks, WT hNPC projections were identified within the pons whereas  $\alpha 9$ -eYFP hNPCs only reached the cerebral peduncles. Similarly, at 7 weeks post-transplantation, WT hNPCs projected axons to the cerebral peduncles while  $\alpha 9$ -eYFP hNPCs only projected as far as the internal capsule (Fig. 9). These results suggest that endogenous levels of integrins are sufficient to induce significant levels of axonal growth likely because the nervous system as well as the transplanted cells are still undergoing development. In further analysis of  $\alpha 9$ -eYFP hNPC grafts, a total of 5.9% (n=1) showed projections within the medullary pyramids. No projections were observed within the pons yet 11.8% (n=2) of grafts showed hNCAM-positive projections within the cerebral peduncles. In 29.4% (n=5) of the grafts, there were projections within the internal capsule whilst in 17.6% (n=3) of the grafts there were projections only within the localized cortical region of the cell bolus. A further 35.3% (n=6) showed no hNCAM-positive projections likely due to the degeneration of the cell bolus.

## **Discussion**

Stem cell research has radically changed regenerative medicine, even more so since the discovery of iPSC technology (Takahashi and Yamanaka, 2006; Takahashi et al., 2007). In addition, we now have valuable insight into the underlying mechanisms that prevent and contribute to regeneration of human axons following CNS injury. Specifically, it has been



shown that the failure of CST axon regeneration is partly due to a loss of anterograde trafficking of growth-promoting proteins including the  $\alpha 9$  integrin subunit as demonstrated in cultured cortical neurons (Franssen et al., 2015). The present study demonstrates that human iPSC-derived NPCs (hNPCs) can increase expression of  $\alpha 9$  integrin within the developing rodent CST. We have carried out extensive characterization of these cells both *in vitro* and following transplantation into the developing sensorimotor cortex. These cells express low levels of the integrin subunits,  $\alpha 9$  and  $\beta 1$ , as well as other growth-promoting proteins including BDNF and TrkB, and the cortical neuron markers Tbr1 and Ctip2. Furthermore, we have generated  $\alpha 9$ -eYFP-expressing hNPCs and demonstrated the functional effect of the  $\alpha 9$  receptor in the presence of TN-C *in vitro*. Following transplantation, hNPCs retained expression of  $\alpha 9$ -eYFP up to 8 weeks post-transplantation, the time point after which the transplanted cells died. After only 2 weeks post-transplantation, we demonstrate that these cells project axons within the endogenous CST reaching as far as the medullary pyramids. Due to the growth-promoting properties of  $\alpha 9$ -eYFP, we initially hypothesized that  $\alpha 9$ -eYFP-expressing hNPCs would project longer axons than WT hNPCs, however the presence of  $\alpha 9$ -eYFP did not result in longer axonal projections *in vivo* compared to WT hNPCs. These results suggest that endogenous levels of integrins are sufficient to induce axonal growth from hNPCs within a developing nervous system.

### Endogenous protein expression in hNPCs

A large proportion of the cultured hNPCs were  $\beta$ III-tubulin-positive confirming a high proportion of neurons within these cultures. Although these cells were pre-programmed to differentiate into cerebral cortical neurons, other cell types such as GFAP-expressing cells were identified within the cell population suggesting that these hNPC cultures may give rise to a small proportion (~5%) of astrocytes. For the purposes of transplantation, the presence of astrocytes may have been advantageous for hNPC survival and integration *in vivo*. For example, astrocytes can promote neuron function and synaptogenesis (Ullian et al., 2004; 2001; Christopherson et al., 2005).

It was also critical that the expression profile of the cells have a cortical neuron phenotype which is important for their development, integration and function (Finger et al., 2002; Arlotta et al., 2005; Sano et al., 2017) and because we were transplanting these cells into the cortex these proteins are required to induce growth within the CST. Specifically, when transplanting cells into the lesioned cortex, research has shown that receiving an area-specific graft can impact the level of integration and regeneration observed (Gaillard et al., 2007; Michelson et al., 2015). Previous literature has shown expression of cortical neuron markers within iPSC-hNPCs, such as Tbr1 and Ctip2 (Shi et al., 2012b), which our results confirm *in vitro*. Specifically, these proteins are involved in projection of neurons from deep-layers of the cortex to sub-cortical targets (Hevner et al., 2001; Arlotta et al., 2005; Chen et al., 2008; McKenna et al., 2011). Of particular interest is Ctip2, which is known to be involved in formation of CST projections during development while perturbation of its expression can result in aberrant CST formation (Chen et al., 2008). Following transplantation, we were also able to confirm hNPCs retained Tbr1 expression over time. Due to a lack of consistency with Ctip2 antibodies *in vivo*, however, we were unable to clearly ascertain Ctip2 expression in hNPCs following transplantation. Evaluating integrin expression *in vitro*, specifically of  $\alpha 9$  and  $\beta 1$  integrin, we have also demonstrated that iPSC-derived hNPCs express low levels of each integrin subunit which can be further enhanced using lentiviral transduction (as shown for  $\alpha 9$  integrin) *in vivo*.

### Overexpressing $\alpha 9$ -eYFP results in increased neurite outgrowth from hNPCs in the presence of TN-C

Overexpression of  $\alpha 9$ -eYFP in hNPCs was successfully achieved using lentivirus. We used full-length  $\alpha 9$  integrin tagged to eYFP to allow the exogenous protein to be detected *in vivo* using an anti-GFP antibody due to a lack of commercially available  $\alpha 9$  integrin antibodies that can detect this integrin subunit *in vivo*. The function of  $\alpha 9$  integrin was analyzed using a neurite outgrowth assay in the presence of TN-C as previously described (Andrews et al., 2009). Our results show neurite outgrowth from  $\alpha 9$ -eYFP hNPCs was significantly greater compared to either fGFP-hNPCs or untransduced WT hNPCs when grown on chicken TN-C (10 $\mu$ g/mL). The  $\alpha 9$ -eYFP hNPCs also had significant outgrowth on low concentrations of human TN-C (1 $\mu$ g/mL) suggesting they may show greater specificity for human TN-C. Interestingly, when the concentration of the human TN-C was increased to 5 $\mu$ g/mL and 10 $\mu$ g/mL, the outgrowth from both the fGFP-hNPCs and WT hNPCs gradually increased compared to their growth on 1 $\mu$ g/mL TN-C, whilst the outgrowth  $\alpha 9$ -eYFP hNPCs plateaued. These results suggest endogenous integrin function is increased as the concentration of human TN-C is increased.

Research has shown embryonic DRGs can increase integrin cell surface expression in the presence of inhibitory CSPGs resulting in integrin-induced outgrowth (Condic et al., 1999). This adaptation is not mimicked with adult neurons, however overexpression of integrin in adult DRG neurons can rescue this effect and result in increased neurite growth (Condic, 2001; Cheah et al., 2016). Our present results suggest that increasing the concentration of human TN-C results in an intrinsic activation of endogenous integrin function. Although other TN-C-binding integrin subunits such as  $\alpha 7$  and  $\alpha 8$  (Mercado et al., 2004) may influence neurite outgrowth, expression of these subunits were not assessed within this study but furthermore they have not been directly linked to neurite outgrowth.

### **Characterization of hNPC transplants in developing rat sensorimotor cortex**

We used a neonatal rat model and transplanted cells between the ages of P0-2 into the predicted layer 5 of the sensorimotor cortex using a developmental rat brain atlas (Altman and Bayer, 1995). Coordinates were optimized with each new litter and all off-target grafts were rejected from study analysis. The coordinates that resulted with the most on-target transplants were: AP 0.0 mm, ML 1.5/-1.5 mm, DV -0.4 mm. A window of opportunity exists at early postnatal ages where the rat immune system is not yet fully developed (Reviewed in Marshall-Clarke et al., 2000; Reviewed in Holsapple et al., 2003) allowing for longer term survival of grafts that ensures analysis, compared to a naïve adult rodent model. Transplantation at this age of transplantation also exploits the advantages of a readily developing nervous system which may facilitate integration and development of cells and their axonal projections. Transplantation of embryonic stem cell (ESC)-derived NPCs and neurons has previously been carried out in the immune-competent rodent neonatal brain with promising results for graft survival and integration (Ideguchi et al., 2010; Denham et al., 2012). For example, transplanted ESC-derived neurons can survive up to 10 weeks following transplant into the neonatal rat striatum (Denham et al., 2012). Similarly, ESC-derived NPCs transplanted into P0-P2 mouse cerebral cortex differentiated into area-specific cortical projection neurons and projected within the host CST within 2-3 weeks after transplantation (Ideguchi et al., 2010). In the present study, transplant survival was highest at earlier time points of 2 and 4 weeks, after which IHC results show an increase in Iba-1 immunoreactivity indicating microglia activity (or infiltration) and an increase in GFAP immunoreactivity indicating astrocytic activity, correlating with a decrease in graft survival in conjunction with a loss of axonal projections. In addition, despite the increase in GFAP expression at the transplant site, there was minimal colocalization between transplanted cells and GFAP immunoreactivity confirming the fact that any transplanted astrocytes (up to 5% total of the cultured hNPCs prior to transplant) did not significantly contribute to the increase in astrocytes over time.

Using human-specific antibodies, the transplanted cell bolus and emanating projections were detected *in vivo*. Specifically, these projections followed the trajectory of the CST through the corona radiata and internal capsule, cerebral peduncle, pons and pyramids of the brain stem. The furthest projections were observed within the pyramids 2 weeks following transplantation, in both WT hNPC and  $\alpha 9$ -eYFP hNPC transplants. By 8 weeks, WT and  $\alpha 9$ -eYFP hNPCs were found to project only as far as the internal capsule. It is unclear whether at this time point, the axons were actively dying back in conjunction with the increased immune response but the increase in microglial activation suggests this may have been the case.

It is important to consider not only the survival and projections of transplanted cells, but also whether the cells integrate and mature *in vivo*. Over a 2 month time period, we would expect the cells to develop and follow a path of maturation, which was assessed by analysing levels of DCX expression. Previously published work on DCX expression following transplantation of mouse embryonic cortical neurons show DCX expression is reduced 2 weeks following transplant into the adult lesioned motor cortex (Ballout et al., 2016). However, rodent cells and human cells follow different time courses of development and maturation. In the human NPCs used in this study, DCX expression was retained from 2 weeks up until 8 weeks *in vivo* suggesting that a large proportion of the cells are maintained in progenitor cell state following grafting. The expression of endogenous DCX was present in the developing rat cortex at 2 weeks at a very low level and by 4 weeks, no DCX expression was detected. This is in line with previously published literature suggesting cell proliferation ends at around P15 (Reviewed in Rice and Barone, 2000). We also assessed vGLUT1 expression in the cultured hNPCs and found it to be expressed throughout the cells. We did not investigate expression following transplantation since the growing axons did not reach their targets in the spinal cord, however this is an area for future investigation.

Another way to analyze hNPC maturation is by assessing myelination. Research has suggested hNPCs must mature and become functional neurons before myelination can occur (Lu et al., 2014). Following immunolabelling for MBP, we were unable to confirm myelination of hNPCs *in vivo* (data not shown) however it may be advantageous for hNPCs to retain an immature cell state when considering the need to extend long distance axonal projections and make new synaptic contacts. Recent research has demonstrated human iPSC-derived cortical neurons (Axol Bioscience) require long-term culture of 20-30 weeks for functional synapses to form (Odawara et al., 2016). Similarly, transplanted human NSCs into the adult immunodeficient rodent lesioned spinal cord showed no signs of maturation until 3 months *in vivo* with oligodendrocyte formation not detected until 1 year post-transplant (Lu et al., 2017). These results highlight the requirement for long-term analysis of human iPSC-NPCs *in vivo* to enable full characterization of integration and maturation.

### **Why do $\alpha 9$ -eYFP hNPC axonal projections not extend further than WT hNPCs *in vivo*?**

Over the period of 8 weeks,  $\alpha 9$ -eYFP hNPC-derived axons did not project as far as the WT hNPC-derived axons (Fig. 9). There are a number of potential reasons which may have impacted on this outcome. Previous literature suggests genetic modification of transplanted cells using lentivirus does not affect cell viability, providing the host is immunocompromised (Behrstock et al., 2008; Fujimoto et al., 2012). Some studies however have reported a significant loss of virally-transduced cells following engraftment into the rodent brain. For example, transplantation of mouse luciferase-expressing NSCs into immune-competent adult mouse brain show 80% of cells undergo apoptotic cell death within the first 24 hours following grafting, with only 1% of cells surviving 14 days following transplant. This was correlated with an increase in hypoxic marker expression within the graft bolus suggesting lack of oxygen and nutrients contributed to reduced cell survival (Reekmans et al., 2012).

As the ligand for  $\alpha 9 \beta 1$  is TN-C, the expression of this ECM protein was analyzed within the endogenous tissue and was found to be localized to the area surrounding the transplant sites following engraftment of hNPCs (Fig. 6). It is unclear whether the localized expression of TN-C at the injection site limited the hNPCs projections from extending long distances. If  $\alpha 9$ -eYFP-hNPCs preferentially project to a TN-C rich environment this may predict their behavior if transplanted into an injury model, where TN-C is upregulated at the injury site (Zhang et al., 1997). Although care was taken during the transplant, the injection itself may have induced a micro-injury that could have impacted upon  $\alpha 9$  integrin activity. The results from the outgrowth assays show  $\alpha 9$ -eYFP was functional prior to transplantation yet proteins associated with CNS injury, including CSPGs and Nogo-A, inactivate integrin activity (Hu and Strittmatter, 2008; Tan et al., 2011). Modulation of the integrin activation state using forced expression of kindlin-1 or human-specific antibodies TS2/16 (Hu and Strittmatter, 2008; Tan et al., 2011; Cheah et al., 2016), can overcome this inhibition and may be required if  $\alpha 9$ -eYFP-hNPCs were to be used for regenerative purposes in the CNS.

### Trafficking of $\alpha 9$ integrin within the CST

We know some integrin subunits, including  $\alpha 9$ , are not efficiently transported into the axons of certain neuron subtypes including adult CST axons (Andrews et al., 2016). Recent work by Franssen *et al* have highlighted that Rab11- and Arf6-mediated trafficking of integrins play a key role in their age-related downregulation as a result of increased retrograde integrin trafficking leading to a reduced presence of integrins within the axonal compartment (Franssen et al., 2015). However, during development, integrins are freely trafficked within axons of the corticospinal tract (Andrews et al., 2016). This observation does not seem to be limited to integrins as other transmembrane proteins, such as TrkB and IGF-IR, are also not transported within the entire length of adult axons such as in the CST (Hollis et al., 2009a, 2009b). In the current study however, we show  $\alpha 9$ -eYFP expression throughout the length of the axonal projections at all time points. As these are early stage developing neurons we would expect trafficking of growth-promoting integrins to be fully functioning in an anterograde direction encouraging neurite outgrowth and axonal elongation. Indeed, our results demonstrate expression of  $\alpha 9$ -eYFP from the cell body through to the tip of extending projections.

### Conclusions

In conclusion, although these iPSC-derived hNPCs were not grafted into a SCI model, there is substantial evidence here and from the literature to suggest these cells could be beneficial following SCI. Crucially however, iPSC-derived hNPCs not only present a potential possibility for targeted cell replacement, but they also provide a tool to study integrin adaptation responses in human neurons following exposure to components of the CNS injury milieu. Our data suggests iPSC-derived hNPCs respond to the presence of environmental TN-C by modifying intrinsic integrin activity, a trait previously demonstrated in embryonic DRGs (Condic et al., 1999). Equally, at this developmental state of early postnatal transplantation, our results suggest that overexpression of growth-promoting integrin receptors are not required to induce axonal outgrowth and elongation and that the endogenous levels of integrin may be sufficient for this purpose.

### Abbreviations

$\alpha 9$	alpha 9 integrin
A-P	Anterior-posterior

$\beta 1$	beta 1 integrin
BCA	Bicinchoninic acid
BDNF	Brain-derived neurotrophic factor
CNS	Central nervous system
CSPGs	Chondroitin sulphate proteoglycans
CST	Corticospinal tract
Ctip2	Chicken ovalbumin upstream promoter transcription factor (COUP-TF) interacting protein 2
D-V	Dorsal-ventral
DCX	Doublecortin
DRG	Dorsal root ganglia
ECM	Extracellular matrix
ESCs	Embryonic stem cells
eYFP	Enhanced yellow fluorescent protein
fGFP	Farnesylated green fluorescent protein
GFAP	Glial fibrillary acid protein
HEK293T	Human embryonic kidney 293 cells expressing SV40 T antigen
hNCAM	Human neural cell adhesion molecule
hNPCs	Human neural progenitor cells
HuNu	Human nuclear antigen
ICC	Immunocytochemistry
IHC	Immunohistochemistry
iPSCs	Induced pluripotent stem cells
M-L	Medial-lateral
MBP	Myelin basic protein
NSCs	Neural stem cells
PLO	Poly-L-ornithine
PFA	Paraformaldehyde
PKC $\gamma$	Protein kinase C gamma
RST	Rubrospinal tract
SCI	Spinal cord injury

Tbr1	T-box brain 1
TN-C	Tenascin-C
Trk/TrkB	Tropomyosin receptor kinase/B
VGlut1	Vesicular glutamate transporter 1
WB	Western blotting
WT	Wild type

## Acknowledgements

We thank Dr. David A. Tumbarello (University of Southampton) for critical reading of the manuscript and experimental advice. We also thank Dr. Yichen Shi (Axol Bioscience) for technical and experimental advice.

## Author Contributions Statement

MA designed the project. LF and MA carried out the experiments and performed data analysis. LF and MA wrote the manuscript.

## Funding

This work was supported by a 600<sup>th</sup> Anniversary PhD studentship from the University of St Andrews to LHF, as well as grants from the RS MacDonald Trust, Northwood Charitable Trust and BBSRC New Investigator Award to MRA.

## Conflict of Interest Statement

The authors have no conflicts of interest to declare.

**Table 1 Primary antibodies used for ICC, IHC and WB**

Antibody	Species	Dilution			Supplier	Cat No.
		ICC	IHC	WB		
BDNF	Polyclonal, rabbit	1:200	----	----	Abcam	AB72439
Ctip2	Polyclonal, rabbit	1:500	----	----	Abcam	Ab70453
DAPI	----	1:10,000	1:10,000	----	Thermo Scientific™	62248
DCX	Polyclonal, rabbit	1:500	1:500	----	Abcam	AB18723
GFAP	Polyclonal, rabbit	1:750	1:500	----	DAKO	ZO33429-2
GFP	Polyclonal, rabbit	1:1,000	1:1,000	1:1,000	Invitrogen	A11122
HuNu	Monoclonal, mouse	----	1:500	----	Millipore	MAB1281
Iba-1	Polyclonal, rabbit	----	1:500	----	Wako	019-19741
MBP	Monoclonal, rabbit	1:50	----	----	Cell signalling	D8X4Q
hNCAM	Monoclonal, mouse	----	1:200	----	Santa Cruz, Insight	SC-106
PKCγ	Polyclonal, rabbit	----	1:200	----	Santa Cruz, Insight	SC-211
Tbr1	Polyclonal, rabbit	1:200	1:400	----	Abcam	AB31940
TN-C	Monoclonal, rabbit	1:250	1:250	----	Abcam	AB108930
TrkB	Polyclonal, rabbit	1:1,000	----	----	Santa Cruz Biotechnology	SC-12

$\alpha$ 9-integrin	Polyclonal, rabbit	1:1,000	----	1:1,000	Thermo Scientific™	PA5-27771
$\beta$ -actin	Monoclonal, mouse	----	----	1:10,000	Sigma	A5441
$\beta$ III-tubulin	Monoclonal, mouse	1:1,000	----	----	Sigma	T8660
$\beta$ III-tubulin	Polyclonal, rabbit	1:1,000	----	----	Covance	MRB-435P
$\beta$ 1-integrin	Monoclonal, mouse	1:500	----	1:1,000	BD Transductions	610467
Vinculin	Monoclonal, mouse	1:200	----	----	Millipore	MAB3574

**Table 2 Tissue analysis time points**

	Tissue collection time points					Total
	2 weeks	4 weeks	6 weeks	7 weeks	8 weeks	
No. of pups with WT hNPC grafts	6	7	4	6	3	26
No. of pups with $\alpha$ 9-hNPC grafts	3	0*	5	4	5	17
Total	9	7	9	10	8	43

\*Only 2 animals received on target injections in this group and have therefore been removed from the analysis.

**Fig. 1 Endogenous expression of integrins, growth-promoting proteins and deep layer cortical neuron markers in human iPSC-derived NPCs.** Endogenous  $\alpha 9$  (A) and  $\beta 1$  (B) integrin subunit expression in WT hNPCs was confirmed using WB, with faint bands observed at approximately 115 kDa ( $\alpha 9$  integrin) or 120 kDa ( $\beta 1$  integrin). This was analyzed alongside retinal pigmented epithelial (RPE) cell lysates as positive controls for both integrin subunits. Blots were also stained with  $\beta$ -actin (approximately 42 kDa) to confirm equal protein loading. ICC analysis further confirmed a low level of endogenous  $\alpha 9$  (C and D) and  $\beta 1$  (E and F) integrin subunit expression within hNPCs. Furthermore, ICC analysis demonstrated hNPCs express TN-C as observed in punctate dots (white arrows, G and H) around the cell nucleus (identified using DAPI). The hNPCs also express a number of growth-promoting proteins including the progenitor cell marker DCX, observed in both the cell body and neurite projections (white arrows, I), which co-labelled with  $\beta$ III-tubulin (white arrows, J). Expression of the neurotrophic factor BDNF (white arrows in K and L) was also identified as well as expression of its receptor, Trk B (white arrows in M and N). The hNPC cultures largely contained  $\beta$ III-tubulin-positive cells (white arrows in O), however some GFAP-positive cells (white arrows in P) were also detected. No MBP-positive cells were detected following 21 days in culture (Q). Expression of deep-layer cortical neuron markers Tbr1 (white arrows, R-T) and Ctip2 (white arrows, U-W) was detected following ICC within the DAPI-stained nucleus of hNPCs. Cells were co-stained with  $\beta$ III-tubulin and DAPI. Scale bar in C-N, T and W = 20  $\mu$ m; O-S, U and V = 100  $\mu$ m.

**Fig. 2 Overexpression of  $\alpha 9$ -eYFP and fGFP using lentivirus.** The overexpression of  $\alpha 9$ -eYFP was detected using ICC with an anti-GFP antibody (white arrows in A) detected within the hNPC projections (white arrows in B) and co-labelled with  $\beta$ III-tubulin. Integrin expression was observed in the cell body and throughout the length of the projections. fGFP expression was identified following hNPC transduction using LV-fGFP detected using GFP and  $\beta$ III-tubulin antibodies (white arrows in C and D). No GFP expression was observed in untransduced WT hNPCs (E and F). Overexpression was confirmed using WB (G) with an anti-GFP antibody, resulting in a band of approximately 140 kDa in lane 2 (LV- $\alpha 9$ -eYFP hNPC lysates) and a band at approximately 30 kDa in lane 3 (LV-fGFP hNPC lysates). No GFP bands were observed in lane 1 containing WT hNPC lysates.  $\beta$ -actin immunoblotting confirmed equal protein loading (approximately 42 kDa). At high magnification,  $\alpha 9$ -eYFP was observed co-localizing to the tip of extended projections in a punctate arrangement (white arrows in H-J). Scale bar in A-F = 50  $\mu$ m, H = 15  $\mu$ m, I = 5  $\mu$ m and J = 2.5  $\mu$ m.

**Fig. 3 Increasing expression of  $\alpha 9$  integrin in iPSC-derived hNPCs promotes neurite outgrowth when grown on TN-C.** Following transduction of hNPCs with either LV- $\alpha 9$ -eYFP or LV-fGFP, cells were grown on either 10  $\mu$ g/mL chicken TN-C (n = 4; A-C, M) or 1  $\mu$ g/mL (n = 3), 5  $\mu$ g/mL (n = 3) or 10  $\mu$ g/mL (n = 3) of human TN-C (D-L, N-P). Cells were fixed after 72 hours in culture and analyzed with ICC using anti- $\beta$ III-tubulin to label neurites (white arrows in A-L) which were measured using NeuronJ. Results were analyzed using one-way ANOVA and expressed as mean  $\pm$  SEM. \* = P < 0.05, \*\* = P < 0.01, \*\*\* = P < 0.001. Scale bar in A-L = 200  $\mu$ m.

**Fig. 4 iPSC-derived hNPCs injected into the sensorimotor cortex extend axons along the intrinsic pyramidal tract, projecting to the brain stem.** Tissue was stained with anti-HuNu



which detected the cell bolus at the injection site which was either dispersed (white arrows in A) or more compact (white arrows in B). The projections were identified using anti-hNCAM and were observed emanating from the bolus through areas of the intrinsic pyramidal tract. These included the internal capsule (C, D and E), the cerebral peduncles, (F and G) and areas of the pyramids (H, J and J\*). To confirm that the fibers observed at 2 weeks post-transplantation within the brain stem were within the pyramidal tract, tissue was stained for PKC $\gamma$ , to detect the pyramids (I, J and J\*). Abbreviations: ic = internal capsule; cp = cerebral peduncle; py = pyramidal tract. Scale bar in A, B, F and G = 250  $\mu$ m, C and H = 200  $\mu$ m; D, E and I = 500  $\mu$ m; J and J\* = 150  $\mu$ m.

**Fig. 5 Rosette formation within HuNu-positive hNPC bolus.** Following transplantation, some hNPCs formed a rosette formation (A-D), typical of neural stem cells. Scale bar A and B = 250  $\mu$ m; C and D = 150  $\mu$ m.

**Fig. 6 Transplantation of human iPSC-derived NPCs results in an increased rat host immune response and expression of TN-C over time.** Tissue surrounding the graft sites were analyzed using IHC for expression of IBA-1 and GFAP alongside either HuNu and hNCAM to identify the graft site. At early time points of 2 weeks (A, C – low magnification; B, D – high magnification), IBA-1 staining is mildly increased compared to control tissue (I). However, over time there is a noticeable increased immune response from microglia surrounding the HuNu-positive graft site (K, M – low magnification; L, N – high magnification) compared to control tissue (S). Similarly, GFAP staining is mildly increased at earlier time point of 2 weeks (E, G – low magnification; F, H – high magnification) compared to control tissue (J). Over time this expression is markedly increased (O, Q – low magnification; P, R – high magnification) compared to control tissue (T). Following transplantation of hNPCs there was a localized upregulation of TN-C at the graft site (U-W). Scale bar in A, C, E and G = 250  $\mu$ m; B, D, F, H, I, J, L, N, P, R, S, T and W = 150  $\mu$ m; K, M, O, Q, U and V = 500  $\mu$ m.

**Fig. 7 iPSC-derived hNPCs express deep-layer cortical neuron markers and progenitor cell markers *in vivo*.** The expression of the cortical neuron marker Tbr1 was analyzed using an antibody against Tbr1 demonstrating that a large proportion of the transplanted hNPCs expressed Tbr1 that co-localized within HuNu-positive cells at 4 weeks (A and B, white arrows). This expression was observed in both WT and  $\alpha$ 9-eYFP hNPCs across all time points (data not shown). Similarly, expression of the progenitor cell marker DCX was retained following transplant shown at 4 weeks (C-E) and at 8 weeks (F-H) post-transplantation within hNCAM-positive projections (white arrows C-H). Scale bar in A-H = 150  $\mu$ m.

**Fig. 8 iPSC-derived hNPCs retain exogenous  $\alpha$ 9 integrin expression up to 8 weeks *in vivo*.** Following IHC analysis with an anti-GFP antibody, exogenous  $\alpha$ 9-eYFP expression was detected from 2 weeks up to 8 weeks. This expression co-localized throughout hNCAM-positive axonal projections (A, C and E). At 2 weeks  $\alpha$ 9-eYFP expression was observed as punctate dots throughout the axons (A and B). Scale bar in A = 5  $\mu$ m; B = 2.5  $\mu$ m; C = 25  $\mu$ m; D = 10  $\mu$ m; E = 50  $\mu$ m; F = 20  $\mu$ m.

**Fig. 9 In the uninjured developing rodent brain, increasing integrin expression does not result in longer projections from iPSC-derived hNPCs.** Transplants from all 45 pups were compared within the two groups (WT and  $\alpha 9$ -eYFP hNPC transplants). The transplant sites for both groups (panels A and C) indicate that both groups received on-target injections across all time points (2 weeks to 8 weeks). When comparing axonal projections from these two groups (panels B, D and E), results show the distance axons of WT hNPCs project are comparable to the distance  $\alpha 9$ -eYFP hNPCs project. At 2 weeks, both groups show projections emanating through the pyramidal tract and both reaching the pyramids. Similarly, at 4 weeks, WT hNPCs projected axons from the cortex to the pons (The  $\alpha 9$ -eYFP hNPCs at 4 weeks have been removed from analysis due to small group size.) At 6 weeks, WT hNPCs were also able to project to the pons whilst  $\alpha 9$ -eYFP hNPCs projected only as far as the cerebral peduncles. Finally, at 7 and at 8 weeks, WT hNPC projections were detected within the cerebral peduncles and  $\alpha 9$ -eYFP hNPCs were detected only as far as the internal capsule.

**Supplementary Fig 1. Overexpression of alpha9 integrin-eYFP localizes to vinculin- and actin-rich focal adhesions in hNPCs.** The overexpression of  $\alpha 9$ -eYFP was detected using ICC with an anti-GFP antibody (A) within the hNPC projections and co-labelled with the focal adhesion marker, vinculin (B) and actin (C); merged in image D. Integrin expression was observed in the cell body and throughout the length of the projections. At high magnification,  $\alpha 9$ -eYFP was observed within extended projections in vinculin- and actin-rich areas (inserts of D; images i and ii). Monochrome images of i and ii are shown in the lower panels. Arrows in i and ii indicate localization of alpha9 integrin to vinculin- and actin-rich focal adhesions; arrowheads in ii indicate localization of alpha9 integrin an actin-rich focal adhesion site. Scale bar in D = 25  $\mu$ m; i and ii = 5  $\mu$ m.

**Supplementary Fig 2. Changes in endogenous expression of beta1 integrin in hNPCs overexpressing  $\alpha 9$ -eYFP by lentivirus.** Endogenous  $\beta 1$  integrin subunit expression was observed in hNPCs using WB, with bands observed at approximately 120 kDa ( $\beta 1$  integrin), analyzed alongside retinal pigmented epithelial (RPE1) cell lysates. Overexpression was confirmed using WB (G) with an anti-GFP antibody, resulting in a band of approximately 140 kDa in lane 2 (LV- $\alpha 9$ -eYFP hNPC lysates). Despite unequal protein loading (shown with  $\beta$ -actin at approximately 42 kDa), there is a subtle enhancement of endogenous  $\beta 1$  integrin expression following overexpression of  $\alpha 9$  integrin as calculated by the normalized integrated density levels (“Relative  $\beta 1$  integrin expression;”  $\beta 1$  integrin versus  $\beta$  actin).

**Supplementary Fig. 3 hNPC grafts were transplanted into the deep layers of the sensorimotor cortex.** On-target hNCAM-positive grafts were detected within the deep layers of the sensorimotor cortex (SMC, A). This region (cortical layers 5 and 6) can be identified by the presence of TBR1-expressing cells indicated by the white arrows in B. The cell bolus was detected using anti-HuNu antibody (C); cc- corpus callosum; SMC - sensorimotor cortex LV- lateral ventricle. Scale bar in A-C = 500  $\mu$ m.


**Supplementary Fig 4. Transplanted hNPCs show minimal to no overlap with GFAP-positive cells *in vivo*.** HuNu-positive hNPCs did not show overlap with GFAP-positive cells following transplantation (A-D). Scale bar A and B = 250  $\mu$ m; C and D = 150  $\mu$ m.

## References

- Altman J, Bayer SA. (1995). Atlas of prenatal rat brain development. Boca Raton, CRC Press.
- Andrews, M.R., Czvitkovich, S., Dassie, E., Vogelaar, C.F., Faissner, A., Blits, B., Gage, F.H., French-Constant, C., Fawcett, J.W. (2009).  $\alpha 9$  Integrin Promotes Neurite Outgrowth on Tenascin-C and Enhances Sensory Axon Regeneration. *J. Neurosci.* 29, 5546–5557. <https://doi.org/10.1523/JNEUROSCI.0759-09.2009>
- Andrews, M.R., Soleman, S., Cheah, M., Tumbarello, D.A., Mason, M.R.J., Moloney, E., Verhaagen, J., Bensadoun, J.-C., Schneider, B., Aebischer, P., Fawcett, J.W. (2016). Axonal localization of integrins in the CNS is neuronal type and age dependent. *eneuro* ENEURO.0029-16.2016. <https://doi.org/10.1523/ENEURO.0029-16.2016>
- Arlotta, P., Molyneaux, B.J., Chen, J., Inoue, J., Kominami, R., Macklis, J.D. (2005). Neuronal Subtype-Specific Genes that Control Corticospinal Motor Neuron Development *In Vivo*. *Neuron* 45, 207–221. <https://doi.org/10.1016/j.neuron.2004.12.036>
- Ballout, N., Frappé, I., Péron, S., Jaber, M., Zibara, K., Gaillard, A. (2016). Development and Maturation of Embryonic Cortical Neurons Grafted into the Damaged Adult Motor Cortex. *Front. Neural Circuits* 10. <https://doi.org/10.3389/fncir.2016.00055>
- Behrstock, S., Ebert, A.D., Klein, S., Schmitt, M., Moore, J.M., Svendsen, C.N. (2008). Lesion-induced increase in survival and migration of human neural progenitor cells releasing GDNF. *Cell Transplant.* 17, 753–762.
- Bi, X., Lynch, G., Zhou, J., Gall, C.M. (2001). Polarized distribution of  $\alpha 5$  integrin in dendrites of hippocampal and cortical neurons. *J. Comp. Neurol.* 435, 184–193.
- Bradbury, E.J., Moon, L.D.F., Popat, R.J., King, V.R., Bennett, G.S., Patel, P.N., Fawcett, J.W., McMahon, S.B. (2002). Chondroitinase ABC promotes functional recovery after spinal cord injury. *Nature* 416, 636–640. <https://doi.org/10.1038/416636a>
- Cheah, M., Andrews, M.R., Chew, D.J., Moloney, E.B., Verhaagen, J., Fässler, R., Fawcett, J.W. (2016). Expression of an Activated Integrin Promotes Long-Distance Sensory Axon Regeneration in the Spinal Cord. *J. Neurosci.* 36, 7283–7297. <https://doi.org/10.1523/JNEUROSCI.0901-16.2016>
- Chen, B., Wang, S.S., Hattox, A.M., Rayburn, H., Nelson, S.B., McConnell, S.K. (2008). The Fezf2–Ctip2 genetic pathway regulates the fate choice of subcortical projection neurons in the developing cerebral cortex. *Proc. Natl. Acad. Sci.* 105, 11382–11387. <https://doi.org/10.1073/pnas.0804918105>
- Christopherson, K.S., Ullian, E.M., Stokes, C.C.A., Mallowney, C.E., Hell, J.W., Agah, A., Lawler, J., Mosher, D.F., Bornstein, P., Barres, B.A. (2005). Thrombospondins are astrocyte-secreted proteins that promote CNS synaptogenesis. *Cell* 120, 421–433. <https://doi.org/10.1016/j.cell.2004.12.020>
- Condic, M.L. (2001). Adult Neuronal Regeneration Induced by Transgenic Integrin Expression. *J. Neurosci.* 21, 4782–4788.
- Condic, M.L., Snow, D.M., Letourneau, P.C. (1999). Embryonic Neurons Adapt to the Inhibitory Proteoglycan Aggrecan by Increasing Integrin Expression. *J. Neurosci.* 19, 10036–10043.
- Denham, M., Parish, C.L., Leaw, B., Wright, J., Reid, C.A., Petrou, S., Dottori, M., Thompson, L.H. (2012). Neurons derived from human embryonic stem cells extend long-distance axonal projections through growth along host white matter tracts after intra-cerebral transplantation. *Front. Cell. Neurosci.* 6. <https://doi.org/10.3389/fncel.2012.00011>
- Finger, J.H., Bronson, R.T., Harris, B., Johnson, K., Przyborski, S.A., Ackerman, S.L. (2002). The netrin 1 receptors Unc5h3 and Dcc are necessary at multiple choice points for the guidance of corticospinal tract axons. *J. Neurosci. Off. J. Soc. Neurosci.* 22, 10346–10356.
- Franssen, E.H.P., Zhao, R.-R., Koseki, H., Kanamarlapudi, V., Hoogenraad, C.C., Eva, R., Fawcett, J.W. (2015). Exclusion of Integrins from CNS Axons Is Regulated by Arf6 Activation and the AIS. *J. Neurosci.* 35, 8359–8375. <https://doi.org/10.1523/JNEUROSCI.2850-14.2015>
- Fujimoto, Y., Abematsu, M., Falk, A., Tsujimura, K., Sanosaka, T., Juliandi, B., Semi, K., Namihira, M., Komiya, S., Smith, A., Nakashima, K. (2012). Treatment of a mouse model of spinal cord injury by transplantation of human induced pluripotent stem cell-derived long-term self-renewing neuroepithelial-like stem cells. *Stem Cells Dayt. Ohio* 30, 1163–1173. <https://doi.org/10.1002/stem.1083>
- Gaillard, A., Prestoz, L., Dumartin, B., Cantereau, A., Morel, F., Roger, M., Jaber, M., 2007. Reestablishment of damaged adult motor pathways by grafted embryonic cortical neurons. *Nat. Neurosci.* 10, 1294–1299. <https://doi.org/10.1038/nn1970>
- Hevner, R.F., Shi, L., Justice, N., Hsueh, Y., Sheng, M., Smiga, S., Bulfone, A., Goffinet, A.M., Campagnoni, A.T., Rubenstein, J.L. (2001). Tbr1 regulates differentiation of the preplate and layer 6. *Neuron* 29, 353–366.
- Hollis, E.R., Jamshidi, P., Löw, K., Blesch, A., Tuszynski, M.H. (2009a). Induction of corticospinal regeneration by lentiviral trkB-induced Erk activation. *Proc. Natl. Acad. Sci.* 106, 7215–7220. <https://doi.org/10.1073/pnas.0810624106>

- Hollis, E.R., Lu, P., Blesch, A., Tuszynski, M.H. (2009b). IGF-I gene delivery promotes corticospinal neuronal survival but not regeneration after adult CNS injury. *Exp. Neurol.* 215, 53–59. <https://doi.org/10.1016/j.expneurol.2008.09.014>
- Holsapple, M.P., West, L.J., Landreth, K.S. (2003). Species comparison of anatomical and functional immune system development. *Birth Defects Res. B. Dev. Reprod. Toxicol.* 68, 321–334. <https://doi.org/10.1002/bdrb.10035>
- Hu, F., Strittmatter, S.M. (2008). The N-Terminal Domain of Nogo-A Inhibits Cell Adhesion and Axonal Outgrowth by an Integrin-Specific Mechanism. *J. Neurosci.* 28, 1262–1269. <https://doi.org/10.1523/JNEUROSCI.1068-07.2008>
- Ideguchi, M., Palmer, T.D., Recht, L.D., Weimann, J.M. (2010). Murine Embryonic Stem Cell-Derived Pyramidal Neurons Integrate into the Cerebral Cortex and Appropriately Project Axons to Subcortical Targets. *J. Neurosci.* 30, 894–904. <https://doi.org/10.1523/JNEUROSCI.4318-09.2010>
- Lu, P., Blesch, A., Tuszynski, M.H. (2001). Neurotrophism without neurotropism: BDNF promotes survival but not growth of lesioned corticospinal neurons. *J. Comp. Neurol.* 436, 456–470. <https://doi.org/10.1002/cne.1080>
- Lu, P., Ceto, S., Wang, Y., Graham, L., Wu, D., Kumamaru, H., Staufenberg, E., Tuszynski, M.H. (2017). Prolonged human neural stem cell maturation supports recovery in injured rodent CNS. *J. Clin. Invest.* 127, 3287–3299. <https://doi.org/10.1172/JCI92955>
- Lu, P., Wang, Y., Graham, L., McHale, K., Gao, M., Wu, D., Brock, J., Blesch, A., Rosenzweig, E.S., Havton, L.A., Zheng, B., Conner, J.M., Marsala, M., Tuszynski, M.H. (2012). Long-Distance Growth and Connectivity of Neural Stem Cells After Severe Spinal Cord Injury. *Cell* 150, 1264–1273. <https://doi.org/10.1016/j.cell.2012.08.020>
- Lu, P., Woodruff, G., Wang, Y., Graham, L., Hunt, M., Wu, D., Boehle, E., Ahmad, R., Poplawski, G., Brock, J., Goldstein, L.S.B., Tuszynski, M.H. (2014). Long-distance axonal growth from human induced pluripotent stem cells after spinal cord injury. *Neuron* 83, 789–796. <https://doi.org/10.1016/j.neuron.2014.07.014>
- Marshall-Clarke, S., Reen, D., Tasker, L., Hassan, J. (2000). Neonatal immunity: how well has it grown up? *Immunol. Today* 21, 35–41. [https://doi.org/10.1016/S0167-5699\(99\)01548-0](https://doi.org/10.1016/S0167-5699(99)01548-0)
- McKenna, W.L., Betancourt, J., Larkin, K.A., Abrams, B., Guo, C., Rubenstein, J.L.R., Chen, B. (2011). Tbr1 and Fezf2 Regulate Alternate Corticofugal Neuronal Identities during Neocortical Development. *J. Neurosci.* 31, 549–564. <https://doi.org/10.1523/JNEUROSCI.4131-10.2011>
- Meijering, E., Jacob, M., Sarria, J.-C.F., Steiner, P., Hirling, H., Unser, M. (2004). Design and validation of a tool for neurite tracing and analysis in fluorescence microscopy images. *Cytom. Part J. Int. Soc. Anal. Cytol.* 58, 167–176. <https://doi.org/10.1002/cyto.a.20022>
- Mercado, M.L.T., Nur-e-Kamal, A., Liu, H.-Y., Gross, S.R., Movahed, R., Meiners, S. (2004). Neurite Outgrowth by the Alternatively Spliced Region of Human Tenascin-C Is Mediated by Neuronal  $\alpha 7 \beta 1$  Integrin. *J. Neurosci.* 24, 238–247. <https://doi.org/10.1523/JNEUROSCI.4519-03.2004>
- Michelsen, K.A., Acosta-Verdugo, S., Benoit-Marand, M., Espuny-Camacho, I., Gaspard, N., Saha, B., Gaillard, A., Vanderhaeghen, P., 2015. Area-Specific Reestablishment of Damaged Circuits in the Adult Cerebral Cortex by Cortical Neurons Derived from Mouse Embryonic Stem Cells. *Neuron* 85, 982–997. <https://doi.org/10.1016/j.neuron.2015.02.001>
- Naldini, L., Blömer, U., Gallay, P., Ory, D., Mulligan, R., Gage, F.H., Verma, I.M., Trono, D. (1996). *In vivo* gene delivery and stable transduction of nondividing cells by a lentiviral vector. *Science* 272, 263–267.
- Nieuwenhuis, B., Haenzi, B., Andrews, M.R., Verhaagen, J., Fawcett, J.W. (2018). Integrins promote axonal regeneration after injury of the nervous system. *Biol. Rev. Camb. Philos. Soc.* <https://doi.org/10.1111/brv.12398>
- Nori, S., Okada, Y., Yasuda, A., Tsuji, O., Takahashi, Y., Kobayashi, Y., Fujiyoshi, K., Koike, M., Uchiyama, Y., Ikeda, E., Toyama, Y., Yamanaka, S., Nakamura, M., Okano, H. (2011). Grafted human-induced pluripotent stem-cell-derived neurospheres promote motor functional recovery after spinal cord injury in mice. *Proc. Natl. Acad. Sci.* 201108077. <https://doi.org/10.1073/pnas.1108077108>
- Odawara, A., Katoh, H., Matsuda, N., Suzuki, I. (2016). Physiological maturation and drug responses of human induced pluripotent stem cell-derived cortical neuronal networks in long-term culture. *Sci. Rep.* 6, srep26181. <https://doi.org/10.1038/srep26181>
- Reekmans, K., De Vocht, N., Praet, J., Fransen, E., Le Blon, D., Hoornaert, C., Daans, J., Goossens, H., Van der Linden, A., Berneman, Z., Ponsaerts, P. (2012). Spatiotemporal evolution of early innate immune responses triggered by neural stem cell grafting. *Stem Cell Res. Ther.* 3, 56. <https://doi.org/10.1186/scrt147>
- Rice, D., Barone, S. (2000). Critical periods of vulnerability for the developing nervous system: evidence from humans and animal models. *Environ. Health Perspect.* 108, 511–533.


- Sano, N., Shimogawa, T., Sakaguchi, H., Ioroi, Y., Miyawaki, Y., Morizane, A., Miyamoto, S., Takahashi, J. (2017). Enhanced Axonal Extension of Subcortical Projection Neurons Isolated from Murine Embryonic Cortex using Neuropilin-1. *Front. Cell. Neurosci.* 11, 123. <https://doi.org/10.3389/fncel.2017.00123>
- Schneider, C.A., Rasband, W.S., Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image analysis [WWW Document]. *Nat. Methods*. <https://doi.org/10.1038/nmeth.2089>
- Schwab, M.E., Strittmatter, S.M. (2014). Nogo limits neural plasticity and recovery from injury. *Curr. Opin. Neurobiol.* 0, 53–60. <https://doi.org/10.1016/j.conb.2014.02.011>
- Shi, Y., Kirwan, P., Livesey, F.J. (2012a). Directed differentiation of human pluripotent stem cells to cerebral cortex neurons and neural networks. *Nat. Protoc.* 7, 1836–1846. <https://doi.org/10.1038/nprot.2012.116>
- Shi, Y., Kirwan, P., Smith, J., Robinson, H.P.C., Livesey, F.J. (2012b). Human cerebral cortex development from pluripotent stem cells to functional excitatory synapses. *Nat. Neurosci.* 15, 477–486. <https://doi.org/10.1038/nn.3041>
- Silver, J., Miller, J.H. (2004). Regeneration beyond the glial scar. *Nat. Rev. Neurosci.* 5, 146–156. <https://doi.org/10.1038/nrn1326>
- Takahashi, K., Tanabe, K., Ohnuki, M., Narita, M., Ichisaka, T., Tomoda, K., Yamanaka, S. (2007). Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 131, 861–872. <https://doi.org/10.1016/j.cell.2007.11.019>
- Takahashi, K., Yamanaka, S. (2006). Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors. *Cell* 126, 663–676. <https://doi.org/10.1016/j.cell.2006.07.024>
- Tan, C.L., Kwok, J.C.F., Patani, R., Chandran, S., Fawcett, J.W. (2011). Integrin activation promotes axon growth on inhibitory CSPGs by enhancing integrin signaling. *J. Neurosci. Off. J. Soc. Neurosci.* 31, 6289–6295. <https://doi.org/10.1523/JNEUROSCI.0008-11.2011>
- Tornero, D., Wattananit, S., Grønning Madsen, M., Koch, P., Wood, J., Tatarishvili, J., Mine, Y., Ge, R., Monni, E., Devaraju, K., Hevner, R.F., Brüstle, O., Lindvall, O., Kokaia, Z. (2013). Human induced pluripotent stem cell-derived cortical neurons integrate in stroke-injured cortex and improve functional recovery. *Brain J. Neurol.* 136, 3561–3577. <https://doi.org/10.1093/brain/awt278>
- Tran, A.P., Warren, P.M., Silver, J. (2018). The Biology of Regeneration Failure and Success After Spinal Cord Injury. *Physiol. Rev.* 98, 881–917. <https://doi.org/10.1152/physrev.00017.2017>
- Tuszynski, M.H., Steward, O. (2012). Concepts and Methods for the Study of Axonal Regeneration in the CNS. *Neuron* 74, 777–791. <https://doi.org/10.1016/j.neuron.2012.05.006>
- Ullian, E.M., Harris, B.T., Wu, A., Chan, J.R., Barres, B.A. (2004). Schwann cells and astrocytes induce synapse formation by spinal motor neurons in culture. *Mol. Cell. Neurosci.* 25, 241–251. <https://doi.org/10.1016/j.mcn.2003.10.011>
- Ullian, E.M., Sapperstein, S.K., Christopherson, K.S., Barres, B.A. (2001). Control of Synapse Number by Glia. *Science* 291, 657–661. <https://doi.org/10.1126/science.291.5504.657>



**Fig. 1** Endogenous expression of integrins, growth-promoting proteins and deep layer cortical neuron markers in human iPSC-derived NPCs.




931 **Figure 2**



932 **Fig. 2 Overexpression of  $\alpha 9$ -eYFP and fGFP using lentivirus.**


950 **Figure 3**



951  
952 **Fig. 3** Increasing expression of  $\alpha 9$  integrin in iPSC-derived hNPCs promotes neurite  
953 outgrowth when grown on TN-C.  
954  
955




**Figure 4**




**Fig. 4 iPSC-derived hNPCs injected into the sensorimotor cortex extend axons along the intrinsic pyramidal tract, projecting to the brain stem.**

**Figure 5**



**Fig. 5 Rosette formation within HuNu-positive hNPC bolus.**


**Figure 6**



**Fig. 6** Transplantation of human iPSC-derived NPCs results in an increased rat host immune response and expression of TN-C over time.




1034 **Figure 7**




**Fig. 7 iPSC-derived hNPCs express deep-layer cortical neuron markers and progenitor cell markers *in vivo*.**

**Figure 8**




**Fig. 8 iPSC-derived hNPCs retain exogenous  $\alpha 9$  integrin expression up to 8 weeks *in vivo*.**

**Figure 9**




**Fig. 9 In the uninjured developing rodent brain, increasing integrin expression does not result in longer projections from iPSC-derived hNPCs.**

# Supplementary Figure 1



**Supplementary Fig 1. Overexpression of alpha9 integrin-eYFP localizes to focal adhesions in hNPCs.**


**Supplementary Figure 2**



**Supplementary Fig 2. Changes in endogenous expression of beta1 integrin in hNPCs overexpressing  $\alpha 9$ -eYFP by lentivirus.**




# Supplementary Figure 3



Supplementary Fig. 3 hNPC grafts were transplanted into the deep layers of the sensorimotor cortex.

**Supplementary Figure 4**



**Supplementary Fig 4. Transplanted hNPCs show minimal to no overlap with GFAP-positive cells *in vivo*.**