


# Substance Use Patterns of Gay and Bisexual Men in the Momentum Health Study

American Journal of Men's Health  
2018, Vol. 12(5) 1759–1773  
© The Author(s) 2018  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/1557988318786872  
journals.sagepub.com/home/jmh  


Eric Abella Roth, PhD<sup>1</sup>, Zishan Cui, MSc<sup>2</sup>, Lu Wang, MSc<sup>2</sup>,  
Heather L. Armstrong, PhD<sup>2,3</sup>, Ashleigh J. Rich, MPH<sup>2</sup>,  
Nathan J. Lachowsky, PhD<sup>2,4</sup>, Paul Sereda, BA<sup>2</sup>, Kiffer G. Card, PhD<sup>2,5</sup>,  
Jody Jollimore, MPP<sup>6</sup>, Terry Howard, MSc PPH<sup>7</sup>, Gbolahan Olarewaju, MSc<sup>2</sup>,  
David M. Moore, MD, MHSc<sup>2,3</sup>, and Robert S. Hogg, PhD<sup>2,8</sup>

## Abstract

Research with male sexual minorities frequently combines gay and bisexual men as Men Who Have Sex with Men or Gay and Bisexual Men. When analyzed separately, bisexual men consistently feature negative health differentials, exemplified by higher substance use levels. This interpretation is not clear-cut because studies may combine bisexual men and women, use different dimensions of sexual orientation to define bisexuality, and/or not consider number of sexual partners as a possible confounding factor. This study conducted separate bivariate and multivariate analyses comparing gay to bisexual Momentum Study participants based on self-identity, sexual attraction, and sexual behavior, while controlling for number of sexual partners and sociodemographic, psychosocial, and sexual behavior measures. The study hypothesized that, regardless of definition, bisexual men feature higher substance use levels compared to gay men. Bivariate analyses revealed significantly higher ( $p < .05$ ) use among bisexual men for multiple substances in all samples. Nonprescription stimulants and heroin were significant in all multivariate logistic regression models. In contrast, all bisexual samples reported lower use of erectile dysfunction drugs and poppers, substances associated with anal sex among gay men. Subsequent analysis linked these results to lower levels of anal sex in all bisexual samples. Bivariate analyses also revealed that bisexual men featured significantly lower educational levels, annual incomes, and Social Support Scales scores and higher Anxiety and Depression Sub-Scale Scores. In summary, findings revealed bisexual men's distinctive substance use, sexual behavior, psychosocial, and sociodemographic profiles, and are important for tailoring specific health programs for bisexual men.

## Keywords

bisexual men, gay men, substance use, behavioral issues, sexual dimensions

Received March 4, 2018; revised May 23, 2018; accepted June 7, 2018

The majority of research with male sexual minorities categorizes them as “gay and bisexual men” (GBM), or “men who have sex with men” (MSM). For example, Kaestle and Ivory's (2012) content analysis of PUBMED literature dating from 1987 to 2007 reported that less than 20% of 348 articles mentioning bisexuality performed separate analyses for bisexuals. While intended to be inclusive, terms like MSM and GBM make it difficult to draw valid conclusions regarding bisexuals or other men who have sex with both men and women while simultaneously skewing data for gay men (Young & Meyer,

2005; Bauer & Brennan, 2013). In particular, conflating gay and bisexual men is an important problem for men's health research, since studies that do separate gay from bisexual men indicate multiple negative health differentials for bisexual men (Friedman et al., 2014; Friedman & Dodge, 2016). Currently, problems in defining and classifying bisexuality in survey data remain in analyzing gay and bisexual men's health parameters separately and/or in comparison to other sexual groups.

Reports of higher substance use levels for bisexual men in comparison to gay men (Shelton, 2017) exemplify



both these health differentials and their associated problems. Analysis of US Multicenter AIDS Cohort Study (MACS) data reported significantly higher polydrug use for bisexual men compared to gay men (Friedman et al., 2014). In other comparisons with gay men, Ibañez, Purcell, Stall, Parsons, and Gomez (2005) identified HIV-positive bisexual men more frequently using injection drugs, while Nakamura, Semple, Strathdee, and Patterson (2011) revealed HIV-positive methamphetamine-using bisexual men with significantly higher use levels for crack, alcohol, marijuana, cocaine, hallucinogens, and heroin. Brennan-Ing, Porter, Seidel, and Kapiak (2014) indicated that bisexual men 50 years of age and over were significantly more likely to use tobacco, cocaine, crack, and heroin in comparison to same-aged gay men, and a comparison of gay and bisexual men from upstate New York identified significantly higher alcohol severity scores for bisexual men (Hequembourg, Parks, Collins, & Hughes, 2015). Finally, a national probability sample determined that bisexual male adolescents use more illegal substances than either gay or straight male youths (Russell, Driscoll, & Truong, 2002).

While the earlier studies denote higher substance use levels for bisexual men compared to gay men, this interpretation can be problematic for at least three reasons. First, some studies combine bisexual men and women in their analysis (Bauer, Flanders, MacLeod, & Ross, 2016; Hequembourg & Dearing, 2013; Ross et al., 2014), negating direct comparisons between gay and bisexual men. Second, studies may use differing criteria to define bisexuality. Miller, André, Ebin, and Besonova (2007, p. 2) define bisexuality as, "... the capacity for emotional, romantic, and/or physical attraction to more than one sex or gender," while stressing that, "... capacity for attraction may or may not manifest itself in terms of sexual interaction." This definition recognizes three dimensions of human sexual orientation: identity, attraction, and behavior (Laumann, 1994). Using differing dimensions as eligibility criteria negates direct comparisons. Thus, while McCabe, Bostwick, Hughes, West, and Boyd (2010) used identity in their analysis of National Epidemiologic Survey on Alcohol and Related Conditions Data, Bowers, Branson, Fletcher, and Reback (2011)

emphasized sexual behavior to study Los Angeles' bisexual men, and Bauer et al. (2016) focused on sexual attraction to examine Canadian bisexual men and women. One proposed solution to this challenge is to use all three dimensions of sexual orientation whenever possible (Bauer & Brennan, 2013; McCabe, Hughes, Bostwick, & Boyd, 2005; Saewyc et al., 2004; Scheer et al., 2003).

The third difficulty is particularly relevant for behavioral bisexuality. Bauer and Brennan (2013) note that in survey data, classification as bisexual necessitates having both male and female partners within a stated time-period. In contrast, lesbians, gays, and heterosexual men and women need only one sexual partner to "define" their sexual orientation. This is an important difference because increased partner number is frequently associated with increased substance use and sexual behavior risk (Cavazos-Rehg et al., 2011; Mercer, Hart, Johnson, & Cassell, 2009). For example, Cavazos-Rehg et al. (2011) discovered that sexual partner numbers increased with substance use intensification, and Armstrong et al. (2018) documented significant associations between poppers, crystal methamphetamine, and MDMA and increasing number of sexual partners. Despite such findings, sexual partner number is often omitted from analyses comparing bisexuals with other groups (Baldwin et al., 2015; Friedman et al., 2014; Hequembourg & Dearing, 2013; Nakamura et al., 2011). Bauer and Brennan (2013) suggest this potential confounding problem can be mitigated by including the number of sexual partners (both male and female) in multivariable models, restricting analyses to survey participants with at least two partners, and/or conducting sensitivity analyses to check for statistically significant differences between study groups.

This study analyzes data generated by the Momentum Health Study. Momentum focused primarily on gay men, and previous analyses have not separated gay from bisexual study participants (Moore et al., 2016; Rich et al., 2016; Roth et al., 2018). Nonetheless, Momentum data have the potential to address the three challenges outlined earlier. Specifically, they pertain to gay and bisexual men only, the study questionnaire considers all three sexual dimensions of sexual orientation, and includes the number of recent sexual partners, both male and female, for

<sup>1</sup>Department of Anthropology, University of Victoria, Victoria, BC, Canada

<sup>2</sup>British Columbia Centre for Excellence in HIV/AIDS, Vancouver, BC, Canada

<sup>3</sup>Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada

<sup>4</sup>School of Public Health and Social Policy, University of Victoria, Victoria, BC, Canada

<sup>5</sup>Faculty of Health Sciences, Simon Fraser University, Vancouver, BC, Canada

<sup>6</sup>Executive Director, Community Based Research Centre for Gay Men's Health, Vancouver, BC, Canada

<sup>7</sup>Community Advisory Board, Momentum Health Study, Vancouver, BC, Canada

<sup>8</sup>Faculty of Health Sciences, Simon Fraser University, Burnaby, British Columbia, Canada

#### Corresponding Author:

Eric Abella Roth, PhD, Department of Anthropology, University of Victoria, Victoria, BC V8W 3P5, Canada.  
Email: ericroth@uvic.ca

all participants. Study goals in analyzing these data are two-fold. The first is to test the hypothesis that compared to gay Momentum Study participants bisexual men in the study have higher substance use patterns, regardless of the definition of bisexuality. The second is to identify psychosocial, demographic, and socioeconomic factors associated with bisexuality in the Momentum sample. These factors are important because as Ebin (2012) observes, it is not bisexuality *per se* that determines substance patterns, but rather how bisexual men and women are stigmatized and marginalized. Shelton (2017, pp. 120–121) lists three related societal factors that negatively affect bisexual men and women's health: (a) invisibility, (b) double discrimination, and (c) insistence that one must be either gay or straight. The first factor results from conflating bisexual and gay men, as discussed earlier, as well as bisexuals labeled as gay or straight based on their partner's gender. Double discrimination means that bisexuals lack recognition and support from society in general as well as from lesbian and/or gay communities (Dodge et al., 2012). The third factor is the belief that bisexuality is merely a transitional stage before individuals inevitably conform to the heterosexuality/homosexuality dichotomy (Friedman et al., 2014). Results of such societal stressors include higher prevalence of mental health and substance use problems for bisexual men and women (Feinstein & Dyer, 2017).

## Methods

### Protocol

Momentum is a prospective cohort study of the health, sexual behavior, and substance use patterns of gay and bisexual men in Vancouver, British Columbia, Canada. Momentum uses respondent-driven sampling (RDS; Heckathorn, 1997) to recruit participants. In this approach, purposefully selected "seeds" sharing similar characteristics with target populations are initial study participants who are subsequently encouraged to recruit additional participants from their social and sexual networks. Seed recruitment was through community agencies with the assistance of Momentum's community advisory board and later via mobile smartphones and gay Website advertisements (Moore et al., 2016). Seeds underwent a short training session of peer-recruitment procedures and received six coupons, hard copy or digitally generated, to offer acquaintances who met the study's eligibility criteria. These included being at least 16 years of age, self-identification as a man (regardless of sex at birth), reporting sex with a man in the past 6 months, ability to complete a questionnaire in English, and residence in the Greater Vancouver Area. Potential participants came to the study's downtown Vancouver

study office where a research assistant insured they met the eligibility criteria and were a seed or had received a study coupon from another study participant. All eligible participants signed an Informed Consent form, received a \$50 honorarium for participation, and a further \$10 for each person they recruited. Alternatively, participants could enter a monthly draw for a \$250 gift card or a 6-monthly draw for a \$2000 travel voucher. RDSCM v. 3.0 software (Ithaca, NY) managed coupons, recorded recruiter-recruit relationships, and tracked compensation and coupon redemption patterns. Every 6 months, study participants completed a computer-assisted self-interview questionnaire and biological tests including point-of-care HIV testing or HIV blood work as appropriate, blood tests for hepatitis C and syphilis, and optional tests for gonorrhea and chlamydia. Participant recruitment covered the period February 2012–February 2015. This study used only baseline data from participants' first study visit. All procedures received human ethics clearances from Simon Fraser University (ID Number 2011s0691), the University of British Columbia (ID Number H11-00691), and the University of Victoria (ID Number 11-459).

### Measures

Dependent variables were responses to questions about sexual orientation, measured by identity, attraction, and behavior. For identity, the study questionnaire asked, "How would you describe your sexual orientation?" with responses including "gay," "bisexual," "queer," "questioning," and "other." Men giving the last three responses ( $n=46$ ) were removed from the identity analysis, leaving a direct comparison between self-identified gay ( $n = 655$ ) and bisexual men ( $n = 73$ ). Responses to the query, "In the past two years who have you had sexual fantasies about?" determined sexual attraction. Respondents who answered "men only" were considered gay ( $n = 541$ ), while bisexuality ( $n = 233$ ) was defined by responses indicting fantasies about men as well as women. Behavioral bisexuality was determined from responses to the question, "In the past two years who have you had sex with?" Behavioral bisexuality was defined as having sex with at least one male and one female during this time ( $n = 114$ ). In contrast, participants having sex only with men were considered gay ( $n = 660$ ).

Independent variables consisted of substance use, sociodemographic characteristics, and psychosocial and sexual behavior measures. Substance use questions asked if respondents used erectile dysfunction drugs, crystal methamphetamine, poppers, heroin, and/or injected drugs (including steroids) in the past 6 months. In addition, composite variables asked about use of hallucinogens (Ecstasy/MDMA, LSD, Ketamine, mushrooms), prescription stimulants (Concerta®, Adderall®, and

Ritalin®), nonprescription stimulants (crack and cocaine), prescription sedatives (GHB, benzodiazepines, and barbiturates), and prescription opioids (Morphine, Codeine, Oxycontin®, Percocet®) in the same time period. Questions pertaining to alcohol use permitted calculation of Alcohol Use Disorder Test scores (AUDIT; Saunders, Aasland, Babor, De la Fuente, & Grant, 1993). These classified participants as: (a) Low Risk Drinkers (AUDIT score 0–7), (b), Hazardous Drinkers (AUDIT score  $\geq 8$ –15), (c) Harmful Drinkers (AUDIT score 16–19), or (d) Alcohol Dependent Drinkers (AUDIT score  $\geq 20$ ). Sociodemographic variables included measures of age, annual income, education, neighborhood, and current health self-assessment. Psychosocial variables consisted of the Hospital Anxiety and Depression Scale (HAD) (Snaith, 2003), divided into the Anxiety Subscale (study  $\alpha = 0.79$ ), plus the Depression Subscale (study  $\alpha = 0.83$ ), and the Social Support Scale (Lubben et al., 2006, study  $\alpha = 0.86$ ). Sexual variables included the total number of sexual partners, both male and female, number of male sex partners, and male anal sex partners reported in the past 6 months.

### Statistical Analysis

Samples were interdependent, as men could be included in one, two, or all three samples. Therefore, analysis followed the McCabe et al. (2005) methodology and conducted separate bivariate and multivariate analyses among each bisexual sample. In the first regard, differences between bisexual and gay men were assessed using Wilcoxon Rank-Sum tests for continuous variables and  $\chi^2$  tests for categorical variables. Subsequent analyses compared bisexual with gay men as the dependent variable in univariate and multivariate logistic regression models using the SAS® v. 9.4 PROC LOGISTIC routine (Cary, NC) (Allison, 2012). Final multivariable models were determined using a backward elimination procedure based on the Akaike Information Criterion (AIC) and Type-III  $p$ -values (Lima et al., 2007). RDS uses social network size estimates and measures of homophily between recruited men and their recruiters to adjust for sampling bias and produce population parameter estimates (Heckathorn, 2002). The RDS program RDSTAT v. 4.9 adjusted raw data, with network size determined by asking participants how many gay and bisexual men in the Vancouver area they would be comfortable giving a study voucher.

## Results

### Descriptive Sample Statistics

The final sample contained 774 gay and bisexual men, including 134 seeds. Table 1 presents raw count data and

percentages, plus RDS-adjusted values and their 95% confidence intervals (CIs) for the total sample. The raw data indicated that 76% ( $n = 585$ ) of the sample self-identified as White, 9.6% ( $n = 74$ ) as Asian, 6.5% ( $n = 50$ ) as Indigenous, and 4.5% ( $n = 35$ ) as Hispanic. A final classification consisted of 30 men (3.9%) who identified as Arab ( $n = 7$ ), Black ( $n = 10$ ), or simply responded as “Other,” with the last designation retained for analysis. More than three-quarters of participants had more than a high school education ( $n = 595$ , 76.9%). Despite the overall high educational level, the majority were in the lowest annual income group,  $< \$30,000$  ( $n = 485$ , 62.7%). The RDS-adjusted percentage of self-reported HIV-positive men was 21% ( $n = 218$ ).

### Bivariate Analysis Results

Tables 2–4 present bivariate analyses results conducted upon RDS-adjusted data respectively for the identity, attraction, and behavior samples. In the identity results presented in Table 2, bisexual men featured significantly higher values for Anxiety ( $p < .001$ ) and Depression ( $p < .001$ ) Sub-Scales, and lower Social Support Scale scores ( $p < .001$ ). Contrary to expectations, bisexual men reported lower number of male and female sex partners ( $p = .020$ ), male sex partners ( $p = .003$ ), and male anal sex partners ( $p = .011$ ). Other sociodemographic differentials included significantly lower annual income ( $p < .001$ ), educational attainment ( $p < .001$ ), and percentages of men self-identifying as White ( $p = .006$ ) for bisexual men. In addition, these men had a different residence pattern, with fewer living in Downtown Vancouver and the suburbs, represented by Greater Vancouver ( $p = .007$ ). Finally, bisexual men in this sample uniformly self-assessed their current health as inferior to gay men ( $p < .001$ ). With respect to substance use patterning, bisexual men reported significantly higher levels of use of prescription ( $p < .001$ ) and nonprescription ( $p < .001$ ) stimulants, crystal methamphetamine ( $p = .001$ ), prescription opioids ( $p < .001$ ), and heroin ( $p < .001$ ). Only in the case of poppers did bisexual men exhibit significantly lower use levels ( $p = .001$ ).

This pattern of higher substance use, combined with negative socioeconomic and health differentials, is repeated for the much larger ( $n = 233$ ) attraction sample results, presented in Table 3. Compared to gay men, bisexual men in this sample again featured significantly higher Anxiety ( $p < .001$ ) and Depression Sub-Scale Scores ( $p < .001$ ), along with lower Social Support Scale scores ( $p < .001$ ), annual incomes ( $p < .001$ ), and educational attainment ( $p < .007$ ), and similarly assessed their health as worse than gay sample members ( $p < .001$ ). Ethnic differences were also significant ( $p < .028$ ), with more bisexual men identifying as Indigenous, Hispanic,

**Table 1.** Total Sample Descriptive Statistics, Raw and RDS-Adjusted, Values Outside the RDS 95% CI in Bold.

| Variable  | N                 | %           | RDS %       | RDS 95% CI          |
|---|-------------------|-------------|-------------|---------------------|
| <b>Age (Median, Q1, Q3)</b>                         | <b>34, 26, 47</b> |             |             |                     |
| <b>Ethnicity</b>                                    |                   |             |             |                     |
| White   | <b>585</b>        | <b>75.6</b> | <b>68.5</b> | <b>[61.1, 74.5]</b> |
| Asian   | 74                | 9.6         | 9.2         | [5.9, 14.8]         |
| Aboriginal  | 50                | 6.5         | 9.7         | [5.1, 15.1]         |
| Hispanic  | 35                | 4.5         | 7.3         | [3.2, 11.7]         |
| Other   | 30                | 3.9         | 5.1         | [2.6, 8.7]          |
| <b>Education</b>                                    |                   |             |             |                     |
| Less than or equal to high school                   | <b>179</b>        | <b>23.1</b> | <b>32.6</b> | <b>[26.9, 39.8]</b> |
| More than high school                               | <b>595</b>        | <b>76.9</b> | <b>67.4</b> | <b>[60.2, 73.1]</b> |
| <b>Neighborhood</b>                                 |                   |             |             |                     |
| Downtown  | 382               | 49.4        | 51.0        | [43.2, 58.2]        |
| Vancouver   | 240               | 31.0        | 30.8        | [24.8, 37.0]        |
| Outside Greater Vancouver                           | 152               | 19.6        | 18.2        | [13.5, 24.6]        |
| <b>Annual Income</b>                                |                   |             |             |                     |
| <\$30,000   | <b>485</b>        | <b>62.7</b> | <b>72.9</b> | <b>[67.6, 78.5]</b> |
| \$30–\$59,999                                       | <b>200</b>        | <b>25.8</b> | <b>18.6</b> | <b>[14.4, 22.7]</b> |
| >\$60,000   | 89                | 11.5        | 8.6         | [5.3, 12.0]         |
| <b>HIV status—Self-report</b>                       |                   |             |             |                     |
| Negative/Unknown                                    | <b>556</b>        | <b>71.8</b> | <b>79.0</b> | <b>[72.1, 85.9]</b> |
| Positive  | <b>218</b>        | <b>28.2</b> | <b>21.0</b> | <b>[14.1, 27.9]</b> |
| <b>Sexual orientation (Identity)</b>                |                   |             |             |                     |
| Gay   | 655               | 84.6        | 80.4        | [76.0, 84.9]        |
| Bisexual  | <b>73</b>         | <b>9.4</b>  | <b>14.7</b> | <b>[10.4, 18.7]</b> |
| Queer/Questioning <sup>a</sup>                      | <b>25</b>         | <b>3.5</b>  | <b>1.5</b>  | <b>[0.7, 2.5]</b>   |
| Other <sup>a</sup>                                  | 19                | 2.5         | 3.4         | [1.6, 5.7]          |
| <b>Sexual orientation (past 2 years—Attraction)</b> |                   |             |             |                     |
| Males only—Gay                                      | 541               | 69.9        | 66.6        | [60.8, 72.1]        |
| Other = Bisexual                                    | 233               | 30.1        | 33.4        | [27.9, 39.2]        |
| <b>Sexual orientation (past 2 years—Behavior)</b>   |                   |             |             |                     |
| Males only = Gay                                    | <b>660</b>        | <b>85.3</b> | <b>77.3</b> | <b>[71.9, 82.9]</b> |
| Other = Bisexual                                    | <b>114</b>        | <b>14.7</b> | <b>22.7</b> | <b>[17.1, 28.1]</b> |

Note. <sup>a</sup>Omitted from analysis of self-identified men. CI = confidence interval; Q1 = first quartile; Q3 = third quartile; RDS = respondent-driven sampling.

and “Other.” As in the identity model, bisexual men overwhelmingly featured higher substance use levels, represented by significantly higher use for prescription ( $p = .001$ ) and nonprescription ( $p < .001$ ) stimulants, crystal methamphetamine ( $p = .020$ ), injection drug use ( $p = .022$ ), hallucinogens ( $p = .014$ ), heroin ( $p < .001$ ) and prescription opioids ( $p = .013$ ). An important exception to this pattern was significantly lower popper use ( $p < .001$ ), also recorded for self-identified bisexual men. As in the identity sample, erectile dysfunction drug use was lower for bisexual than gay men but was statistically significant ( $p = .023$ ).

Table 4 presents bivariate analysis results for men who were behaviorally bisexual within the past 2 years. As before, overall results depict bisexual men with

negative socioeconomic and psychosocial differentials and distinctly higher substance use levels. Behaviorally bisexual men had significantly higher Anxiety ( $p = .005$ ) and Depression Sub-Scale Scores ( $p < .001$ ), and lower Social Support Scale Scores ( $p < .001$ ) relative to gay sample members. Similar to the other samples, behaviorally bisexual men featured lower educational attainment ( $p < .001$ ) and annual income distribution ( $p < .001$ ), and assessed their current health as worse than gay men ( $p < .001$ ). Behaviorally bisexual men had more Indigenous, Hispanic and “Other” ethnic members ( $p < .001$ ) than gay sample members, and a significantly different residence pattern ( $p = .001$ ). Despite being twice as large as the identity sample and one-half the size of the attraction sample, this group had very

**Table 2.** Results for Bivariate RDS-Adjusted Data Analysis, *Identify Model* (Bisexual Men = 73, Gay Men = 655).

| Continuous variable                     | Gay men |    |    | Bisexual men |    |    | Probability |
|---|---------|----|----|--------------|----|----|-------------|
|   | MD      | Q1 | Q3 | MD           | Q1 | Q3 |             |
| <b>Age</b>                              | 34      | 26 | 47 | 39           | 28 | 50 | .088        |
| <b>HAD anxiety</b>                      | 8       | 5  | 11 | 9            | 7  | 12 | .015        |
| <b>HAD depression</b>                   | 4       | 2  | 6  | 6            | 3  | 9  | <.001       |
| <b>Social support</b>                   | 10      | 8  | 12 | 8            | 5  | 10 | <.001       |
| <b>Male and female sex partners P6M</b> | 5       | 2  | 13 | 3            | 2  | 8  | .020        |
| <b>Male sex partners P6M</b>            | 5       | 2  | 12 | 2            | 1  | 4  | .003        |
| <b>Male sex partners P6M</b>            | 3       | 1  | 9  | 1            | 10 | 3  | .011        |

| Categorical Variable              | Gay men  |      | Bisexual men |      | Probability |
|-----------------------------------|----------|------|--------------|------|-------------|
|                                   | <i>n</i> | %    | <i>n</i>     | %    |             |
| <b>Annual income</b>              |          |      |              |      | <.001       |
| <\$30,000                         | 387      | 66.6 | 62           | 89.9 |             |
| \$30,000–\$59,999                 | 185      | 22.6 | 7            | 6.0  |             |
| >\$60,000                         | 83       | 10.8 | 4            | 4.1  |             |
| <b>Ethnicity</b>                  |          |      |              |      | .006        |
| White                             | 490      | 70.1 | 57           | 66.2 |             |
| Asian                             | 71       | 10.3 | 2            | 3.3  |             |
| Indigenous                        | 41       | 8.3  | 4            | 8.9  |             |
| Hispanic                          | 31       | 7.7  | 4            | 7.6  |             |
| Other                             | 22       | 3.6  | 7            | 14.0 |             |
| <b>Education</b>                  |          |      |              |      | <.001       |
| Less than or equal to high school | 136      | 25.8 | 37           | 53.8 |             |
| More than high school             | 519      | 74.2 | 36           | 46.2 |             |
| <b>Neighborhood</b>               |          |      |              |      | .007        |
| Downtown                          | 333      | 52.3 | 34           | 47.8 |             |
| Vancouver                         | 192      | 25.9 | 44           | 39.3 |             |
| Greater Vancouver                 | 130      | 21.8 | 15           | 12.9 |             |
| <b>Current Health</b>             |          |      |              |      | <.001       |
| Excellent                         | 120      | 19.6 | 4            | 2.0  |             |
| Very good                         | 267      | 36.9 | 22           | 25.6 |             |
| Good                              | 196      | 30.4 | 27           | 45.4 |             |
| Fair                              | 60       | 10.4 | 15           | 17.3 |             |
| Poor                              | 10       | 2.4  | 3            | 4.1  |             |
| DK                                | 2        | 0.3  | 2            | 2.7  |             |

| Categorical variable          | Gay men  |      | Bisexual men |      | Probability |
|-------------------------------|----------|------|--------------|------|-------------|
|                               | <i>n</i> | %    | <i>n</i>     | %    |             |
| <b>Non-Rx stimulants P6M</b>  |          |      |              |      | <.001       |
| No                            | 479      | 72.0 | 39           | 45.5 |             |
| Yes                           | 176      | 28.0 | 34           | 54.5 |             |
| <b>Injection drug use P6M</b> |          |      |              |      | .147        |
| No                            | 595      | 91.6 | 59           | 87.4 |             |
| Yes                           | 60       | 8.4  | 14           | 13.6 |             |
| <b>Rx stimulants P6M</b>      |          |      |              |      | <.001       |
| No                            | 620      | 94.7 | 65           | 85.8 |             |
| Yes                           | 35       | 5.3  | 8            | 14.2 |             |

(continued)

**Table 2. (continued)**

| Categorical variable                  | Gay men  |      | Bisexual men |      | Probability |
|---------------------------------------|----------|------|--------------|------|-------------|
|                                       | <i>n</i> | %    | <i>n</i>     | %    |             |
| <b>Crystal meth P6M</b>               |          |      |              |      |             |
| No                                    | 532      | 82.6 | 50           | 68.5 | .001        |
| Yes                                   | 132      | 17.4 | 23           | 31.5 |             |
| <b>Rx sedatives P6M</b>               |          |      |              |      |             |
| No                                    | 517      | 81.3 | 57           | 74.2 | .082        |
| Yes                                   | 138      | 18.7 | 16           | 25.8 |             |
| <b>Hallucinogens P6M</b>              |          |      |              |      |             |
| No                                    | 452      | 72.4 | 50           | 71.3 | .801        |
| Yes                                   | 203      | 27.6 | 23           | 18.7 |             |
| <b>Heroin P6M</b>                     |          |      |              |      |             |
| No                                    | 641      | 98.0 | 60           | 85.6 | <.001       |
| Yes                                   | 14       | 2.0  | 13           | 14.4 |             |
| <b>Rx opioids P6M</b>                 |          |      |              |      |             |
| No                                    | 602      | 91.8 | 59           | 80.0 | <.001       |
| Yes                                   | 53       | 8.2  | 14           | 20.0 |             |
| <b>Poppers P6M</b>                    |          |      |              |      |             |
| No                                    | 401      | 64.5 | 51           | 80.4 | .001        |
| Yes                                   | 254      | 35.5 | 22           | 19.6 |             |
| <b>Erectile dysfunction drugs P6M</b> |          |      |              |      |             |
| No                                    | 491      | 78.8 | 58           | 86.1 | .071        |
| Yes                                   | 164      | 21.2 | 15           | 13.9 |             |
| <b>Other drugs P6M</b>                |          |      |              |      |             |
| No                                    | 590      | 90.1 | 63           | 93.6 | .391        |
| Yes                                   | 65       | 9.9  | 10           | 6.4  |             |
| <b>AUDIT scores</b>                   |          |      |              |      |             |
| Low risk 0–7                          | 394      | 63.3 | 39           | 56.9 | .558        |
| Hazardous 8–15                        | 173      | 23.9 | 21           | 27.7 |             |
| Harmful 16–19                         | 46       | 6.9  | 7            | 8.3  |             |
| Dependence ≥20                        | 38       | 5.9  | 5            | 7.0  |             |

Note. Referent = gay men. P6M = past 6 months. MD = median; Q1 = first quartile; Q3 = third quartile; HAD = hospital anxiety and depression scale; RDS = respondent-driven sampling.

strong similarities to both, reporting significantly higher use of injection drugs ( $p = .049$ ), prescription ( $p < .001$ ) and nonprescription stimulants ( $p < .001$ ), crystal methamphetamine ( $p = .004$ ), prescription sedatives ( $p < .001$ ), and heroin ( $p < .001$ ). This sample also had a significantly higher AUDIT score distribution ( $p < .001$ ), with a larger proportion of men classified as Alcohol Dependent (gay men = 5.2%, bisexual men = 17.7%). Important exceptions to the overall pattern of higher substance use for bisexual men were once again represented by significantly lower scores for poppers ( $p < .001$ ) and slightly lower erectile dysfunction drug use ( $p = .571$ ).

### Logistic Regression Results

Overall, bivariate analyses revealed distinctive pattern of substance use, psychosocial and socioeconomic and

demographic variables associated with bisexuality, no matter how defined within the overall Momentum sample. These patterns were maintained in subsequent univariate and multivariate logistic regression analyses, with Table 5 presenting statistically significant ( $p < .05$ ) variables in the final multivariable model. These revealed significantly lower Social Support Scale scores for all three samples, lower annual income distributions for the identity and behavior models, and significantly lower education attainment for the identity model. In contrast, in all three models, bisexual men assessed their current health status as inferior to gay men, and the small “Other” ethnic group was statistically significant in the identity and behavior samples, while the also numerically small Hispanic sample was significant in the behavior model. Multivariate results again indicated higher substance use levels for bisexual men, with heroin and nonprescription stimulants significantly higher in all samples, along with

**Table 3.** Results for RDS-Adjusted Bivariate Data Analysis, Attraction Model (Bisexual Men = 233, Gay Men = 541).

| Continuous variable                   | Gay men |    |    | Bisexual men |    |    | Probability |
|---------------------------------------|---------|----|----|--------------|----|----|-------------|
|                                       | MD      | Q1 | Q3 | MD           | Q1 | Q3 |             |
| <b>Age</b>                            | 34      | 26 | 47 | 33           | 25 | 46 | .472        |
| <b>HAD anxiety</b>                    | 8       | 5  | 10 | 9            | 6  | 12 | <.001       |
| <b>HAD depression</b>                 | 4       | 2  | 6  | 5            | 2  | 7  | <.001       |
| <b>Social support</b>                 | 10      | 8  | 12 | 9            | 7  | 11 | <.001       |
| <b>Male + female sex partners P6M</b> | 5       | 2  | 14 | 4            | 2  | 10 | .748        |
| <b>Male sex partner P6M</b>           | 5       | 2  | 12 | 4            | 2  | 10 | .946        |
| <b>Male anal sex partner P6M</b>      | 4       | 1  | 9  | 2            | 1  | 5  | .487        |

| Categorical variable              | Gay men  |      | Bisexual men |      | Probability |
|-----------------------------------|----------|------|--------------|------|-------------|
|                                   | <i>n</i> | %    | <i>n</i>     | %    |             |
| <b>Annual income</b>              |          |      |              |      |             |
| <\$30,000                         | 318      | 65.5 | 167          | 81.0 | <.001       |
| \$30,000–\$59,999                 | 153      | 22.9 | 47           | 14.0 |             |
| >\$60,000                         | 70       | 11.6 | 19           | 5.1  |             |
| <b>Ethnicity</b>                  |          |      |              |      |             |
| White                             | 406      | 70.3 | 179          | 68.6 | .028        |
| Asian                             | 60       | 10.7 | 14           | 5.3  |             |
| Indigenous                        | 35       | 8.4  | 15           | 10.8 |             |
| Hispanic                          | 22       | 6.9  | 13           | 8.2  |             |
| Other                             | 14       | 3.8  | 12           | 7.1  |             |
| <b>Education</b>                  |          |      |              |      |             |
| Less than or equal to high school | 113      | 25.9 | 66           | 35.2 | .007        |
| More than high school             | 428      | 74.1 | 167          | 64.8 |             |
| <b>Neighborhood</b>               |          |      |              |      |             |
| Downtown                          | 273      | 51.6 | 109          | 47.8 | .177        |
| Vancouver                         | 165      | 26.8 | 75           | 39.3 |             |
| Greater Vancouver                 | 103      | 21.5 | 49           | 12.9 |             |
| <b>Current health</b>             |          |      |              |      |             |
| Excellent                         | 100      | 20.4 | 28           | 8.6  | <.001       |
| Very good                         | 228      | 38.7 | 75           | 26.0 |             |
| Good                              | 154      | 29.8 | 89           | 40.7 |             |
| Fair                              | 51       | 9.4  | 30           | 16.7 |             |
| Poor                              | 6        | 1.3  | 9            | 5.8  |             |
| DK                                | 2        | 0.4  | 2            | 2.2  |             |
| <b>Rx stimulants P6M</b>          |          |      |              |      |             |
| No                                | 513      | 96.0 | 211          | 89.9 | .001        |
| Yes                               | 28       | 4.0  | 22           | 10.1 |             |

| Categorical                   | Gay men  |      | Bisexual men |      | Probability |
|-------------------------------|----------|------|--------------|------|-------------|
|                               | <i>n</i> | %    | <i>n</i>     | %    |             |
| <b>Non-Rx stimulants P6M</b>  |          |      |              |      |             |
| No                            | 399      | 73.7 | 149          | 56.6 | <.001       |
| Yes                           | 142      | 26.3 | 84           | 43.4 |             |
| <b>Injection drug use P6M</b> |          |      |              |      |             |
| No                            | 490      | 90.6 | 201          | 87.0 | .022        |
| Yes                           | 51       | 9.4  | 32           | 13.0 |             |
| <b>Crystal meth P6M</b>       |          |      |              |      |             |
| No                            | 427      | 82.4 | 177          | 75.2 | .020        |
| Yes                           | 114      | 17.6 | 56           | 24.8 |             |

(continued)



**Table 3. (continued)**

| Categorical                           | Gay men  |      | Bisexual men |      | Probability |
|---------------------------------------|----------|------|--------------|------|-------------|
|                                       | <i>n</i> | %    | <i>n</i>     | %    |             |
| <b>Rx sedatives P6M</b>               |          |      |              |      |             |
| No                                    | 423      | 80.7 | 179          | 74.3 | .040        |
| Yes                                   | 118      | 19.3 | 54           | 25.7 |             |
| <b>Hallucinogens P6M</b>              |          |      |              |      |             |
| No                                    | 373      | 73.9 | 152          | 65.3 | .014        |
| Yes                                   | 168      | 26.1 | 82           | 34.7 |             |
| <b>Heroin P6M</b>                     |          |      |              |      |             |
| No                                    | 530      | 98.1 | 213          | 90.8 | <.001       |
| Yes                                   | 11       | 1.9  | 20           | 8.2  |             |
| <b>Rx opioids P6M</b>                 |          |      |              |      |             |
| No                                    | 496      | 91.2 | 204          | 85.3 | .013        |
| Yes                                   | 45       | 8.8  | 29           | 14.7 |             |
| <b>Poppers P6M</b>                    |          |      |              |      |             |
| No                                    | 322      | 62.4 | 161          | 77.2 | <.001       |
| Yes                                   | 219      | 37.6 | 72           | 22.8 |             |
| <b>Erectile dysfunction drugs P6M</b> |          |      |              |      |             |
| No                                    | 403      | 77.8 | 184          | 84.8 | .023        |
| Yes                                   | 138      | 22.2 | 49           | 15.2 |             |
| <b>Other drugs P6M</b>                |          |      |              |      |             |
| No                                    | 483      | 90.3 | 210          | 93.7 | .115        |
| Yes                                   | 58       | 9.7  | 23           | 6.3  |             |
| <b>AUDIT scores</b>                   |          |      |              |      |             |
| Low risk 0–7                          | 328      | 64.8 | 133          | 56.5 | .073        |
| Hazardous 8–15                        | 137      | 22.3 | 67           | 27.9 |             |
| Harmful 16–19                         | 41       | 6.4  | 15           | 5.6  |             |
| Dependence ≥20                        | 32       | 6.5  | 16           | 9.9  |             |

Note. Referent = gay men. MD = median; Q1 = first quartile; Q3 = third quartile; P6M = past 6 months; HAD = hospital anxiety and depression scale; RDS = respondent-driven sampling.

prescription opioids and AUDIT scores in the behavior model, and hallucinogens in the attraction model.

The two important exceptions to the pattern of higher substance use remained in the multivariate models, with erectile dysfunction drug use significantly lower in the identity analysis and popper use significantly lower in all three analyses. These results differ from previously reported patterns for gay men, in which erectile dysfunction drugs and poppers are respectively associated with insertive and receptive anal sex behavior (Fisher, Reynolds, & Napper, 2010; Rich et al., 2016). This study's results parallel reports of lower bisexual men's popper use compared to gay men (Bowers et al., 2011; Brennan-Ing et al., 2014). Lower popper and erectile dysfunction drug levels suggested lower anal sex frequency for bisexual men. RDS-adjusted measures denoting the number of gay and bisexual men reporting no anal sex in the past 6 months for each sample allowed testing of this suggestion. Figure 1 demonstrates that all bisexual samples reported higher levels for no anal sex.  $\chi^2$  analysis revealed that these differences were significant for the

identity ( $p = .016$ ) and attraction ( $p = .006$ ), but not the behavior ( $p = .105$ ) samples. We anticipate further testing this patterning using Momentum event-level data (Rich et al., 2016).

## Discussion

Studies with gay and bisexual men are frequently analyzed and reported under the combined rubric "Men Who Have Sex with Men," or conflated as "Gay and Bisexual Men." When studies do distinguish between male sexual minorities, results point to higher substance use levels among bisexual men. This interpretation is weakened when analyses combine bisexual men and women, use differing dimensions of human sexuality to define bisexuality, and do not consider possible differences in number of sexual partners. In an attempt to avoid these pitfalls, this study analyzed data from Momentum Health Study participants, conducting separate analyses for bisexuality defined by self-identity, attraction to men and women, and sex with both men and women. Bivariate univariable

**Table 4.** Results for Bivariate RDS-Adjusted Data Analysis, Behavior Model (Bisexual Men = 114, Gay Men = 660).

| Continuous variable                   | Gay men |    |    | Bisexual men |    |    | Probability |
|---------------------------------------|---------|----|----|--------------|----|----|-------------|
|                                       | MD      | Q1 | Q3 | MD           | Q1 | Q3 |             |
| <b>Age</b>                            | 34      | 26 | 48 | 33           | 25 | 44 | .101        |
| <b>HAD anxiety</b>                    | 8       | 5  | 11 | 9            | 6  | 12 | .005        |
| <b>HAD depression</b>                 | 4       | 2  | 6  | 5            | 3  | 7  | <.001       |
| <b>Social support</b>                 | 10      | 8  | 12 | 9            | 6  | 11 | <.001       |
| <b>Male + female sex partners P6M</b> | 5       | 2  | 13 | 5            | 2  | 10 | .138        |
| <b>Male sex partners P6M</b>          | 5       | 2  | 13 | 3            | 2  | 6  | .253        |
| <b>Male anal sex partners P6M</b>     | 3       | 1  | 9  | 2            | 1  | 5  | .676        |

| Categorical variable              | Gay men  |      | Bisexual men |      | Probability |
|-----------------------------------|----------|------|--------------|------|-------------|
|                                   | <i>n</i> | %    | <i>n</i>     | %    |             |
| <b>Annual income</b>              |          |      |              |      |             |
| <\$30,000                         | 395      | 66.4 | 90           | 88.6 | <.001       |
| \$30,000–\$59,999                 | 184      | 22.9 | 16           | 14.0 |             |
| ≥\$60,000                         | 81       | 10.7 | 8            | 7.0  |             |
| <b>Ethnicity</b>                  |          |      |              |      |             |
| White                             | 500      | 72.6 | 85           | 56.6 | <.001       |
| Asian                             | 71       | 10.3 | 3            | 2.6  |             |
| Indigenous                        | 41       | 7.9  | 9            | 7.9  |             |
| Hispanic                          | 26       | 3.9  | 9            | 7.9  |             |
| Other                             | 22       | 3.3  | 8            | 7.0  |             |
| <b>Education</b>                  |          |      |              |      |             |
| Less than or equal to high school | 135      | 25.9 | 44           | 41.7 | <.001       |
| More than high school             | 525      | 74.1 | 70           | 58.3 |             |
| <b>Neighborhood</b>               |          |      |              |      |             |
| Downtown                          | 336      | 52.8 | 46           | 42.7 | .001        |
| Vancouver                         | 195      | 26.1 | 45           | 40.5 |             |
| Greater Vancouver                 | 129      | 21.4 | 23           | 16.8 |             |
| <b>Current health</b>             |          |      |              |      |             |
| Excellent                         | 116      | 19.5 | 12           | 4.7  | <.001       |
| Very good                         | 275      | 36.9 | 28           | 24.7 |             |
| Good                              | 192      | 30.5 | 51           | 44.8 |             |
| Fair                              | 67       | 11.2 | 14           | 13.6 |             |
| Poor                              | 8        | 1.5  | 7            | 8.4  |             |
| DK                                | 2        | 0.3  | 2            | 3.7  |             |

| Categorical variable          | Gay men  |      | Bisexual men |      | Probability |
|-------------------------------|----------|------|--------------|------|-------------|
|                               | <i>n</i> | %    | <i>n</i>     | %    |             |
| <b>Injection drug use P6M</b> |          |      |              |      |             |
| No                            | 598      | 91.5 | 93           | 86.4 | .049        |
| Yes                           | 62       | 8.5  | 21           | 13.6 |             |
| <b>Non-Rx stimulants P6M</b>  |          |      |              |      |             |
| No                            | 489      | 74.3 | 59           | 42.8 | <.001       |
| Yes                           | 171      | 25.7 | 55           | 57.2 |             |
| <b>Rx stimulants P6M</b>      |          |      |              |      |             |
| No                            | 623      | 95.6 | 101          | 87.8 | <.001       |
| Yes                           | 37       | 4.4  | 13           | 12.2 |             |
| <b>Crystal meth P6M</b>       |          |      |              |      |             |
| No                            | 542      | 82.0 | 80           | 70.2 | .004        |
| Yes                           | 136      | 18.0 | 34           | 29.8 |             |

(continued)

**Table 4. (continued)**

| Categorical variable                  | Gay men  |      | Bisexual men |      | Probability |
|---------------------------------------|----------|------|--------------|------|-------------|
|                                       | <i>n</i> | %    | <i>n</i>     | %    |             |
| <b>Rx sedatives P6M</b>               |          |      |              |      |             |
| No                                    | 521      | 81.5 | 81           | 66.8 | <.001       |
| Yes                                   | 139      | 18.5 | 33           | 33.2 |             |
| <b>Hallucinogens P6M</b>              |          |      |              |      |             |
| No                                    | 453      | 72.4 | 71           | 66.7 | .142        |
| Yes                                   | 207      | 27.6 | 43           | 33.3 |             |
| <b>Heroin P6M</b>                     |          |      |              |      |             |
| No                                    | 645      | 97.7 | 98           | 87.3 | <.001       |
| Yes                                   | 15       | 2.3  | 16           | 12.7 |             |
| <b>Rx opioids P6M</b>                 |          |      |              |      |             |
| No                                    | 610      | 92.0 | 90           | 77.7 | <.001       |
| Yes                                   | 50       | 8.0  | 24           | 22.3 |             |
| <b>Poppers P6M</b>                    |          |      |              |      |             |
| No                                    | 403      | 63.7 | 80           | 81.4 | <.001       |
| Yes                                   | 257      | 36.3 | 34           | 18.6 |             |
| <b>Erectile dysfunction drugs P6M</b> |          |      |              |      |             |
| No                                    | 497      | 79.7 | 90           | 81.6 | .571        |
| Yes                                   | 163      | 20.3 | 24           | 18.4 |             |
| <b>Other drugs P6M</b>                |          |      |              |      |             |
| No                                    | 596      | 91.2 | 97           | 92.1 | .701        |
| Yes                                   | 64       | 8.8  | 17           | 7.9  |             |
| <b>AUDIT Scores</b>                   |          |      |              |      |             |
| Low risk 0–7                          | 401      | 65.4 | 59           | 48.8 | <.001       |
| Hazardous 8–15                        | 172      | 23.5 | 32           | 26.3 |             |
| Harmful 16–19                         | 47       | 5.9  | 9            | 7.2  |             |
| Dependence ≥20                        | 34       | 5.2  | 14           | 17.7 |             |

Note. MD = median; Q1 = first quartile; Q3 = third quartile; P6M = past 6 months; HAD = hospital anxiety and depression scale; RDS = respondent-driven sampling.

analyses indicated overall higher substance use levels for all bisexual samples, and all multivariable analyses revealed that bisexual men specifically featured significantly higher non-prescription stimulant and heroin use. Prescription opioid use also was significant in the identity and behavior models.

These results support the hypothesis that no matter how defined, bisexual Momentum participants feature higher substance use patterns. This interpretation is strengthened by noting the different sample sizes, from the small ( $n = 73$ ) self-identity sample to the largest attraction ( $n = 233$ ) sample, as well as by analyses controlling for sexual partner number in the multivariate models. Important exceptions to this pattern were lower popper and erectile dysfunction drug use for bisexual men, which subsequent analysis linked to lower levels of anal sex compared to gay men. These findings indicate that bisexual men's substance patterns differ not only in use levels, but also in terms of sexual context. Specifically, they suggest that bisexual men's substance use patterns may differ from the strong association between sex and

substances known as “Chemsex” or “Party and Play” for gay men (Card et al., 2018; Weatherburn, Hickson, Reid, Torres-Rueda, & Bourne, 2016).

This study's second goal was to identify socioeconomic, demographic and psychosocial variables associated with bisexuality. Bivariate analyses revealed that all bisexual samples featured significantly lower educational levels, Social Support Scale scores, and annual incomes, combined with significantly higher Anxiety and Depression Sub-Scale scores. Social Support Scale scores remained significant in all multivariate models. This may be particularly important since previous studies noted bisexual men's feeling of not belonging to either the general population or the gay community (Dodge et al., 2012; Feinstein & Dyar, 2017; Friedman & Dodge, 2016). Furthermore, finding higher Anxiety and Depression Sub-Scale scores, coupled with lower Social Support Scale scores, annual incomes, and educational levels, suggests a possible syndemic effect for bisexual men's health, as suggested by Friedman and Dodge (2016). However, our cross-sectional data preclude attribution of


**Table 5.** Results from Multivariable Models.

| Variable                                 | Identity ( <i>n</i> = 73)<br>Referent = Gay men |                           | Attraction ( <i>n</i> = 233)<br>Referent = Men only |                           | Behavior ( <i>n</i> = 114)<br>Referent = Men only |                      |
|--|---|---------------------------|---|---------------------------|---|----------------------|
|  | aOR   | 95% CI                    | aOR   | 95% CI                    | aOR   | 95% CI               |
| <b>Social support</b>                    | <b>0.87</b>                                     | <b>[0.81, 0.96]</b>       | <b>0.93</b>   | <b>[0.88, 0.98]</b>       | <b>0.92</b>                                       | <b>[0.86, 0.98]</b>  |
| <b>Annual income</b>                     |   |                           |   |                           |   |                      |
| <\$30,000                                | Ref.  |                           | Not selected <sup>a</sup>                           |                           | Ref.  |                      |
| \$30,000–\$59,999                        | <b>0.32</b>                                     | <b>[0.13, 0.80]</b>       |   |                           | <b>0.45</b>                                       | <b>[0.23, 0.89]</b>  |
| >\$60,000                                | 0.67  | [0.23, 1.96]              |   |                           | 0.52  | [0.20, 1.36]         |
| <b>Ethnicity</b>                         |   |                           |   |                           |   |                      |
| White                                    | Ref.  |                           | Not selected <sup>a</sup>                           |                           | Ref.  |                      |
| Asian                                    | 0.52  | [0.16, 1.63]              |   |                           | 0.67  | [0.26, 1.72]         |
| Indigenous                               | 0.40  | [0.14, 1.02]              |   |                           | 1.21  | [0.60, 2.41]         |
| Hispanic                                 | 1.55  | [0.64, 3.76]              |   |                           | <b>5.14</b>                                       | <b>[2.59, 10.19]</b> |
| Other                                    | <b>9.52</b>                                     | <b>[3.84, 23.62]</b>      |   |                           | <b>5.18</b>                                       | <b>[2.31, 11.66]</b> |
| <b>Education</b>                         |   |                           |   |                           |   |                      |
| Less than or equal to high school        | Ref.  |                           | Not selected <sup>a</sup>                           |                           | Not selected <sup>a</sup>                         |                      |
| More than high school                    | <b>0.52</b>                                     | <b>[0.31, 0.89]</b>       |   |                           |   |                      |
| <b>Current health</b>                    |   |                           |   |                           |   |                      |
| Excellent                                | Ref.  |                           | Ref.  |                           | Ref.  |                      |
| Very good                                | <b>9.03</b>                                     | <b>[1.98, 41.09]</b>      | 1.39  | [0.80, 2.41]              | <b>3.09</b>                                       | <b>[1.31, 7.30]</b>  |
| Good                                     | <b>17.39</b>                                    | <b>[3.91, 77.29]</b>      | <b>2.71</b>   | <b>[1.58, 4.63]</b>       | <b>6.02</b>                                       | <b>[2.61, 13.89]</b> |
| Fair                                     | <b>11.34</b>                                    | <b>[2.35, 54.73]</b>      | <b>2.66</b>   | <b>[1.39, 5.14]</b>       | <b>2.89</b>                                       | <b>[1.17, 5.60]</b>  |
| Poor                                     | <b>9.46</b>                                     | <b>[1.50, 59.80]</b>      | <b>5.69</b>   | <b>[1.94, 16.70]</b>      | <b>12.15</b>                                      | <b>[3.28, 45.01]</b> |
| <b>Non-Rx stimulants P6M<sup>a</sup></b> |   |                           |   |                           |   |                      |
| No                                       | Ref.  |                           | Ref.  |                           | Ref.  |                      |
| Yes                                      | <b>1.73</b>                                     | <b>[1.01, 2.96]</b>       | <b>1.55</b>   | <b>[1.07, 2.25]</b>       | <b>2.09</b>                                       | <b>[1.31, 3.33]</b>  |
| <b>Rx stimulants P6M</b>                 |   |                           |   |                           |   |                      |
| No                                       | Ref.  |                           | Not selected <sup>a</sup>                           |                           | Not selected <sup>a</sup>                         |                      |
| Yes                                      | <b>3.19</b>                                     | <b>[1.27, 7.59]</b>       |   |                           |   |                      |
| <b>Heroin</b>                            |   |                           |   |                           |   |                      |
| No                                       | Ref.  |                           | Ref.  |                           | Ref.  |                      |
| Yes                                      | <b>4.56</b>                                     | <b>[1.68, 12.46]</b>      | <b>3.91</b>   | <b>[1.67, 9.18]</b>       | <b>4.45</b>                                       | <b>[1.91, 10.38]</b> |
| <b>Rx opioids</b>                        |   |                           |   |                           |   |                      |
| No                                       |   | Not selected <sup>a</sup> |   | Not selected <sup>a</sup> | Ref.  |                      |
| Yes                                      |   |                           |   |                           | <b>2.31</b>                                       | <b>[1.28, 4.18]</b>  |
| <b>Poppers</b>                           |   |                           |   |                           |   |                      |
| No                                       | Ref.  |                           | Ref.  |                           | Ref.  |                      |
| Yes                                      | <b>0.47</b>                                     | <b>[0.22, 0.99]</b>       | <b>0.41</b>   | <b>[0.28, 0.61]</b>       | <b>0.35</b>                                       | <b>[0.22, 0.57]</b>  |

| Variable                          | Identity ( <i>n</i> = 73)<br>Referent = Gay men |                     | Attraction ( <i>n</i> = 233)<br>Referent = Men only |                     | Behavior ( <i>n</i> = 114)<br>Referent = Men only |                     |
|-----------------------------------|---|---------------------|---|---------------------|---|---------------------|
|                                   | aOR   | 95% CI              | aOR   | 95% CI              | aOR   | 95% CI              |
| <b>Erectile dysfunction drugs</b> |   |                     |   |                     |   |                     |
| No                                | Ref.  |                     | Not selected <sup>a</sup>                           |                     | Not selected <sup>a</sup>                         |                     |
| Yes                               | <b>0.47</b>                                     | <b>[0.22, 0.99]</b> |   |                     |   |                     |
| <b>Hallucinogens P6M</b>          |   |                     |   |                     |   |                     |
| No                                | Not selected <sup>a</sup>                       |                     | Ref.  |                     | Not selected <sup>a</sup>                         |                     |
| Yes                               |   |                     | <b>1.71</b>   | <b>[1.14, 2.57]</b> |   |                     |
| <b>AUDIT scores</b>               |   |                     |   |                     |   |                     |
| Low risk 0–7                      |   |                     |   |                     | Ref.  |                     |
| Hazardous 8–15                    | Not selected <sup>a</sup>                       |                     | Not selected <sup>a</sup>                           |                     | 1.27  | [0.77, 2.09]        |
| Harmful 16–19                     |   |                     |   |                     | 1.13  | [0.48, 2.67]        |
| Dependence ≥20                    |   |                     |   |                     | <b>3.44</b>                                       | <b>[1.63, 7.25]</b> |

Note. Statistically significant ( $p < .05$ ) variables are in bold. aOR = adjusted odds ratio; Ref. = referent.

<sup>a</sup>Not selected by AIC.



**Figure 1.** RDS-adjusted percentages reporting no anal sex in past 6 months.

causation. Future longitudinal research, using both quantitative and qualitative approaches, could fruitfully focus on identifying underlying motives, contexts, and associations in bisexual men's substance use levels and patterns in relation to these factors. In particular, mediation analysis may help define possible relationships between psychosocial factors, social isolation, and substance use, while qualitative studies could search for substance use motivations, as previously done for gay men (Weatherburn, Hickson, Reed, Torres-Rueda, & Bourne, 2016).

This study has limitations. While adjusting data for RDS sampling, we make no claim that the result is a representative sample, and therefore results are inapplicable to other locales. Further, as in all studies based on self-reports of substance use and sexual behavior, data may suffer from social desirability bias. In addition, substance use data constituted categorical "yes/no" responses and did not indicate use frequency. We could not therefore distinguish between episodic substance use and more serious substance use disorders. Finally, having sex with a man in the past 6 months as an eligibility criterion meant our sample did not include bisexual men attracted to both men and women or who self-identify as bisexual but did not have sex with a man during this period. Instead, we focused on identifying and analyzing separately bisexual and gay men's health parameters in a sample that previously lumped both together under the rubric Men Who Have Sex with Men (Moore et al., 2016) or Gay and Bisexual Men (Roth et al., 2018).

Despite these caveats, while focusing on substance use, this study found that Momentum Health Study bisexual participants also featured distinctive sexual behavior, psychosocial, and sociodemographic patterns in comparison to their gay counterparts. These results are particularly important in relation to calls for intervention and education patterns specifically tailored for bisexual men

(Friedman & Dodge, 2016; Shelton, 2017) and illustrate the benefits of separating bisexual from gay men's data, rather than conflating them under the rubric of "gay and bisexual men" and/or "Men Who Have Sex with Men."

### Acknowledgments

The authors are grateful for the assistance and involvement of Momentum Health Study participants, office staff and community advisory board, as well as our community partner agencies, the Health Initiative for Men, YouthCo HIV and Hep C Society, and the Positive Living Society of BC.

### Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Funding for Momentum is through the National Institute on Drug Abuse (Grant # R01DA031055-01A1) and the Canadian Institutes for Health Research (Grant # MOP-107544, 143342, PJT-153139). A CANFAR/CTN Postdoctoral Fellowship Award supported NJL. Scholar Awards from the Michael Smith Foundation for Health Research (#5209, #16863) support DMM and NJL. HLA is supported by a Postdoctoral Fellowship Award from the Canadian Institutes of Health Research (Grant # MFE-152443). AJR is supported by a Frederick Banting and Charles Best Doctoral Research Award from the Canadian Institutes of Health Research (#379361).

### References

- Allison, P. (2012). *Logistic regression using SAS*. Cary, NC: SAS Press.
- Armstrong, H., Roth, E., Rich, A., Lachowsky, N., Cui, Z., Sereda, P., ... Hogg, R. (2018). Associations between sexual partner number and HIV risk behaviors: Implications for HIV prevention efforts in a Treatment as Prevention (TasP) environment. *AIDS Care*, 1–8. [Epub ahead of print].
- Baldwin, A., Dodge, B., Schick, V., Hubach, R. D., Bowling, J., Malebranche, D., ... Fortenberry, J. D. (2015). Sexual self-identification among behaviorally bisexual men in the Midwestern United States. *Archives of Sexual Behavior*, 44(7), 2015–2026.
- Bauer, G., & Brennan, D. (2013). The problem with 'behavioral bisexuality': Assessing sexual orientation in survey research. *Journal of Bisexuality*, 13(2), 148–165.
- Bauer, G., Flanders, C., MacLeod, M., & Ross, L. (2016). Occurrence of multiple mental health or substance use outcomes among bisexuals: A respondent-driven sampling study. *BMC Public Health*, 16(1), 497. doi:http://dx.doi.org/10.1186/s12889-016-3173-z
- Bowers, R., Branson, J., Fletcher, J., & Reback, C. (2011). Differences in substance use and sexual partnering between

- men who have sex with men, men who have sex with men and women and transgender women. *Culture, Health & Sexuality*, 13(06), 629–642.
- Brennan-Ing, M., Porter, K., Seidel, L., & Karpiak, S. (2014). Substance use and sexual risk differences among older bisexual and gay men with HIV. *Behavioral Medicine*, 40(3), 108–115.
- Card, K., Armstrong, H., Carter, A., Cui, Z., Wang, L., Zhu, J., ... Roth, E. (2018). A latent class analysis of substance use and culture among gay, bisexual and other men who have sex with men. *Culture, Health & Sexuality*, 1–6. [Epub ahead of print].
- Cavazos-Rehg, P., Krauss, M., Spitznagel, E., Schootman, M., Cottler, L., & Jean Bierut, L. (2011). Number of sexual partners and associations with initiation and intensity of substance use. *AIDS & Behavior*, 15, 869–874.
- Dodge, B., Schnarrs, P., Reece, M., Goncalves, G., Martinez, O., Nix, R., ... Fortenberry, J. (2012). Community involvement among behaviourally bisexual men in the Midwestern USA: Experiences and perceptions across communities. *Culture, Health & Sexuality*, 14(9), 1095–1110.
- Ebin, J. (2012). Why bisexual health? *Journal of Bisexuality*, 12(2), 168–177.
- Feinstein, B., & Dyer, C. (2017). Bisexuality, minority stress and health. *Current Sex Research*, 9, 42–49.
- Fisher, D., Reynolds, G., & Napper, L. (2010). Use of crystal meth, Viagra and sexual behaviour. *Current Opinion in Infectious Diseases*, 23, 53–56.
- Friedman, M. R., & Dodge, B. M. (2016). The role of syndemic in explaining health disparities among bisexual men: A blueprint for a theoretically informed perspective. In E. R. Wright & N. Carnes (Eds.), *Understanding the HIV/AIDS epidemic in the United States* (pp. 71–98). New York, NY: Springer International Publishing.
- Friedman, M. R., Stall, R., Silvestre, A. J., Mustanski, B., Shoptaw, S., Surkan, P. J., ... Plankey, M. W. (2014). Stuck in the middle: Longitudinal HIV-related disparities among men who have sex with men and women (MSMW). *Journal of Acquired Immune Deficiency Syndromes*, 66(2), 213–220.
- Heckathorn, D. (1997). Respondent-driven sampling: A new approach to the study of hidden populations. *Social Problems*, 44, 174–199.
- Heckathorn, D. (2002). Respondent-driven sampling II: Deriving valid population estimates from chain-referral samples of hidden populations. *Social Problems*, 49(1), 1–34.
- Hequembourg, A., & Dearing, R. (2013). Exploring shame, guilt, and risky substance use among sexual minority men and women. *Journal of Homosexuality*, 60(4), 615–638.
- Hequembourg, A., Parks, K., Collins, R., & Hughes, L. (2015). Sexual assault risks among gay and bisexual men. *The Journal of Sex Research*, 52(3), 282–295.
- Ibañez, G., Purcell, D., Stall, R., Parsons, J., & Gómez, C. (2005). Sexual risk, substance use, and psychological distress in HIV-positive gay and bisexual men who also inject drugs. *AIDS*, 19, S49–S55.
- Kaestle, C., & Ivory, A. (2012). A forgotten sexuality: Content analysis of bisexuality in the medical literature over two decades. *Journal of Bisexuality*, 12(1), 35–48.
- Laumann, E. (1994). *The social organization of sexuality: Sexual practices in the United States*. Chicago, IL: University of Chicago Press.
- Lima, V., Geller, J., Bangsberg, D., Patterson, T., Daniel, M., Kerr, T., ... Hogg, R. (2007). The effect of adherence on the association between depressive symptoms and mortality among HIV-infected individuals first initiating HAART. *AIDS*, 21, 1175–1183.
- Lubben, J., Blozik, E., Gillmann, G., Iliffe, S., von Renteln Kruse, W., Beck, J., & Stuck, A. (2006). Performance of an abbreviated version of the Lubben Social Network Scale among three European community-dwelling older adult populations. *The Gerontologist*, 46(4), 503–513.
- McCabe, S., Bostwick, W., Hughes, T., West, B., & Boyd, C. (2010). The relationship between discrimination and substance use disorders among lesbian, gay, and bisexual adults in the United States. *American Journal of Public Health*, 100(10), 1946–1952.
- McCabe, S., Hughes, T., Bostwick, W., & Boyd, C. (2005). Assessment of difference in dimensions of sexual orientation: Implications for substance use research in a college-age population. *Journal of Studies on Alcohol*, 66(5), 620–629.
- Mercer, C. H., Hart, G. J., Johnson, A. M., & Cassell, J. A. (2009). Behaviourally bisexual men as a bridge population for HIV and sexually transmitted infections? Evidence from a national probability survey. *International Journal of STD & AIDS*, 20(2), 87–94.
- Miller, M., André, A., Ebin, J., & Bessonova, L. (2007). *Bisexual health: An introduction and model practices for HIV/STI prevention programming*. New York, NY: National Gay and Lesbian Task Force Policy Institute, the Fenway Institute at Fenway Community Health, and BiNet USA.
- Moore, D., Cui, Z., Lachowsky, N., Raymond, H., Roth, E., Rich, A., ... Montaner, J. (2016). HIV community viral load and factors associated with elevated viremia among a community-based sample of Men Who Have Sex With Men in Vancouver, Canada. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 72(1), 87–95.
- Nakamura, N., Semple, S., Strathdee, S., & Patterson, T. (2011). HIV risk profiles among HIV-positive, methamphetamine-using men who have sex with both men and women. *Archives of Sexual Behavior*, 40(4), 793–801.
- Rich, A., Lachowsky, N., Cui, Z., Sereda, P., Lal, A., Moore, D., Hogg, R., & Roth, E. (2016). Event-level analysis of anal sex roles and sex drug use among gay and bisexual men in Vancouver, British Columbia, Canada. *Archives of Sexual Behavior*, 44, 389–397.
- Ross, L., Bauer, G., MacLeod, M., Robinson, M., MacKay, J., & Dobinson, C. (2014). Mental health and substance use among bisexual youth and non-youth in Ontario, Canada. *PLoS One*, 9(8), e101604.
- Roth, E., Cui, Z., Rich, A., Lachowsky, N., Sereda, P., Card, K., ... Hogg, R. (2018). Seroadaptive strategies of Vancouver gay and bisexual men in a treatment as prevention environment. *Journal of Homosexuality*, 65(4), 524–539.
- Russell, S. T., Driscoll, A. K., & Truong, N. (2002). Adolescent same-sex romantic attractions and relationships: Implications for substance use and abuse. *American Journal of Public Health*, 92(2), 198–202.

- Saewyc, E., Bauer, G., Skay, C., Bearinger, L., Resnick, M., Reis, E., & Murphy, A. (2004). Measuring sexual orientation in adolescent health surveys: Evaluation of eight school-based surveys. *Journal of Adolescent Health, 35*, 345.e1–345.e15.
- Saunders, J., Aasland, O., Babor, T., De la Fuente, J., & Grant, M. (1993). Development of the Alcohol Use Disorders Identification Test (AUDIT). WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction, 88*, 791–804.
- Scheer, S., Parks, C., McFarland, W., Page-Shafer, K., Delgado, V., Ruiz, J., ... Klausner, J. (2003). Self-reported sexual identity, sexual behaviors and health risks: Examples from a population-based survey of young women. *Journal of Lesbian Studies, 7*, 69–83.
- Shelton, M. (2017). *Fundamentals of LGBT substance use disorders: Multiple identities, multiple challenges*. New York, NY: Harrington Park Press.
- Snaith, R. (2003). The hospital depression and anxiety scale. *Health and Quality of Life Outcomes, 1*(1), 1.
- Weatherburn, P., Hickson, F., Reid, D., Torres-Rueda, S., & Bourne, A. (2017). Motivations and values associated with combining sex and illicit drugs ('chemsex') among gay men in South London: Findings from a qualitative study. *Sexually Transmitted Infections, 93*(3), 206–212.
- Young, R., & Meyer, I. (2005). The trouble with "MSM" and "WSW": Erasure of the sexual-minority person in public health discourse. *American Journal of Public Health, 95*(7), 1144–1149.