



Article

Hexaborate(2−) and Dodecaborate(6−) Anions as Ligands to Zinc(II) Centres: Self-Assembly and Single-Crystal XRD Characterization of $[\text{Zn}\{\kappa^3\text{O-B}_6\text{O}_7(\text{OH})_6\}(\kappa^3\text{N-dien})]\cdot 0.5\text{H}_2\text{O}$ (dien = $\text{NH}(\text{CH}_2\text{-CH}_2\text{NH}_2)_2$), $(\text{NH}_4)_2[\text{Zn}\{\kappa^2\text{O-B}_6\text{O}_7(\text{OH})_6\}_2(\text{H}_2\text{O})_2]\cdot 2\text{H}_2\text{O}$ and $(1,3\text{-pnH}_2)_3[(\kappa^1\text{N-H}_3\text{N}\{\text{CH}_2\}_3\text{NH}_2)\text{Zn}\{\kappa^3\text{O-B}_{12}\text{O}_{18}(\text{OH})_6\}]_2\cdot 14\text{H}_2\text{O}$ (1,3-pn = 1,3-diaminopropane)

Mohammed A. Altahan ^{1,†}, Michael A. Beckett ^{1,*} , Simon J. Coles ² and Peter N. Horton ²¹ School of Natural Sciences, Bangor University, Bangor LL57 2UW, UK; chs030@bangor.ac.uk² Chemistry, University of Southampton, Southampton SO17 1BJ, UK; S.J.Coles@soton.ac.uk (S.J.C.); P.N.Horton@soton.ac.uk (P.N.H.)

* Correspondence: m.a.beckett@bangor.ac.uk; Tel.: +44-1248-382-378

† Current address: Chemistry Department, College of Science, University of Thi-Qar, Nasiriyah, Iraq.

Received: 27 February 2019; Accepted: 23 March 2019; Published: 27 March 2019



Abstract: Two zinc(II) hexaborate(2−) complexes, $[\text{Zn}\{\kappa^3\text{O-B}_6\text{O}_7(\text{OH})_6\}(\kappa^3\text{N-dien})]\cdot 0.5\text{H}_2\text{O}$ (dien = $\text{NH}(\text{CH}_2\text{CH}_2\text{NH}_2)_2$) (1) and $(\text{NH}_4)_2[\text{Zn}\{\kappa^2\text{O-B}_6\text{O}_7(\text{OH})_6\}_2(\text{H}_2\text{O})_2]\cdot 2\text{H}_2\text{O}$ (2), and a zinc(II) dodecaborate(6−) complex, $(1,3\text{-pnH}_2)_3[(\kappa^1\text{N-H}_3\text{N}\{\text{CH}_2\}_3\text{NH}_2)\text{Zn}\{\kappa^3\text{O-B}_{12}\text{O}_{18}(\text{OH})_6\}]_2\cdot 14\text{H}_2\text{O}$ (1,3-pn = 1,3-diaminopropane) (3), have been synthesized and characterized by single-crystal XRD studies. The complexes crystallized through self-assembly processes, from aqueous solutions containing 10:1 ratios of $\text{B}(\text{OH})_3$ and appropriate Zn(II) amine complex: $[\text{Zn}(\text{dien})_2](\text{OH})_2$, $[\text{Zn}(\text{NH}_3)_4](\text{OH})_2$, and $[\text{Zn}(\text{pn})_3](\text{OH})_2$. The hexaborate(2−) anions in 1 and 2 are coordinated to octahedral Zn(II) centres as tridentate (1) or bidentate ligands (2) and the dodecaborate(6−) ligand in 3 is tridentate to a tetrahedral Zn(II) centre.

Keywords: dodecaborate(6−); hexaborate(2−); oxidoborate; polyborate; self-assembly; X-ray structure; zinc(II) complex

1. Introduction

There are more than two hundred known borate (polyborate) minerals, and many more known synthetic polyborates [1–3]. Borates are generally comprised of cationic moieties partnered with anionic units containing boron, oxygen, and in many cases hydroxyl hydrogen. Oxidoborates (or hydroxyoxidoborates) are the more appropriate terms, but the term borate (or polyborate) has been used for many years and will be used in this manuscript. Borates are a class of compounds with rich structural diversity [4–7], and have been synthesized by solvothermal methods or from aqueous solution by the addition of $\text{B}(\text{OH})_3$ to a solution containing the appropriate templating cation [7]. Polyborate salts obtained from aqueous solution usually contain discrete, isolated or insular hydroxyl anions, whilst polyborate salts prepared via solvothermal methods are often more condensed and contain anionic polymeric 1-D chains, 2-D layers or 3-D networks with a variety of framework building blocks [1,7]. Salts formed from aqueous solution often contain the pentaborate(1−)

$[\text{B}_5\text{O}_6(\text{OH})_4]^-$ anion since this anion is structurally well suited to forming crystalline supramolecular lattices, which are held together by strong H-bond interactions [8–11]. We have developed a strategy to overcome pentaborate(1–) salt formation by utilizing more highly charged ($> +1$) metal complex cations with ligands having the potential to form multiple H-bond interactions to template crystallization from aqueous solution of polyborate salts of unusual structures. In this context we have isolated two novel polyborate anions: $[\text{B}_7\text{O}_9(\text{OH})_6]^{3-}$ [12] and $[\text{B}_8\text{O}_{10}(\text{OH})_6]^{2-}$ [13]. We have also recently started to investigate Zn(II)/polyborate chemistry and have been able to isolate an insular bi-Zn(II) complex containing a rare dodecaborate(6–) anion [14] and two polymeric 1-D coordination chains with hexaborate(2–) ligands bridging Zn(II) centres [15]. There are a number of other structural reports on polyborate/Zn(II) chemistry [16–23], including the industrially important $\text{Zn}[\text{B}_3\text{O}_4(\text{OH})_3]$ [24].

In this manuscript we describe the synthesis and XRD structures of two new Zn(II)/hexaborate(2–) complexes: $[\text{Zn}\{\kappa^3\text{O}-\text{B}_6\text{O}_7(\text{OH})_6\}(\kappa^3\text{N-dien})]\cdot 0.5\text{H}_2\text{O}$ (dien = $\text{NH}(\text{CH}_2\text{CH}_2\text{NH}_2)_2$) (1) and $(\text{NH}_4)_2[\text{Zn}\{\kappa^2\text{O}-\text{B}_6\text{O}_7(\text{OH})_6\}_2(\text{H}_2\text{O})_2]\cdot 2\text{H}_2\text{O}$ (2). We also report a Zn(II)/dodecaborate(6–) complex $(1,3\text{-pnH}_2)_3[\kappa^1\text{N}-\text{H}_3\text{N}(\text{CH}_2)_3\text{NH}_2]\text{Zn}\{\kappa^3\text{O}-\text{B}_{12}\text{O}_{18}(\text{OH})_6\}_2\cdot 14\text{H}_2\text{O}$ (1,3-pn = 1,3-diaminopropane) (3). All three complexes are insular and the hexaborate(2–) ligand is tridentate in 1, whereas in 2 it is bidentate to octahedral Zn(II) centres. The dodecaborate(6–) ligand in 3 is tridentate to a tetrahedral Zn(II) centre. The structures of these two anions are drawn schematically in Figure 1.





Figure 1. The (a) hexaborate(2–) anion, $[\text{B}_6\text{O}_7(\text{OH})_6]^{2-}$, observed in 1 and 2; and (b) dodecaborate(6–) anion, $[\text{B}_{12}\text{O}_{18}(\text{OH})_6]^{6-}$, observed in 3. These diagrams show the location of formal Lewis charges.

2. Results and Discussion

2.1. Synthesis and Characterization

Compounds 1, 2 and 3 were prepared in moderate yield through crystallization from aqueous solution initially containing $\text{B}(\text{OH})_3$ and $[\text{Zn}(\text{dien})_2](\text{OH})_2$, $[\text{Zn}(\text{NH}_3)_4](\text{OH})_2$ or $[\text{Zn}(\text{pn})_3](\text{OH})_2$ for 1, 2 and 3, respectively. The hydroxide salts were prepared in situ from the corresponding sulphate salts by the addition of $\text{Ba}(\text{OH})_2$ and removal of precipitated BaSO_4 (Scheme 1).



Scheme 1. Synthesis of Zn(II) hexaborate(2−) and dodecaborate(6−) complexes (dien = $\text{NH}(\text{CH}_2\text{CH}_2\text{NH}_2)_2$, pn = 1,3-diaminopropane).

Compounds **1**, **2** and **3** are formed through self-assembly processes. B(OH)_3 , when dissolved in aqueous solution at moderate to high pH, exists not as boric acid but as a dynamic combinatorial library (DCL) [25,26] of a variety of polyborate anions which are in rapid equilibria [27,28]. Likewise, Zn(II) complexes are labile [29], and a DCL of Zn(II)/amine species are also present in the solution. The products crystallize from solution maximizing energetically favourable solid-state interactions, including coordination bonds, Coulombic attractions, H-bonding and steric effects [30,31].

Compounds **1**, **2** and **3** were characterized spectroscopically (NMR and IR), by thermal DSC/TGA analysis and by single-crystal XRD studies (Section 2.2). They all gave satisfactory bulk elemental analysis.

The thermal TGA/DSC data obtained for **1–3** (see Supplementary Materials) were consistent with the structures determined by single-crystal X-ray diffraction studies (see below) and can be interpreted by multi-step decomposition processes. For **1** this involved loss of interstitial water (<190 °C), further loss of water with cross-condensation of hexaborate(2−) ligands (190–380 °C) and finally oxidation and/or evaporation of the organic dien ligand (380–650 °C) to leave an anhydrous zinc borate $\text{ZnB}_6\text{O}_{10}$ (= $\text{ZnO} \cdot 3\text{B}_2\text{O}_3$) as a glassy residue. Glassy solids with masses consistent with $\text{ZnB}_{12}\text{O}_{19}$ (= $\text{ZnO} \cdot 6\text{B}_2\text{O}_3$) were obtained as the final residues for both **2** and **3** since the initial starting Zn/B ratio was 1:12. The thermal decomposition of **3** followed a similar pattern to **1**. Compound **2** had a TGA trace consistent with loss of initial interstitial water (<110 °C), loss of ammonia (110–250 °C), and final condensation of hexaborate(2−) anions (250–500 °C). Similar thermal behaviour has been observed in other metal polyborate species [12,13,24,32–35], including 1-D zinc hexaborate(1−) coordination polymers $[\text{Zn}(\text{en})\{\text{B}_6\text{O}_7(\text{OH})_6\}] \cdot 2\text{H}_2\text{O}$ and $[\text{Zn}(\text{pn})\{\text{B}_6\text{O}_7(\text{OH})_6\}] \cdot 1.5\text{H}_2\text{O}$ [15]. Magnetic susceptibility χ_m data for **1–3** were $\sim -200 \times 10^{-6} \text{ cm}^3 \cdot \text{mol}^{-1}$ and typical for diamagnetic zinc(II) complexes.

IR spectra can be used to characterize polyborate species since characteristic B–O stretches are generally strong and often diagnostic [36]. Hexaborate(2−) ions, which are never “isolated” and usually found coordinated tridentate to metal centres, have been reported to show such bands at $\sim 953(\text{m}) \text{ cm}^{-1}$ and $808(\text{s}) \text{ cm}^{-1}$. Compound **1** displayed bands at 950(m), 861(m) and 806(s) whilst **2** showed bands at 953(m), 904 (s) and 857(m). Thus, the strong band usually observed at 808 cm^{-1} was absent in **2** and replaced by a strong band at 904 cm^{-1} . This may be a reflection on the unusual centrosymmetric bidentate hexaborate(2−) coordination mode observed in **2**. The IR spectrum of **3** showed peaks at 1047(s), 952(m), 902(s) and 855(m), and there were corresponding absorptions in the reported spectrum of $[(\text{H}_3\text{NCH}_2\text{CH}_2\text{NH}_2)\text{Zn}\{\text{B}_{12}\text{O}_{18}(\text{OH})_6\}]\text{Zn}(\text{en})(\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2) \cdot 8\text{H}_2\text{O}$ [14], which also contains a coordinated dodecaborate(6−) ion. Possible diagnostic absorption bands for this anion have not been described before.

Compounds **1–3** were all insoluble in organic solvents but “dissolved” with decomposition in aqueous solution. ^1H , ^{11}B spectra of these solutions were obtained in D_2O , as were the ^{13}C spectra of **1** and **3**. The ^1H and ^{13}C spectra showed peaks consistent with the organics present and the ^1H spectra additionally displayed at H_2O /exchangeable hydrogen peak (H_2O , NH , BOH) at ~ 4.8 ppm. ^{11}B spectra of **1–3** all showed a single signal at a $+17.4$, $+15.9$ and $+14.0$ ppm, respectively. These signals are all downfield of those calculated [10] (at infinite dilution) for the boron/charge ratio of three ($+13.8$) for a hexaborate($2-$) system, and two ($+11.0$) for the dodecaborate($6-$) ions. This assumes fast $\text{B}(\text{OH})_3/[\text{B}(\text{OH})_4]^-$ exchange [27,28] and is also associated with the pH of the solution. The influence of the zinc(II) ions may also be important here by reducing the effective charge at boron.

2.2. X-ray Diffraction Studies

The structures of **1**, **2** and **3** were determined by single-crystal XRD methods. Crystal data are given in the experimental section and all XRD data are available as Supplementary Materials.

Compounds **1** and **2** both contained the hexaborate($2-$) anion coordinated to a Zn(II) centre and the structures of **1** and **2**, showing their atomic numbering schemes, are shown in Figures 2 and 3, respectively. The anionic complex in **2** was centrosymmetric with the asymmetric unit comprising of half the anion with the zinc($2+$) ion on the inversion centre. Compound **1** was a neutral zinc(II) complex with 0.5 waters of crystallization. The neutral Zn(II) complex, $[\text{Zn}\{\text{B}_6\text{O}_7(\text{OH})_6\}(\text{dien})]$, contained a tridentate ($\kappa^3\text{N}$) dien ligand and a tridentate ($\kappa^3\text{O}$) hexaborate($2-$) ligand. Compound **1** was disordered with two heavy atoms (O10, C4) of the ligand, and associated hydrogen atoms, split in a 1:1 ratio. One position also had an associated water of crystallization (O21). Compound **2** was a salt comprised of $[\text{NH}_4]^+$ cations, $[\text{Zn}\{\text{B}_6\text{O}_7(\text{OH})_6\}_2(\text{H}_2\text{O})_2]^{2-}$ anions and interstitial H_2O molecules. Both hexaborate($2-$) ligands in **2** were bidentate ($\kappa^2\text{O}$) and the coordinated H_2O molecules were *trans*. The Zn–O (hexaborate) distances in **2** { $2.0692(9)$ Å (O11) and $2.1208(9)$ Å (O12)} were within the range of distances observed for **1** { $2.0612(11)$ – $2.1864(10)$ Å} despite the change in coordination mode of the hexaborate($2-$) ligand. The Zn–O (H_2O) distance in **2** was $2.1292(9)$ Å (O21), and the three Zn–N (dien) distances in **1** ranged from $2.1283(14)$ – $2.1473(15)$ Å. The angles about the Zn(II) centres were $82.56(5)$ – $100.26(5)^\circ$ and $166.45(5)$ – $175.22(5)^\circ$ for **1**, and $87.90(3)$ – $92.10(3)^\circ$ and 180.00° for **2**. These angles and distances were consistent with previous reported octahedral complexes of Zn(II) with O and N donor ligands [37]. Bond lengths (B–O) and OBO and BOB bond angles associated with the hexaborate($2-$) ligands in both **1** and **2** were very similar. For example, bond lengths to the central pyramidal O^+ ($1.5154(18)$ – $1.5231(18)$ Å, **1**; $1.5053(15)$ – $1.5247(16)$ Å, **2**) > other bond lengths to four coordinate borons ($1.4407(19)$ – $1.4791(19)$ Å, **1**; $1.4413(18)$ – $1.4889(15)$ Å, **2**) > bond-lengths to three coordinate borons ($1.362(2)$ – $1.418(4)$ Å, **1**; $1.3570(17)$ – $1.3793(17)$ Å, **2**) and consistent with distances and angles previously reported specifically for hexaborate($2-$) complexes [15,32,38,39] and related polyborate systems [8–24,32–36,38–40].




Figure 2. Molecular structure of $[\text{Zn}\{\kappa^3\text{O-B}_6\text{O}_7(\text{OH})_6\}(\kappa^3\text{N-dien})]\cdot 0.5\text{H}_2\text{O}$ (dien = $\text{NH}(\text{CH}_2\text{CH}_2\text{NH}_2)_2$) (1) showing atomic labelling.




Figure 3. Molecular structure of the asymmetric unit of $(\text{NH}_4)_2[\text{Zn}\{\kappa^2\text{O-B}_6\text{O}_7(\text{OH})_6\}(\text{H}_2\text{O})_2]\cdot 2\text{H}_2\text{O}$ (2), showing atomic labelling.

H-bonding interactions are commonly observed in most polyborate solid-state structures. They were observed at many locations in the solid-state structures of **1** and **2** and must be partly responsible for the self-assembly of these structures from their constituents. Compound **1** showed H-bond interactions between the neutral complexes as well as these complexes and the water of crystallization. Compound **2** showed H-bond cation/anion and anion/H₂O interactions. The energetically favourable reciprocal R₂²(8) (Etter [41] nomenclature) O8H8→O3*, O8*H8*→O3 linked hexaborate(2−) units in **1**. There were also unusual R₂²(6) (O9H9→O12*H12*→O4) and R₂²(8) (N2H2→O8* and O13H13→O2*) arrangements between neighbouring hexaborate units in **1**; the latter ring included Zn(1). Compound **2** also had two energetically favourable reciprocal R₂²(8) interactions between neighbouring hexaborate(2−) units (O13H13→O6*, O13*H13*→O6 and O8H8→O3*, O8*H8*→O3). There was also an unusual intramolecular H-bond in **2** between the coordinated H₂O molecule and the hexaborate(2−) ligand (O21H21A→O13) as part of an intramolecular R₁¹(8) system incorporating the Zn1 centre (Figure 4). The coordinated H₂O also H-bonded to a neighbouring hexaborate O21H21B→O2*. O13 is the hexaborate hydroxyl oxygen atom that fulfilled the role as third coordination donor atom in **1** and in other tridentate hexaborate complexes. In this particular local environment of **2**, the energetics of forming this H-bond and the H₂O–Zn coordination bond must outweigh the energetics of a simple borate O–Zn coordinate bond. O13H13 also H-bonded to a neighbouring hexaborate (O13H13→O6*). Full details of these H-bond interactions are given in the Supplementary Materials.




Figure 4. The intramolecular O21H21A→O13 H-bond interaction in **2**. [d(O21–H21) 0.87 Å, d(H21–O13) 1.79 Å; d(O21⋯O13) 2.6446(13) Å; angle O21H21O13, 169.7°] which is part of two R₁¹(8) rings, incorporating Zn–O coordinate bonds (symmetry $i = 2 - x, 1 - y, 2 - z$).

Compound **3** was an ionic compound comprised of [H₃N(CH₂)₃NH₃]²⁺ cations and [(H₃N(CH₂)₃NH₂)ZnB₁₂O₁₈(OH)₆]^{3−} anions, with the anions containing the dodecaborate(6−) ligand coordinated κ³O to a tetrahedral Zn(II) centre which also had a monoprotonated monodentate

κ^1N -H₃N(CH₂)₃NH₂ ligand. There were also seven waters of crystallization per Zn(II) centre. A diagram of the structure is shown in Figure 5.




Figure 5. Diagram of (1,3-pnH₂)₃[(κ^1N -H₃N(CH₂)₃NH₂)Zn{ κ^3O -B₁₂O₁₈(OH)₆}]₂·14H₂O (1,3-pn = 1,3-diaminopropane) (**3**) showing atomic labelling.

The Zn–O (dodecaborate) distances in **3** {1.9592(18) Å (O3)–1.9717(18) Å (O1)} were shorter than those observed for **1** or **2**, reflecting tetrahedral vs. octahedral coordination geometries. The Zn1N1 distance was 2.006(2) Å, and internuclear angles about Zn1 ranged from 103.43(7)–117.86(9)°. These data are very similar to those of the closely related di-Zn(II) complex [(NH₃CH₂CH₂NH₂)Zn{B₁₂O₁₈(OH)₆}Zn(en)(NH₂CH₂CH₂NH₃)] [**14**] that features 1,2-diaminoethane. The dodecaborate(6−) anion (Figure 1b) is comprised of six boroxole rings fused so as to produce a larger central {B₆O₆} ring, with each boron atom within this ring carrying a formal negative charge due to their four-coordinate nature. This anion was first reported in 1990 in the structure of Ag₆[B₁₂O₁₈(OH)₆]·3H₂O [**42**]. The dodecaborate(6−) anion in **3** is closely related to the deprotonated structures found in Na₈[B₁₂O₂₀(OH)₄] [**43**] and Zn₆[B₁₂O₂₄] [**44**]. The central ring oxygen atoms alternate up and down on different sides of the central ring and are ideally set-up to bind tridentate to a metal centre. The dodecaborate(6−) anion has been previously observed to coordinate in a tridentate mode in the following compounds: [(NH₃CH₂CH₂NH₂)Zn{B₁₂O₁₈(OH)₆}–Zn(en)(NH₂CH₂CH₂NH₃)] [**14**], Na₂Cs₄Ba₂[B₁₂O₁₈(OH)₆](OH)₄ [**45**], K₇[(BO₃)Mn{B₁₂O₁₈(OH)₆}]·H₂O [**46**] and K₇[(BO₃)Zn{B₁₂O₁₈(OH)₆}]·H₂O [**47**].

The six four-coordinate boron atoms had B–O distances ranging between 1.441(3)–1.506(3) Å and their O–B–O angles ranged from 106.3(2)–112.1(2)°. The remaining six borons of the anion were three-coordinate and had significantly shorter B–O distances {1.351(3)–1.386(3) Å} and larger O–B–O angles {115.4(2)–123.1(2)°}. These bond lengths are similar to those observed for [(NH₃CH₂CH₂NH₂)Zn(B₁₂O₁₈(OH)₆)Zn(en)(NH₂CH₂CH₂NH₃)] [14], other similarly fused boroxole systems [14,45–47] and the hexaborate(2–) complexes **1** and **2**.

The hydroxyl hydrogen atom, the amino hydrogen atoms of the protonated 1,3-diaminopropane ions and ligands and the waters of crystallization form numerous H-bond interactions and they were presumably responsible—at least in part—for this remarkable self-assembly from mono-boron species. There are numerous cation/anion H-bond interactions, and three of the six potential dodecaborate hydroxyl interactions are R₂²(8): O20H20→O15*, O23H23→O9* and O24H24→O17*, with only the latter reciprocal. “Simple” inter-borate H-bonds originate from O19H19 and O22H22 whilst O21H21 has a non-borate interaction and H-bonds to an H2O (O31). This configuration contrasts with that of [(NH₃CH₂CH₂NH₂)Zn{B₁₂O₁₈(OH)₆}Zn(en)(NH₂CH₂CH₂NH₃)], where all six were involved in R₂²(8) interactions. However, a structural motif that is similar to that found in [(NH₃CH₂CH₂NH₂)Zn{B₁₂O₁₈(OH)₆}Zn(en)(NH₂CH₂CH₂NH₃)] is that amino hydrogen atoms of the uncoordinated nitrogen (N2) of the H₃N(CH₂)₃NH₂ ligand H-bond and link with dodecaborate(6–) units of adjacent complexes. Full details of these H-bond interactions are in the Supplementary Materials.

3. Experimental

3.1. General

All chemicals were obtained from commercial sources. Combustion analysis (CHN) were obtained from OEA laboratories Ltd. in Callington, Cornwall, UK. NMR spectra were obtained on a Bruker Avance spectrometer (Bruker, Coventry, UK) (in D₂O) operating at 400.1 MHz (¹H), 100.6 MHz (¹³C) or 128.4 MHz (¹¹B) with data reported as δ (ppm) with positive chemical shifts to a high frequency of tetramethylsilane (TMS) (¹H, ¹³C) and BF₃·OEt₂ (¹¹B). FTIR spectra were obtained on a PerkinElmer 100 FTIR spectrometer (PerkinElmer, Seer Green, UK) as KBr pellets. TGA/DSC analyses were undertaken in air on an SDT Q600 V4.1 Build 59 instrument (New Castle, DE, USA), using Al₂O₃ crucibles between 10–800 °C with a ramp temperature rate of 10 °C·min^{−1}.

3.2. Synthesis, Spectroscopic, Analytical and Crystallographic data for **1**

A solution of NH(CH₂CH₂NH₂)₂ (dien) (2.16 mL, 20 mmol) in H₂O (5 mL) was added to a solution of ZnSO₄·H₂O (1.79 g, 10 mmol) in H₂O (10 mL). The reaction mixture was stirred at room temperature for 60 min before the addition of Ba(OH)₂·8H₂O (3.15 g, 10 mmol) in H₂O (25 mL). This mixture was rapidly stirred for a further 30 min. The white precipitate of BaSO₄ was removed by filtration and B(OH)₃ (6.18 g, 10 mmol) dissolved in H₂O (50 mL) was added to the filtrate, which was further stirred at room temperature for 3 h. The volume of this solution was reduced to 20 mL by gentle evaporation in a warm water bath. The concentrated solution was left for 10 days in NMR tubes for crystallization and yielded colourless crystals of [Zn(dien){B₆O₇(OH)₆}]·0.46H₂O (**1**) (1.9 g, 42%). Mp ≥ 300 °C. Anal. Calc.: C = 10.5%, H = 4.4%, N = 9.2%. Found: C = 10.7%, H = 4.1%, N = 9.3%. NMR. ¹H/ppm: 2.5 (m, 8H), 4.8 (s, 37H, NH₂, H₂O, OH). ¹³C/ppm: 38.10. ¹¹B/ppm: 17.4. IR (KBr/cm^{−1}): 3549(s), 3384(s), 1642(m), 1442(s), 1427(s), 1362(s), 1249(m), 1193(s) 1108(s), 1028(s), 951(m), 861(m), 808(m). TGA: 100–190 °C, loss of 0.46 interstitial H₂O 2.5 (1.8 calc.); 190–380 °C, condensation of polyborate with loss of three further H₂O 15.2% (13.7% calc.); 380–650 °C, oxidation of dien 38.5% (36.3% calc.); residual ZnB₆O₁₀ 61.5% (63.4% calc.). Magnetic susceptibility: χ_m = −210 × 10^{−6} cm³·mol^{−1}.

Crystal data: C₄H_{19.91}B₆N₃O_{13.5}Zn, M_r = 456.46, monoclinic, C2/c (No. 15), a = 26.0212(3) Å, b = 9.15620(10) Å, c = 13.6318(2) Å, β = 99.5800(10)°, α = γ = 90°, V = 3202.55(7) Å³, T = 100(2) K, Z = 8,

$Z' = 1$, $\mu(\text{Mo K}\alpha) = 1.613 \text{ mm}^{-1}$, 18390 reflections measured, 3651 unique ($R_{int} = 0.0241$) which were used in all calculations. The final wR_2 was 0.0666 (all data) and R_1 was 0.0240 ($I > 2\sigma(I)$).

3.3. Synthesis, Spectroscopic, Analytical and Crystallographic Data for 2

A solution of NH_3 (35%, 2.4 mL, 36 mmol) was added dropwise to a solution of $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$ (1.08 g, 6 mmol) in H_2O (15 mL). The addition of $\text{Ba}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$ (1.89 g, 6 mmol) in H_2O (35 mL) followed by rapid stirring for 15 min resulted in a precipitate of BaSO_4 which was removed by filtration. $\text{B}(\text{OH})_3$ (3.71 g, 60 mmol) dissolved in H_2O (30 mL) was added to the filtrate which was further stirred at room temperature for 30 min. The volume of this solution was reduced to 5 mL by gentle evaporation on a warm water bath and the concentrated solution was left for 3 days in NMR tubes for crystallization and yielded colourless crystals of $[\text{NH}_4]_2[\text{Zn}\{\text{B}_6\text{O}_7(\text{OH})_6\}_2(\text{H}_2\text{O})_2] \cdot 2\text{H}_2\text{O}$ (**2**) (2.1 g, 48%). $\text{Mp} \geq 300 \text{ }^\circ\text{C}$. Anal. Calc.: H = 3.8%, N = 3.8%. Found: H = 4.0%, N = 3.7%. NMR: ^{11}B /ppm: 15.9. IR (KBr/ cm^{-1}): 3212(s), 1400(s), 1357(s), 1048(s), 953(m), 904(m), 857(m). TGA: 100–110 $^\circ\text{C}$, loss of 4 interstitial/coordinated H_2O 10.2% (9.9% calc.); 110–250 $^\circ\text{C}$, loss of 2 NH_3 15.5% (14.8% calc.); 250–500 $^\circ\text{C}$, condensation of polyborate with loss of six further H_2O 31.1 (29.6 calc.); residual $\text{ZnB}_6\text{O}_{19}$ 68.9% (68.2% calc.). Magnetic susceptibility: $\chi_m = -290 \times 10^{-6} \text{ cm}^3 \cdot \text{mol}^{-1}$.

Crystal data: $\text{B}_{12}\text{H}_{28}\text{N}_2\text{O}_{30}\text{Zn}$, $M_r = 731.33$, triclinic, $P-1$ (No. 2), $a = 7.4831(2) \text{ \AA}$, $b = 7.8551(2) \text{ \AA}$, $c = 11.0111(3) \text{ \AA}$, $\alpha = 108.065(2)^\circ$, $\beta = 95.020(2)^\circ$, $\gamma = 90.118(2)^\circ$, $V = 612.68(3) \text{ \AA}^3$, $T = 100(2) \text{ K}$, $Z = 1$, $Z' = 0.5$, $\mu(\text{Mo K}\alpha) = 1.138 \text{ mm}^{-1}$, 16475 reflections measured, 2799 unique ($R_{int} = 0.0314$) which were used in all calculations. The final wR_2 was 0.0559 (all data) and R_1 was 0.0212 ($I > 2\sigma(I)$).

3.4. Synthesis, Spectroscopic, Analytical and Crystallographic Data for 3

A solution of $\text{NH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$ (1,3-pn) (2.52 mL, 30 mmol) in H_2O (10 mL) was added to a solution of $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$ (1.79 g, 10 mmol) in H_2O (10 mL). The reaction mixture was stirred at room temperature for 60 min before the addition of $\text{Ba}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$ (3.15 g, 10 mmol) in H_2O (25 mL). This mixture was rapidly stirred for a further 30 min. The white precipitate of BaSO_4 was removed by filtration and $\text{B}(\text{OH})_3$ (6.18 g, 10 mmol) dissolved in H_2O (50 mL) was added to the filtrate, which was further stirred at room temperature for 30 min. The volume of this solution was reduced to 5 mL by gentle evaporation in a warm water bath. The product was collected by filtration and carefully washed with cold H_2O followed by CH_3COCH_3 , and then dried at 40 $^\circ\text{C}$ for 1 h to yield colourless crystals of $[\text{H}_3\text{N}(\text{CH}_2)_3\text{NH}_3]_3[\text{H}_3\text{N}(\text{CH}_2)_3\text{NH}_2]\text{ZnB}_{12}\text{O}_{18}(\text{OH})_6]_2 \cdot 14\text{H}_2\text{O}$ (**3**) (4.1 g, 46%). $\text{Mp} \geq 300 \text{ }^\circ\text{C}$. Anal. Calc.: C = 10.0%, H = 5.9%, N = 7.8%. Found: C = 9.7%, H = 5.2%, N = 7.8%. NMR. ^1H /ppm: 1.93 (p, 10H, CH_2), 3.01 (t, 20H, CH_2) 4.8 (s, 68H, NH_2 , H_2O , OH). ^{13}C /ppm: 26.9, 37.6. ^{11}B /ppm: 14.0. IR (KBr/ cm^{-1}): 3405(s), 3263(s), 1644(m), 1532(m), 1352(s), 1151(m) 1047(s), 952(m), 902(s), 855(m). TGA: 100–190 $^\circ\text{C}$, loss of 14 interstitial H_2O 14.1% (13.9% calc.); 190–350 $^\circ\text{C}$, condensation of polyborate with loss of six further H_2O 6.9 (6.0 calc.); 350–800 $^\circ\text{C}$, oxidation of organics 22.8% (22.0% calc.); residual $\text{Zn}_2\text{B}_{24}\text{O}_{38}$ 56.6% (55.4% calc.). p-XRD: d-spacing (Å)/(%) rel. int.): 9.98(36), 9.44 (100), 8.50 (54), 8.08 (35), 6.93 (43). Magnetic susceptibility: $\chi_m = -180 \times 10^{-6} \text{ cm}^3 \cdot \text{mol}^{-1}$.

Crystal data: $\text{C}_{7.5}\text{H}_{49}\text{B}_{12}\text{N}_5\text{O}_{31}\text{Zn}$, $M_r = 900.60$, triclinic, $P-1$ (No. 2), $a = 9.3681(2) \text{ \AA}$, $b = 10.6910(2) \text{ \AA}$, $c = 19.2746(4) \text{ \AA}$, $\alpha = 82.954(2)^\circ$, $\beta = 76.156(2)^\circ$, $\gamma = 68.655(2)^\circ$, $V = 1744.44(7) \text{ \AA}^3$, $T = 100(2) \text{ K}$, $Z = 2$, $Z' = 1$, $\mu(\text{Mo K}\alpha) = 0.821 \text{ mm}^{-1}$, 38,867 reflections measured, 7958 unique ($R_{int} = 0.0389$) which were used in all calculations. The final wR_2 was 0.1053 (all data) and R_1 was 0.0425 ($I > 2\sigma(I)$).

3.5. X-ray Crystallography

Single-crystal X-ray crystallography was undertaken at the Engineering and Physical Sciences Research Council (EPSRC) National Crystallography service at the University of Southampton, (Southampton, UK). Suitable crystals of **1**, **2** and **3** were selected and mounted on a MITIGEN holder in perfluoroether oil on a Rigaku FRE+ equipped with HF Varimax confocal mirrors and an AFC12 goniometer and HG Saturn 724+ detector diffractometer. The crystals were kept at $T = 100(2) \text{ K}$ during data collection. Using *Olex2* [48], the structures were solved with the *ShelXT* [49] structure solution

program using the Intrinsic Phasing solution method. The models were then refined with *ShelXL* [50] using least squares minimisation. Cambridge Crystallographic Data Centre (CCDC) 1898912 (1), 1898913 (2), 1898914 (3) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via <http://www.ccdc.cam.ac.uk/conts/retrieving.html> (or from CCDC, 12 Union Road, Cambridge CB2 1EZ, UK; Fax: +44 1223 336033; email deposit@ccdc.ac.uk).

4. Conclusions

The strategy of using more highly charged cationic labile transition-metal complexes to template self-assembly (by crystallization) of polyborate anions from alkaline aqueous solutions originally containing $B(OH)_3$ has resulted in the synthesis of three new zinc polyborate complexes in moderate yields (40–50%). These complexes contain either hexaborate(2−) or dodecaborate(6−) ligands and are stabilized by Zn–O coordinate bonds. The solid-state structures are further stabilized by multiple intramolecular and/or intermolecular H-bond interactions which are prevalent in polyborate structures.

Supplementary Materials: The following are available online at <http://www.mdpi.com/2304-6740/7/4/44/s1>. TGA and single-crystal XRD data. Cif and checkcif files.

Author Contributions: M.A.B. conceived the experiments; M.A.A. synthesized and characterized the complexes and grew the single crystals; P.N.H. and S.J.C. solved the crystal structures; M.A.B. wrote the paper with contributions from all co-authors.

Funding: This research received no external funding.

Acknowledgments: We thank the EPSRC for the use of the X-ray Crystallographic Service (NCS, Southampton, UK).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Schubert, D.M.; Knobler, C.B. Recent studies of polyborate anions. *Phys. Chem. Glasses Eur. J. Glass Sci. Technol. B* **2009**, *50*, 71–78.
2. Grice, J.D.; Burns, P.C.; Hawthorne, F.C. Borate minerals II. A hierarchy of structures based upon the borate fundamental building block. *Can. Mineral.* **1999**, *37*, 731–762.
3. Becker, P. A contribution to borate crystal chemistry: Rules for the occurrence of polyborate anion types. *Z. Kristallogr.* **2001**, *216*, 523–533. [[CrossRef](#)]
4. Heller, G. A survey of structural types of borates and polyborates. *Top. Curr. Chem.* **1986**, *131*, 39–98.
5. Belokoneva, E.L. Borate crystal chemistry in terms of the extended OD theory: Topology and symmetry analysis. *Crystallogr. Rev.* **2005**, *11*, 151–198. [[CrossRef](#)]
6. Christ, C.L.; Clark, J.R. A crystal-chemical classification of borate structures with emphasis on hydrated borates. *Phys. Chem. Miner.* **1977**, *2*, 59–87. [[CrossRef](#)]
7. Beckett, M.A. Recent Advances in crystalline hydrated borates with non-metal or transition-metal complex cations. *Coord. Chem. Rev.* **2016**, *323*, 2–14. [[CrossRef](#)]
8. Wiebcke, M.; Freyhardt, C.C.; Felsche, J.; Engelhardt, G. Clathrates with three-dimensional host structures of hydrogen bonded pentaborate $[B_5O_6(OH)_4]^-$ ions: Pentaborates with the cations NMe_4^+ , NEt_4^+ , $NPhMe_3^+$ and $pipH^+$ ($pipH^+$ = piperidinium). *Z. Naturforsch.* **1993**, *48b*, 978–985. [[CrossRef](#)]
9. Visi, M.Z.; Knobler, C.B.; Owen, J.J.; Khan, M.I.; Schubert, D.M. Structures of self-assembled nonmetal borates derived from α,ω -diaminoalkanes. *Cryst. Growth Des.* **2006**, *6*, 538–545. [[CrossRef](#)]
10. Beckett, M.A.; Coles, S.J.; Davies, R.A.; Horton, P.N.; Jones, C.L. Pentaborate(1−) salts templated by substituted pyrrolidinium cations: Synthesis, structural characterization, and modelling of solid-state H-bond interactions by DFT calculations. *Dalton Trans.* **2015**, *44*, 7032–7040. [[CrossRef](#)] [[PubMed](#)]
11. Beckett, M.A.; Bland, C.C.; Horton, P.N.; Hursthouse, M.B.; Varma, K.S. Supramolecular structures containing “isolated” pentaborate anions and non-metal cations: Crystal structures of $[Me_3NCH_2CH_2OH][B_5O_6(OH)_4]$ and $[4-Mepy_4-MepyH][B_5O_6(OH)_4]$. *J. Organomet. Chem.* **2007**, *692*, 2832–2838. [[CrossRef](#)]

12. Altahan, M.A.; Beckett, M.A.; Coles, C.J.; Horton, P.N. A new polyborate anion $[B_7O_9(OH)_6]^{3-}$: Self-assembly, XRD and thermal properties of *s-fac*- $[Co(en)_3][B_7O_9(OH)_6] \cdot 9H_2O$. *Inorg. Chem. Commun.* **2015**, *59*, 95–98. [[CrossRef](#)]
13. Altahan, M.A.; Beckett, M.A.; Coles, C.J.; Horton, P.N. A new decaoxidoctaborate(2−) anion, $[B_8O_{10}(OH)_6]^{2-}$: Synthesis and characterization of $[Co(en)_3][B_5O_6(OH)_4][B_8O_{10}(OH)_6] \cdot 5H_2O$ (en = 1,2-diaminoethane). *Inorg. Chem.* **2015**, *54*, 412–414. [[CrossRef](#)]
14. Altahan, M.A.; Beckett, M.A.; Coles, C.J.; Horton, P.N. Transition-metal complexes with oxidoborates. Synthesis and XRD characterization of $[H_3NCH_2CH_2NH_2]Zn\{\kappa^3O,O',O''-B_{12}O_{18}(OH)_6-\kappa^1O''\}Zn(en)(NH_2CH_2CH_2NH_3)] \cdot 8H_2O$: A neutral bimetallic zwitterionic polyborate system containing the “isolated” dodecaborate(6−) anion. *Pure Appl. Chem.* **2018**, *90*, 625–632. [[CrossRef](#)]
15. Altahan, M.A.; Beckett, M.A.; Coles, C.J.; Horton, P.N. Two 1-D Coordination Polymers containing Zinc(II) Hexaborates: $[Zn(en)\{B_6O_7(OH)_6\}] \cdot 2H_2O$ (en = 1,2-diaminoethane) and $[Zn(pn)\{B_6O_7(OH)_6\}] \cdot 1.5H_2O$ (pn = (+/−) 1,2-diaminopropane). *Crystals* **2018**, *8*, 470. [[CrossRef](#)]
16. Wang, G.-M.; Sun, Y.-Q.; Yang, G.-Y. Synthesis and crystal structures of two new pentaborates. *J. Solid State Chem.* **2005**, *178*, 729–735. [[CrossRef](#)]
17. He, Y.; Yang, J.; Xi, C.-Y.; Chen, J.-S. Solvothermal synthesis and crystal structure of $Zn(en)_3B_5O_7(OH)_3$. *Chem. Res. Chin. Univ.* **2006**, *22*, 271–273. [[CrossRef](#)]
18. Jiang, H.; Yang, B.-F.; Wang, G.-M. $[Zn(dap)_3][Zn(dap)B_5O_8(OH)_2]_2$: A novel organic-inorganic hybrid chain-like zincoborate made up of $[B_5O_8(OH)_2]^{3-}$ and $[Zn(dap)]^{2+}$ linkers. *J. Clust. Sci.* **2017**, *28*, 1421–1429. [[CrossRef](#)]
19. Wei, L.; Sun, A.-H.; Xue, Z.-Z.; Pan, J.; Wang, G.-M.; Wang, Y.-X.; Wang, Z.-H. Hydrothermal synthesis and structural characterization of a new hybrid zinc borate, $[Zn(dap)_2][B_4O_6(OH)_2]$. *J. Clust. Sci.* **2017**, *28*, 1453–1462. [[CrossRef](#)]
20. Zhao, P.; Cheng, L.; Yang, G.Y. Synthesis and characterization of a new organic-inorganic hybrid borate $[Zn(dab)_{0.5}(dab')_{0.5}(B_4O_6(OH)_2)] \cdot H_2O$. *Inorg. Chem. Commun.* **2012**, *20*, 138–141. [[CrossRef](#)]
21. Paul, A.V.; Sachidananda, K.; Natarajan, S. $[B_4O_9H_2]$ cyclic borate units as the building unit in a family of zinc borate structures. *Cryst. Growth Des.* **2010**, *10*, 456–464. [[CrossRef](#)]
22. Pan, R.; Chen, C.-A.; Yang, B.-F. Two new octaborates constructed of two different sub-clusters and supported by metal complexes. *J. Clust. Sci.* **2017**, *28*, 1237–1248. [[CrossRef](#)]
23. Zhao, P.; Lin, Z.-E.; Wei, Q.; Cheng, L.; Yang, G.-Y. A pillared-layered zincoborate with an anionic network containing unprecedented zinc oxide chains. *Chem. Commun.* **2014**, *50*, 3592–3594. [[CrossRef](#)]
24. Schubert, D.M.; Alam, F.; Visi, M.Z.; Knobler, C.B. Structural characterization and chemistry of an industrially important zinc borate, $Zn[B_3O_4(OH)_3]$. *Chem Mater.* **2003**, *15*, 866–871. [[CrossRef](#)]
25. Sola, J.; Lafuente, M.; Atcher, J.; Alfonso, I. Constitutional self-selection from dynamic combinatorial libraries in aqueous solution through supramolecular interactions. *Chem. Commun.* **2014**, *50*, 4564–4566. [[CrossRef](#)] [[PubMed](#)]
26. Corbett, P.T.; Leclaire, J.; Vial, L.; West, K.R.; Wietor, J.-L.; Sanders, J.K.M.; Otto, S. Dynamic combinatorial chemistry. *Chem. Rev.* **2006**, *106*, 3652–3711. [[CrossRef](#)] [[PubMed](#)]
27. Salentine, G. High-field ^{11}B NMR of alkali borate. Aqueous polyborate equilibria. *Inorg. Chem.* **1983**, *22*, 3920–3924. [[CrossRef](#)]
28. Anderson, J.L.; Eyring, E.M.; Whittaker, M.P. Temperature jump rate studies of polyborate formation in aqueous boric acid. *J. Phys. Chem.* **1964**, *68*, 1128–1132. [[CrossRef](#)]
29. Taube, H. Rates and mechanisms of substitutions in inorganic complexes in aqueous solution. *Chem. Rev.* **1952**, *50*, 69–126. [[CrossRef](#)]
30. Dunitz, J.D.; Gavezzotti, A. Supramolecular synthons: Validation and ranking of intermolecular interaction energies. *Cryst. Growth Des.* **2012**, *12*, 5873–5877. [[CrossRef](#)]
31. Desiraju, G.R. Supramolecular synthons in crystal engineering—A new organic synthesis. *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 2311–2327. [[CrossRef](#)]
32. Altahan, M.A.; Beckett, M.A.; Coles, S.J.; Horton, P.N. Synthesis and characterization of polyborates template by cationic copper(II) complexes: Structural (XRD), spectroscopic, thermal (TGA/DSC) and magnetic properties. *Polyhedron* **2017**, *135*, 247–257. [[CrossRef](#)]
33. Wang, G.-M.; Sun, Y.-Q.; Yang, G.-Y. Synthesis and crystal structures of three new borates templated by transition-metal complexes in situ. *J. Solid State Chem.* **2006**, *179*, 1545–1553. [[CrossRef](#)]

34. Yang, Y.; Wang, Y.; Zhu, J.; Liu, R.-B.; Xu, J.; Meng, C.-C. A new mixed ligand copper pentaborate with square-like, rectangular-like and ellipse-like channels formed via hydrogen bonds. *Inorg. Chim. Acta* **2011**, *376*, 401–407. [[CrossRef](#)]
35. Liu, Z.-H.; Zhang, J.-J.; Zhang, W.-J. Synthesis, crystal structure and vibrational spectroscopy of a novel mixed ligands Ni(II) pentaborate $[\text{Ni}(\text{C}_4\text{H}_{10}\text{N}_2)(\text{C}_2\text{H}_8\text{N}_2)_2][\text{B}_5\text{O}_6(\text{OH})_4]_2$. *Inorg. Chim. Acta* **2006**, *359*, 519–524. [[CrossRef](#)]
36. Li, J.; Xia, S.; Gao, S. FT-IR and Raman spectroscopic study of hydrated borates. *Spectrochim. Acta* **1995**, *51A*, 519–532. [[CrossRef](#)]
37. Archibald, S.J. Zinc. In *Comprehensive Coordination Chemistry II*, 2nd ed.; McCleverty, J.A., Meyer, T.J., Eds.; Elsevier: Amsterdam, The Netherlands, 2003; Volume 6, pp. 1147–1251.
38. Natarajan, S.; Klein, W.; Panthofer, M.; Wuellen, L.V.; Jansen, M. Solution mediated synthesis and structure of the first anionic bis(hexaborato)-zincate prepared in the presence of organic base. *Z. Anorg. Allg. Chem.* **2003**, *629*, 959–962. [[CrossRef](#)]
39. Jemai, N.; Rzaigui, S.; Akriche, S. Piperazine-1,4-dium bis(hexahydroxidoheptaaxidohexaborato- $\kappa^3\text{O},\text{O}',\text{O}''$) cobaltate(II) hexahydrate. *Acta Cryst.* **2014**, *E70*, m167–m169. [[CrossRef](#)] [[PubMed](#)]
40. Beckett, M.A.; Hibbs, D.E.; Hursthouse, M.B.; Malik, K.M.A.; Owen, P.; Varma, K.S. *cyclo*-Boratrissiloxane and *cyclo*-diboratetrasiloxane derivatives and their reactions with amines: Crystal and molecular structure of $(p\text{-BrC}_6\text{H}_4\text{BO})_2(\text{Ph}_2\text{SiO})_2$. *J. Organomet. Chem.* **2000**, *595*, 241–247. [[CrossRef](#)]
41. Etter, M.C. Encoding and decoding hydrogen-bond patterns of organic chemistry. *Acc. Chem. Res.* **1990**, *23*, 120–126. [[CrossRef](#)]
42. Skakibaie-Moghadam, M.; Heller, G.; Timper, U. Die kristallstruktur von $\text{Ag}_6[\text{B}_{12}\text{O}_{18}(\text{OH})_6] \cdot 3\text{H}_2\text{O}$ einen neuen dokekaborat. *Z. Kristallogr.* **1990**, *190*, 85. [[CrossRef](#)]
43. Menchetti, M.; Sabelli, C. A new borate polyanion in the structure of $\text{Na}_8[\text{B}_{12}\text{O}_{20}(\text{OH})_4]$. *Acta Cryst.* **1979**, *B35*, 2488–2493. [[CrossRef](#)]
44. Choudhury, A.; Neeraj, S.; Natarajan, S.; Rao, C.N.R. An open-framework zincoborate formed by $\text{Zn}_6\text{B}_{12}\text{O}_{24}$ clusters. *J. Chem. Soc. Dalton Trans.* **2002**, 1535–1538. [[CrossRef](#)]
45. Zhang, T.-J.; Pan, R.; He, H.; Yang, B.-F.; Yang, G.-Y. Solvothermal synthesis and structure of two new boranes containing $[\text{B}_7\text{O}_9(\text{OH})_5]^{2-}$ and $[\text{B}_{12}\text{O}_{18}(\text{OH})_6]^{6-}$ clusters. *J. Clust. Sci.* **2016**, *27*, 625. [[CrossRef](#)]
46. Zhang, H.-X.; Zhang, J.; Zheng, S.T.; Yang, G.-Y. $\text{K}_7\{(\text{BO}_3)\text{Mn}[\text{B}_{12}\text{O}_{18}(\text{OH})_6]\} \cdot \text{H}_2\text{O}$: First manganese borate based on covalently linked $\text{B}_{12}\text{O}_{18}(\text{OH})_6$ clusters and BO_3 units via Mn^{2+} cations. *Inorg. Chem. Commun.* **2004**, *7*, 781–783. [[CrossRef](#)]
47. Rong, C.; Jiang, J.; Li, Q.-L. Synthesis and transitional metal borates $\text{K}_7\{(\text{BO}_3)\text{Zn}[\text{B}_{12}\text{O}_{18}(\text{OH})_6]\} \cdot \text{H}_2\text{O}$ and quantum chemistry study. *Chinese J. Inorg. Chem.* **2012**, *28*, 2217–2222.
48. Dolomanov, O.V.; Bourhis, L.J.; Gildea, R.J.; Howard, J.A.K.; Puschmann, H. *Olex2*: A complete structure solution, refinement and analysis program. *J. Appl. Cryst.* **2009**, *42*, 339–341. [[CrossRef](#)]
49. Sheldrick, G.M. *ShelXT*-intergrated space-group and crystal structure determination. *Acta Cryst.* **2015**, *A71*, 3–8. [[CrossRef](#)]
50. Sheldrick, G.M. Crystal structure refinement with *ShelXL*. *Acta Cryst.* **2015**, *C27*, 3–8. [[CrossRef](#)]

