

SCIENTIFIC DATA

OPEN

DATA DESCRIPTOR

Worldwide measurements of bioturbation intensity, ventilation rate, and the mixing depth of marine sediments

Martin Solan ¹, Ellie R. Ward ¹, Ellen L. White¹, Elizabeth E. Hibberd¹, Camilla Cassidy ¹, Jasmin M. Schuster ^{1,2}, Rachel Hale ¹ & Jasmin A. Godbold ¹

The activities of a diverse array of sediment-dwelling fauna are known to mediate carbon remineralisation, biogeochemical cycling and other important properties of marine ecosystems, but the contributions that different seabed communities make to the global inventory have not been established. Here we provide a comprehensive georeferenced database of measured values of bioturbation intensity (Db, $n = 1281$), burrow ventilation rate (q , $n = 765$, 47 species) and the mixing depth (L , $n = 1780$) of marine soft sediments compiled from the scientific literature (1864–2018). These data provide reference information that can be used to inform and parameterise global, habitat specific and/or species level biogeochemical models that will be of value within the fields of geochemistry, ecology, climate, and palaeobiology. We include metadata relating to the source, timing and location of each study, the methodology used, and environmental and experimental information. The dataset presents opportunity to interrogate current ecological theory, refine functional typologies, quantify uncertainty and/or test the relevance and robustness of models used to project ecosystem responses to change.

Background & Summary

Marine sediments are known to harbour significant levels of biodiversity that play a key role in biogeochemical cycling, carbon storage and the regulation of climate active gases^{1–3}, but the geographic contribution of extant benthic communities is not well constrained at large scales⁴. Quantifying the extent, timing and way in which organisms transport particles and pore water fluids⁵ has received a considerable amount of attention⁶, yet few attempts to seek universalities, generalities, and particularities have taken place that can inform the architecture of global biogeochemical models^{7–10}. Variations in the intensity of faunal mediation in relation to changing conditions that alter species interactions¹¹, community structure¹² and environmental setting^{13–15} are well-known and mean that the contributions of individual species and/or definable communities cannot be applied universally¹⁶, yet these sources of variation are not generally incorporated into modelling frameworks¹⁷. Indeed, most models are parameterised with broad functional descriptors or selected values of bioturbation that oversimplify or misrepresent temporal and spatial variation in the mediating role of biota^{18–20}, largely because comprehensive compilations of such biological information are not readily available²¹. For these reasons, the treatment of key processes can differ greatly between models such that simulated ecosystem outcomes commonly misalign with ecosystem properties measured at local to regional scales¹⁷, frustrating efforts to accurately project the effects and consequences of environmental change²².

Descriptions of how infaunal invertebrates mediate ecosystem properties are common in the literature and have largely become synonymous with particle displacement and burrow ventilation²³, although alternative descriptors have been considered and emphasised²⁴. As sediment particle reworking often consists of a series of small particle displacement events, standard practice has been to treat the resulting vertical profile of mixing in

¹Ocean and Earth Science, National Oceanography Centre Southampton, University of Southampton, Waterfront Campus, European Way, Southampton, SO14 3ZH, UK. ²Department of Ocean Sciences, Memorial University of Newfoundland, St. John's, Newfoundland, NL, Canada. These authors contributed equally: Martin Solan and Ellie R. Ward. Correspondence and requests for materials should be addressed to M.S. (email: m.solan@soton.ac.uk)




Fig. 1 The geographical location of reported bioturbation parameter values. (a) D_b , the biodiffusion coefficient, (b) L , the sediment mixing depth, (c) q_1 , the ventilation rate for named macro-invertebrate species or for mixed communities taken during active bouts of ventilation and (d) q_2 , the ventilation rate for named macro-invertebrate species or for mixed communities estimated over extended periods of time. Data points may represent multiple observations at that locality. The boundaries of ecoregion domains and divisions³³ (dark blue shading) and provinces³⁴ (light blue shading) are indicated.

Realm	D_b	L	q_1	q_2
Arctic	68	45	0	0
Central IndoPacific	17	48	0	0
Eastern IndoPacific	29	4	0	7
Not allocated (Polar)	4	4	0	0
Not allocated (Temperate)	12	17	0	0
Not allocated (Tropical)	36	21	0	0
Southern Ocean	58	17	0	0
Temperate Australasia	3	5	3	13
Temperate Northern Atlantic	661	1312	206	553
Temperate Northern Pacific	272	122	17	45
Temperate South America	32	25	0	6
Temperate Southern Africa	1	1	0	0
Tropical Atlantic	33	41	0	0
Tropical Eastern Pacific	36	88	0	0
Western IndoPacific	19	30	0	0
Grand total	1281	1780	226	624

Table 1. Number of observations for bioturbation intensity (D_b), mixing depth (L) and ventilation rate (q_1 and q_2) listed by marine realm.

an analogous way to that of diffusive heat transport, calculating a biodiffusion coefficient (D_b , $\text{cm}^2 \text{year}^{-1}$) that describes the rate at which the variance of the location of a particle tracer changes over time within the sediment profile²⁵. Similarly, as the active transfer of fluid by infaunal organisms may be orders of magnitude greater (volumetrically) than particle reworking²⁶, the non-diffusive exchange of pore-water solutes with over-lying water is routinely examined²⁷, but these data have not previously been collated in an accessible archive. The combined effect of particulate and fluid transport on sediment biogeochemical processes is reflected in the vertical colour transition (from brown to olive green/black) of the sediment profile²⁸, dictated by the transition from iron (oxyhydr)oxides at the surface to black sulphidic phases at depth²⁹ that correlate with a variety of environmental drivers³⁰. Hence, regions of high reflectance (brown) in an image represent a well-mixed region of sediment and provide a reasonable approximation of the mixing depth³¹.

Here, motivated by the need to relate changes in ecosystem properties to local heterogeneity rather than global mean conditions^{17,32}, we have collated the extensive repository of information that exists in the primary scientific literature concerning how faunal communities redistribute sediment particles, ventilate their burrows and effect

Method	Db	L	q1	q2
¹³⁷ Cs	43	10	0	0
¹⁴ C	2	59	0	0
²¹⁰ Pb	551	429	0	0
²²² Ra	10	10	0	0
²²⁸ Th	12	0	0	0
²³⁴ Th	423	46	0	0
²³⁵ Th	4	1	0	0
²³⁹²⁴⁰ Pu	34	16	0	0
³² Si	3	3	0	0
⁷ Be	39	8	0	0
calculated	0	71	0	0
Chla	71	7	0	0
Bromide	0	0	10	155
Caesium	0	0	0	4
clearance	0	0	0	34
Doppler	0	0	14	23
dye	0	0	19	28
Eh	0	109	0	0
Em	0	0	95	46
estimate	0	0	1	0
glassbeads	19	0	0	0
hydraulic	0	0	6	6
luminescence	0	2	0	0
luminophores	57	21	0	0
model	0	0	0	6
OrgC	0	1	0	0
oxygen	0	0	6	6
permeability	0	0	1	0
pet	0	0	0	7
piv	0	0	7	2
pressure	0	0	48	265
radon	0	0	11	0
SPI	5	833	0	0
TCO ₂	0	0	0	3
tekbeads	8	8	0	0
thermistor	0	0	6	38
uranine	0	0	2	0
visual	0	104	0	0
xray	0	42	0	0
Grand total	1281	1780	226	623

Table 2. Number of observations for bioturbation intensity (Db), mixing depth (L) and ventilation rate (q1 and q2) categorised against method of quantification. Method definitions are listed as Descriptor categories S1 and S2 in in solan_etal_suppl_info_v3.docx⁴¹.

the depth to which mixing typically occurs in relation to their physical location. Our hope is that the inherent spatial and temporal heterogeneity shown within these data will be embraced by modellers, statisticians and ecologists and contribute to the development of next generation biogeochemical models that can better inform conservation and management strategies.

Methods

We searched the Thomson Reuters Web of Science collection (<http://www.webofknowledge.com>, accessed 07/03/2019) using a 'General Search' across all databases with the search term (i) *bioturbation*, (ii) *sediment profile imag**, and (iii) *bioirrigation OR burrow ventilation* in the titles and key words of all document types, in all languages, for the publication years 1864 to 2018. Citation returns were manually searched for reported values of the sediment mixing depth (L, cm)^{30,31}, the biodiffusion coefficient (D_b , $\text{cm}^2 \text{year}^{-1}$) estimated from models of sediment particle reworking^{6,25}, and the rate of ventilation (q , $\text{ml h}^{-1} \text{ind.}^{-1}$) for named macro-invertebrate species or mixed communities. These data for L and D_b supersede records collated elsewhere⁷⁻¹⁰ and include observations from the older literature (pre-1970) cited by the authors of the returns from our search.

	Db	L	q1	q2
Season				
Spring	295	332	81	125
Summer	266	607	28	115
Autumn	245	233	3	9
Winter	70	31	41	70
Multiple	68	72	1	78
Total:	944	1275	154	397
Depth category				
0–50 m	277	956	225	599
50–200 m	281	276	0	16
200–1000 m	245	145	0	9
1000–4000 m	303	251	0	0
4000–6000 m	175	152	0	0
>6000 m	0	0	0	0
Total:	1281	1780	225	624
Class				
Bivalvia	—	—	9 [4]	22 [5]
Echinoidea	—	—	5 [2]	10 [1]
Ophiuroidea	—	—	0 [0]	9 [1]
Malacostraca	—	—	37 [9]	89 [11]
Polychaeta	—	—	157 [11]	427 [21]
Community	—	—	18 [n/a]	67 [n/a]
Total:			226	624

Table 3. Number of observations for bioturbation intensity (Db), mixing depth (L) and ventilation rate (q1 and q2) for season and depth category. For q, the number of experimental observations are listed against taxonomic class. The number of species considered within each taxonomic class are indicated in square brackets.

For each unique record, we collated associated environmental metadata (latitude, longitude, water depth, sedimentation rate), information on the methodology used, and details about the timing (year, season, month) and ecoregion (following accepted biogeographical typologies)^{33,34} from the original publication, personal communication with the corresponding author and/or from third party sources of information. Where specific values were not presented in the original publication and had to be derived, values were extracted from graphical summaries using *Web Plot Digitiser* (<https://automeris.io/WebPlotDigitizer/>). When the location of a study was not provided, latitude and longitude coordinates and/or water depth were retrieved from Google Earth (<http://earth.google.com/>) and manually cross referenced with site descriptions within the source publication. Following standard practice⁹, the seasonal offset between Northern (NH) and Southern (SH) hemisphere was corrected by attributing a nominal season to each study: Spring, April–June in the NH or October–December in the SH; Summer, July–September in the NH or January–March in the SH; Autumn, October–December in the NH or April–June in the SH; or Winter, January–March in the NH or July–September in the SH. Due to variations in seasonal timing at any given latitude, the scheme is not necessarily representative of geographical clines in forcing. Data collected from multiple months or unspecified periods are also included. The methodology used to generate each record includes 21 techniques for L and Db (reviewed in ref.³⁵) and 18 techniques for q (reviewed in ref.³⁶).

As species ventilation behaviour varies over time³⁷, we distinguish ventilation measurements based solely on active bouts of ventilation (q1, an indication of peak activity) from those estimated over extended periods of time that span rest periods (q2, a more representative indication of species contribution). Similarly, in recognising that experimental configuration^{24,38} and the geometry of the sediment–water interface³⁹ can influence species behaviour, our database includes information on aquaria dimensions. Given the time span of the studies under consideration, species nomenclature has been standardised in line with the *World Register of Marine Species*⁴⁰.

Data Records

Data records are available via an unrestricted repository hosted by Harvard Dataverse⁴¹. Data represent reported values for the biodiffusion coefficient (D_b , $\text{cm}^2 \text{year}^{-1}$; Fig. 1a) and/or the sediment mixing depth (L, cm; Fig. 1b) for specific locations and can be found in `solan_etal_DbL.csv`. Separately, the data set also includes volumetric ventilation flow rates (q , $\text{ml h}^{-1} \text{ind.}^{-1}$) for named macro-invertebrate species or mixed communities taken during active bouts of ventilation (q1, Fig. 1c) and/or estimated over extended periods of time (q2, Fig. 1d) for specific locations. These can be found in `solan_etal_q.csv`. The number of records within the dataset are listed to ecoregion (Table 1), method of quantification (Table 2) and by season and water depth (Table 3). Table 3 also includes the number of experimental observations of q1 and q2 listed against taxonomic class. A summary of the definitions for the descriptors (=column headings) used in the Db and L (Descriptor categories S1) and q (Descriptor categories S2) datasets are documented separately in `solan_etal_suppl_info_v3.docx`⁴¹.

Technical Validation

The data has been collated from the peer-reviewed literature (Data Source S1 in solan_et_al_suppl_info_v3.docx)⁴¹ and has undergone rigorous quality control prior to publication. Each individual record (unique identification number) in the dataset is traceable to the point of origin (data source identification number)⁴¹.

Usage Notes

We have included all reported values from the literature without prejudice or downstream processing steps. Reporting errors and updates of the data will be periodically issued. Users should use the latest version of the data listed (under the 'versions' tab) at Harvard Dataverse⁴¹ and maintained at *Bioturbation Online* (<http://bioturbation.online>). This contribution is based on data release 3.0. There are no limitations on the use of these data.

References

- Solan, M. *et al.* Extinction and ecosystem function in the marine benthos. *Science* **306**, 1177–1180 (2004).
- Martin, R. E. Secular increase in nutrient levels through the Phanerozoic: implications for productivity, biomass and diversity of the marine biosphere. *Palaios* **11**, 209–219 (1996).
- Covich, A. P. *et al.* The role of biodiversity in the functioning of freshwater and marine benthic ecosystems. *Bioscience* **54**, 767–775 (2004).
- Wei, C. L. *et al.* Global patterns and predictions of seafloor biomass using random forests. *Plos One* **5**, e15323, <https://doi.org/10.1371/journal.pone.0015323> (2010).
- Pearson, T. H. Functional group ecology in soft-sediment marine benthos: The role of bioturbation. *Oceanogr. Mar. Biol. Ann. Rev.* **39**, 233–267 (2001).
- Meysman, F. J. R., Boudreau, B. & Middelburg, J. J. Relations between local, non-local, discrete and continuous models of bioturbation. *J. Mar. Res.* **61**, 391–410 (2003).
- Boudreau, B. P. Is burial velocity a master parameter for bioturbation? *Geochim. Cosmochim. Acta* **58**, 1243–1249 (1994).
- Boudreau, B. P. Mean mixed depth of sediments: The wherefore and the why. *Limnol. Oceanogr.* **43**, 524–526 (1998).
- Teal, L. R., Bulling, M. T., Parker, E. R. & Solan, M. Global patterns of bioturbation intensity and mixed depth of marine soft sediments. *Aquatic Biol.* **2**, 207–218 (2008).
- Lecroart, P. *et al.* Bioturbation, short-lived radioisotopes, and the tracer-dependence of biodiffusion coefficients. *Geochim. Cosmochim. Acta* **74**, 6049–6063 (2010).
- Godbold, J. A. & Solan, M. Long-term effects of warming and ocean acidification are modified by seasonal variation in species responses and environmental conditions. *Phil. Trans. R. Soc. B.* **368**, UNSP 20130186 (2013).
- Wohlgemuth, D., Solan, M. & Godbold, J. A. Specific arrangements of species dominance can be more influential than evenness in maintaining ecosystem process and function. *Sci. Rep.* **6**, <https://doi.org/10.1038/srep39325> (2016).
- Berelson, W. M. *et al.* Modelling bio-irrigation rates in the sediments of Port Phillip Bay. *Mar. Freshwater Res.* **50**, 573–579 (1999).
- Morys, C., Forster, S. & Graf, G. Variability of bioturbation in various sediment types and on different spatial scales in the southwestern Baltic Sea. *Mar. Ecol. Prog. Ser.* **557**, 31–49 (2016).
- Wohlgemuth, D., Solan, M. & Godbold, J. A. Species contributions to ecosystem process and function can be population dependent and modified by biotic and abiotic setting. *Proc. R. Soc. B.* **284**, 20162805 (2017).
- Murray, F., Douglas, A. & Solan, M. Species that share traits do not necessarily form distinct and universally applicable functional effect groups. *Mar. Ecol. Prog. Ser.* **516**, 23–34 (2014).
- Snelgrove, P. *et al.* Global carbon cycling on a heterogeneous seafloor. *Trends Ecol. Evol.* **33**, 96–105 (2018).
- Butenschoen, M. *et al.* ERSEM 15.06: a generic model for marine biogeochemistry and the ecosystem dynamics of the lower trophic levels. *Geosci. Model Dev.* **9**, 1293–1339 (2016).
- Yakushev, E. V. *et al.* Bottom RedOx Model (BROM v.1.1): a coupled benthic–pelagic model for simulation of water and sediment biogeochemistry. *Geosci. Model Dev.* **10**, 453–482 (2017).
- Lessin, G. *et al.* Modelling marine sediment biogeochemistry: Current knowledge gaps, challenges, and some methodological advice for advancement. *Front. Mar. Sci.* **5**, 19, <https://doi.org/10.3389/fmars.2018.00019> (2018).
- Tyler, E. H. M. *et al.* Extensive gaps and biases in our knowledge of a well-known fauna: implications for integrating biological traits into macroecology. *Glob. Ecol. Biogeogr.* **21**, 922–934 (2012).
- Sweetman, A. K. *et al.* Major impacts of climate change on deep-sea benthic ecosystems. *Elem. Sci. Anth.* **5**, 4, <https://doi.org/10.1525/elementa.203> (2017).
- Kristensen, E. *et al.* What is bioturbation? The need for a precise definition for fauna in aquatic sciences. *Mar. Ecol. Prog. Ser.* **446**, 285–302 (2012).
- Hale, R., Mavrogordato, M. N., Tolhurst, T. J. & Solan, M. Characterizations of how species mediate ecosystem properties require more comprehensive functional effect descriptors. *Sci. Rep.* **4**, <https://doi.org/10.1038/srep06463> (2014).
- Crank, J. *The mathematics of diffusion* (Oxford University Press, 1975).
- Dornoffer, T. M., Waldbusser, G. G. & Meile, C. Modeling lugworm irrigation behavior effects on sediment nitrogen cycling. *Mar. Ecol. Prog. Ser.* **534**, 121–134 (2015).
- Riisgård, H. U. & Larsen, P. S. Water pumping and analysis of flow in burrowing zoobenthos: an overview. *Aquatic Ecol.* **39**, 237–258 (2005).
- Lyle, M. The brown-green color transition in marine sediments: a marker of the Fe(III)–Fe(II) redox boundary. *Limnol. Oceanogr.* **2**, 1026–1033 (1983).
- Statham, P. J. *et al.* Extending the applications of sediment profile imaging to geochemical interpretations using colour. *Cont. Shelf Res.*, <https://doi.org/10.1016/j.csr.2017.12.001> (2017).
- Teal, L. R., Parker, E. R. & Solan, M. Sediment mixed layer as a proxy for benthic ecosystem process and function. *Mar. Ecol. Prog. Ser.* **414**, 27–40 (2010).
- Germano, J. D., Rhoads, D. C., Valente, R. M., Carey, D. A. & Solan, M. The use of Sediment Profile Imaging (SPI) for environmental impact assessments and monitoring studies: lessons learned from the past four decades. *Oceanogr. Mar. Biol. Ann. Rev.* **49**, 235–298 (2011).
- Bates, A. E. *et al.* Biologists ignore ocean weather at their peril. *Nature* **560**, 299–301 (2018).
- Bailey, R. G. *Ecoregions: The ecosystem geography of the oceans and continents* (New York: Springer, 1998).
- Spalding, M. D. *et al.* Marine ecoregions of the world: a bioregionalization of coastal and shelf areas. *Bioscience* **57**, 573–583 (2007).
- Maire, O. *et al.* Quantification of sediment reworking rates in bioturbation research: a review. *Aquatic Biol.* **2**, 219–238 (2008).
- Roskosch, A., Hupfer, M., Nutzmann, G. & Lewandowski, J. Measurement techniques for quantification of pumping activity of invertebrates in small burrows. *Fundam. Appl. Limnol.* **178**, 89–110 (2011).
- Timmerman, K., Banta, G. T. & Glud, R. N. Linking *Arenicola marina* irrigation behavior to oxygen transport and dynamics in sandy sediments. *J. Mar. Res.* **64**, 915–938 (2006).
- Bulling, M. T. *et al.* Species effects on ecosystem processes are modified by faunal responses to habitat composition. *Oecologia* **158**, 511–520 (2008).

39. Meysman, F. J. R. *et al.* Quantifying biologically and physically induced flow and tracer dynamics in permeable sediments. *Biogeosciences* **4**, 627–646 (2007).
40. Horton, T. *et al.* World Register of Marine Species. VLIZ, <https://doi.org/10.14284/170> (2017).
41. Solan, M. A global database of measured values of benthic invertebrate bioturbation intensity, ventilation rate, and the mixed depth of marine soft sediments. *Harvard Dataverse*, <https://doi.org/10.7910/DVN/GBELFW> (2018).

Acknowledgements

M.S., J.A.G. and R.H. were supported by Work Package 2 of the Shelf Sea Biogeochemistry Programme (NE/K001906/1, 2011–2017), jointly funded by the Natural Environment Research Council (NERC) and the Department for Environment, Food and Rural Affairs (Defra).

Author Contributions

M.S. initiated the database, contributed to data collection and wrote the manuscript. E.R.W. and M.S. coordinated data collection, were responsible for quality control and constructed the database. M.S., E.R.W., E.L.W., E.E.H., C.C., J.S., R.H. and J.A.G. contributed to data collection. R.H. and J.A.G. contributed to database design and reviewed and provided critical commentary on the manuscript prior to submission.

Additional Information

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

The Creative Commons Public Domain Dedication waiver <http://creativecommons.org/publicdomain/zero/1.0/> applies to the metadata files associated with this article.

© The Author(s) 2019