



**Small Vessel Disease Pathological Changes in  
Neurodegenerative and Vascular Dementias concomitant  
with Autonomic Dysfunction**

Journal:	<i>Brain Pathology</i>
Manuscript ID	BPA-19-02-RA-044.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Hase, Yoshiki; Newcastle University, Institute of Neuroscience Polvikoski, Tuomo; Newcastle Upon Tyne Hospitals NHS Foundation Trust, Cellular Pathology Department; Newcastle University, Institute of Neuroscience Firbank, Micheal; Newcastle University, Institute of Neuroscience Craggs, Lucinda; Newcastle University, Institute of Neuroscience; Hawthorne, Emily; Newcastle University, Institute of Neuroscience Platten, Charlotte; Newcastle University, Institute of Neuroscience Stevenson, William; Newcastle University, Institute of Neuroscience Deramecourt, Vincent; Univ Lille Nord de France, Histology and Pathology Department Ballard, Clive; University of Exeter, School of Medicine Kenny, RoseAnne; Trintiy College Institute for Neuroscience, Neuroscience Perry, Robert; Newcastle University, Institute of Neuroscience Ince, Paul; University of Sheffield, Department of Neuroscience Carare, Roxana; University of Southampton, Clinical Neurosciences Allan, Louise; University of Exeter, Institute of Health Research Horsburgh, Karen; University of Edinburgh, Neuroscience; Kalaria, Raj; Newcastle University, Institute of Neuroscience
Keywords:	Alzheimer's disease, Autonomic Dysfunction, Dementia with Lewy Bodies, Parkinson's Disease with Dementia, Small Vessel Disease, Vascular Dementia

SCHOLARONE™  
Manuscripts

1  
2  
3 **Small Vessel Disease Pathological Changes in Neurodegenerative and Vascular Dementias**  
4 **concomitant with Autonomic Dysfunction**  
5  
6  
7

8 Yoshiki Hase,<sup>1</sup> Tuomo M Polvikoski,<sup>1</sup> Michael J Firbank,<sup>1</sup> Lucinda JL Craggs,<sup>1</sup> Emily  
9 Hawthorne,<sup>1</sup> Charlotte Platten,<sup>1</sup> William Stevenson,<sup>1</sup> Vincent Deramecourt,<sup>2</sup> Clive Ballard,<sup>3</sup>  
10 Rose Anne Kenny,<sup>4</sup> Robert H Perry,<sup>1</sup> Paul Ince,<sup>5</sup> Roxana O Carare,<sup>6</sup> Louise M Allan,<sup>3</sup> Karen  
11 Horsburgh,<sup>7</sup> and Raj N Kalaria,<sup>1\*</sup>  
12  
13  
14  
15  
16  
17

- 18 1. Institute of Neuroscience, Newcastle University, Newcastle upon Tyne, United Kingdom  
19 2. University Lille Nord de France, Histology and Pathology Department, Lille University  
20 Hospital, Lille, France  
21  
22 3. School of Medicine, University of Exeter, Exeter, United Kingdom  
23  
24 4. Department of Medical Gerontology, Trinity College Dublin, Dublin, Ireland  
25  
26 5. Sheffield Institute for Translational Neuroscience, University of Sheffield, Sheffield, United  
27 Kingdom  
28  
29 6. Clinical and Experimental Sciences Unit, Faculty of Medicine, University of Southampton,  
30 Southampton, United Kingdom  
31  
32 7. Centre for Neuroregeneration, University of Edinburgh, Little France Crescent. Edinburgh.  
33 United Kingdom  
34  
35  
36  
37

38 **Running title:** Autonomic Dysfunction, Cerebrovascular pathology and Dementia  
39  
40  
41

42  
43 \* Corresponding author: Professor Raj N. Kalaria  
44

45 Neurovascular Research Group, Institute of Neuroscience, Newcastle University, Campus for  
46 Ageing and Vitality, Newcastle upon Tyne, NE4 5PL, United Kingdom; Tel: 0191 208 1352;  
47 Fax: 0191 208 1301; E-mail: [raj.kalaria@ncl.ac.uk](mailto:raj.kalaria@ncl.ac.uk)  
48  
49

50  
51 Number of words: 4,563 words (excluding abstract, tables, figure legends, references, and  
52 supplemental data); Abstract; 296 words; Number of Figures: 3; Number of Tables: 2; Number  
53 of references 50  
54  
55  
56  
57

## Abstract

We performed a clinicopathological study to assess the burden of small vessel disease (SVD) type of pathological changes in elderly demented subjects, who had clinical evidence of autonomic dysfunction, either carotid sinus hypersensitivity or orthostatic hypotension or both or had exhibited unexpected repeated falls. Clinical and neuropathological diagnoses in 112 demented subjects comprised dementia with Lewy bodies (DLB), Parkinson's disease with dementia (PDD), Alzheimer's disease (AD), Mixed dementia (mostly AD-DLB) and vascular dementia (VaD). Of these, 12 DLB subjects had no recorded unexpected falls in life and therefore no evidence of concomitant autonomic dysfunction. A further 17 subjects were assessed as ageing controls without significant pathology or signs of autonomic dysfunction. We quantified brain vascular pathological changes and determined severities of neurodegenerative lesions including  $\alpha$ -synuclein pathology. We found moderate-severe vascular changes and high vascular pathology scores ( $P < 0.01$ ) in all neurodegenerative dementias and as expected in VaD compared to similar age controls. Arteriolosclerosis, perivascular spacing and microinfarcts were frequent in the basal ganglia and frontal white matter (WM) across all dementias whereas small infarcts ( $< 5$  mm) were restricted to VaD. In a sub-set of demented subjects, we found that vascular pathology scores were correlated with WM hyperintensity volumes determined by MRI in life ( $P < 0.02$ ). Sclerotic index values were increased by  $\sim 50\%$  in both the WM and neocortex in all dementias compared to similar age controls. We found no evidence for increased  $\alpha$ -synuclein deposition in subjects with autonomic dysfunction. Our findings suggest greater SVD pathological changes occur in the elderly diagnosed with neurodegenerative dementias including DLB and who acquire autonomic dysfunction. SVD changes may not necessarily manifest in clinically overt symptoms but they likely confound motor or cognitive dysfunction. We propose autonomic dysfunction promotes chronic cerebral hypoperfusion to impact upon ageing-related neurodegenerative processes and characterise their end-stage clinical syndromes.

## Keywords

Alzheimer's disease, Autonomic Dysfunction, Dementia, Dementia with Lewy Bodies, Microvascular Pathology, Mixed Dementia, Parkinson's Disease with Dementia, Small Vessel Disease, Vascular Dementia

**Abbreviations**

AD, Alzheimer's disease; CSH, carotid sinus hypersensitivity; CSM, carotid sinus massage; DLB, dementia with Lewy bodies; H&E, Haematoxylin and Eosin; OH, orthostatic hypotension; PD, Parkinson's disease; PDD, Parkinson's disease with dementia; SI, sclerotic index; SVD, small vessel disease; WM, white matter; WMH, white matter hyperintensity.

For Peer Review Only

## Introduction

Autonomic dysfunction occurs commonly after 50 years of age. At least 40% of the elderly with Dementia with Lewy bodies (DLB), Parkinson's disease with dementia (PDD), vascular dementia (VaD) and Alzheimer disease (AD) will exhibit some form of autonomic dysfunction. They will invariably develop carotid sinus hypersensitivity (CSH), orthostatic hypotension (OH) or neurocardiovascular instability, associated with hypotension and bradycardia (2, 7, 23). CSH involving an excessive fall in arterial pressure (53) is reported to occur in 35-58% of elderly with AD and DLB (24). OH develops in nearly 50% of Parkinsonians (16, 18, 43, 49, 50) and may be present in over 90% of PDD patients (2). OH is an independent risk for incident dementia (9, 55) and often accompanied by disabling symptoms, impaired quality of life, dizziness, nausea, sweating, and loss of consciousness (1, 4, 32).

The vascular hypothesis of autonomic dysfunction proposes that recurrent episodic hypotension results in chronic cerebral hypoperfusion, especially in the frontal lobe. This in turn causes anoxic damage and leads to impaired cognitive function (45). While the true burden of asymptomatic autonomic dysfunction is unknown (31), it is crucial to protect the ageing brain from fluctuations between hyper- and hypotension, which are known to be associated with white matter (WM) changes (58). We previously showed increased WM hyperintensities (WMHs) (24) in DLB patients with autonomic dysfunction. WMHs are also associated with a postural drop in blood pressure in PD (39). Here, we tested the hypothesis that there is increased small vessel disease (SVD) or microvascular pathology in patients with clinical evidence of autonomic dysfunction or in those with repeated unexplained falls or syncope and progress to dementia. We performed clinicopathological studies to explore the extent of SVD pathological changes (51) in patients exhibiting signs of autonomic dysfunction and had developed dementia.

## Materials and Methods

### *Study design and subjects*

This study involved the assessment of a total of 129 subjects. Demographic, clinical and pathological details of all the subjects are given in Tables 1 and 2. Initially, we screened clinical records and post-mortem reports filed between 1992 and 2017. Of the 350 records screened, we

1  
2  
3 identified 77 demented subjects, who were assessed for autonomic function in the Falls and  
4 Syncope Service (FASS) at the Royal Victoria Infirmary, Newcastle upon Tyne and had had a  
5 post-mortem (Table 1). We then found 23 additional dementia cases with clinical records  
6 reporting repeated unexplained falls prior to death. This information was used as a proxy for the  
7 presence of autonomic dysfunction (1, 2). In total, we identified 100 demented subjects with  
8 direct clinical or reported evidence for autonomic dysfunction (Table 2). The unexplained falls  
9 in these individuals were deemed to be for reasons other than arthritis, osteoporosis or joint  
10 disease. We also identified an additional 12 DLB cases with no reported unexplained falls prior  
11 to death (Table 2). This group with likely no or minimal autonomic dysfunction was compared  
12 with those DLB subjects, who had clear evidence of autonomic dysfunction or repeated  
13 unexplained falls. We could not find any cases diagnosed with PDD, AD, VaD or mixed  
14 dementia and without evidence of autonomic dysfunction and had come to post-mortem.  
15 Furthermore, we identified 17 similar age controls with no evidence of reported falls, syncope or  
16 autonomic dysfunction (Table 2). We could not find any control subjects, who were clinically  
17 assessed in the FASS for autonomic function or had falls or syncope and had had a post-mortem.  
18 There were also no dementia cases tested in the FASS without autonomic dysfunction and had  
19 had an autopsy.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33 Patients suspected of autonomic dysfunction with recurrent symptoms of syncope or falls  
34 had evidence of CSH or OH or both (Table 1). CSH was diagnosed by an exaggerated  
35 haemodynamic response to longitudinal carotid sinus massage (CSM). The exaggerated  
36 response was defined as >3 seconds of asystole (cardioinhibitory CSH) or a fall in systolic blood  
37 pressure in excess of 30 mmHg whilst supine or 50 mmHg whilst in the upright position  
38 (vasodepressor CSH) during 5 seconds of carotid sinus stimulation by longitudinal massage over  
39 the carotid sinus (24, 28, 29, 37). OH was defined as >30 mmHg drop in systolic blood pressure  
40 after an active stand test (Head-up tilt test) (11, 54). Patients diagnosed with OH had undergone  
41 CSM and were confirmed not meet the clinical diagnostic criteria for CSH. The mean interval  
42 between the time of autonomic function assessment and death was 1.92 years with a range from  
43 0 to 7 years. Almost 80% of the subjects died 3 years after they were assessed in the FASS. The  
44 neuropsychometric assessments included the revised Cambridge Cognition Examination  
45 (CAMCOG) battery and Mini-Mental State Examination (MMSE). Sub-scores for cognitive  
46 domains including memory, orientation and other domains of executive function were generated  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 from the total scores (3). The suspected clinical diagnosis of dementia was confirmed at  
4 monthly clinicopathological consensus meetings where clinicians met with the pathologists to  
5 designate a final diagnosis in accordance with current criteria including the diagnostic and  
6 statistical manual of mental disorders, 4th Edition (DSM-IV).  
7  
8  
9

10  
11 Study subjects were participants of the Newcastle longitudinal prospective dementia  
12 series. They had a clinical diagnosis of either DLB, PDD, mixed dementia (Mixed), AD or VaD.  
13 VaD subjects were participants of the Newcastle Cognitive Function After Stroke study (3).  
14 Age-matched control subjects aged >70 years were either part of previous prospective studies or  
15 based on unrelated brain donations to the Newcastle Brain Tissue Resource (NBTR). They were  
16 included as controls if they had not been diagnosed with autonomic dysfunction, cognitive  
17 impairment or any neurological or psychiatric illness. Ethics approval was granted by local  
18 research ethics committees of the Newcastle upon Tyne NHS Foundation Hospitals Trust.  
19 Permission for use of brains for post-mortem research was also granted by consent from the  
20 participants themselves or next-of-kin or family member. Brain tissues were retained in and  
21 obtained from the NBTR.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

### 33 *Magnetic Resonance Imaging*

34  
35 A total of 42 subjects who had post-mortem had undergone brain MRI in life. T2-weighted MRI  
36 images (T2WI) were obtained in each case, and WMH volumes (vWMHs) estimated using  
37 previously validated in house software (12). The mean time from MRI scans to death (years  $\pm$   
38 SEM) for all the subjects was  $6.4 \pm 0.6$  years. For each group this was as follows: DLB  $3.6 \pm 0.7$   
39 years; PDD  $4.3 \pm 1.0$  years; Mixed  $4.6 \pm 0.6$  years; AD  $8.1 \pm 1.1$  years and VaD  $6.4 \pm 1.3$  years.  
40 For controls, without autonomic dysfunction it was  $12.2 \pm 1.4$  years (n=5). The percentage of  
41 WMH volume per total brain volume (%vWMH/vTB) in disease cases was calculated. We then  
42 performed correlation analysis between WMH volume and vascular pathology scores from all  
43 the demented subjects with autonomic dysfunction (Figure 2).  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

### 54 *Brain tissues and Neuropathological analysis*

1  
2  
3 Neuropathological assessment was carried out as described previously (3). Briefly,  
4 haematoxylin and eosin (H&E) staining was used for assessment of structural integrity and  
5 infarcts, Nissl and Luxol Fast blue staining for cellular patterns and myelin loss, Bielschowsky's  
6 silver impregnation and amyloid- $\beta$  for Consortium to Establish a Registry for Alzheimer's  
7 Disease (CERAD) rating of neuritic plaques, Gallays stain for neuritic pathology, and tau  
8 immunohistochemistry for Braak staging of neurofibrillary tangles. Large sections from the  
9 frontal, temporal (including hippocampal formation), parietal and occipital lobes, caudate-  
10 putamen, brain stem including midbrain, pons and medulla oblongata, and cerebellum were  
11 examined from sampled regional blocks as identified in the Newcastle brain map (20, 42). The  
12 clinical diagnosis of DLB and PDD were confirmed according to established criteria (30). The  
13 clinical diagnosis of AD was confirmed on evidence of significant Alzheimer's-type pathology  
14 incorporating Braak stages V–VI, moderate-severe CERAD (26) and high ABC scores per  
15 National Institute of Aging-Alzheimer's Association guidelines (36), in the general absence of  
16 significant vascular pathology (Table 2). The clinical diagnosis of vascular dementia (VaD) was  
17 made when there were multiple or cystic infarcts, lacunae, border-zone infarcts, microinfarcts  
18 and small vessel disease, and pathologically confirmed as Braak stage  $\leq$  IV (20, 21). Mixed AD  
19 and VaD case was classified when there was sufficient degree of pathology to reach Braak V–VI  
20 and significant vascular pathology (6). We also included cases with Mixed dementia, who met  
21 two or more neuropathological diagnostic criteria for DLB, PDD and AD.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

37 Vascular pathology scores were derived from the presence of vascular lesions/pathologies  
38 in brain areas, including the frontal lobe at the level of the olfactory bulbs, temporal lobe at level  
39 of the anterior hippocampus, and basal ganglia at level of mamillary body. Lesions including  
40 arteriosclerosis, cerebral amyloid angiopathy (CAA), perivascular haemosiderin deposition,  
41 perivascular space dilatation in the deep and juxtacortical WM, myelin loss, and cortical micro  
42 ( $<0.5$  mm), small ( $<5$  mm) and large ( $>0.5$  cm) infarcts were recorded with increasing severity  
43 resulting in greater scores (10). These measures were compatible with the recently established  
44 vascular cognitive impairment neuropathology consortium criteria (46). The vascular pathology  
45 scores were determined as detailed previously (10) with  $>95\%$  agreement between scoring  
46 performed by VD and RNK. CAA was scored using the system described previously (25).  
47 Tissues from control subjects had occasional ageing-related pathology and were classified as 'no  
48 pathological diagnosis' (Table 2). Except for neuropathological examination (TP, RHP, PI and  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 RNK), all subsequent morphological analyses were always undertaken under operator blinded  
4 conditions. Samples were identified with coded sequential numbers. In addition, at least two of  
5 both positive and negative controls were included to monitor the quality of staining.  
6  
7  
8

#### 9 10 11 *Assessment of vascular sclerotic index (SI)*

12  
13 H&E stained coronal frontal lobe (Brodmann area 9) sections were analysed to assess SI in a  
14 total of ~4,000 brain vessels, essentially as described previously (8, 56). Twenty images from  
15 each entire depth of the cerebral cortex and the deep WM were randomly captured using a bright  
16 field microscope (Leitz DIALUX 20, Leica) with a 10x objective lens coupled to a lumenera  
17 infinity digital camera (Lumenera Corporation, Canada). Using VasCalc software (56), internal  
18 diameter and external diameter of each vessel was measured in three opposing axis. Sclerotic  
19 index (SI) was calculated using the following formula:  $SI = (\text{External diameter} - \text{Internal}$   
20  $\text{diameter}) / (\text{External diameter})$ . Three SI values were calculated from three measured axis in  
21 each vessel and averaged. The inter-rater reliability between the assessors was 0.9 either DV  
22 and RK (SVD pathology scores) or YH and RK (SI scores).  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

#### 34 *Statistical analysis*

35  
36 Statistical analysis was carried out using SPSS (IBM, version 23.0, IBM Corporation, Armonk,  
37 NY, USA) with the level of significance set at  $P < 0.05$ . First, distribution of values was tested  
38 using the Shapiro-Wilk test followed by one-way analysis of variance (one-way ANOVA) with  
39 post-hoc Tukey's tests for normally distributed values or Kruskal-Wallis H tests for non-  
40 normally distributed values to compare data amongst autonomic dysfunction plus dementia and  
41 control group. Student's t-test or Mann-Whitney U test was used for normally or non-normally  
42 distributed data respectively to assess the differences between cortex and WM data. Spearman's  
43 rho correlation was used to assess the relationship between WM hyperintensity volumes and  
44 vascular pathology. Pearson's correlation was used to assess the correlation between vascular  
45 pathology and microinfarcts/microbleeds. Anonymized data will be shared by request from any  
46 qualified investigator. All the data we have generated in this study have been included in this  
47 paper.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

## Results

### *Clinical features in the autonomic dysfunction plus dementia subjects*

Tables 1 and 2 provide the mean age and gender distributions of all the dementia subjects with relevant clinical manifestations. We noted that the mean age of PDD subjects was marginally lower compared to controls and other autonomic dysfunction plus dementia subjects ( $\dagger P < 0.01$ ) (Table 1 and 2). Of the 77 dementia subjects tested in the FASS, 29 (38%) had evidence of both OH and CSH (Table 1). The total CAMCOG and MMSE scores indicated all subjects had evidence of dementia at least 6 months prior to death. There were no differences in CAMCOG memory and executive sub-scores amongst the autonomic dysfunction plus dementia groups (Table 1). We further noted that on average 55% of the autonomic dysfunction plus dementia subjects exhibited hypertension and often had more than one other vascular disease risk factor including diabetes mellitus (DM), ischaemic heart disease (IHD) and smoking (Table 1). The frequency of hypertension in DLB without falls subjects was 33%, which was not different compared with that in dementia subjects who had autonomic dysfunction (Chi-square test).

### *Cerebrovascular Pathological changes in Dementia with Autonomic Dysfunction*

Table 2 provides details of clinically tested autonomic dysfunction subjects and ageing controls assessed for SVD or microvascular pathology. None of the control subjects had reported unexplained falls in life or had been tested in the FASS. Subjects with dementia exhibited the expected pathology defining each type of disease. The Mixed and AD subjects had the highest amount of neurofibrillary tangle (Braak stage) and neuritic or amyloid  $\beta$  plaque burden (Table 2). The DLB, PDD and VaD groups had relatively lower burdens of AD-related hallmark pathologies and there was even less in ageing controls ( $*P < 0.01$ ;  $**P < 0.01$ ) (Table 2). In addition, nigral Lewy body pathology and neuronal loss were characteristically most severe in DLB, PDD and Mixed dementia subjects (Table 2). Of the 13 VaD subjects clinically diagnosed with autonomic dysfunction, 3 had some degree of Lewy body pathology (Table 2).

VaD subjects expectedly exhibited the highest vascular and WM pathology scores (10). However, remarkably all neurodegenerative dementia groups including DLB, PDD, Mixed and

AD had high vascular pathology scores compared to ageing controls ( $\ddagger P < 0.01$ ) (Table 2). AD subjects had the highest ABC scores (36) and the greatest frequency of moderate to severe CAA (Table 2). The types of vascular changes seen in subcortical regions in the autonomic dysfunction plus dementia subjects predominantly constituted those akin to SVD or microangiopathy, which included arteriolosclerosis, intimal thickening, fibroid necrosis, hyalinisation, perivascular spacing, microinfarcts, WM rarefaction and cerebral amyloid angiopathy (10) (Figure 1A, a-f). WM scores incorporating demyelination (17) were also greater in the dementia cases compared to controls. Variable calcification was present in numerous vessels within the extrapyramidal regions particularly in and vicinity of the globus pallidus (Figure 1B, a-d) that we also observed, albeit to a lesser degree in normal ageing controls. The diameters of the calcified vessels ranged 40 to 160  $\mu\text{m}$ . However, distribution of the vascular pathology scores in different brain regions indicated that higher scores were evident in the frontal lobe and the basal ganglia (Figure 1C). There was high predilection for microinfarcts in the frontal lobe WM as well as the basal ganglia across all the dementias.

Further verification of our observations on clinically tested subjects was evident from retrospective assessment of DLB subjects, who had had no apparent autonomic dysfunction (Table 2) and of demented subjects who had had repeated unexplained falls in life (Table 2). The vascular pathology features and scores in most of the dementia subjects were similar or higher compared to those who were clinically tested in the FASS (Table 2). We also could not find differences in SVD features in demented individuals whether they were clinically tested in the FASS or experienced unexplained falls as proxy for autonomic dysfunction. However, the key observation was that DLB subjects without reported unexplained falls or autonomic dysfunction, had lower vascular pathology scores compared to dementia subjects who had autonomic dysfunction. The vascular pathology scores in DLB subjects without autonomic dysfunction were also not different from ageing controls ( $P = 0.236$ ) (Figure 1C). We also found a positive correlation between severity of CAA and haemosiderin deposition (proxy for microbleeds) ( $r = 0.88$ ,  $P = 0.01$ ) but not between CAA and microinfarcts ( $P > 0.05$ ). We further noted that the degree of arteriolosclerosis was strongly correlated with numbers of microinfarcts ( $r = 0.87$ ,  $P = 0.012$ ) (Table 2).

### *Correlation between Vascular pathological changes and WMH volumes upon MRI*

To strengthen the evidence that WMHs associated with the various dementias in life (Figure 2A) reflected greater SVD or microvascular pathology, we tested the relationship between vWMHs determined by T2WI MRI and vascular pathology scores evaluated after post-mortem in those subjects who exhibited autonomic dysfunction. We found there was a positive correlation ( $\rho=0.40$ ,  $P=0.016$ ) between vWMH and vascular pathology scores across the dementias (Figure 2B). However, there was no clear relationship between vWMH and any of the other neurodegenerative disease markers such as tau or  $\alpha$ -synuclein pathology ( $P>0.05$ ).

### *Sclerotic Index and autonomic dysfunction plus dementia*

We then quantified the degree of arteriolosclerosis using the SI (Figure 3A) in the deep WM and entire depth of the cortex of the frontal lobe. The SI was consistently higher by 40-50% in all of the autonomic dysfunction plus dementia subjects compared to controls in both the frontal cortex and the WM (\*\* $P<0.01$  and †† $P<0.01$  respectively) (Figure 3B). Dementia subjects exhibited higher SI values in the cortex compared to the WM, particularly in AD subjects († $P<0.05$ ), whereas controls showed similar SI values in both the cortex and WM (Figure 3B). However, it was notable that DLB subjects without falls exhibited similar SI values as ageing controls and lower SI values, particularly in the WM compared to all dementia with evidence of autonomic dysfunction (‡ $P<0.035$ ) (Figure 3B).

## **Discussion**

We previously reported that deep WMHs correlated with autonomic dysfunction in DLB subjects (24). In this study, we provide robust evidence for the presence of significantly greater degree of SVD pathology including WM changes in dementia subjects with clinical evidence of concomitant autonomic dysfunction. The remarkable finding is that irrespective of common dementia type whether DLB, PDD, Mixed dementia, AD or VaD they all exhibited greater SVD type of pathological changes in the presence of autonomic dysfunction. This was corroborated by low vascular pathology scores and SI values in DLB subjects without evidence of autonomic

1  
2  
3 dysfunction. Whereas the highest burden of SVD pathology (51) consisting of microinfarcts,  
4 lacunar infarcts, severe arteriolosclerosis, WM rarefaction and perivascular spacing was present  
5 in VaD, it was rather surprising that subjects diagnosed with neurodegenerative dementias  
6 including PDD, DLB and AD exhibited these similar microvascular changes particularly severe  
7 arteriolosclerosis and perivascular spacing. SI values were higher by 45-50% signifying  
8 moderate-severe arteriolosclerosis in dementia cases with autonomic dysfunction.  
9  
10 Arteriolosclerosis and microinfarcts are strongly associated with cognitive impairment (5, 19,  
11 52), and were consistent features across all dementias. Vascular pathology scores, particularly in  
12 the WM were also increased and correlated with vWMHs in those demented subjects with  
13 autonomic dysfunction. These observations collectively suggest arterial wall changes and  
14 thickening may impact on blood rheology to impair cerebral blood flow and perfusion (20).  
15  
16  
17  
18  
19  
20  
21  
22

23 With reference to other features of SVD such as CAA, except in AD, dementia subjects  
24 with and without autonomic dysfunction exhibited similar CAA pathology to ageing controls.  
25 This further suggested that CAA was not necessarily a factor in the observed arteriosclerotic  
26 differences in demented subjects with and without autonomic dysfunction. While, we observed  
27 strong correlations between arteriolosclerosis and microinfarcts and between CAA and  
28 haemosiderin deposition or microbleeds, we did not observe correlations between CAA and  
29 microinfarcts (40, 47). The latter is probably because we had low numbers of cases in the  
30 analysis and had included VaD subjects, who showed relatively low CAA but high burden of  
31 microinfarction. The frequency of hypertension in 'DLB without falls' group was not different  
32 from other dementia groups. Moreover, DLB subjects without falls exhibited similar SI values  
33 as ageing controls but had lower SI values, particularly in the WM, compared to all dementias  
34 with concomitant of autonomic dysfunction. This suggests hypertension per se was not  
35 necessarily a factor in the observed differences in SVD pathology in demented subjects with and  
36 without autonomic dysfunction.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 **Large vessel disease may also contribute to SVD type of changes in the WM and to**  
49 **microinfarcts (20, 51, 57). In comparison to the neurodegenerative dementias, VaD subjects bear**  
50 **greater degrees of large vessel disease i.e. intracranial and extracranial atherosclerotic disease.**  
51 **In a recent study (14), we noted that regardless of severity of carotid artery stenosis, most of the**  
52 **infarcts were small (<5mm in size) and most were located in the cerebral cortex. In a previous it**  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 was further noted that cerebral atherosclerosis was not only expectedly associated with cystic  
4 infarcts but importantly microinfarcts (57). It would appear microinfarcts arise from a  
5 combination of mechanisms including artery-to-artery embolism, cardiogenic embolism and  
6 haemodynamic impairment that likely involve microemboli.  
7  
8  
9

10  
11 In comparison to ageing controls with no recorded autonomic dysfunction and DLB  
12 subjects without reported falls, overall our findings indicate SVD pathological changes are  
13 common in elderly subjects with autonomic dysfunction irrespective of the type of dementia they  
14 acquire. This may compound effects of co-existing pathologies including neurofibrillary or  $\alpha$ -  
15 synuclein pathology or amyloid deposits, which albeit may occur even in cases of pure  
16 autonomic failure such as idiopathic OH or Bradbury-Eggleston syndrome (pure autonomic  
17 failure causing idiopathic OH) (ROC and RNK, unpublished observations). It seems reasonable  
18 to conclude that cerebral SVD changes contribute to the clinical syndrome in elderly subjects  
19 who develop dementia and exhibit autonomic dysfunction. While these ageing-related SVD  
20 changes may not result in overt neurological signs or cerebrovascular events except in VaD, they  
21 likely collectively contribute to the declining brain health of the elderly with autonomic  
22 dysfunction.  
23  
24  
25  
26  
27  
28  
29  
30  
31

32 Our findings underscore that vascular protection is important in patients who develop any  
33 type of dementia and are symptomatic or even asymptomatic for autonomic dysfunction. We  
34 enrolled symptomatic autonomic dysfunction plus obviously demented subjects. However, some  
35 patients with neurodegenerative diseases and dementia develop autonomic dysfunction prior to  
36 motor and cognitive symptoms. A previous study reported that up to 40% of PD patients with  
37 autonomic dysfunction were 'asymptomatic', and even 'asymptomatic' autonomic dysfunction  
38 caused higher prevalence of falls, worsened activities of daily life (ADL) and affected their  
39 quality of life (QOL) (31). In a follow up study (32), the severity of autonomic symptoms  
40 progressed by 20% over the course of a year and was independently associated with impairment  
41 in ADLs and QOL. Symptomatic and asymptomatic OH were both associated with increased  
42 falls and health care utilization. Thus, 'asymptomatic' autonomic dysfunction has almost similar  
43 detrimental effects to being symptomatic. Importantly, if 'asymptomatic' autonomic dysfunction  
44 could be established clinically at early stages of cognitive impairment, relevant treatment or  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 intervention(s) has the potential to prevent further cognitive decline in patients with  
4 neurodegenerative dementias.  
5  
6

7 **While our study provides good evidence to support our hypothesis in that autonomic**  
8 **dysfunction is linked to alterations in the cerebral vasculature of subjects with neurodegenerative**  
9  
10 diseases, it is unclear whether autonomic dysregulation is a consequence or an instigator of  
11 dementia. A recent study (38), provided novel evidence for autonomic dysfunction in mild  
12 cognitive impairment. This was associated with orthostatic blood pressure dysregulation and  
13 suggests that dysautonomia is an early feature in the development of cognitive dysfunction.  
14 CSH is an important risk factor for hypertension, IHD and systemic atherosclerosis (48). In our  
15 cohort, autonomic dysfunction plus dementia subjects displayed frequent vascular risk factors.  
16 These may contribute to atherosclerosis in the carotid arteries, affecting the baroreceptors (27)  
17 and possibly intracranial cerebral arteries to cause deep WM changes (15, 24) and worsen  
18 cognitive function. Alternatively, CSH may independently cause chronic or episodic  
19 hypotension promoting abnormal perfusion with resultant changes in cerebral small vessels and  
20 the WM leading to cognitive decline. While previous studies have confirmed the association  
21 between features of autonomic dysfunction e.g. OH and selective cognitive deficits, albeit in PD,  
22 they have rejected the hypothesis that this is underlined by the development of cerebrovascular  
23 disease (43). The underlying mechanism could be mediated by development of cerebrovascular  
24 disease induced by chronic or episodic hypoperfusion but until now the extent of brain vascular  
25 load in PD or DLB patients with OH was not elucidated.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 Another hypothesis is that generalised brain  $\alpha$ -synuclein pathology or focal  $\alpha$ -  
40 synucleinopathy in the sympathetic ganglia relates to autonomic dysfunction and dementia.  
41 Clinical and post-mortem neuropathological findings in patients with PD and DLB, both  
42 clinically diagnosed as pure autonomic failure (PAF) revealed that neurodegeneration in PD and  
43 DLB may originate from outside the CNS in autonomic postganglionic neurons (22).  
44 Involvement of postganglionic sympathetic nerves including the cardiac sympathetic nerves were  
45 predominant in PD and PAF (41), suggesting that the cardiac but not the medullary sympathetic  
46 neurones are affected in people with PD and PAF. Cardioinhibitory CSH in patients with Lewy  
47 body pathology was also related to severity of the cardiac sympathetic neurons rather than in  
48 CNS (44). Furthermore, severity of autonomic dysfunction was associated with cognitive  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 dysfunction in patients with PD (50). These reports strongly support our findings in that PDD  
4 and DLB, which exhibit extensive brain Lewy body pathology, showed similar SVD pathologies  
5 and cognitive dysfunction as those cases with autonomic dysfunction plus dementia without  
6 significant Lewy body pathology. Thus, extensive Lewy body pathologies in the brain may not  
7 directly affect SVD pathological changes to lead to dementia but autonomic dysfunction related  
8 cerebral hypoperfusion likely contributes to microvascular pathology, WM changes and  
9 cognitive decline in patients with Lewy body pathology. It is plausible, however, that the  
10 observed SVD pathological changes result from neuronal damage in the brainstem nuclei or  
11 disruption of the monoaminergic neuronal system (27, 35). While further investigation is  
12 necessary, we observed alpha-synuclein pathology in sympathetic ganglia of some VaD subjects,  
13 suggesting autonomic dysfunction in VaD could partly result from alpha-synuclein pathology in  
14 the sympathetic nervous system.

15  
16  
17 We emphasise that one of the main limitations of our study is that we were not able to  
18 find and compare groups of dementia subjects who were tested in the FASS, did not have  
19 evidence of autonomic dysfunction and had come to post-mortem. As suggested by several  
20 clinical studies (cited above), high numbers of dementia patients may be 'asymptomatic' for  
21 autonomic dysfunction and it is likely that they bear considerable burden of SVD pathology. We  
22 previously demonstrated the spectrum of vascular pathology in different dementias including  
23 VaD, AD and DLB (10). In that previous post-mortem cohort, while we were not aware  
24 whether any of the patients had autonomic dysfunction or were asymptomatic for it, on the whole  
25 they tended to exhibit relatively low SVD pathology scores. Another limitation is that we did  
26 not quantify microvascular pathology in all the regions of the brain used to score vascular  
27 pathology. Such analysis is profoundly cumbersome but in accord with our previous hypothesis  
28 (13), we concentrated on the deep WM of the frontal lobe and incorporating the centrum  
29 semiovale. Finally, in terms of elucidating some mechanistic approaches we did not have  
30 enough frozen tissues from cases without clear evidence of autonomic dysfunction to  
31 substantiate study power and enable biochemical assays of myelin degrading proteins or  
32 gliovascular unit markers to demonstrate consequences of microvascular damage (13, 17).

33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Autonomic functions are regulated in the brainstem, e.g. medulla and its nuclei (34). In  
patients with CSH, tau pathology was evident in the medullary baroreflex nuclei (35) and



1  
2  
3 medullary vessels were degenerated in patients with multiple system atrophy (MSA) (33).  
4  
5 Microvascular changes in the brainstem also relate to or are caused by autonomic dysfunction. A  
6  
7 future expanded analysis of SVD pathological changes involving other brain regions, particularly  
8  
9 in the brainstem in this cohort may provide more evidence to elucidate the pathophysiological  
10  
11 mechanisms and impact of autonomic dysfunction on microvascular changes in dementia  
12  
13 patients.

### 14 **Conclusions**

15  
16 In summary, autonomic dysfunction irrespective of final diagnosis of dementia type is associated  
17  
18 with consistent higher burden of SVD pathological changes, particularly in the deep WM. These  
19  
20 SVD changes do not necessarily manifest into overt clinical signs but likely add to the dementia  
21  
22 syndrome. Our findings implicate that autonomic dysfunction is an important risk factor of  
23  
24 chronic cerebral hypoperfusion, which may differentially affect neurodegenerative processes to  
25  
26 produce the end-stage pathology. We propose the importance of assessing autonomic  
27  
28 dysfunction in patients with dementia, particularly at an early stage. Treatment for detrimental  
29  
30 autonomic dysfunction (even when it is asymptomatic) implementing stricter control of vascular  
31  
32 risk factors, pacemakers, life-style intervention or pharmacological approaches might be  
33  
34 effective therapeutic strategies to reduce SVD during ageing and prevent or slow down cognitive  
35  
36 decline in patients with autonomic dysfunction plus dementia, in addition to reducing their  
37  
38 immediate risk of falls.  
39  
40 .

### 41 **Acknowledgements**

42  
43 We are grateful to the patients, families, and clinical house staff for their cooperation in the  
44  
45 investigation of this study. We also appreciate the cooperation of the NBTR directors and staff  
46  
47 in assisting us with this study. We are thankful to Janet Slade and Arthur Oakley for the expert  
48  
49 technical assistance and for assisting us in managing and screening the cohort. Our work is  
50  
51 supported by grants from Alzheimer's Research UK (ARUK, PG2013-022) and the Medical  
52  
53 Research Council (MRC, G0500247), Newcastle Centre for Brain Ageing and Vitality (BBSRC,  
54  
55 EPSRC, ESRC and MRC, LLHW). Tissue for this study was collected by the Newcastle Brain  
56  
57 Tissue Resource, which is funded in part by a grant from the UK MRC (G0400074), by the  
58  
59  
60

1  
2  
3 Newcastle NIHR Biomedical Research Centre in Ageing and Age Related Diseases award to the  
4 Newcastle upon Tyne Hospitals NHS Foundation Trust, and by a grant from the Alzheimer's  
5 Society and ART as part of the Brains for Dementia Research Project. Dr. Hase was supported  
6 by SENSHIN Medical Research Foundation, Osaka Japan and the Great Britain Sasakawa  
7 Foundation, London, UK and reports no disclosures.  
8  
9  
10  
11  
12  
13  
14

### 15 **Data Availability**

16  
17 The data that support the findings of this study are available on request from the corresponding  
18 author. The data are not publicly available due to privacy or ethical restrictions.  
19  
20  
21  
22  
23

### 24 **Author Contributions**

25 YH and RNK conceived the study and wrote the first drafts of the manuscript. YH, EH, CP,  
26 LJLC, WS, MJF, TP and RNK performed or contributed to different aspects of the quantitative  
27 analysis. CB, RAK, LMA were responsible for original set up and clinical evaluation of the  
28 cohorts. TMP, VD, RHP, PI and RNK provided the pathological diagnosis. YH, TMP, MJF,  
29 ROC, KH, LMA and RNK contributed to critically revising the manuscript for important  
30 intellectual content, and all approved the final version of the manuscript for submission.  
31  
32  
33  
34  
35  
36  
37  
38

### 39 **Conflict of interests**

40  
41 The authors have no disclosures or conflicts of interest in relation to this manuscript.  
42  
43  
44  
45  
46  
47

### 48 **Ethical Approval**

49 Ethical approvals were granted by local research ethics committees of the Newcastle upon Tyne  
50 Foundation Hospitals Trust. Permission for use of brains for post-mortem research was also  
51 granted by consent from next-of-kin or family. All the brain tissues were retained in and  
52 obtained from the Newcastle Brain Tissue Resource.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review Only

## References

1. Allan L, McKeith I, Ballard C, Kenny RA (2006) The prevalence of autonomic symptoms in dementia and their association with physical activity, activities of daily living and quality of life. *Dement Geriatr Cogn Disord.*22(3):230-7.
2. Allan LM, Ballard CG, Rowan EN, Kenny RA (2009) Incidence and prediction of falls in dementia: a prospective study in older people. *PLoS One.*4(5):e5521.
3. Allan LM, Rowan EN, Firbank MJ, Thomas AJ, Parry SW, Polvikoski TM, O'Brien JT, Kalaria RN (2011) Long term incidence of dementia, predictors of mortality and pathological diagnosis in older stroke survivors. *Brain.*134(Pt 12):3716-27.
4. Arnold AC, Shibao C (2013) Current concepts in orthostatic hypotension management. *Curr Hypertens Rep.*15(4):304-12.
5. Arvanitakis Z, Leurgans SE, Barnes LL, Bennett DA, Schneider JA (2011) Microinfarct pathology, dementia, and cognitive systems. *Stroke.*42(3):722-7.
6. Ballard C, McKeith I, O'Brien J, Kalaria R, Jaros E, Ince P, Perry R (2000) Neuropathological substrates of dementia and depression in vascular dementia, with a particular focus on cases with small infarct volumes. *Dement Geriatr Cogn Disord.*11(2):59-65.
7. Ballard C, Shaw F, McKeith I, Kenny R (1998) High prevalence of neurovascular instability in neurodegenerative dementias. *Neurology.*51(6):1760-2.
8. Craggs LJ, Hagel C, Kuhlenbaeumer G, Borjesson-Hanson A, Andersen O, Viitanen M, Kalimo H, McLean CA, Slade JY, Hall RA, Oakley AE, Yamamoto Y, Deramecourt V, Kalaria RN (2013) Quantitative vascular pathology and phenotyping familial and sporadic cerebral small vessel diseases. *Brain Pathol.*23(5):547-57.
9. Cremer A, Soumare A, Berr C, Dartigues JF, Gabelle A, Gosse P, Tzourio C (2017) Orthostatic hypotension and risk of incident dementia: results from a 12-year follow-up of the three-city study cohort. *Hypertension.*70(1):44-9.
10. Deramecourt V, Slade JY, Oakley AE, Perry RH, Ince PG, Maurage CA, Kalaria RN (2012) Staging and natural history of cerebrovascular pathology in dementia. *Neurology.*78(14):1043-50.
11. Fedorowski A, Burri P, Melander O (2009) Orthostatic hypotension in genetically related hypertensive and normotensive individuals. *J Hypertens.*27(5):976-82.

12. Firbank MJ, Minett T, O'Brien JT (2003) Changes in DWI and MRS associated with white matter hyperintensities in elderly subjects. *Neurology*.61(7):950-4.
13. Hase Y, Horsburgh K, Ihara M, Kalaria RN (2017) White matter degeneration in vascular and other ageing-related dementias. *J Neurochem*.10.1111/jnc.14271.
14. Hase Y, Polvikoski TM, Ihara M, Hase M, Zafar R, Stevenson W, Allan LM, Ennaceur A, Horsburgh K, Gallart-Palau X, Sze SK, Kalaria RN (2019) Carotid artery disease in post-stroke survivors and effects of enriched environment on stroke pathology in a mouse model of carotid artery stenosis. *Neuropathol Appl Neurobiol*.10.1111/nan.12550.
15. Hollander M, Bots ML, Del Sol AI, Koudstaal PJ, Witteman JC, Grobbee DE, Hofman A, Breteler MM (2002) Carotid plaques increase the risk of stroke and subtypes of cerebral infarction in asymptomatic elderly: the Rotterdam study. *Circulation*.105(24):2872-7.
16. Hommel A, Faber MJ, Weerkamp NJ, van Dijk JG, Bloem BR, Koopmans RT (2016) Prevalence and prescribed treatments of orthostatic hypotension in institutionalized patients with Parkinson's disease. *J Parkinsons Dis*.6(4):805-10.
17. Ihara M, Polvikoski TM, Hall R, Slade JY, Perry RH, Oakley AE, Englund E, O'Brien JT, Ince PG, Kalaria RN (2010) Quantification of myelin loss in frontal lobe white matter in vascular dementia, Alzheimer's disease, and dementia with Lewy bodies. *Acta Neuropathol*.119(5):579-89.
18. Isaacson SH, Skettini J (2014) Neurogenic orthostatic hypotension in Parkinson's disease: evaluation, management, and emerging role of droxidopa. *Vasc Health Risk Manag*.10:169-76.
19. Kalaria RN (2012) Cerebrovascular disease and mechanisms of cognitive impairment: evidence from clinicopathological studies in humans. *Stroke*.43(9):2526-34.
20. Kalaria RN (2016) Neuropathological diagnosis of vascular cognitive impairment and vascular dementia with implications for Alzheimer's disease. *Acta Neuropathol*.131(5):659-85.
21. Kalaria RN, Kenny RA, Ballard CG, Perry R, Ince P, Polvikoski T (2004) Towards defining the neuropathological substrates of vascular dementia. *J Neurol Sci*.226(1-2):75-80.
22. Kaufmann H, Nahm K, Purohit D, Wolfe D (2004) Autonomic failure as the initial presentation of Parkinson disease and dementia with Lewy bodies. *Neurology*.63(6):1093-5.
23. Kenny RA, Richardson DA (2001) Carotid sinus syndrome and falls in older adults. *Am J Geriatr Cardiol*.10(2):97-9.

- 1  
2  
3 24. Kenny RA, Shaw FE, O'Brien JT, Scheltens PH, Kalaria R, Ballard C (2004) Carotid  
4 sinus syndrome is common in dementia with Lewy bodies and correlates with deep white matter  
5 lesions. *J Neurol Neurosurg Psychiatry*.75(7):966-71.  
6  
7  
8 25. Love S, Chalmers K, Ince P, Esiri M, Attems J, Kalaria R, Jellinger K, Yamada M,  
9 McCarron M, Minett T, Matthews F, Greenberg S, Mann D, Kehoe PG (2015) Erratum:  
10 Development, appraisal, validation and implementation of a consensus protocol for the  
11 assessment of cerebral amyloid angiopathy in post-mortem brain tissue. *Am J Neurodegener*  
12 *Dis*.4(2):49.  
13  
14  
15  
16 26. Lowe J, Kalaria RN (2015) *Dementia*. In: Greenfield's Neuropathology, Love S PA,  
17 Ironside J, Budka H, (ed.), *Chapter 50*, pp. 1001-55, CRC Press: London.  
18  
19 27. McDonald C, Newton JL, Burn DJ (2016) Orthostatic hypotension and cognitive  
20 impairment in Parkinson's disease: Causation or association? *Mov Disord*.31(7):937-46.  
21  
22 28. McIntosh SJ, Kenny RA (1994) Carotid sinus syndrome in the elderly. *J R Soc*  
23 *Med*.87(12):798-800.  
24  
25 29. McIntosh SJ, Lawson J, Kenny RA (1994) Heart rate and blood pressure responses to  
26 carotid sinus massage in healthy elderly subjects. *Age Ageing*.23(1):57-61.  
27  
28 30. McKeith IG, Dickson DW, Lowe J, Emre M, O'Brien JT, Feldman H, Cummings J, Duda  
29 JE, Lippa C, Perry EK, Aarsland D, Arai H, Ballard CG, Boeve B, Burn DJ, Costa D, Del Ser T,  
30 Dubois B, Galasko D, Gauthier S, Goetz CG, Gomez-Tortosa E, Halliday G, Hansen LA, Hardy  
31 J, Iwatsubo T, Kalaria RN, Kaufer D, Kenny RA, Korczyn A, Kosaka K, Lee VM, Lees A,  
32 Litvan I, Lodos E, Lopez OL, Minoshima S, Mizuno Y, Molina JA, Mukaetova-Ladinska EB,  
33 Pasquier F, Perry RH, Schulz JB, Trojanowski JQ, Yamada M (2005) Diagnosis and  
34 management of dementia with Lewy bodies: third report of the DLB Consortium.  
35 *Neurology*.65(12):1863-72.  
36  
37 31. Merola A, Romagnolo A, Rosso M, Lopez-Castellanos JR, Wissel BD, Larkin S,  
38 Bernardini A, Zibetti M, Maule S, Lopiano L, Espay AJ (2016) Orthostatic hypotension in  
39 Parkinson's disease: Does it matter if asymptomatic? *Parkinsonism Relat Disord*.33:65-71.  
40  
41 32. Merola A, Romagnolo A, Rosso M, Suri R, Berndt Z, Maule S, Lopiano L, Espay AJ  
42 (2017) Autonomic dysfunction in Parkinson's disease: A prospective cohort study. *Mov*  
43 *Disord*.10.1002/mds.27268.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
33. Miller VM, Kalaria RN, Hall R, Oakley AE, Kenny RA (2007) Medullary microvessel degeneration in multiple system atrophy. *Neurobiol Dis.*26(3):615-22.
34. Miller VM, Kenny RA, Oakley AE, Hall R, Kalaria RN, Allan LM (2009) Dorsal motor nucleus of vagus protein aggregates in Lewy body disease with autonomic dysfunction. *Brain Res.*1286:165-73.
35. Miller VM, Kenny RA, Slade JY, Oakley AE, Kalaria RN (2008) Medullary autonomic pathology in carotid sinus hypersensitivity. *Neuropathol Appl Neurobiol.*34(4):403-11.
36. Montine TJ, Phelps CH, Beach TG, Bigio EH, Cairns NJ, Dickson DW, Duyckaerts C, Frosch MP, Masliah E, Mirra SS, Nelson PT, Schneider JA, Thal DR, Trojanowski JQ, Vinters HV, Hyman BT, National Institute on A, Alzheimer's A (2012) National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease: a practical approach. *Acta Neuropathol.*123(1):1-11.
37. Munro NC, McIntosh S, Lawson J, Morley CA, Sutton R, Kenny RA (1994) Incidence of complications after carotid sinus massage in older patients with syncope. *J Am Geriatr Soc.*42(12):1248-51.
38. Nicolini P, Ciulla MM, Malfatto G, Abbate C, Mari D, Rossi PD, Pettenuzzo E, Magrini F, Consonni D, Lombardi F (2014) Autonomic dysfunction in mild cognitive impairment: evidence from power spectral analysis of heart rate variability in a cross-sectional case-control study. *PLoS One.*9(5):e96656.
39. Oh YS, Kim JS, Lee KS (2013) Orthostatic and supine blood pressures are associated with white matter hyperintensities in Parkinson disease. *J Mov Disord.*6(2):23-7.
40. Okamoto Y, Yamamoto T, Kalaria RN, Senzaki H, Maki T, Hase Y, Kitamura A, Washida K, Yamada M, Ito H, Tomimoto H, Takahashi R, Ihara M (2012) Cerebral hypoperfusion accelerates cerebral amyloid angiopathy and promotes cortical microinfarcts. *Acta Neuropathol.*123(3):381-94.
41. Orimo S, Oka T, Miura H, Tsuchiya K, Mori F, Wakabayashi K, Nagao T, Yokochi M (2002) Sympathetic cardiac denervation in Parkinson's disease and pure autonomic failure but not in multiple system atrophy. *J Neurol Neurosurg Psychiatry.*73(6):776-7.
42. Perry RH, Oakley AE (1993) *Newcastle Brain Map*. In: Neuropsychiatric Disorders, Roberts G, Leigh P, Weinberger D, (eds.), *Chapter 1*, pp. 1-10, Wolfe Publishing: London.

- 1  
2  
3 43. Pilleri M, Facchini S, Gasparoli E, Biundo R, Bernardi L, Marchetti M, Formento P,  
4 Antonini A (2013) Cognitive and MRI correlates of orthostatic hypotension in Parkinson's  
5 disease. *J Neurol*.260(1):253-9.  
6  
7  
8 44. Polvikoski T, Kalaria RN, Perry R, Miller V, Kenny RA (2006) Carotid sinus  
9 hypersensitivity associated with focal alpha-synucleinopathy of the autonomic nervous system. *J*  
10 *Neurol Neurosurg Psychiatry*.77(9):1064-6.  
11  
12 45. Sambati L, Calandra-Buonaura G, Poda R, Guaraldi P, Cortelli P (2014) Orthostatic  
13 hypotension and cognitive impairment: a dangerous association? *Neurol Sci*.35(6):951-7.  
14  
15 46. Skrobot OA, Attems J, Esiri M, Hortobagyi T, Ironside JW, Kalaria RN, King A,  
16 Lammie GA, Mann D, Neal J, Ben-Shlomo Y, Kehoe PG, Love S (2016) Vascular cognitive  
17 impairment neuropathology guidelines (VCING): the contribution of cerebrovascular pathology  
18 to cognitive impairment. *Brain*.139(11):2957-69.  
19  
20 47. Soontornniyomkij V, Lynch MD, Mermash S, Pomakian J, Badkoobehi H, Clare R,  
21 Vinters HV (2010) Cerebral microinfarcts associated with severe cerebral beta-amyloid  
22 angiopathy. *Brain Pathol*.20(2):459-67.  
23  
24 48. Tsioufis CP, Kallikazaros IE, Toutouzas KP, Stefanadis CI, Toutouzas PK (2002)  
25 Exaggerated carotid sinus massage responses are related to severe coronary artery disease in  
26 patients being evaluated for chest pain. *Clin Cardiol*.25(4):161-6.  
27  
28 49. Velseboer DC, de Haan RJ, Wieling W, Goldstein DS, de Bie RM (2011) Prevalence of  
29 orthostatic hypotension in Parkinson's disease: a systematic review and meta-analysis.  
30 *Parkinsonism Relat Disord*.17(10):724-9.  
31  
32 50. Verbaan D, Marinus J, Visser M, van Rooden SM, Stiggelbout AM, van Hilten JJ (2007)  
33 Patient-reported autonomic symptoms in Parkinson disease. *Neurology*.69(4):333-41.  
34  
35 51. Vinters HV, Zarow C, Borys E, Whitman JD, Tung S, Ellis WG, Zheng L, Chui HC  
36 (2018) Review: Vascular dementia: clinicopathologic and genetic considerations. *Neuropathol*  
37 *Appl Neurobiol*.44(3):247-66.  
38  
39 52. Westover MB, Bianchi MT, Yang C, Schneider JA, Greenberg SM (2013) Estimating  
40 cerebral microinfarct burden from autopsy samples. *Neurology*.80(15):1365-9.  
41  
42 53. Wieling W, Krediet CT, Solari D, de Lange FJ, van Dijk N, Thijs RD, van Dijk JG,  
43 Brignole M, Jardine DL (2013) At the heart of the arterial baroreflex: a physiological basis for a  
44 new classification of carotid sinus hypersensitivity. *J Intern Med*.273(4):345-58.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
54. Wieling W, Schatz IJ (2009) The consensus statement on the definition of orthostatic hypotension: a revisit after 13 years. *J Hypertens.*27(5):935-8.
55. Wolters FJ, Mattace-Raso FU, Koudstaal PJ, Hofman A, Ikram MA, Heart Brain Connection Collaborative Research G (2016) Orthostatic hypotension and the long-term risk of dementia: a population-based study. *PLoS Med.*13(10):e1002143.
56. Yamamoto Y, Ihara M, Tham C, Low RW, Slade JY, Moss T, Oakley AE, Polvikoski T, Kalaria RN (2009) Neuropathological correlates of temporal pole white matter hyperintensities in CADASIL. *Stroke.*40(6):2004-11.
57. Zheng L, Vinters HV, Mack WJ, Zarow C, Ellis WG, Chui HC (2013) Cerebral atherosclerosis is associated with cystic infarcts and microinfarcts but not Alzheimer pathologic changes. *Stroke.*44(10):2835-41.
58. Zuccala G, Onder G, Pedone C, Carosella L, Pahor M, Bernabei R, Cocchi A, Group G-OS (2001) Hypotension and cognitive impairment: selective association in patients with heart failure. *Neurology.*57(11):1986-92.

Table 1: Clinical features in the autonomic dysfunction plus dementia subjects assessed in the FASS

Variable	All dementias	DLB	PDD	Mixed	AD	VaD
Number of subjects	77	16	13	17	18	13
Age, years, mean (range)	82.3 (64-98)	79.6 (69-96) <sup>‡</sup>	72.8 (64-81) <sup>†</sup>	83.0 (72-93)	87.8 (76-96)	88.1 (75-98)
OH / CSH / OH and CSH, number	58, 42, 29	12, 10, 7	11, 5, 3	12, 11, 8	14, 11, 9	9, 5, 2
Total CAMCOG score (/100), mean (range)	54.2 (20-80)	53.8 (24-80)	61.1 (39-74)	51.0 (23-68)*	44.0 (20-73)	61.2 (25-80)
Memory sub-score (/27), mean (range)	13.7 (4-23)	14.9 (8-21)	16.2 (9-23)	13.5 (4-23)	10.0 (5-14)	14.1 (4-22)
Executive sub-score (/28), mean (range)	11.2 (2-19)	12.1 (4-23)	9.6 (5-18)	11.2 (7-15)	12.3 (10-17)	10.6 (2-19)
MMSE score (/30), mean (range)	13.1 (0-24)	13.8 (6-20)	14.2 (3-20)	11.9 (2-19)	8.5 (0-16)	17.2 (8-24) <sup>†</sup>
Hypertension, number (%)	42 (55%)	11 (69%)	7 (54%)	7 (41%)	9 (50%)	9 (69%)
Systolic BP, mean (range)	146 (101-200)	149 (122-187)	152 (101-200)	142 (109-186)	141 (110-165)	146 (131-172)
Diastolic BP, mean (range)	76 (51-120)	74 (60-86)	82 (57-120)	74 (53-98)	72 (51-95)	76 (52-92)
Other VRF, number (smoking / DM / IHD)	32/5/15	7/0/4	6/1/3	6/1/3	5/3/2	8/0/3

All dementia subjects had clinical evidence of autonomic dysfunction. All dementias group represent all causes of dementia including DLB, PDD, Mixed, AD and VaD; Mixed dementia: 7 =AD+DLB, 6 =AD+VaD, 3=AD+DLB+VaD and 1 = PD+DLB+VaD.

Age, <sup>†</sup> $P<0.01$  vs Control, DLB, Mixed, AD and VaD; <sup>‡</sup> $P<0.01$  vs AD and VaD;  $P=0.057$ , Mixed vs AD;  $P=0.053$ , Mixed vs VaD; OH, orthostatic hypotension; CSH, carotid sinus hypersensitivity; 'OH / CSH / OH and CSH, number' indicates number of cases who had OH / CSH / both OH and CSH in each group. For example, in DLB group, OH / CSH / OH and CSH are described as 12, 10, 7.

1  
2  
3 This means 12 cases in DLB had OH, 10 subjects confirmed to have CSH and 7 had evidence to have both OH and CSH (OH and  
4 CSH overlapped in some cases); Total CAMCOG score, \* $P=0.034$  vs PDD, \* $P=0.028$  vs VaD; CAMCOG executive and memory  
5 subscore, n.s. between groups; MMSE score, † $P=0.041$  vs DLB, † $P=0.021$  vs Mixed, † $P=0.003$  vs AD; Hypertension, number (%),  
6 n.s. between groups (Chi-square test); Systolic and Diastolic BP, n.s. between groups. Clinical features other than age and gender in  
7 Controls and DLB without falls cases were not completely available as they were not assessed at the falls and syncope clinic (FASS).

8  
9  
10 Abbreviations: AD, Alzheimer's disease; BP, blood pressure; CAMCOG, revised Cambridge Cognition Examination; DLB,  
11 Dementia with Lewy Bodies; DM, diabetes mellitus; F, female; IHD, ischaemic heart disease; M, male; Mixed, Mixed dementia;  
12 MMSE, Mini-Mental State Examination; PDD, Parkinson's disease with dementia; VaD, Vascular dementia; VRF, vascular risk  
13 factor; WML, white matter lesion.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Table 2

## Neuropathological features of all dementia cases and similar age controls in the study

Variable	Controls	DLB without Falls	All dementias	DLB	PDD	Mixed†	AD	VaD
Number of subjects	17	12	100	24	18	22	18	18
Age, years, mean (range)	86.9 (72-99)	80.0 (74-90)*	82.0 (64-98)	79.6 (69-96)*	74.6 (64-89)†	83.4 (72-93)	87.8 (76-96)	85.8 (71-98)
Gender, number (F/M)	13 / 4	1 / 11	52 / 48	11 / 13	9 / 9	16 / 6	9 / 9	7 / 11
Autonomic Dysfunction (n=)§	0	0	100	24	18	22	18	18
Brak Stage, mean (range)	1.9 (0-4)	3.0 (0-6)	3.3 (0-6)	2.2 (0-4)	2.0 (0-4)	4.8 (2-6)*	5.6 (5-6)**	1.6 (0-4)
CERAD, mean (range)	0.5 (0-2)	1.2 (0-2)	1.7 (0-3)	1.5 (0-3)¶	0.5 (0-2)	2.6 (0-3)*	2.9 (2-3)**	0.8 (0-3)
Alzheimer's Disease Neuropathologic Changes; A, B, C (mean)#	A0.5, B1.2, C0.5	A1.2, B1.6, C1.2	A1.5, B1.8, C1.7	A0.9, B1.3, C1.5	A0.4, B1.1, C0.5	A2.5, B2.6, C2.6	A3, B3, C3	A0.6, B1.2, C0.8
Vascular pathology score, mean (range)#	6.7 (0-10)‡	8.1 (5-10)§	11.4 (5-18)	9.9 (5-13)	10.8 (7-17)	11.9 (7-15)	11.2 (6-16)	13.6 (10-18)¶
White matter lesion score, mean (range)	0.5 (0-2)§	1.6 (1-3)	2.2 (0-3)	2.0 (1-3)	2.1 (1-3)	2.1 (0-3)	1.8 (0-3)	2.9 (2-3)¶¶
White matter/Vascular lesions, moderate - severe (%)	17.6**	41.7*	94	83	94	91	72	100
Lewy body pathology, number	0 / 0	2 / 7	16 / 36	3 / 19	5 / 10	5 / 7	0 / 1	3 / 0

1														
2														
3		(limbic/neocortical)												
4														
5														
6		SN neuronal loss (n=),												
7		(none/mild/moderate/ severe)	N/A	3/1/3/5	19/25/28/24	0/4/1/8	0/1/4/12	3/5/9/4	7/8/2/0	9/7/2/0				
8														
9														
10		CAA, total (%)	17.6	8.3	18.6	8.3	5.6	18.2	66.7**	27.8				
11														
12		CAA, moderate-severe	5.8	8.3	10.1	0.0	5.6	9.1	38.9**	16.7				
13		(%)												
14														
15		Arteriolo sclerosis,												
16		moderate-severe (%)	5.9**	25.0	60.6	45.9	61.1	63.4	61.1	66.7				
17														
18		Microinfarct, total (%)	35.3**	25.0**	81.1	79.1	88.9	77.3	66.7	100*				
19														
20														
21		Microinfarct, moderate-	0.0*	0.0*	30.5	25.0	33.3	27.2	15.6	76.9**				
22		severe (%)												
23														
24		Haemosiderin deposition	0.0	0.0	15.0	8.3	11.1	9.1	33.3**	16.7				
25		(microbleeds), total (%)												
26														
27		Brain only necropsy (%)	100	25	72.0	70.8	88.9	95.4	66.7	33.3				
28														
29		Intracranial MCA												
30		stenosis, moderate-severe	N/A	16.7	14.0	4.2	5.6	0.0	11.1	90.9				
31		(%)												
32		Extracranial ICA												
33		stenosis, moderate-severe	N/A	0.0	16.8	N/A	0.0	N/A	11.1	100				
34		(%)												
35		$\alpha$ -Synuclein pathology in												
36		the sympathetic ganglia,												
37		moderate-severe (n=)	N/A	N/A	5	N/A	1	N/A	N/A	4				
38														
39														
40														
41														
42														
43														
44														
45														
46														
47														

§All dementia subjects had clinical symptoms or evidence of autonomic dysfunction. #Mixed dementia group represents: 7=AD+DLB, 7=AD+VaD, 4=AD+DLB+VaD, 3=DLB+VaD and 1=PD+DLB+VaD.

Age, † $P<0.01$  vs Control, DLB, Mixed, AD and VaD; ‡ $P<0.01$  vs AD and VaD; & $P<0.01$  vs AD and VaD; Braak Stage, \*\* $P<0.01$  vs Control, DLB without Falls, DLB, PDD and VaD; \*\* $P<0.05$  vs Mixed; \* $P<0.01$  vs Control, DLB and PDD; \* $P<0.01$  vs DLB without Falls and VaD; CERAD, \*\* $P<0.01$  vs Control, DLB, PDD and VaD; \* $P<0.01$  vs Control, DLB, PDD and VaD;  $\phi P<0.05$  vs Control;  $\psi P<0.01$  vs PDD; #Alzheimer's Disease Neuropathologic Changes (36); #Vascular Pathology Score (10), † $P<0.01$  vs DLB, PDD, Mixed, AD and VaD; & $P<0.01$  vs PDD, Mixed, AD and VaD; † $P<0.05$  vs DLB without Falls, DLB and PDD; White matter lesion score; § $P<0.01$  vs DLB without Falls, DLB, PDD, Mixed, AD and VaD; † $P<0.01$  vs DLB without Falls, DLB, PDD, Mixed and AD; White matter/Vascular lesions, \*\* $P<0.01$  vs Control, DLB, PDD, Mixed, AD and VaD; \* $P<0.05$  vs Control, DLB, PDD, Mixed, AD and VaD (Chi-square test); Lewy body pathology, only the number of limbic/neocortical cases are shown. Fifteen Controls, 7 Mixed, 15 AD and 14 VaD cases had no Lewy body pathology. Two controls, 1 DLB, 2 Mixed and 1 AD showed Lewy body pathology in brain stem. Data were not available for 1 case in DLB and PDD groups due to limited autopsy; Neuronal loss in the substantia nigra, data was not available for 1 case in each group other than VaD due to limited autopsy. CAA total (%) and CAA moderate-severe (%), \*\* $P<0.01$  (Chi-square test); Arteriolosclerosis moderate-severe (%), \*\* $P<0.01$  (Chi-square test); Microinfarct total (%) and CAA moderate-severe (%), \*\* $P<0.01$ , \* $P<0.05$  (Chi-square test); Microbleeds total (%), \*\* $P<0.01$  (Chi-square test).

Abbreviations: AD, Alzheimer's disease; CAA, cerebral amyloid angiopathy; CERAD, Consortium to Establish a Registry for Alzheimer's Disease; DLB, Dementia with Lewy Bodies; F, female; ICA, internal carotid artery; M, male; Mixed, MCA, middle cerebral artery; Mixed dementia; PDD, Parkinson's disease with dementia; SN, substantia nigra; VaD, Vascular dementia; WML, white matter lesion.

## Figure Legends

**Figure 1: Spectrum of SVD pathology (subcortical cerebrovascular lesions) in autonomic dysfunction plus dementia subjects.** **A (a-f)**, Representative images (H&E staining) of different vascular pathologies in frontal WM (**a-d**) and basal ganglia (**e and f**) observed in autonomic dysfunction plus DLB and PDD subjects. **A (a)**, vascular wall thickening with perivascular spacing; **A (b)**, perivascular spacing; **A (c)**, small infarct (arrows); **A (d)**, perivascular microinfarct; **A (e)**, microvascular hyalinosis with perivascular spacing; **A (f)**, rarefaction (arrows). **B (a-d)**, Representative images (H&E stain) of microvascular changes and variable calcification observed in DLB and PDD subjects. **B (a-d)**, microvascular hyalinosis; **B (b, c and d)**, calcification in the vessel wall. **C**, Bar chart showing the total vascular pathology scores and distribution of individual vascular pathology scores in different regions of the brain. All dementias with autonomic dysfunction groups showed consistently higher total ( $\dagger P < 0.008$ ) and individual vascular pathology scores compared to controls, particularly in both the frontal lobe and the basal ganglia ( $**P < 0.016$  and  $\dagger\dagger P < 0.000$  respectively). In the temporal lobe, DLB, Mixed, AD and VaD subjects showed higher vascular pathology scores compared to controls ( $\ddagger P < 0.019$ ). Vascular pathology scores in the hippocampus remained same across all dementias and control subjects. High scores were evident in VaD subjects who also exhibited lacunar infarcts. Subjects with dementia also exhibited microinfarcts particularly in the frontal WM and basal ganglia with scores of 5/6 and 3/4 respectively (10). Total and individual vascular pathology scores were not different between controls and DLB without Falls. Scale bar represents 100  $\mu\text{m}$  (A, a-f) and 20  $\mu\text{m}$  (B, a-d).

**Figure 2: Relationship between WM hyperintensity volumes and vascular pathology in autonomic dysfunction.** **A (a-f)**, Axial views of T2-weighted brain MRI images at the level of basal ganglia and corona radiata to illustrate degrees of vWMHs in different dementias (**b-f**) in comparison to a control subject without autonomic dysfunction (**a**). The mean percentage of vWMH per total brain volume and vascular pathology scores in the controls (n=5) was 10.4% and 8.8 respectively (**a**). **B**, Correlational analysis between percentage of vWMH per total brain volume and vascular pathology scores of demented subjects with autonomic dysfunction.

1  
2  
3 Spearman's rho correlation analysis revealed that vascular pathology score exhibited a positive  
4 correlation with % vWMH per total brain volume ( $\rho=0.40$ ,  $P=0.016$ ).  
5  
6  
7  
8

9  
10 **Figure 3: Assessment of vascular sclerotic index in brain vessels.** **A**, Representative profile  
11 of a mildly hyalinised microvessel indicating the dimensions used to determine the sclerotic  
12 index (SI). The dotted lines mark inner diameters of the vessel. The normal lines indicate  
13 external diameters. Scale bar represents 50  $\mu\text{m}$ . **B**, Histogram showing SI values in the WM and  
14 the cortex in controls and autonomic dysfunction plus dementia groups. All dementias with  
15 autonomic dysfunction groups showed consistently higher mean SI values compared to controls  
16 in both the WM and the cortex (\*\* $P<0.005$  and †† $P<0.005$  respectively). SI values were not  
17 different between controls and DLB without Falls in both the WM and the cortex. In the WM,  
18 DLB without Falls showed lower SI values compared with all dementia with autonomic  
19 dysfunction groups (‡ $P<0.035$ ). In autonomic dysfunction plus dementia subjects, higher SI was  
20 evident in the cortex compared with WM, particularly in AD subjects (¶ $P=0.019$ ).  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60






Figure 1\_revised\_Hase et al\_2019




Figure 2

194x283mm (300 x 300 DPI)




Figure 3

169x237mm (300 x 300 DPI)