

# BOLIB 2019: Bilevel Optimization LIBrary of test problems version 2

Shenglong Zhou, Alain B. Zemkoho, and Andrey Tin

**Abstract** This paper presents the Bilevel Optimization LIBrary of the test problems (BOLIB—for short), which contains a collection of test problems, with continuous variables, to help support the development of numerical solvers for bilevel optimization. The first version of the library, available at [arxiv.org/abs/1812.00230](https://arxiv.org/abs/1812.00230), is made of 124 examples of nonlinear bilevel optimization problems. This version contains 173 examples with 138 nonlinear (including the 124 from the first version), 24 linear, and 11 simple bilevel optimization problems. This expansion of BOLIB further establishes it as the largest bilevel optimization library of test problems. Moreover, as the library is computation-enabled with the MATLAB m-files of all the examples, it provides a uniform basis for testing and comparing algorithms. The library, together with all the related codes, is freely available at [biopt.github.io/bolib](https://biopt.github.io/bolib).

## 1 Introduction

The bilevel optimization problem can take the form

$$\begin{aligned} \min_{x,y} F(x, y) \\ \text{s.t. } G(x, y) \leq 0, \\ y \in S(x) := \arg \min_y \{f(x, y) \mid g(x, y) \leq 0\}, \end{aligned} \tag{1}$$

---

Shenglong Zhou  
School of Mathematics, University of Southampton, University Road, SO17 1BJ Southampton,  
United Kingdom, e-mail: [shenglong.zhou@soton.ac.uk](mailto:shenglong.zhou@soton.ac.uk)

Alain B. Zemkoho  
School of Mathematics, University of Southampton, University Road, SO17 1BJ Southampton,  
United Kingdom, e-mail: [a.b.zemkoho@soton.ac.uk](mailto:a.b.zemkoho@soton.ac.uk)

Andrey Tin  
School of Mathematics, University of Southampton, University Road, SO17 1BJ Southampton,  
United Kingdom, e-mail: [a.tin@soton.ac.uk](mailto:a.tin@soton.ac.uk)

where the functions  $G : \mathbb{R}^{n_x} \times \mathbb{R}^{n_y} \rightarrow \mathbb{R}^{n_G}$  and  $g : \mathbb{R}^{n_x} \times \mathbb{R}^{n_y} \rightarrow \mathbb{R}^{n_g}$  define the upper-level and lower-level constraints, respectively. As for  $F : \mathbb{R}^{n_x} \times \mathbb{R}^{n_y} \rightarrow \mathbb{R}$  and  $f : \mathbb{R}^{n_x} \times \mathbb{R}^{n_y} \rightarrow \mathbb{R}$ , they denote the upper-level and lower-level objective functions, respectively. The set-valued map  $S : \mathbb{R}^{n_x} \rightrightarrows \mathbb{R}^{n_y}$  represents the optimal solution/argminimum mapping of the lower-level problem. Further recall that problem (1) as a whole is often called upper-level problem.

Our aim is to propose a computation-enabled library of test problems to help accelerate the development of numerical methods for bilevel programs in the form (1). For the sake of clarity, note that the bilevel optimization problems in this library, that we classify into the following three categories, only involve continuous variables:

- *Nonlinear bilevel programs*, which are problems in the form (1) with at least one of the functions involved being nonlinear.
- *Linear bilevel programs* are problems in the form (1) with functions  $F$ ,  $f$ , and all the components of  $G$  and  $g$  being linear.
- *Simple bilevel programs* (term coined in [13]) are optimization problems where the feasible set is partly defined by the optimal solution set of a second optimization problem. But unlike in (1), the lower-level problem is not a parametric optimization problem. More precisely, a simple bilevel optimization has the form:

$$\begin{aligned} \min_y F(y) \\ \text{s.t. } G(y) \leq 0, \\ y \in S := \arg \min_y \{f(y) \mid g(y) \leq 0\}, \end{aligned} \quad (2)$$

where, similarly to (1),  $G : \mathbb{R}^{n_y} \rightarrow \mathbb{R}^{n_G}$  and  $g : \mathbb{R}^{n_y} \rightarrow \mathbb{R}^{n_g}$  describe the upper-level and lower-level constraints, respectively, while the real-valued function  $F$  (resp.  $f$ ), defined  $\mathbb{R}^{n_y}$ , represents the upper-level (resp. lower-level) objective function. The expression ‘‘simple bilevel program’’ is used in [32] to refer to bilevel optimization problems of the form (1), where  $y$  (resp.  $x$ ) is not involved in the upper-level (resp. lower-level) constraints.

The main contributions of the library are three-fold. First, BOLIB provides MATLAB codes for 173 examples, including 138 nonlinear, 24 linear, and 11 simple bilevel programs, ready to be used to test numerical algorithms. Secondly, it puts together the true or best known solutions and the corresponding references for all the examples included. Hence, can serve as a benchmark platform for numerical accuracy evaluation for methods designed to solve problem (1). Thirdly, all examples as well as their gradients and Hessians are programmed and stored in the MATLAB m-files. Thus, facilitating the use of the examples and corresponding derivatives in the implementation of numerical methods, where such information is necessary.

To the best of our knowledge, this is the largest library of test examples for bilevel optimization, especially for the nonlinear class of the problem. It includes bilevel optimization problems from Colson’s BIPA [11], Leyffer’s MacMPEC [35], as well as from Mitsos and Barton’s technical report [39]. We would like to emphasize that the fundamental objective that we hope to achieve with BOLIB is the acceleration of numerical software development for bilevel optimization, as it is our opinion that

the level of expansion of applications of the problem has outpaced the development rate for numerical solvers, especially for the nonlinear class of the problem.

In the next section, we describe the library with details on the inputs and outputs of the codes, as well as some useful insights on the examples. In the subsequent section, a guideline is given on how to access the library.

## 2 Description of the library

This section describes the structure of the library, while focusing on the inputs and outputs of each example, as well as the list of all examples together with their true or best known solutions and corresponding references. Before we proceed, note that each `m-file` contains information about the corresponding example, which include the first and second order derivatives of the input functions. For the upper-level objective function  $F : \mathbb{R}^{n_x} \times \mathbb{R}^{n_y} \rightarrow \mathbb{R}$ , these derivatives are defined as follows

$$\begin{aligned} \nabla_x F(x, y) &= \begin{bmatrix} \nabla_{x_1} F \\ \vdots \\ \nabla_{x_{n_x}} F \end{bmatrix} \in \mathbb{R}^{n_x}, \\ \nabla_{xx}^2 F(x, y) &= \begin{bmatrix} \nabla_{x_1 x_1}^2 F & \cdots & \nabla_{x_{n_x} x_1}^2 F \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 x_{n_x}}^2 F & \cdots & \nabla_{x_{n_x} x_{n_x}}^2 F \end{bmatrix} \in \mathbb{R}^{n_x \times n_x}, \\ \nabla_{xy}^2 F(x, y) &= \begin{bmatrix} \nabla_{x_1 y_1}^2 F & \cdots & \nabla_{x_{n_x} y_1}^2 F \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 y_{n_x}}^2 F & \cdots & \nabla_{x_{n_x} y_{n_x}}^2 F \end{bmatrix} \in \mathbb{R}^{n_y \times n_x}. \end{aligned} \quad (1)$$


Similar expressions are valid for  $\nabla_y F(x, y) \in \mathbb{R}^{n_y}$ ,  $\nabla_{yy}^2 F(x, y) \in \mathbb{R}^{n_y \times n_y}$ , and the lower-level objective function  $f$ . As the constraint functions are vector-valued, we use the following notations to refer to derivative information in the context of the upper-level constraint function  $G : \mathbb{R}^{n_x} \times \mathbb{R}^{n_y} \rightarrow \mathbb{R}^{n_G}$ , for instance:

$$\begin{aligned}
\nabla_x G(x, y) &= \begin{bmatrix} \nabla_x G_1 \\ \vdots \\ \nabla_x G_{n_G} \end{bmatrix} = \begin{bmatrix} \nabla_{x_1} G_1 & \cdots & \nabla_{x_{n_x}} G_1 \\ \vdots & \ddots & \vdots \\ \nabla_{x_1} G_{n_G} & \cdots & \nabla_{x_{n_x}} G_{n_G} \end{bmatrix} \in \mathbb{R}^{n_G \times n_x}, \\
\nabla_{xx}^2 G(x, y) &= \begin{bmatrix} \nabla_{xx}^2 G_1 \\ \vdots \\ \nabla_{xx}^2 G_{n_G} \end{bmatrix} = \begin{bmatrix} \nabla_{x_1 x_1}^2 G_1 & \cdots & \nabla_{x_{n_x} x_1}^2 G_1 \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 x_{n_x}}^2 G_1 & \cdots & \nabla_{x_{n_x} x_{n_x}}^2 G_1 \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 x_1}^2 G_{n_G} & \cdots & \nabla_{x_{n_x} x_1}^2 G_{n_G} \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 x_{n_x}}^2 G_{n_G} & \cdots & \nabla_{x_{n_x} x_{n_x}}^2 G_{n_G} \\ \nabla_{x_1 y_1}^2 G_1 & \cdots & \nabla_{x_{n_x} y_1}^2 G_1 \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 y_{n_y}}^2 G_1 & \cdots & \nabla_{x_{n_x} y_{n_y}}^2 G_1 \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 y_1}^2 G_{n_G} & \cdots & \nabla_{x_{n_x} y_1}^2 G_{n_G} \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 y_{n_y}}^2 G_{n_G} & \cdots & \nabla_{x_{n_x} y_{n_y}}^2 G_{n_G} \end{bmatrix} \in \mathbb{R}^{(n_G n_x) \times n_x}, \\
\nabla_{xy}^2 G(x, y) &= \begin{bmatrix} \nabla_{xy}^2 G_1 \\ \vdots \\ \nabla_{xy}^2 G_{n_G} \end{bmatrix} = \begin{bmatrix} \nabla_{x_1 y_1}^2 G_1 & \cdots & \nabla_{x_{n_x} y_1}^2 G_1 \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 y_{n_y}}^2 G_1 & \cdots & \nabla_{x_{n_x} y_{n_y}}^2 G_1 \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 y_1}^2 G_{n_G} & \cdots & \nabla_{x_{n_x} y_1}^2 G_{n_G} \\ \vdots & \ddots & \vdots \\ \nabla_{x_1 y_{n_y}}^2 G_{n_G} & \cdots & \nabla_{x_{n_x} y_{n_y}}^2 G_{n_G} \end{bmatrix} \in \mathbb{R}^{(n_G n_y) \times n_x}.
\end{aligned} \tag{2}$$

Similar formulas are also valid for  $\nabla_y G(x, y) \in \mathbb{R}^{n_G \times n_y}$ ,  $\nabla_{yy}^2 G(x, y) \in \mathbb{R}^{n_G n_y \times n_y}$ , and the lower-level constraint  $g$ . It is important to emphasize that in the context of the constraints,  $\nabla_x G(x, y) \in \mathbb{R}^{1 \times n_x}$ , for example, is a row vector when  $n_G = 1$ . However,  $\nabla_x F(x, y) \in \mathbb{R}^{n_x}$  and  $\nabla_x f(x, y) \in \mathbb{R}^{n_x}$  are column vectors.

## 2.1 Inputs and outputs

The BOLIBver2 folder (see Section 3 on how to access the library), which contains all the library material, includes 3 sub-folders named `Nonlinear`, `Linear`, and `Simple`. In the `Nonlinear` subfolder, there are 138 MATLAB m-files. Each one specifies a nonlinear bilevel optimization test example, named by a combination of authors' surnames, year of publication, and when necessary, the order of the example in the corresponding reference. For example, as in following figure (showing a partial list of the examples), `AiyoshiShimizu1984Ex2.m` stands for Example 2 in the paper by Aiyoshi and Shimizu published in 1984 [1]. However, for a few examples (`DesignCentringP1`, `NetworkDesignP1`, etc.), the problem naming is based on previous use in the literature and therefore could help to easily recognize them.



In folder `Linear`, there are 24 MATLAB m-files defining 24 liner bilevel optimization test examples. The rule of naming each example is same as in the nonlinear case. Similarly, folder `Simple` contains 11 simple bilevel optimization test examples.

Now we describe the inputs and outputs of the m-file of a given example. Each file has the function handle named in the following way:

$$w = \text{example\_name}(x, y, \text{keyf}, \text{keyxy}). \quad (3)$$

For the inputs, we have

$$\begin{aligned} x &\in \mathbb{R}^{n_x}, \quad y \in \mathbb{R}^{n_y}, \\ \text{keyf} &\in \{ 'F', 'G', 'f', 'g' \}, \\ \text{keyxy} &\in \{ [], 'x', 'y', 'xx', 'xy', 'yy' \}, \end{aligned}$$

where `'F'`, `'G'`, `'f'`, and `'g'`, respectively stand for the four functions involved in (1). `'x'` and `'y'` represent the first order derivative with respect to  $x$  and  $y$ , respectively. Finally, `'xx'`, `'xy'`, and `'yy'` correspond to the second order derivative of the function  $F, G, f$ , and  $g$ , with respect to  $xx, xy$ , and  $yy$ , respectively.

For the outputs,  $w = \text{example\_name}(x, y, \text{keyf})$  or  $w = \text{example\_name}(x, y, \text{keyf}, [])$  returns the function value of `keyf` while  $w = \text{example\_name}(x, y, \text{keyf}, \text{keyxy})$  can additionally return the first or second order derivative of `keyf` w.r.t. the choice of `keyxy` as described above. We summarize all the scenarios in Table 1:

**Table 1** Input–output scenarios from the m-files containing the examples

keyf/keyxy	[]	'x'	'y'	'xx'	'xy'	'yy'
'F'	$F(x, y)$	$\nabla_x F(x, y)$	$\nabla_y F(x, y)$	$\nabla_{xx}^2 F(x, y)$	$\nabla_{xy}^2 F(x, y)$	$\nabla_{yy}^2 F(x, y)$
'G'	$G(x, y)$	$\nabla_x G(x, y)$	$\nabla_y G(x, y)$	$\nabla_{xx}^2 G(x, y)$	$\nabla_{xy}^2 G(x, y)$	$\nabla_{yy}^2 G(x, y)$
'f'	$f(x, y)$	$\nabla_x f(x, y)$	$\nabla_y f(x, y)$	$\nabla_{xx}^2 f(x, y)$	$\nabla_{xy}^2 f(x, y)$	$\nabla_{yy}^2 f(x, y)$
'g'	$g(x, y)$	$\nabla_x g(x, y)$	$\nabla_y g(x, y)$	$\nabla_{xx}^2 g(x, y)$	$\nabla_{xy}^2 g(x, y)$	$\nabla_{yy}^2 g(x, y)$

For the dimension of  $w$  in each scenario, see (1)–(2). If  $n_G = 0$  (or  $n_g = 0$ ), all outputs related to  $G$  (or  $g$ ) should be empty, namely,  $w = []$ . To further clarify the outputs, let us look at some specific usage:

- $w = \text{example\_name}(x, y, 'F')$  or  $w = \text{example\_name}(x, y, 'F', [])$  returns the function value of  $F$ , i.e.,  $w = F(x, y)$ ; this is similar for  $G$ ,  $f$ , and  $g$ ;
- $w = \text{example\_name}(x, y, 'F', 'x')$  returns the partial derivative of  $F$  with respect to  $x$ , i.e.,  $w = \nabla_x F(x, y)$ ;
- $w = \text{example\_name}(x, y, 'G', 'y')$  returns the Jacobian matrix of  $G$  with respect to  $y$ , i.e.,  $w = \nabla_y G(x, y)$ ;
- $w = \text{example\_name}(x, y, 'f', 'xy')$  returns the Hessian matrix of  $f$  with respect to  $xy$ , i.e.,  $w = \nabla_{xy}^2 f(x, y)$ ;
- $w = \text{example\_name}(x, y, 'g', 'yy')$  returns the second order derivative of  $g$  with respect to  $yy$ , i.e.,  $w = \nabla_{yy}^2 g(x, y)$ .

We now use two examples to illustrate the definitions above. The first one is nonlinear while the second one is a simple bilevel program.

### Example

Shimizu et al. (1997), see [50], considered the bilevel program (1) with

$$\begin{aligned} F(x, y) &:= (x - 5)^2 + (2y + 1)^2, \\ f(x, y) &:= (y - 1)^2 - 1.5xy, \\ g(x, y) &:= \begin{bmatrix} -3x + y + 3 \\ x - 0.5y - 4 \\ x + y - 7 \end{bmatrix}. \end{aligned}$$

Here, we have dimensions  $n_x = 1$ ,  $n_y = 1$ ,  $n_G = 0$ , and  $n_g = 3$ . The m-file is named by ShimizuEtal1997a (i.e., `example_name = ShimizuEtal1997a`) and was coded in MATLAB as it can be seen in Table 2. If we are given some inputs (as in left column of the table below), then ShimizuEtal1997a will return us corresponding results as in the right column:

Inputs	Outputs
<code>x = 4</code>	<code>x = 4</code>
<code>y = 0</code>	<code>y = 0</code>
<code>F = ShimizuEtal1997a(x, y, 'F')</code>	<code>F = 2</code>
<code>Fx = ShimizuEtal1997a(x, y, 'F', 'x')</code>	<code>Fx = -2</code>
<code>Gy = ShimizuEtal1997a(x, y, 'G', 'y')</code>	<code>Gy = []</code>
<code>fx = ShimizuEtal1997a(x, y, 'f', 'xy')</code>	<code>fx = -1.5</code>
<code>gy = ShimizuEtal1997a(x, y, 'g', 'yy')</code>	<code>gy = [0; 0; 0]</code>

**Table 2** Matlab code for ShimizuEtal1997a.

---

```
function w=ShimizuEtal1997a(x,y,keyf,keyxy)
if nargin<4 || isempty(keyxy)
    switch keyf
    case 'F'; w = (x-5)^2+(2*y+1)^2;
    case 'G'; w = [];
    case 'f'; w = (y-1)^2-1.5*x*y;
    case 'g'; w = [-3*x+y+3; x-0.5*y-4; x+y-7];
    end
else
    switch keyf
    case 'F'
        switch keyxy
        case 'x' ; w = 2*(x-5);
        case 'y' ; w = 4*(2*y+1);
        case 'xx'; w = 2;
        case 'xy'; w = 0;
        case 'yy'; w = 8;
        end
    case 'G'
        switch keyxy
        case 'x' ; w = [];
        case 'y' ; w = [];
        case 'xx'; w = [];
        case 'xy'; w = [];
        case 'yy'; w = [];
        end
    case 'f'
        switch keyxy
        case 'x' ; w = -1.5*y;
        case 'y' ; w = 2*(y-1)-1.5*x;
        case 'xx'; w = 0;
        case 'xy'; w = -1.5;
        case 'yy'; w = 2;
        end
    case 'g'
        switch keyxy
        case 'x' ; w = [-3; 1; 1];
        case 'y' ; w = [ 1;-0.5; 1];
        case 'xx'; w = [ 0; 0; 0];
        case 'xy'; w = [ 0; 0; 0];
        case 'yy'; w = [ 0; 0; 0];
        end
    end
end
end
end
```

---

**Example**

Franke et al. (2018), see [73], considered the bilevel program (1) with

$$\begin{aligned} F(y) &:= -y_2, \\ f(y) &:= y_3, \\ g(y) &:= \begin{bmatrix} y_1^2 - y_3 \\ y_1^2 + y_2^2 - 1 \\ -y_3 \end{bmatrix}. \end{aligned}$$

Here, we have dimensions  $n_x = 0$ ,  $n_y = 3$ ,  $n_G = 0$ , and  $n_g = 3$ . The m-file is named by `FrankeEta12018Ex513` (i.e., `exmaple_name = FrankeEta12018Ex513`) and is equally coded in MATLAB as described in Table 3.

It is worth mentioning that despite the lack of variable  $x$  in the latter example, we still treat it as an input, for the sake of unifying the inputs of the function handle as in (3). Hence, for all the simple bilevel optimization examples, we input  $x$  as a scalar. In this way,  $x$  has no impact on the example itself.

**2.2 Useful details on the examples**

The details related to each example presented in the BOLIB library are in a column of Table 4 below. As we mentioned before, those examples are classified into 3 categories: nonlinear, linear and simple bilevel optimisation test examples. The first column of the table provides the list of problems, as they appear in the `Examples` subfolder of the `BOLIBver2` folder. The second column gives the reference in the literature where the example might have first appeared. The third column combines the labels corresponding to the nature of the functions involved in (1). Precisely, “N” and “L” will be used to indicate whether the functions  $F$ ,  $G$ ,  $f$ , and  $g$  are nonlinear (N) or linear (L), while “O” is used to symbolize that there is either no function  $G$  or  $g$  present in problem (1). Then follows the column with  $n_x$  and  $n_y$  for the upper and lower-level variable dimensions, as well as  $n_G$  (resp.  $n_g$ ) to denote the the number of components of the upper (resp. lower)-level constraint function. On the other hand,  $F^*$  and  $f^*$  denote the best known optimal upper and lower-level objective function values, respectively, according to the reference that is listed in the last column.

Note that examples `Zlobec2001b` and `MitsosBartonEx32` have no optimal solutions. There are 4 examples involving parameters; i.e., `CalamaiVicente1994a` with  $\rho \geq 1$  (its  $F^*$  and  $f^*$  listed in the table are under  $\rho = 1$ , other cases can be found in [7]), `HenrionSurowiec2011` with  $c \in \mathbb{R}$ , `IshizukaAiyoshi1992a` with  $M > 1$  and `RobustPortfolioP1` with  $\delta \in [1, +\infty]$  (its  $F^*$  and  $f^*$  listed in the



**Table 3** Matlab code for FrankeEtal2018Ex513.

---

```

function w=FrankeEtal2018Ex513(x,y,keyf,keyxy)
if nargin<4 || isempty(keyxy)
    switch keyf
    case 'F'; w = -y(2);
    case 'G'; w = [];
    case 'f'; w = y(3);
    case 'g'; w = [y(1)^2-y(3); y(1)^2+y(2)^2-1; -y(3)];
    end
else
    switch keyf
    case 'F'
        switch keyxy
        case 'x' ; w = 0;
        case 'y' ; w = [0; -1; 0];
        case 'xx'; w = 0;
        case 'xy'; w = zeros(3,1);
        case 'yy'; w = zeros(3,3);
        end
    case 'G'
        switch keyxy
        case 'x' ; w = [];
        case 'y' ; w = [];
        case 'xx'; w = [];
        case 'xy'; w = [];
        case 'yy'; w = [];
        end
    case 'f'
        switch keyxy
        case 'x' ; w = 0;
        case 'y' ; w = [0; 0; 1];
        case 'xx'; w = 0;
        case 'xy'; w = zeros(3,1);
        case 'yy'; w = zeros(3,3);
        end
    case 'g'
        switch keyxy
        case 'x' ; w = zeros(3,1);
        case 'y' ; w = [2*y(1) 0 -1; 2*y(1) 2*y(2) 0; 0 0 -1];
        case 'xx'; w = zeros(3,1);
        case 'xy'; w = zeros(9,1);
        case 'yy'; w = [2 0 0;0 0 0;0 0 0;2 0 0; 0 2 0;zeros(4,3)];
        end
    end
end
end
end

```

---

table are under  $\delta = 2$ ). Dimensions  $n_x, n_y, n_G$  or  $n_g$  of examples `OptimalControl`, `RobustPortfolioP1`, `RobustPortfolioP2`, and `ShehuEtal2019Ex42` can be altered to get problems of different sizes, as necessary.

Table 4: List of bilevel programs with related labels and known solutions.

Example name	RefI	$F-G-f-g$	$n_x$	$n_y$	$n_G$	$n_g$	$F^*$	$f^*$	RefII
<b>Nonlinear bilevel programs</b>									
AiyoshiShimizu1984Ex2	[1]	L-L-N-L	2	2	5	6	5	0	[1]
AllendeStill2013	[2]	N-L-N-N	2	2	5	2	1	-0.5	[2]
AnEtal2009	[3]	N-L-N-L	2	2	6	4	2251.6	565.8	[3]
Bard1988Ex1	[4]	N-L-N-L	1	1	1	4	17	1	[4]
Bard1988Ex2	[4]	N-L-N-L	4	4	9	12	-6600	54	[11]
Bard1988Ex3	[4]	N-N-N-N	2	2	3	4	-12.68	-1.02	[8]
Bard1991Ex1	[5]	L-L-N-L	1	2	2	3	2	12	[5]
BardBook1998Ex832	[6]	N-L-L-L	2	2	4	7	0	5	
CalamaiVicente1994a	[7]	N-O-N-L	1	1	0	3	0	0	[7]
CalamaiVicente1994b	[7]	N-O-N-L	4	2	0	6	0.3125	-0.4063	[7]
CalamaiVicente1994c	[7]	N-O-N-L	4	2	0	6	0.3125	-0.4063	[7]
CalveteGale1999P1	[9]	L-L-L-N	2	3	2	6	-29.2	0.31	[9, 23]
ClarkWesterberg1990a	[10]	N-L-N-L	1	1	2	3	5	4	[47]
Colson2002BIPA1	[11]	N-L-N-L	1	1	3	3	250	0	
Colson2002BIPA2	[11]	N-L-N-L	1	1	1	4	17	2	[8]
Colson2002BIPA3	[11]	N-L-N-L	1	1	2	2	2	24.02	[8]
Colson2002BIPA4	[11]	N-L-N-L	1	1	2	2	88.79	-0.77	[8]
Colson2002BIPA5	[11]	N-L-N-N	1	2	1	6	2.75	0.57	[8]
Dempe1992a	[12]	L-N-N-N	2	2	1	2	×	×	
Dempe1992b	[12]	N-O-N-N	1	1	0	1	31.25	4	[8]
DempeDutta2012Ex24	[14]	N-O-N-N	1	1	0	1	0	0	[14]
DempeDutta2012Ex31	[14]	L-N-N-N	2	2	4	2	-1	4	[14]
DempeEtal2012	[15]	L-L-N-L	1	1	2	2	-1	-1	[15]
DempeFranke2011Ex41	[16]	N-L-N-L	2	2	4	4	5	-2	[16]
DempeFranke2011Ex42	[16]	N-L-N-L	2	2	4	3	2.13	-3.5	[16]
DempeFranke2014Ex38	[17]	L-L-N-L	2	2	4	4	-1	-4	[17]
DempeLohse2011Ex31a	[18]	N-O-N-L	2	2	0	4	-5.5	0	[18]
DempeLohse2011Ex31b	[18]	N-O-N-L	3	3	0	5	-12	0	
DeSilva1978	[19]	N-O-N-L	2	2	0	4	-1	0	[8]
FalkLiu1995	[20]	N-O-N-L	2	2	0	4	-2.1962	0	[8]
FloudasEtal2013	[21]	L-L-N-L	2	2	4	7	0	200	[52]
FloudasZlobec1998	[22]	N-L-L-N	1	2	2	6	1	-1	[23, 39]
GumusFloudas2001Ex1	[23]	N-L-N-L	1	1	3	3	2250	197.75	[39]
GumusFloudas2001Ex3	[23]	L-L-N-L	2	3	4	9	-29.2	0.31	[39]
GumusFloudas2001Ex4	[23]	N-L-N-L	1	1	5	2	9	0	[39]
GumusFloudas2001Ex5	[23]	L-L-N-N	1	2	2	6	0.19	-7.23	[39]
HatzEtal2013	[24]	L-O-N-L	1	2	0	2	0	0	[24]
HendersonQuandt1958	[25]	N-L-N-L	1	1	2	1	-3266.7	-711.11	[25]
HenrionSurowiec2011	[26]	N-O-N-O	1	1	0	0	$-c^2/4$	$-c^2/8$	[27]
IshizukaAiyoshi1992a	[28]	N-L-L-L	1	2	1	5	0	-M	[28]
KleniatiAdjiman2014Ex3	[29]	L-L-N-L	1	1	2	2	-1	0	[29]
KleniatiAdjiman2014Ex4	[29]	N-N-N-N	5	5	13	11	-10	-3.1	[29]
LamparSagrat2017Ex23	[30]	L-L-N-L	1	2	2	2	-1	1	[30]

LamparSagrat2017Ex31	[31]	N-L-L-L	1	1	1	1	1	0	[31]
LamparSagrat2017Ex32	[31]	N-O-N-O	1	1	0	0	0.5	0	[31]
LamparSagrat2017Ex33	[31]	N-L-L-L	1	2	1	3	0.5	0	[31]
LamparSagrat2017Ex35	[31]	N-L-L-L	1	1	2	3	0.8	-0.4	[31]
LucchettiEtal1987	[33]	N-L-N-L	1	1	2	2	0	0	[33]
LuDebSinha2016a	[34]	N-L-N-O	1	1	4	0	1.14	1.69	[34]
LuDebSinha2016b	[34]	N-L-N-O	1	1	4	0	0	1.66	[34]
LuDebSinha2016c	[34]	N-L-N-O	1	1	4	0	1.12	0.06	[34]
LuDebSinha2016d	[34]	L-N-L-N	2	2	11	3	×	×	
LuDebSinha2016e	[34]	N-L-L-N	1	2	6	3	×	×	
LuDebSinha2016f	[34]	L-N-N-O	2	1	9	0	×	×	
MacalHurter1997	[36]	N-O-N-O	1	1	0	0	81.33	-0.33	[36]
Mirrlees1999	[38]	N-O-N-O	1	1	0	0	1	0.06	[38]
MitsosBarton2006Ex38	[39]	N-L-N-L	1	1	4	2	0	0	[39]
MitsosBarton2006Ex39	[39]	L-L-N-L	1	1	3	2	-1	-1	[39]
MitsosBarton2006Ex310	[39]	L-L-N-L	1	1	2	2	0.5	-0.1	[39]
MitsosBarton2006Ex311	[39]	L-L-N-L	1	1	2	2	-0.8	0	[39]
MitsosBarton2006Ex312	[39]	N-L-N-L	1	1	2	2	0	0	[39]
MitsosBarton2006Ex313	[39]	L-L-N-L	1	1	2	2	-1	0	[39]
MitsosBarton2006Ex314	[39]	N-L-N-L	1	1	2	2	0.25	-0.08	[39]
MitsosBarton2006Ex315	[39]	L-L-N-L	1	1	2	2	0	-0.83	[39]
MitsosBarton2006Ex316	[39]	L-L-N-L	1	1	2	2	-2	0	[39]
MitsosBarton2006Ex317	[39]	N-L-N-L	1	1	2	2	0.19	-0.02	[39]
MitsosBarton2006Ex318	[39]	N-L-N-L	1	1	2	2	-0.25	0	[39]
MitsosBarton2006Ex319	[39]	N-L-N-L	1	1	2	2	-0.26	0	[39]
MitsosBarton2006Ex320	[39]	N-L-N-L	1	1	2	2	0.31	-0.08	[39]
MitsosBarton2006Ex321	[39]	N-L-N-L	1	1	2	2	0.21	-0.07	[39]
MitsosBarton2006Ex322	[39]	N-L-N-N	1	1	2	3	0.21	-0.07	[39]
MitsosBarton2006Ex323	[39]	N-N-L-N	1	1	3	3	0.18	-1	[39]
MitsosBarton2006Ex324	[39]	N-L-N-L	1	1	2	2	-1.75	0	[39]
MitsosBarton2006Ex325	[39]	N-N-N-N	2	3	6	9	-1	-2	[39]
MitsosBarton2006Ex326	[39]	N-N-N-L	2	3	7	6	-2.35	-2	[39]
MitsosBarton2006Ex327	[39]	N-N-N-N	5	5	13	13	2	-1.1	[39]
MitsosBarton2006Ex328	[39]	N-N-N-N	5	5	13	13	-10	-3.1	[39]
MorganPatrone2006a	[40]	L-L-N-L	1	1	2	2	-1	0	[40]
MorganPatrone2006b	[40]	L-O-N-L	1	1	0	4	-1.25	0	[40]
MorganPatrone2006c	[40]	L-O-N-L	1	1	0	4	-1	-0.25	[40]
MuuQuy2003Ex1	[41]	N-L-N-L	1	2	2	3	-2.08	-0.59	[41]
MuuQuy2003Ex2	[41]	N-L-N-L	2	3	3	4	0.64	1.67	[41]
NieEtal2017Ex34	[42]	L-L-N-N	1	2	2	2	2	0	[42]
NieEtal2017Ex52	[42]	N-N-N-N	2	3	5	2	-1.71	-2.23	[42]
NieEtal2017Ex54	[42]	N-N-N-N	4	4	3	2	-0.44	-1.19	[42]
NieEtal2017Ex57	[42]	N-N-N-N	2	3	5	2	-2	-1	[42]
NieEtal2017Ex58	[42]	N-N-N-N	4	4	3	2	-3.49	-0.86	[42]
NieEtal2017Ex61	[42]	N-N-N-N	2	2	5	1	-1.02	-1.08	[42]
Outrata1990Ex1a	[43]	N-O-N-L	2	2	0	4	-8.92	-6.05	[43]
Outrata1990Ex1b	[43]	N-O-N-L	2	2	0	4	-7.56	-0.58	[43]
Outrata1990Ex1c	[43]	N-O-N-L	2	2	0	4	-12	-112.71	[43]
Outrata1990Ex1d	[43]	N-O-N-L	2	2	0	4	-3.6	-2	[43]
Outrata1990Ex1e	[43]	N-O-N-L	2	2	0	4	-3.15	-16.29	[43]

Outrata1990Ex2a	[43]	N-L-N-L	1	2	1	4	0.5	-14.53	[43]
Outrata1990Ex2b	[43]	N-L-N-L	1	2	1	4	0.5	-4.5	[43]
Outrata1990Ex2c	[43]	N-L-N-L	1	2	1	4	1.86	-10.93	[43]
Outrata1990Ex2d	[43]	N-L-N-N	1	2	1	4	0.92	-19.47	[43]
Outrata1990Ex2e	[43]	N-L-N-N	1	2	1	4	0.90	-14.94	[43]
Outrata1993Ex31	[44]	N-L-N-N	1	2	1	4	1.56	-11.67	[44]
Outrata1993Ex32	[44]	N-L-N-N	1	2	1	4	3.21	-20.53	[44]
Outrata1994Ex31	[45]	N-L-N-N	1	2	2	4	3.21	-20.53	[45]
OutrataCervinka2009	[46]	L-L-N-L	2	2	1	3	0	0	[46]
PaulaviciusEtal2017a	[47]	N-L-N-L	1	1	4	2	0.25	0	[47]
PaulaviciusEtal2017b	[47]	L-L-N-L	1	1	4	2	-2	-1.5	[47]
SahinCircic1998Ex2	[48]	N-L-N-L	1	1	2	3	5	4	[48]
ShimizuAiyoshi1981Ex1	[49]	N-L-N-L	1	1	3	3	100	0	[49]
ShimizuAiyoshi1981Ex2	[49]	N-L-N-L	2	2	3	4	225	100	[49]
ShimizuEtal1997a	[50]	N-O-N-L	1	1	0	3	×	×	
ShimizuEtal1997b	[50]	N-L-N-L	1	1	2	2	2250	197.75	[50]
SinhaMaloDeb2014TP3	[51]	N-N-N-N	2	2	3	4	-18.68	-1.02	[51]
SinhaMaloDeb2014TP6	[51]	N-L-N-L	1	2	1	6	-1.21	7.62	[51]
SinhaMaloDeb2014TP7	[51]	N-N-N-L	2	2	4	4	-1.96	1.96	[51]
SinhaMaloDeb2014TP8	[51]	N-L-N-L	2	2	5	6	0	100	[51]
SinhaMaloDeb2014TP9	[51]	N-O-N-L	10	10	0	20	0	1	[51]
SinhaMaloDeb2014TP10	[51]	N-O-N-L	10	10	0	20	0	1	[51]
TuyEtal2007	[52]	N-L-L-L	1	1	2	3	22.5	-1.52	[52]
Vogel2002	[53]	N-L-N-L	1	1	2	1	1	-2	[53]
WanWangLv2011	[54]	N-O-L-L	2	3	0	8	10.63	-0.5	[54]
YeZhu2010Ex42	[55]	N-L-N-L	1	1	2	1	1	-2	[55]
YeZhu2010Ex43	[55]	N-L-N-L	1	1	2	1	1.25	-2	[55]
Yezza1996Ex31	[56]	N-L-N-L	1	1	2	2	1.5	-2.5	[56]
Yezza1996Ex41	[56]	N-O-N-L	1	2	0	2	0.5	2.5	[56]
Zlobec2001a	[57]	N-O-L-L	1	2	0	3	-1	-1	[57]
Zlobec2001b	[57]	L-L-L-N	1	1	2	4	no	solution	[57]
DesignCentringP1	[71]	N-N-N-N	3	6	3	3	×	×	
DesignCentringP2	[71]	N-N-N-N	4	6	5	3	×	×	
DesignCentringP3	[71]	N-N-N-N	6	6	3	3	×	×	
DesignCentringP4	[71]	N-N-N-N	4	6	3	12	×	×	
NetworkDesignP1	[11]	N-L-N-L	5	5	5	6	300.5	419.8	[8]
NetworkDesignP2	[11]	N-L-N-L	5	5	5	6	142.9	81.95	[8]
OptimalControl	[37]	N-N-N-L	2	$n_y$	3	$2n_y$	×	×	
RobustPortfolioP1	[71]	L-N-N-N	N+1	N	N+3	N+1	1.15	0	[71]
RobustPortfolioP2	[71]	L-N-N-N	N+1	N	N+3	N+1	1.15	0	[71]
TollSettingP1	[11]	N-L-N-L	3	8	3	18	-7	12	[8]
TollSettingP2	[11]	N-L-N-L	3	18	3	38	-4.5	32	[8]
TollSettingP3	[11]	N-L-N-L	3	18	3	38	-3.5	32	[8]
TollSettingP4	[11]	N-O-N-L	2	4	0	8	-4	14	[8]
TollSettingP5	[11]	N-O-N-L	1	4	0	8	-2.5	14	[8]
<b>Linear bilevel programs</b>									
AnandalinghamWhite1990	[58]	L-L-L-L	1	1	1	6	-49	15	[58]
Bard1984a	[75]	L-L-L-L	1	1	1	5	28/9	-60/9	[75]
Bard1984b	[75]	L-L-L-L	1	1	1	5	-37.6	1.6	[75]
Bard1991Ex2	[5]	L-L-L-L	1	2	1	5	-1	-1	[5]

BardFalk1982Ex2	[59]	L-L-L-L	2	2	2	5	-3.25	-4	[59]
Ben-AyedBlair1990a	[76]	L-L-L-L	1	2	2	4	-2.5	-5	[76]
Ben-AyedBlair1990b	[76]	L-L-L-L	1	1	1	4	-6	5	[76]
BialasKarwan1984a	[77]	L-L-L-L	1	2	1	7	-2	-0.5	[77]
BialasKarwan1984b	[77]	L-L-L-L	1	1	1	6	-11	11	[77]
CandlerTownsley1982	[60]	L-L-L-L	2	3	2	6	-29.2	3.2	[60]
ClarkWesterberg1988	[61]	L-O-L-L	1	1	0	3	-37	14	[61]
ClarkWesterberg1990b	[10]	L-L-L-L	1	2	2	5	-13	-4	[10]
GlackinEtal2009	[62]	L-L-L-L	2	1	3	3	6	0	[62]
HaurieSavardWhite1990	[78]	L-O-L-L	1	1	0	4	27	-3	[78]
HuHuangZhang2009	[63]	L-L-L-L	1	2	1	5	-76/9	-41/9	[63]
LanWenShihLee2007	[64]	L-L-L-L	1	1	1	7	-85.09	50.17	[64]
LiuHart1994	[67]	L-L-L-L	1	1	1	4	-16	4	[67]
MershaDempe2006Ex1	[65]	L-L-L-L	1	1	1	5	×	×	[65]
MershaDempe2006Ex2	[65]	L-L-L-L	1	1	2	2	-20	-6	[65]
TuyEtal1993	[79]	L-L-L-L	2	2	3	4	-3.25	-6	[79]
TuyEtal1994	[80]	L-L-L-L	2	2	3	3	6	0	[80]
TuyEtal2007Ex3	[52]	L-L-L-L	10	6	12	13	-467.46	-11.62	[52]
VisweswaranEtal1996	[68]	L-L-L-L	1	1	1	5	28/9	-60/9	[68]
WangJiaoLi2005	[66]	L-L-L-L	1	2	2	2	-1000	-1	[66]
<b>Simple bilevel programs</b>									
FrankeEtal2018Ex53	[73]	N-L-N-L	0	2	4	4	1	1	[73]
FrankeEtal2018Ex511	[73]	N-O-L-L	0	3	0	4	3	0	[73]
FrankeEtal2018Ex513	[73]	L-O-L-N	0	3	0	3	-1	0	[73]
FrankeEtal2018Ex521	[73]	L-O-L-N	0	2	0	3	-1	0	[73]
MitsosBarton2006Ex31	[39]	L-L-L-L	0	1	2	2	1	-1	[39]
MitsosBarton2006Ex32	[39]	L-L-L-L	0	1	3	2	no	solution	[39]
MitsosBarton2006Ex33	[39]	L-L-N-N	0	1	2	3	-1	1	[39]
MitsosBarton2006Ex34	[39]	L-L-N-L	0	1	2	2	1	-1	[39]
MitsosBarton2006Ex35	[39]	L-L-N-L	0	1	2	2	0.5	-1	[39]
MitsosBarton2006Ex36	[39]	L-L-N-L	0	1	2	2	-1	-1	[39]
ShehuEtal2019Ex42	[72]	N-O-N-O	0	$n_y$	0	0	×	×	

It is worth pointing out that some examples involve equalities constraints in the upper or lower-level problems. As only 8% of the BOLIB problems have such constraints, we preserve the uniformity in the structure of the codes by converting equalities constraints into inequalities. For the sake of clarity, we list all the examples with equality constraints below.

Table 5: List of bilevel programs with equalities constraints.

Example name	Ref.	$F-G-H-f-g-h$	$n_x$	$n_y$	$n_G$	$n_H$	$n_g$	$n_h$
DempeDutta2012Ex31	[14]	L-L-N-N-N-O	2	2	2	1	2	0
DempeFranke2011Ex41	[16]	N-L-L-N-L-O	2	2	2	1	4	0
DempeFranke2011Ex42	[16]	N-L-L-N-L-O	2	2	2	1	3	0
Zlobec2001b	[57]	L-L-O-L-L-N	1	1	2	0	2	1
NetworkDesignP1	[11]	N-L-O-N-O-L	5	5	5	0	0	3

NetworkDesignP2	[11]	N-L-O-N-O-L	5	5	5	0	0	3
OptimalControl	[37]	N-N-O-N-L-L	2	$n_y$	3	0	$n_y$	$\frac{1}{2}n_y$
RobustPortfolioP1	[71]	L-N-L-N-N-O	N+1	N	N+1	1	N+1	0
RobustPortfolioP2	[71]	L-N-L-N-N-O	N+1	N	N+1	1	N+1	0
TollSettingP1	[11]	N-L-O-N-L-L	3	8	3	0	8	5
TollSettingP2	[11]	N-L-O-N-L-L	3	18	3	0	18	10
TollSettingP3	[11]	N-L-O-N-L-L	3	18	3	0	18	10
TollSettingP4	[11]	N-O-O-N-L-L	2	4	0	0	2	4
TollSettingP5	[11]	N-O-O-N-L-L	1	4	0	0	2	4

### 3 How to access the library?

The BOLIB library (versions 1 and 2) can be accessed through the dedicated website [biopt.github.io/bolib](https://biopt.github.io/bolib). Under this link, you will find the zipped folder named BOLIBver2 containing all the relevant files for the version of the library presented in this paper. The folder contains the subfolder named `Examples`, which contains all the m-files with the codes of the examples, as described in the previous section. The pdf file named `Formulas` collects all the mathematical formulas of all the examples in this library. To start with the library, it is advised to consult the `readme` file for some useful instructions on how to use it.

**Acknowledgements** The work of the first and second authors is partly funded by the EPSRC Grant EP/P022553/1. The third author's work is partly funded by the University of Southampton's Presidential Scholarship. We thank Dr Patrick Mehrlitz (Brandenburgische Technische Universität Cottbus-Senftenberg) for the `OptimalControl` example and related MATLAB files.

### References

1. E. Aiyoshi and K. Shimizu, A solution method for the static constrained Stackelberg problem via penalty method, *IEEE Transactions on Automatic Control*, 29, 1111-1114, 1984.
2. G.B. Allende and G. Still, Solving bilevel programs with the KKT-approach. *Mathematical Programming*, 138(1-2), 309-332, 2013.
3. L.T.H. An, P.D. Tao, N.N. Canh and N.V. Thoai, DC programming techniques for solving a class of nonlinear bilevel programs, *Journal of Global Optimization*, 44(3), 313-337, 2009.
4. J.F. Bard, Convex two-level optimization, *Mathematical Programming*, 40, 15-27, 1988.
5. J.F. Bard, Some properties of the bilevel programming problem, *Journal of Optimization Theory and Applications*, 68(2), 371-378, 1991.
6. J.F. Bard, *Practical bilevel optimization: Algorithms and applications*, Dordrecht: Kluwer Academic Publishers, 1998.
7. P.H. Calamai and L.N. Vicente, Generating quadratic bilevel programming test problems, *ACM Transactions on Mathematical Software*, 20(1), 103-119, 1994.
8. B. Colson, P. Marcotte and G. Savard, A trust-region method for nonlinear bilevel programming: algorithm and computational experience, *Computational Optimization and Applications*, 30(3), 211-227, 2005.

9. H.I. Calvete and C. Galé, The bilevel linear/linear fractional programming problem, *European Journal of Operational Research*, 114(1), 188-197, 1999.
10. P.A. Clark and A.W. Westerberg, Bilevel programming for steady-state chemical process design—I. Fundamentals and algorithms. *Computers & Chemical Engineering*, 14 (1): 87-97, 1990.
11. B. Colson, BIPA (Bilevel Programming with Approximation Methods): Software guide and test problems, Technical report, 2002.
12. S. Dempe, A necessary and a sufficient optimality condition for bilevel programming problems, *Optimization*, 25, 341-354, 1992.
13. S. Dempe, N. Dinh, and J. Dutta, Optimality conditions for a simple convex bilevel programming problem, In *Variational analysis and generalized differentiation in optimization and control*, pp. 149-161, Springer, New York, NY, 2010.
14. S. Dempe and J. Dutta, Is bilevel programming a special case of a mathematical program with complementarity constraints? *Mathematical programming*, 131(1-2), 37-48, 2012.
15. S. Dempe, B. Mordukhovich, and A.B. Zemkoho, Sensitivity analysis for two-level value functions with applications to bilevel programming, *SIAM Journal on Optimization*, 22(4), 1309-1343, 2012.
16. S. Dempe and S. Franke, An algorithm for solving a class of bilevel programming problems, Preprint 2011-04, TU Bergakademie, 2011.
17. S. Dempe and S. Franke, Solution algorithm for an optimistic linear stackelberg problem, *Computers & Operations Research*, 41, 277-281, 2014.
18. S. Dempe and S. Lohse, Dependence of bilevel programming on irrelevant data, Dekan der Fakultät für Mathematik und Informatik, 2011.
19. A.H. De Silva, Sensitivity formulas for nonlinear factorable programming and their application to the solution of an implicitly defined optimization model of US crude oil production, Ph.D. thesis, George Washington University, 1978.
20. J.E. Falk and J. Liu, On bilevel programming, Part I: general nonlinear cases, *Mathematical Programming*, 70(1), 47-72, 1995.
21. C.A. Floudas, P.M. Pardalos, C. Adjiman, W.R. Esposito, Z.H. Gümmüs, S.T. Harding, J.L. Klepeis, C.A. Meyer, and C.A. Schweiger, *Handbook of test problems in local and global optimization*, Springer, 1999.
22. C.A. Floudas and S. Zlobec, Optimality and duality in parametric convex lexicographic programming, In *Multilevel Optimization: Algorithms and Applications*, pp. 359-379, Springer, 1998.
23. Z.H. Gumus and C.A. Floudas, Global optimization of nonlinear bilevel programming problems, *Journal of Global Optimization*, 20(1), 1-31, 2001.
24. K. Hatz, S. Leyffer, J.P. Schlöder, and H.G. Bock, Regularizing bilevel nonlinear programs by lifting, Technical Report, Argonne National Laboratory, USA, 2013.
25. J.M. Henderson and R.E. Quandt, *Microeconomic theory: a mathematical approach*, McGraw-Hill: New York, 1958.
26. R. Henrion and T. Surowiec, On calmness conditions in convex bilevel programming, *Applicable Analysis*, 90(6), 951-970, 2011.
27. F. Facchinei, H. Jiang, and L. Qi, A smoothing method for mathematical programs with equilibrium constraints, *Mathematical programming*, 85(1), 107-134, 1999.
28. Y. Ishizuka and E. Aiyoshi, Double penalty method for bilevel optimization problems, *Annals of Operations Research*, 34(1), 73-88, 1992.
29. P.-M. Kleniati and C.S. Adjiman, Branch-and-sandwich: a deterministic global optimization algorithm for optimistic bilevel programming problems, Part II: Convergence analysis and numerical results, *Journal of Global Optimization*, 60(3), 459-481, 2014.
30. L. Lampariello and S. Sagratella, Numerically tractable optimistic bilevel problems, *Optimization-Online*, 2017.
31. L. Lampariello and S. Sagratella, A bridge between bilevel programs and nash games, *Journal of Optimization Theory and Applications*, 174(2), 613-635, 2017.
32. G.H. Lin, M. Xu and J.Y. Jane, On solving simple bilevel programs with a nonconvex lower level program. *Mathematical Programming*, 144(1-2), 277-305, 2014.

33. R. Lucchetti, F. Mignanego, and G. Pieri, Existence theorems of equilibrium points in Stackelberg games with constraints, *Optimization*, 18(6), 857-866, 1987.
34. Z.C. Lu, K. Deb, and A. Sinha, Robust and reliable solutions in bilevel optimization problems under uncertainties, COIN Report 2016026, Retrived on 19 November 2017 from <http://www.egr.msu.edu/kdeb/papers/c2016026.pdf>.
35. S. Leyffer, MacMPEC: AMPL collection of MPECs, Argonne National Laboratory, 2000. Available at <https://wiki.mcs.anl.gov/leyffer/index.php/MacMPEC>.
36. C.M. Macal and A.P.Hurter, Dependence of bilevel mathematical programs on irrelevant constraints, *Computers and Operations Research*, 24(12), 1129-1140, 1997.
37. P. Mehlitz and G. Wachsmuth, Weak and strong stationarity in generalized bilevel programming and bilevel optimal control, *Optimization*, 65(5), 907-935, 2016.
38. J.A. Mirrlees, The theory of moral hazard and unobservable behaviour: Part I, *The Review of Economic Studies*, 66(1), 3-21, 1999.
39. A. Mitsos and P.I. Barton, A Test Set for Bilevel Programs, Technical report, MIT, 2006.
40. J. Morgan and F. Patrone, Stackelberg problems: Subgame perfect equilibria via tikhonov regularization, *Advances in dynamic games*, 209-221, 2006.
41. L.D. Muu and N.V. Quy, A global optimization method for solving convex quadratic bilevel programming problems, *Journal of Global Optimization*, 26(2), 199-219, 2003.
42. J.W. Nie, L. Wang, and J.J. Ye, Bilevel polynomial programs and semidefinite relaxation methods, *SIAM Journal on Optimization*, 27(3), 1728-1757, 2017.
43. J.V. Outrata, On the numerical solution of a class of Stackelberg problems, *Zeitschrift für Operations Research*, 34(4), 255-277, 1990.
44. J.V. Outrata, Necessary optimality conditions for Stackelberg problems, *Journal of Optimization Theory and Applications*, 76(2), 305-320, 1993.
45. J.V. Outrata, On optimization problems with variational inequality constraints, *SIAM Journal on optimization*, 4(2), 340-357, 1994.
46. J.V. Outrata and M. Cervinka, On the implicit programming approach in a class of mathematical programs with equilibrium constraints, *Control and Cybernetics*, 38(4B), 1557-1574, 2009.
47. R. Paulavicius, P.M. Kleniati, and C.S. Adjiman, BBASBL: Branch-And-Sandwich BiLevel solver. Implementation and computational study with the BASBLib test set, *Computers and Chemical Engineering*, 132, 106609, 2020.
48. K.H. Sahin and A.R. Ciric, A dual temperature simulated annealing approach for solving bilevel programming problems. *Computers and Chemical Engineering*, 23(1), 11-25, 1998.
49. K. Shimizu and E. Aiyoshi, A new computational method for Stackelberg and min-max problems by use of a penalty method, *IEEE Transactions on Automatic Control*, 26(2), 460-466, 1981.
50. K. Shimizu, Y. Ishizuka, and J.F. Bard, *Nondifferentiable and two-level mathematical programming*, Kluwer Academic Publishers, 1997.
51. A. Sinha, P. Malo, and K. Deb, An improved bilevel evolutionary algorithm based on quadratic approximations, In *Evolutionary Computation (CEC), 2014 IEEE Congress*, pp. 1870-1877, 2014.
52. H. Tuy, A. Migdalas, and N.T. Hoai-Phuong, A novel approach to bilevel nonlinear programming, *Journal of Global Optimization*, 38(4), 527-554, 2007.
53. S. Vogel, *Zwei-Ebenen-Optimierungsaufgaben mit nichtkonvexer Zielfunktion in der unteren Ebene*, PhD thesis, Department of Mathematics and Computer Science, TU Bergakademie Freiberg, Freiberg, Germany, 2012.
54. Z.P. Wan, G.M. Wang, and Y.B. Lv, A dual-relax penalty function approach for solving nonlinear bilevel programming with linear lower level problem, *Acta Mathematica Scientia*, 31(2), 652-660, 2011.
55. J.J. Ye and D.L. Zhu, New necessary optimality conditions for bilevel programs by combining the mpec and value function approaches, *SIAM Journal on Optimization*, 20(4), 1885-1905, 2010.
56. A. Yezaa, First-order necessary optimality conditions for general bilevel programming problems, *Journal of Optimization Theory and Applications*, 89(1), 189-219, 1996.



57. S. Zlobec, Bilevel programming: optimality conditions and duality, In: Floudas C., Pardalos P. (eds) *Encyclopedia of Optimization*, pp. 180-185, Springer, 2001.
58. A. Anandalingham and D.J. White, A solution method for the linear static Stackelberg problem using penalty functions, *IEEE Transactions on automatic control*, 35(10), 1170-1173, 1990
59. J. F. Bard and J.E. Falk, An explicit solution to the multi-level programming problem, *Computers and Operations Research*, 9, 77-100, 1982.
60. W. Candler, R. Townsley, A linear two-level programming problem, *Computers and Operations Research*, 9, 59-76, 1982.
61. P.A. Clark and A.W. Westerberg, A note on the optimality conditions for the bilevel programming problem, *Naval Research Logistics Quarterly*, 35(5), 413-418, 1988.
62. J. Glackin, J.G. Ecker, and M. Kupferschmid, Solving bilevel linear programs using multiple objective linear programming, *Journal of Optimization Theory and Applications*, 140(2), 197-212, 2009.
63. T. Hu, B. Huang, X. Zhang, A neural network approach for solving linear bilevel programming problem, In *Proceedings of the 6th ISNN Conference, AISC*, 56, 649-658, 2009.
64. K.M. Lan, U.P. Wen, H.S. Shih, and E.S. Lee, A hybrid neural network approach to bilevel programming problems, *Applied Mathematics Letters*, 20(8), 880-884, 2007.
65. A.G. Mersha and S. Dempe, Linear bilevel programming with upper level constraints depending on the lower level solution, *Applied Mathematics and Computation* 180, 247-254, 2006.
66. Y. Wang, Y.C. Jiao, and H. Li, A evolutionary algorithm for solving nonlinear bilevel programming based on a new constraint-handling scheme, *IEEE Transactions on Systems, Man and Cybernetics part C*, 35(2), 221-232, 2005.
67. Y.-H. Liu and S.M. Hart, Characterizing an optimal solution to the linear bilevel programming problem, *European Journal of Operational Research*, 73(1), 164-166, 1994.
68. V. Visweswaran, C.A. Floudas, M.G. Ierapetritou, and E.N. Pistikopoulos, A decomposition-based global optimization approach for solving bi-level linear and quadratic programs, In: C.A. Floudas and P.M. Pardalos (eds) *State of the Art in Global Optimization. Nonconvex Optimization and Its Applications*, vol 7, pp 139-162, Springer, 1996.
69. M. Labbé, P. Marcotte, and G. Savard, A bilevel model of taxation and its applications to optimal highway, *Management Science*, 44, 1595-1607, 1998.
70. P. Gritzmann and V. Klee, On the complexity of some basic problems in computational convexity. I. Containment problems, *Discrete Mathematics*, 136, 129-174, 1994.
71. O. Stein and G. Still, Solving semi-infinite optimization problems with interior point techniques, *SIAM Journal on Control and Optimization*, 42(3), 769-688, 2003.
72. Y. Shehu, P.T. Vuong, and A.B. Zemkoho, An inertial extrapolation method for convex simple bilevel optimization, *Optimization Methods and Software*, in press, available online at DOI: [10.1080/10556788.2019.1619729](https://doi.org/10.1080/10556788.2019.1619729)
73. S. Franke, P. Mehlitz, and M. Pilecka, Optimality conditions for the simple convex bilevel programming problem in Banach spaces, *Optimization*, 67(2), 237-268, 2018.
74. S. Dempe, F. Harder, P. Mehlitz, and G. Wachsmuth, Solving inverse optimal control problems via value functions to global optimality, *Journal of Global Optimization*, 74(2), 297-325, 2019.
75. J. Bard, Optimality conditions for the bilevel programming problem, *Naval Research Logistics Quarterly*, 31, 13-26, 1984.
76. O. Ben-Ayed and C. Blair, Computational difficulties of bilevel linear programming, *Operations Research*, 38, 556-560, 1990.
77. W. Bialas and M. Karwan, Two-level linear programming, *Management Science*, 30, 1004-1020, 1984.
78. A. Haurie, G. Savard, and D. White, A note on: An efficient point algorithm for a linear two-stage optimization problem, *Operations Research*, 38, 553-555, 1990.
79. H. Tuy, A. Migdalas, and P. Värbrand, A global optimization approach for the linear two-level program, *Journal of Global Optimization*, 3(1), 1-23, 1993.
80. H. Tuy, A. Migdalas and P. Värbrand, A quasiconcave minimization method for solving linear two-level programs, *Journal of Global Optimization*, 4, 243-263, 1994.