



Using marine deposits to understand terrestrial human environments: 6000-year old hyperpycnal flash-flood events and their implications

R. Helen Farr^{a,*}, Alan Clapham^b, Eduard G. Reinhardt^c, Joseph I. Boyce^c, Shawn Collins^c, John Robb^b

^a Archaeology, University of Southampton, SO17 1BF, UK

^b McDonald Institute for Archaeological Research, Department of Archaeology, University of Cambridge, CB2 3DZ, UK

^c School of Geography and Earth Sciences, McMaster University, Hamilton, Canada L8S 4K1



ARTICLE INFO

Keywords:

Submerged prehistory
Hyperpycnal floods
Final Neolithic
Calabria

ABSTRACT

Offshore sedimentary deposits preserved in submarine canyons in southern Calabria, Italy, provide evidence for mid-Holocene (ca. 6000 BP) erosion and the transport of organic-rich floodplain sediments by flash-flood (hyperpycnal) flows from the San Pasquale River. Marine geophysical surveys (bathymetry, side-scan sonar) and diver reconnaissance revealed an unusual offshore peat deposit containing plant macrofossils representing local habitats, potentially reflecting human modified landscapes. The 20–30 cm thick organic deposit included a large tree trunk, well-preserved seeds, leaves, sticks and other delicate organics. Textural and microfossil (foraminifera) analysis of associated sediments (sands and muds) indicate that these deposits resulted from hyperpycnal flood events that were deposited as sediment gravity flows within gullies on the canyon margins. Whilst the value of studying *in situ* submerged prehistoric landscapes is well documented, we demonstrate that reworked floodplain deposits preserved in offshore environments can provide useful palaeoenvironmental information that may not be preserved in terrestrial settings. The botanic archive preserved in submarine flood deposits at San Pasquale affords a unique insight into the local environment in which people lived during the Final Neolithic.

1. Introduction

Coastlines structure human life: they afford access to marine resources, allow travel and inter-regional contact, concentrate population, and furnish points of reference in cultural landscapes (Robb and Farr, 2005:26). The study of coastal prehistoric activity can contribute to our understanding of several central issues, including Mesolithic forager presence, the spread of the Neolithic, and ancient and modern agriculture, subsistence and inter-regional exchange (i.e. Bailey et al., 2017:8; Bell, 2007:3), but in many regions, our knowledge of past coastal environments is limited. This is due to a number of environmental and taphonomic factors that result in poor preservation of archaeological and palaeoenvironmental records; these may include the geomorphology of the coast, tectonics, erosion and sedimentation, and historic and contemporary coastal modification. Knowledge of past changes in coastal environments is also skewed by the questions that we have traditionally been interested in and the types of archaeology we have done. For example, coastal archaeology in the Mediterranean has focussed overwhelmingly upon ports and harbours. Whilst the

archaeology of submerged landscapes has gained interest in the last decade, it has focussed on regions of greatest potential for preservation (Bailey and Flemming, 2008; Benjamin et al., 2011:290; Erlandson, 2001).

The underwater archaeological potential of submerged landscapes on continental shelves is now widely recognised (Bailey and Sakellariou, 2012; Bailey et al., 2017). Previous work has tended to focus on tectonically stable regions where gradual shelving or shallow continental shelves have been affected by postglacial marine transgression (see Flemming et al., 2017 and chapters therein). These tend to be regions with the greatest lateral displacement of palaeocoastlines, where inundation has created potential for *in situ* preservation of terrestrial sites or landforms. Locally steep shelf areas on tectonically active coastlines are, in contrast, generally regarded as of low potential for the preservation of submerged prehistory because of geodynamic conditions (Flemming et al., 2017:4) and their potential has been under explored.

In this paper, we report on the discovery of mid-Holocene (ca. 6000 BP) floodplain sediments preserved in offshore environments on the

* Corresponding author at: Archaeology, University of Southampton, Highfield SO17 1BF, UK.

E-mail address: r.h.farr@soton.ac.uk (R.H. Farr).

<https://doi.org/10.1016/j.jasrep.2019.102176>

Received 22 July 2019; Accepted 16 December 2019

Available online 20 February 2020

2352-409X/ © 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. A) Location of Bova Marina study area in southern Calabria. Generalized shelf and offshore bathymetry also shown (contour interval 250 m). B) Geophysical survey area (6 km²) showing sonar survey track lines (200 kHz side-scan, single-beam bathymetry).

tectonically active coast of southern Calabria, Italy (Fig. 1). Floodplain sediments were deposited by episodic flash-flood events within deep (> 400 m) submarine canyons formed at the mouth of the San Pasquale River (Fig. 1) and contain a rich palaeobotanical archive of Late Neolithic coastal environments. These findings challenge previous assumptions that tectonically active shelf areas have limited potential as palaeoenvironmental archives. We show, in contrast, that the unique preservation of botanical remains in flash-flood deposits (hyperpycnites) can provide palaeoenvironmental data that are otherwise unavailable from terrestrial sediment archives. Submarine canyons are a common feature of the shelf in southern Italy (Casalbore et al., 2011; Ridente et al., 2014) and in other tectonically active coastlines in the Mediterranean. Our results indicate that these settings should not be overlooked as potential sources for palaeoenvironmental and palaeoecological data.

2. Study area

2.1. Geological setting

The San Pasquale valley is located 3 km to the east of the town of Bova Marina, on the south coast of Calabria, Italy (Fig. 1). The geology

of the region is dominated by the Aspromonte massif, a plateau of folded metamorphic rocks, which rise steeply from the coast to a height of ca. 2000 masl (metres above sea-level) (Robb, 1997; Robb and Van Hove, 2003). The San Pasquale River drains from the uplands towards the coast, along a narrow alluvial valley, with steep sided slopes. The San Pasquale is a short (7–8 km), low-sinuosity braided river with coarse gravels and cobbles that typifies the river systems of Calabria and Sicily (Casalbore et al., 2011). Close to the coast (~5 km), the San Pasquale incises through the Early Miocene Stilo Capo d'Orlando Formation, which is composed of conglomerates and marine mudstones containing abundant planktic (Globigerinides and *Orbulina*) and benthic foraminifera (Cavazza and DeCelles, 1993; Patterson et al., 1995). Incision through these poorly indurated, highly erodible mudstones has resulted in steep valley side slopes, with smaller gullies along the margin (Fig. 2).

The coastline at San Pasquale is a narrow littoral plain backed by coastal foothills (Brondi et al., 2003; Antonioli et al., 2017:346). It has a narrow, steeply shelving shelf, which is tectonically active, whilst the ephemeral rivers (*fiumare*) transport a high sediment load from the nearby mountain drainages (for Sicilian comparison see Casalbore et al., 2011). The coastal foothills are dissected by wide alluvial valleys occupied by *fiumare*, which are subject to seasonal flash-flooding during




Fig. 2. A) Digital elevation model (DEM) showing topography of San Pasquale River valley (SPRV) and submarine canyon (elevation in meters relative to sea level). Submarine canyon reaches a depth of > 300 m within 1 km of the shoreline. Outcrops of Holocene and older sediments were discovered in several steep gullies on the eastern margin of the SPRV submarine canyon. B) Bathymetry map for 6 km² inshore area between Bova Marina and Palizzi Marina. Red contour shows the approximate position of the 6000 BP shoreline (~7 m depth) based on sea-level curve in Fig. 3; estimated shoreline position does not take into account subsequent sedimentation. C) Detailed map showing location of outcrops and diver waypoints on the eastern San Pasquale valley canyon margin.

the wet season. The San Pasquale River is one such *fiumara*; like most, today it has been canalised and is seasonally dry.

2.2. Vegetation

The southern Calabrian landscape is dominated by Mediterranean species that vary with altitude. The coast is the driest zone with as little

as 600 mm of annual precipitation, whilst ca.2000 mm fall in the highlands of the Aspromonte (Le Pera and Sorriso-Valvo, 2000). The broad valleys of the larger rivers are primarily used for agriculture, including grain, vines and bergamot orchards. The higher denuded slopes of the clay terraces have little economic use today but as occasional pasture. Coastal vegetation is dominated by *macchia* or *maquis*, Mediterranean scrub, which includes many aromatics such as mints,




Fig. 3. Modeled sea-level curve for Calabria and Sicily from Lambeck et al. (2004) showing the approximate depth for the 6000 BP shoreline.

thyme and oregano, interspersed with olive and figs and other small trees such as *Quercus ilex* (holm oak) and the non-native *Opuntia ficus-indica* (prickly pear cacti). Whilst a typically dry southern Mediterranean vegetation today, it has not always been this way. Historically, the higher mountains inland have had dense conifer and hardwood forests; in 1847 the author and artist Edward Lear described the hillsides around Bova as covered in dense woodland (Lear, 2003:34). Coastal development in the last century has caused dramatic changes to the local environment, as have reclamation of marshes (*bonifica*) and aridification. The San Pasquale valley itself has a long history of land use and has been modified by the construction of agricultural terraces.

2.3. Archaeological and palaeoenvironmental context

Despite its long record of human occupation and central Mediterranean position, there has been little archaeological research focused on the prehistory of southern Calabria. As such, extensive archaeological research along the San Pasquale coast formed part of a broader study of the prehistoric landscape of the region around the *Comune di Bova* in southern Calabria by the Bova Marina Archaeological Project. Whilst evidence for the Palaeolithic and Mesolithic in the region remains sparse, a number of Neolithic (8000–5500 BP), Copper Age (5500–4400 BP) and Bronze Age (4400–3000 BP) sites have been identified and excavated since the start of the project in 1997 (Robb, 2002; Robb and Van Hove, 2003; Robb, 2004; Robb and Michelaki, 2007; Robb, 2007). As part of *Magna Grecia*, this area is better known for its Classical archaeology. The Roman site of Deri is located near the modern San Pasquale River mouth; the Classical Greek site of Mazza stands upon a hill immediately overlooking it.

In the Bova area, prehistoric materials have been found at all elevations from the coast to the mountains. Two Neolithic sites have been excavated: Penitenzeria, a Stentinello settlement (7500–7000 BP) with a short but intense period of occupation and, Umbro, a nearby rock-shelter, which revealed sporadic activity across the Neolithic from the Stentinello, Early to Middle Neolithic, to the Diana Late Neolithic phase (7700–6000 BP). The rockshelter at Umbro was re-used briefly at the end of the Copper Age (5000–4500 BP) (see Robb, 2002; Robb and Van Hove, 2003; Robb, 2004; Robb and Michelaki, 2007; Robb, 2007). A

number of Bronze Age sites are also known, with two excavated (Umbro, 3800–3500 BP), and Sant'Aniceto, 3100–2900 BP). Both excavated Neolithic sites contained local pottery, undecorated, Impressed and Stentinello ware, alongside a small sample of imported ware, polished stone axes, flint, chert and obsidian tools. This indicates knowledge of local raw material sources within the landscape (Michelaki et al., 2012; Michelaki et al., 2015) alongside participation in wider networks of trade and exchange within Calabria and across the Straits of Messina to Sicily and the Aeolian islands (Farr, 2003, 2006; Robb and Farr, 2005:28). Little archaeobotanical or faunal material was preserved. However, that which was recovered reflected a typical Neolithic economy including domesticated animals: sheep/goat, cow, pig and dog, and cereal remains including three carbonised grains of *Hordeum vulgare* (barley). No local proxy palaeoenvironmental record exists for the Calabrian coast.

In the San Pasquale valley, there is also surface evidence for Neolithic occupation, although sites may either have been lost in areas where clay subsoil has been exposed by erosion, or, covered by alluvium of up to 4 m deep in the valley bottom directly adjacent to the San Pasquale River. Surface finds include Impressed Ware pottery about 500 m from the stream, and an obsidian fragment close to the stream mouth.

Evidence of wider regional Holocene climate and palaeoecological variations are limited to pollen records captured in marine cores that only preserve a limited selection of pollen due to wind-blown bias (Antonoli et al., 2017; Di Rita et al., 2018), or inland lacustrine records (Follieri et al., 1998; Allen et al., 2000 Joannin et al., 2012). From the available data, Antonoli et al. (2017:365) suggest the Italian coast would have been dominated by *Pinus* in the highlands and deciduous *Quercus* along lower elevations on the coast. Sicilian records show a general trend toward open vegetation with Mediterranean evergreen shrubs such as *Pistacia* developing into dense maquis (Mediterranean shrub) between 7000 cal BP and 5000 cal BP (Noti et al., 2009; Tinner et al., 2009). This ties to a general shift from a cool wet climate 8000–7000 BP to warmer and drier conditions 7000–3600 BP (Le Pera and Sorriso-Valvo, 2000).

As part of the Bova Marina Project's research into the evolution of the coastal landscape, a geomorphological survey was undertaken along the coast. This included a preliminary offshore diver survey in 2005 to identify geological outcrops along the coast between San Pasquale and Bova Marina (see Fig. 2).

During the survey a dense, thick (20–30 cm) organic unit consisting of seeds and sticks, was found at –27mbsl (metres below sea level) (05WP12 BS4; Figs. 2, 3 and 5). A small stick from this deposit at –27 m was dated by AMS to 6190–5940 cal BP, (05WP12 BS4, Beta-207156, Beta Analytic, Florida, U.S.A.; Fig. 8). This date corresponds to the Final Neolithic in this region, characterised by Diana style pottery. No archaeological remains were identified during the survey, but the unusual nature of the deposit and its possible sea-level implications instigated further geological studies.

3. Materials and methods

3.1. Geophysical and diver surveys

In 2008 and 2009, marine geophysical mapping and underwater surveys using SCUBA were conducted across a 6 km² inshore area between Bova Marina and Palizzi Marina (Fig. 1B). The geophysical survey was collected using a Knudsen 320BP single-beam 200 kHz echosounder and side-scan sonar system with sonar transducers side-mounted on a 5-m inflatable boat. Survey navigation and data positioning were provided by an onboard differential-GPS. A total of 250 line-km of single-beam bathymetry data were acquired in water depths of up to 400 m and corrected for transducer draft and diurnal tidal variations. Bathymetry data were then interpolated using a minimum curvature algorithm (Sonnenburg and Boyce, 2008) with 10 m grid cells




Fig. 4. A) Side-scan mosaic showing outcrops exposed in eastern canyon wall at waypoints 08WP2 and 08WP11 (location of image shown in Fig. 2C). B) Portion of side-scan swath showing exposed strata on northwest face of 'knowle' formed by erosion of gullies into canyon margin. At waypoint 08WP11, conglomerate beds overlie Early Miocene mudstones and are draped by Holocene muds containing abundant *Posidonia* roots and fragments (see Fig. 9).

to produce a colour-shaded bathymetric model (Fig. 2A, B). Bathymetry data were combined with available digital topographic data (NASA Shuttle Radar Topography Mission, 3 s global coverage) to produce a continuous land-sea digital elevation model. Side-scan sonar swaths were collected in water depths of < 30 m and mosaiced (Fig. 4A) to

assist in interpretation of the submarine canyon geomorphology and to identify underwater outcrops for coring and sampling (Fig. 4B).

Diver underwater surveys in 2008 and 2009 identified and mapped several well-exposed stratigraphic exposures (08 waypoint numbers) along the eastern margin of the San Pasquale submarine canyon. Cores




Fig. 5. Well preserved organics from bulk sample at 05WP12. A) *Medicago polymorpha*, toothed medick, B) *Picea* cone (or *Pinus* ?), C) *Quercus* sp. acorn in its cupule, D) charcoal fragments, E) twig, F) *Cerastoderma glaucum*, an estuarine or lagoonal bivalve.

were collected in several stepped outcrops (09WP1) near the organic bed found in 2005 (05WP12) (Figs. 2, 6, 7 and 9). A total of 6 sediment cores were taken along an easterly transect in 25–27 m water depth (Fig. 6) for sediment and microfossil analysis (Figs. 7 and 11). Bulk samples of exposed organic material were also collected at several locations for palaeobotanical analysis and radiocarbon dating (BS2,3,4) (Figs. 7 & 8). All waypoints and coring sites were located using a surface deployed hand-held GPS and depths were measured using a tidally calibrated depth gauge.

3.2. Sedimentology and micropalaeontology

The cores were logged and sampled at 1 cm intervals for particle

size and microfossil analysis. Laser-particle size analysis used standard techniques (Beckman Coulter LS 230; Donato et al., 2009). Loss-on-ignition was measured following the procedure in Heiri et al. (2001).

Micropalaeontological samples (3–4 cm³) were sieved using a 45-μm screen to remove the muds then oven dried at 35 °C and further sieved at 250-μm sieve for microfossil identification. Samples were examined using a binocular dissecting microscope (×60) to identify and quantify the number of hematite-stained planktic foraminifera that originated from the Miocene mudstones within the San Pasquale River valley (specimens/cm³). Many of the specimens also showed diagenetic alteration including overgrowths, infills, and fragmentation. During the microfossil investigation, gypsum crystals were also observed and quantified.




Fig. 6. Detail of bathymetry map (see Fig. 2) showing locations of core transect at locality 09WP1 and relationship with site 05WP12. In 2005, an organic bed - containing well-preserved tree branches and plant macrofossils was exposed at 05WP12 but was subsequently buried and could not be relocated in 2009. The core transect at 09WP1 is located in the same gully but on the eastern side.

3.3. Palaeoecology

In addition to core samples, large ($> 500 \text{ cm}^3$) bulk samples (BS2, 3) were taken from organic beds exposed within the terraced outcrop profile (and also in cores) for archaeobotanical analysis and radiocarbon dating (Figs. 6–8). These samples were dominated by water-logged remains of plant macrofossils with remarkable levels of preservation. Three samples ranging from 400 to 600 mm were analysed (see Table 1 Supplementary material). To extract the plant remains, the samples were soaked in hot water in order to disaggregate the matrix and free the organic material. They were then passed through a series of geological sieves with mesh diameters of 1 mm, 0.5 mm and 0.3 mm. The separate fractions were examined under a low-powered stereomicroscope ($\times 8\text{--}\times 56$) and the plant remains picked out. The remains were identified using the author's (AC) reference collection and Cappers et al. (2006).

4. Results

4.1. Geophysical survey

Bathymetric surveys revealed a network of northwest to northeast-oriented submarine canyons that are aligned with the general trend of the San Pasquale and Sidero river valleys (Fig. 2). The San Pasquale canyon is up to 1 km in width and descends to a water depth of $> 400 \text{ m}$ within 1.5 km of the shoreline. The canyon headwall is defined by arcuate slump scars in 8–10 m water depth, about 200–250 m from the shoreline. The eastern margin of the San Pasquale canyon is incised by 10–20 m deep erosional gullies that have a northwest-southeast orientation, indicating possible structural control (Fig. 2B). The gullies expose the Miocene mudstone bedrock and overlying unconsolidated sediments (Fig. 4). Submarine canyons of similar scale and morphology have been documented on the Sicilian coast and have been attributed to retrogressive slope failure as a result of erosional scour by hyperpycnal flows generated during flash-flooding of *fiumare* (Casalbore et al.,




Fig. 7. A) Core transect at 09WP01 showing stratigraphy and lithofacies in submarine gully fill (location shown in Fig. 6). B) Shallow shelf stratigraphy as exposed in outcrops closer to the San Pasquale River, showing stratigraphic relationships with Neogene bedrock. Contact between *Posidonia* muds and underlying framework supported (fluvial?) conglomerate is interpreted as a marine flooding surface (MFS).

2011). Mass failures can also be triggered by cyclic wave loading of the seabed and ground shaking during earthquakes (Ridente et al., 2014).

Fig. 4 shows side-scan images of a portion of the northeastern margin of the San Pasquale submarine canyon. Here, slope failure and the erosion of gullies has exposed indurated Miocene mudstones and overlying conglomerates. Conglomerates are capped by up to 1–2 m of Holocene marine muds. Areas of low backscatter within gullies indicate the presence of less reflective, fine grained sediments, likely recording failure and downslope transport of Holocene muds into the submarine canyons (Fig. 4).

4.2. Stratigraphy

4.2.1. Shallow shelf

The outcrop stratigraphy in shallow shelf areas (–15 to 22 m) exposed closer to the head of the submarine canyon consisted of Miocene mudstones overlain by a conglomerate of variable thickness that is likely Quaternary in age and likely alluvial in origin, but was modified through the Holocene marine transgression (08WP2, 11) (Fig. 7). The mudstones were compact and semi-consolidated and more erosion resistant than overlying unconsolidated recent Holocene muds and sand deposits. Outcrops of the conglomerate showed some evidence of marine bioencrustation (e.g. barnacles, serpulids), but the encrustation




Fig. 8. Radiocarbon ages and OxCal calibrated age range. See Fig. 7 for stratigraphic position and calibration methods.

could have also occurred more recently with outcrop exposure. In some locations, the conglomerate unit was missing (08WP7,1), or very thin (08WP10). Overlying the conglomerate, was a *Posidonia* rooted peat, containing shell fragments, pebbles and/or interbeds of Holocene marine muddy sand. Large vertical outcrops of conglomerate (> 5 m thick) were also found in deeper locations (08WP13) with beds dipping ~15–20° from horizontal to the W-SW. The conglomerates are likely part of the Early Miocene Stilo Capo d'Orlando Formation, but could also be of Quaternary age (Cavazza and Celles, 1993, Patterson et al., 1995).

4.2.2. Gully sediment fill

The stepped outcrops at 09WP1 extended from ~25 to 27 m depth and exposed mud and sand units containing distinct organic beds (~20–30 cm) that were laterally discontinuous (Figs. 7 and 9). The beds were generally dipping ~15–20° towards the W-NW following the general trend of gullies that are eroded into the eastern margin of the San Pasquale valley submarine canyon (Fig. 2). The edges of sections showed prominent “pop-up” ridges suggesting that exposure of strata on the valley walls was due to a submarine failure and mass wasting events. The valleys appear to have formed within eroded Stilo Capo d'Orlando Fm. units, a large vertical face and promontory was found ~250 m NW of 05WP12 that had similar characteristics to that of 08WP13. At the base of the valley section a large tree trunk with a root ball was found imbedded in intact basal beds of muddy sand that had a similar depth to the organic bed found in 2005 (05WP12) (Fig. 10). The upper exposed surface of the tree was bored with *Teredo navalis*, but the borings did not look recent as the upper boring holes were eroded and exposed. The root ball had embedded cobbles and pebbles indicating it had been eroded from a river-bank/floodplain environment. Whilst we do not have a radiocarbon date for the wood, it was located at a similar stratigraphic level as the organic bed at 05WP12 (6000–6200 BP). The radiocarbon age for the upper and lower organic beds in the valley fill indicate approximately one meter of deposition over a maximum duration of 300 years but sedimentation rates may have been considerably faster as the ages of the organic beds are very similar (Fig. 8).

4.3. Core analyses

Within the cores, a number of event horizons can be seen suggesting episodic events of deposition of organic material within turbidity plumes (Figs. 7 and 11). Particle size analysis revealed sequences of coarsening sand fining upwards to layers of fine silt, indicative of turbidity currents or storm deposits (Drake, 1999; Sommerfield and Nittrouer, 1999). The organic facies are unsorted and include coarser grained sedimentary material and rounded pebbles, occasional shells, together with charcoal and plant debris capped with fine silt layers. In all the cores, except F36, smaller event horizons are also evident in

addition to the larger turbidity flows. These are indicative of minor flood events from the river mouth; they carry less debris and have fewer microfossils indicating a more localised provenance for the sediment.

The micropalaeontological analysis of the cores indicates the sediment source. Hematite stained planktic foraminifera including undistinguished globigerinids and *Orbulina* that were often fragmented (Patterson et al., 1995; Reinhardt et al., 2000) (Fig. 12). These specimens originated from weathered outcrops along the San Pasquale River valley and were transported directly from that source through flood events or through storms on to the shallow shelf. In addition, some of the tests showed gypsum overgrowths, but individual crystals and clusters were also found that may have formed in floodplain sediments or within the Miocene mudstones (Fig. 12). There was a tendency towards increased hematite stained foraminifera and gypsum content with increased grain-size and organic content. The delicate fragile nature of the gypsum and the floodplain organic matter, suggests periods of direct transport from source to sink, with little residence time on the shallow shelf.

4.4. Palaeoecology

Three samples from the organic strata were studied for plant remains. These samples were dominated by waterlogged remains of plants including seeds, leaves and other plant parts (Table 1 Supplementary material). A mosaic of habitats including woodland, aquatic/wetland, and terrestrial open and coastal species is represented.

All three samples consisted of very compacted organic material with little or no inorganic material. Radiocarbon dates were taken on material from samples 2 and 3 (6280–6010 cal BP and 6180–5900 cal BP respectively) (see Fig. 8). All samples contained identifiable woody remains of twigs and larger branches. Leaf fragments, including those of ferns (*Pteridophyte*) and mosses (*Musci*), and other plant remains such as seeds were present (see Table 1 Supplementary material). Readily identifiable remains of oak (*Quercus* sp.) acorns and their cupules were present in all three samples. Whole pine cones were present in the bulk samples, as were a few cone bracts in samples 1–3. Charcoal was recorded in all three samples and in some cases was of sufficient size to be identifiable. In addition to charcoal, some plant remains were preserved by charring, including remains of barley (*Hordeum vulgare*).

5. Discussion

5.1. Origin of terrestrial organic beds

The extraordinary preservation of unsorted organic matter, including not only twigs and branches but also the fine component of leaves, seeds and wood (Fig. 5), is unusual under normal sedimentary




Fig. 9. A) and B) Prominent stepped ('terraced') outcrop profile formed by differential erosion of interbedded sand, mud and organic layers. Prominent ridges or 'pop-ups' at the edge of the beds suggests a prior sediment failure event (failure scarp). C) Organic beds were compressed and laterally thinned along the exposure. D) Photograph showing the dense concentration of organics, including seeds and branches in 20 cm thick organic layer in core F43 (see Figs. 7 and 11 for further detail).




Fig. 10. A) Tree trunk embedded in muddy sand at the bottom of the valley fill (F48; Fig. 7). Trunk was compressed and flattened. B) The outer surface of the trunk was bored by *Teredo navalis*, but the borings were partially eroded, exposing the burrow interior, indicating they were not recent. C) The tree root ball was intact and contained cobbles and pebbles, suggesting it was likely eroded during a flood event and transported downslope by sediment gravity flows.

deposition on shelves; in most events where organic material is transported into the marine system, the fine sediments (i.e. fine silt, mud) remains in suspension and is dispersed. As such it was initially hypothesised that the organic sediments were deposited in a rapid sedimentary event such as a rapid depositional landslide – potentially related to an earthquake or tsunami event – or to meteorological or hydrological factors, such as heavy and prolonged precipitation (Kyoji and Canuti, 2009). A rapid mass wasting event, or series of events, is also suggested by the presence of preserved tree trunks and larger branches within cobble-rich deposits.

The detailed core analysis indicates rapid depositional sedimentary events of varying scale, whilst the lack of post depositional bioturbation further supports the interpretation of rapid burial. The unsorted nature of the sediment deposits suggests the organic material was deposited

within a debris flow. As the energy of the turbulent flow waned, the finer sediment settled out of suspension, rapidly capping the organic horizon and thus preventing the finer organic material from being washed away. It is the rapid burial within this anoxic environment that led to the excellent levels of organic preservation. These sedimentary sequences are likely to have been deposited by hyperpycnal flows triggered by flash-flood events.

In the offshore zone, hyperpycnal flows can carry large concentrations of silt clay and mud in addition to organic matter; they are important sediment transport mechanisms that move continental sediments out into deep water (Lamb and Mohrig, 2009; Felix et al., 2006; Mulder et al., 2003; Piper and Normark, 2009). However, the importance of these episodic events for the transportation of sediment onto the continental shelf has been relatively understudied (Khan et al.,




Fig. 11. Core analyses including particle-size, loss on ignition (LOI) and abundance of hematite stained foraminifera and gypsum clusters.

2005; Syvitski and Morehead, 1999; Traykovski et al., 2000; Wheatcroft and Borgeld, 2000) and has never been considered in relation to potential archaeological preservation. Offshore density flows, such as turbidity currents, occur when sediment laden water flows down slope (Lamb and Mohrig, 2009; Mulder and Syvitski, 1995; Mulder et al., 2003). Hyperpycnal flows are a form of density flow that occur on the shelf when there is an increase in both fluvial flow discharge and sediment load. When density of the sediment-laden water coming from the fluvial outflow exceeds the density of the target basin water, the flow will plunge, resulting in a hyperpycnal flow. Lamb et al. (2008) determined that approximately 40 g/l of suspended sediment is required to produce a large enough density contrast to produce a plunging flow. Whilst the concentrations of suspended sediment needed to generate a turbidity current can vary depending on salinity and environment (Felix et al., 2006), a significant source of sediment and outflow needs to be available to produce a hyperpycnal flow. The resulting sedimentary deposits, hyperpycnites, can be specifically linked to flash-flood events, and can reflect flood duration and flow velocity (Mulder et al., 2003).

The environment of the San Pasquale valley fits well with this interpretation. The steep sided valley and seasonal storms mean that the *fiumara* is prone to flash-flooding and increased outflow. In this region active tectonics and the potential for mass wasting events, such as landslides and coastal slumping, mean that sediment loads can be high, generating the conditions necessary for hyperpycnal flows to form. The organic horizons may have occurred rapidly as a result of one specific

event or represent a series of events more chronologically distinct. AMS radiocarbon dating suggests these events occurred at some point between 6280 and 5900 BP (Fig. 8). If these horizons represented modern sedimentary events carrying recently eroded prehistoric material from another source, it would be unlikely the organic component would be so well preserved; equally, intermixing of wash-in material is unlikely. The high level of preservation indicates that the organic material must have been rapidly transported and buried. As such we can conclude that these event horizons represent isolated flash-flood and hyperpycnal mud flows that occurred in prehistory, thus providing a window into the nature of the landscape and palaeoenvironment of the San Pasquale valley at that time.

The Mediterranean area has a climate that consists of warm wet winters and dry hot summers. Rains in the winter can be very heavy and a large amount of rain can fall in a very short time (Grove and Rackham, 2003). These deluges can saturate the ground very quickly, especially late in the rainy season when the soil is already close to water holding capacity. When this happens, the water runs off over the surface and can carry the soil and the vegetation with it down the river valley in a torrent. If incorporated into an offshore mudflow, or hyperpycnal flood, these rafts of vegetation and organic material are quickly deposited and buried in fine mud.

Recently, Casalbore et al. (2011) documented the effects of flash-flood hyperpycnal flows generating shallow water landslides at *fiumare* mouths in the western Messina Straits, documenting sediment transport patterns. Although they did not document any significant terrestrial




Fig. 12. SEM images showing: A) gypsum crystals on an *Orbulina* fragment, B) Gypsum clusters with planktic foraminifera test, C) Miocene planktic foraminifera showing oxy-hydroxide coating and interior diagenetic calcite or gypsum. D) – F) details of gypsum crystals.

organic concentrations, they did document an eroded tree offshore, similar to the tree with preserved root-ball documented in our study.

5.2. Terrestrial palaeoecology

The presence of charred material and charcoal could be interpreted as resulting from human activity in the area, or, may have been caused by natural fire such as that induced by lightning strikes. However, the charred cereal remains, in the form of a rachis fragment (sample 1) and a single grain (sample 2) of barley (*Hordeum vulgare*) suggest some local human activity. Grain remains often become carbonised during plant processing; *Hordeum vulgare* was also found on the nearby Neolithic sites of Umbro and Penitenzeria.

The presence of moss remains (leaves and stems) and fern pinnule fragments suggests that there were damp, shaded areas, and it is most likely that these remains are associated with the tree species such as oak and possibly pine (*Pinus* sp). The presence of alder (*Alnus glutinosa*), silver birch (*Betula pendula*) and hazel (*Corylus avellana*) suggests the presence of deciduous woodland, with some areas being wet enough to support alder (Clapham, 1999). It is most likely that the slopes and edges of the San Pasquale valley had areas of deciduous woodland, with some pine on the higher drier slopes. This was bordered towards the river by a wetland consisting of common reed (*Phragmites australis*), reed-mace (*Typha* sp), sedges (*Cyperus* sp), grey club-rush (*Schoenoplectus tabernaemontani*), rushes (*Juncus* sp) and spike-rush (*Eleocharis* sp). Aquatic species including fool's water-cress (*Apium nodiflorum*) and soft hornwort (*Ceratophyllum submersum*) were growing in the more open water away from the fringe of wetland plants.

The largest number of plant taxa in the samples were of open habitats (Fig. 13). This type of habitat is abundant within the Mediterranean landscape and the taxa present here could well represent a wide range of different types of open habitat including bare ground (and cultivation), such as the medicks (*Medicago* sp), fumitory (*Fumaria* sp), and pheasant's-eye (*Adonis* sp). Other open habitats may have consisted of shrubs and undershrubs such as heather (*Erica* sp), and




Fig. 13. Overall percentage of taxa per habitat type in botanical samples BS2 and BS3 (see Fig. 7) with detailed identification and quantities provided in Supplementary Material Table 1.

rosemary (*Rosmarinus* sp), other areas may have been covered with grassland with a mixture of grasses and other wild-flowers (see for example: Blamey and Grey-Wilson, 1993; Stace, 2010).

The location of this mosaic of open land is difficult to assess. It is most likely that the river in the past, as now, may well have been seasonal, or with a perennial small volume of flowing water. The river channel may have been braided with gravel and sand islands on which many of these plant taxa could grow, creating a very varied openly vegetated landscape. Coastal environments were also present and indicated by a small number of taxa in the samples, including sea beet (*Beta vulgaris*), sea holly (*Eryngium maritimum*), and radish (*Raphanus raphanistrum*) and salt marshes (*Limonium* sp).

5.3. Archaeological implications

Whilst steep coastal shelves are less likely to preserve submerged sites *in situ*, they can still preserve secondary deposits that contribute to our understanding of the local palaeoenvironment and aid reconstruction of coastal landscapes; enriching our interpretation of terrestrial archaeology. Indeed, it is the very instability of these coastlines that provides excellent conditions for preservation; sedimentary and tectonic events cause rapid erosion and deposition that can create excellent preservation environments for organic material in the marine environment. In addition, the ongoing dynamism of these landscapes mean that these archives are not simply buried in deep sediment but have the potential to be rapidly exposed on the seabed by further sedimentary or tectonic events, making them detectable and accessible. These active coastal zones, whilst not the traditional geomorphological setting for the preservation of submerged landscape archaeology or *in situ* palaeoenvironmental deposits, can hold transported sedimentary archives that are significant for understanding prehistoric coastal landscapes and land use.

In the case of the southern Italian Neolithic, but also more widely in the Mediterranean, our knowledge of the botanical environment is limited. With the exception of rare waterlogged sites, such as La Marmotta (Fugazzola Delpino et al., 1993) and a few pollen cores (Magny et al., 2011; Tinner et al., 2009; Noti et al., 2009), our knowledge of the plants surrounding Neolithic sites is restricted to preserved remains, often charred, excavated from terrestrial sites. This skews discussions of palaeoecology in the Mediterranean Neolithic to domesticated species, particularly grains, or to tree cover associated with climate reconstructions from the pollen records. Yet when attempting to understand Neolithic life, it is clear that such a limited environmental reconstruction affects our understanding of the past. To understand the local vegetation is to provide context to activity and diet, to bring the landscape to life with meals, perfumes, colour and flavour. It changes how we visualise and interpret the past. This is particularly relevant to understanding the Neolithic, where a nuanced understanding of local landscape may help to better comprehend subsistence changes, pastoralism and the introduction of agriculture.

Furthermore, the dynamic nature of these coastlines reminds us that landscapes are active, ever changing processes, rather than static backdrops for prehistoric activity or archaeological research. Destructive flash-flooding and erosion are common around the Mediterranean today; especially on the steeply shelving central Mediterranean coasts (Casalbore et al., 2011). Such coastal flooding is often blamed upon modern deforestation and overgrazing, which lead to the loss of the plant cover securing the soil, and thus to subsequent erosion. When combined with increased coastal development (Antonoli et al., 2017:345), landslides and flash-flood events become a serious geohazard for modern urban planning and Integrated Coastal Zone Management. The potential *longue durée* to these events is overlooked. Within the archaeological record, markers for specific weather events are hard to identify; the archaeological record is usually better suited to reconstructions on the scale of climate and patterns of temperature and rainfall. Therefore, contexts that help us understand re-occurring weather events, and sedimentary events that shaped the landscape, and lives of the people within it in prehistory, are of particular interest. If our goal is to understand the coastal zone and to integrate the onshore-offshore narratives, the potential of these steeply shelving, tectonically active coastal landscapes should not be overlooked.

In addition to adding to our knowledge of the general palaeoecology of the San Pasquale valley at the end of the Neolithic, these samples tantalizingly suggest a human-modified landscape. The presence of barley, oats and legumes suggests human presence close by. Meanwhile, many of the species present are typical of those colonising the margins of human-disturbed environments (i.e. spotted and toothed medicks, sainfoin, fumitories, fat hen and clover) and this may reflect land

clearance. When viewed as a whole, these samples seem to capture the beginnings of change within the San Pasquale valley, from the natural vegetation of the gallery forests along the river course, to one that is adapting to continuous disturbance, most likely, due to nearby human activity, resulting in the production of maquis and garrigue that is characteristic of the region today. It is also noted, how many of the taxa recovered within these samples represent plants that have human uses, for food (brambles, figs, capers, oats, barley, legumes, wild carrot, chicory, rosemary, mints, oregano, basil, hops), medicinal purposes: vervain, valerian, wormwood, black nightshade, soapwort and St John's Wort, or other practical purposes (flax and reeds). Indeed, this is a rich assemblage which, even without directly associated archaeological material, suggests a human-created landscape at or close to the mouth of the San Pasquale River; it gives us unique evidence complementing normal finds-based archaeological knowledge of landscapes. Such a location would have combined a perennial source of fresh water, access to a rich range of resources from multiple terrestrial, littoral and marine environments, and a potentially useful inlet for launching and landing boats. Yet the flash-flooding that carried this raft of vegetation out to sea, and rapidly buried it on the shelf, also attests the potential dangers of living in such a location, in prehistory as much as today.

It is within the Aspromonte foothills, on high ground or rocky promontories overlooking the sea, where the majority of prehistoric finds have been made (Robb et al., 2008). The foothills of the Aspromonte may have been an attractive place to live in prehistory, with fresh water springs, cooler air, raw materials and good visibility. However, the lack of archaeological evidence in the fertile river valleys and coastal corridor has been considered surprising, especially with the evidence that maritime trade and coastal contact was important (evidenced by the high concentration of Liparian obsidian and Sicilian flint in the lithic record) (Farr, 2006). If seasonal flash-flood events, and landslides were indeed features of these *fiumare* and steep shelving coastline, did prehistoric communities choose higher ground for their settlements in the past? Or, has any direct evidence of occupation been eroded? According to the Italian landslide inventory I. F. F. I. project (Guerrieri et al., 2008) extremely rapid debris flows account for 14.9% of landslides in Italy and identification of these geohazards plays an important role in urban planning and protection of cultural heritage.

From our series of short cores, it is not possible to comment on the potential frequency of offshore hyperpycnal events; more recent organic deposits may have been eroded and older events may be buried within the offshore plateau. However, flash-floods are common in the region today but are attributed to coastal development, changes in land use, aridification and variations in precipitation patterns in relation to current climate change. The existence of an offshore flood deposit predating historic land use intensification and coastal development is significant. Could flash-floods in prehistory also relate to modification of the land, increased pastoralism and land clearance? The palaeoecological data recovered from these samples suggests human activity in the valley and the introduction of human modifications to this landscape. Further coring onshore will enable flood sequences to be identified and enable the palaeo frequency of these events to be analysed. This in turn could help resolve questions surrounding the lack of preserved prehistoric sites directly on the coast.

6. Conclusions

This research is significant for a number of reasons beyond elucidating our understanding of the coastal geology and dynamics of the shelf. Due to the narrow shelf, uplift and the active tectonic nature of the coast, this region would not be targeted for maritime archaeological survey for submerged landscapes. However, the very active nature of this coastal landscape has preserved organic remains that can inform us about several interesting issues: the local palaeoenvironment, through species that are rarely preserved on land; probable cultivation and land clearance; the nature of the river valley and coastal zone in the past;

and finally, the *longue durée* of flash-flooding in the region that predates the historic records of land clearance from which it was believed to have resulted.

The preserved organic deposits enrich our understanding of the local palaeoenvironment and the nature of the landscape in the Final Neolithic for this region. Whilst the known terrestrial sites preserve very little organic material, these deposits reveal a rich mosaic of local habitats, much damper than today with a wide variety of taxa, including the local weeds, herbs and wild flowers that are rarely preserved in the terrestrial record. Additionally, these deposits provide potential evidence for land clearance and agriculture that are supported by the archaeological record on nearby sites. The longstanding archaeological survey in the area provides us with the archaeological context of what people were doing within this landscape through the Neolithic and into the Bronze Age (Robb, 2002) but little is known about the Final Neolithic and Copper Age in this region. Whilst not an archaeological site, these organic deposits have archaeological relevance for this period, suggesting ongoing activity within this local landscape, but also revealing the dynamic nature of this landscape.

Viewing the offshore palaeoenvironmental data alongside the regional archaeological narrative begins to blur the boundaries between 'maritime' and terrestrial data, and the division between coastal landscapes and submerged prehistory. In a landscape like that of southern Calabria, the coastal zone is highly dynamic; appreciating this dynamism in the past helps us to understand how people lived within their landscape in prehistory. The environment shapes people's activity, movement and perception of time (Ingold, 2000:189). Rather than determining human activity the environment can be seen to add texture (Evans, 2003:45), providing an additional dimension to the narratives we tell about the past (Thomas, 1996:91). This is particularly true in the dynamic coastal environments of southern Calabria. Therefore, to understand prehistoric activity, and the presence or absence of archaeological material in the coastal zone, it is necessary to understand the specific local coastal dynamics alongside narratives of regional and global environmental change.

This research highlights the value of shelf sedimentary archives for understanding local onshore environments and serves as a reminder that landscapes, whether terrestrial or submerged, are geomorphologically active and ever changing, rather than static backdrops for archaeological research.

Acknowledgements

This research has been funded by the Arts and Humanities Research Council (Grant AH/E00783X/1) and the McDonald Institute for Research, Cambridge (Farr and Robb) and Natural Sciences and Engineering Research Council of Canada (Boyce and Reinhardt).

We thank Professor Martin Jones for his guidance with the archaeobotanical material and Dr Rob Scaife for his support and analysis of samples for potential pollen preservation; Dr Gianna Ayala of the University of Sheffield, Dr Fabio Filianoti and the Mediterranean University of Reggio Calabria; Daniel Riordan, Dan MacKay and Jeremy Gabriel for diving assistance and logistics; the Soprintendenza di Reggio Calabria for their continued support for the Bova Marina Archaeological Project; and our colleagues on the Bova Marina Archaeological Project (Meredith Chesson, Glenn Dunaway, Lin Foxhall, Paula Lazarus, Kostalena Michelaki, and David Yoon) for their help and advice during fieldwork.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jasrep.2019.102176>.

References

- Allen, J.R.M., Watts, W.A., Huntley, B., 2000. Weichselian palynostratigraphy, palaeo-vegetation and palaeoenvironment: the record from Lago Grande di Monticchio, southern Italy. *Quat. Int.* 73–74, 91–110. [https://doi.org/10.1016/S1040-6182\(00\)00067-7](https://doi.org/10.1016/S1040-6182(00)00067-7).
- Antonoli, F., Chiocci, F.L., Anzidei, M., Capotondi, L., Casalbore, D., Magri, D., Silenzi, S., 2017. The Central Mediterranean. In: Flemming, N.C., Harff, J., Moura, D., Burgess, A., Bailey, G.N. (Eds.), *Submerged Landscapes of the European Continental Shelf: Quaternary Palaeoenvironments*. COST European Cooperation in Science and Technology. Wiley Blackwell, Oxford, pp. 341–376.
- Bailey, G., Sakellariou, D. & members of the SPLASHCOS network, 2012. SPLASHCOS: Submerged Prehistoric Archaeology and Landscapes of the Continental Shelf. Antiquity Project Gallery 86(334): <http://www.antiquity.ac.uk/projgall/sakellariou334/>.
- Bailey, G.N., Flemming, N.C., 2008. Archaeology of the continental shelf: marine resources, submerged landscapes and underwater archaeology. *Quat. Sci. Rev.* 27 (23–24), 2153–2165. <https://doi.org/10.1016/j.quascirev.2008.08.012>.
- Bailey, G.N., Harff, J., Sakellariou, D., 2017. Archaeology and palaeolandscapes of the continental shelf: an introduction. In: Bailey, G.N., Harff, J., Sakellariou, D. (Eds.), *Under the Sea: Archaeology and the Palaeolandscapes of the Continental Shelf*. Coastal Research library. Springer, Switzerland, pp. 1–20. <https://doi.org/10.1007/978-3-319-53160-1>.
- Bell, M. 2007. Prehistoric Coastal Communities: The Mesolithic of western Britain. CBA Research Report 149, Council for British Archaeology, York.
- Benjamin, J., Bonsall, C., Pickard, C., Fischer, A., 2011. (Eds.) *Submerged Prehistory*. Oxbow, Oxford.
- Blamey, M., Grey-Wilson, C., 1993. *Mediterranean Wild Flowers*. Harper Collins, London.
- Brondi, A., Cicero, A.M., Magaletti, E., Giovanardi, F., Scarpato, A., Silvestri, C., Spada, E., Casazza, G., 2003. Italian coastal typology for the European water framework directive. In: Özhan, E. (Ed.), *Proceedings of the Sixth International Conference on the Mediterranean Coastal environment*. MEDCOAST E 03, 7–11 October 2003, Ravenna, pp. 1179–1188.
- Cappers, R.T.J., Bekker, R.M., Jans, J.E.A., 2006. *Digital Seed Atlas of the Netherlands*. Barkhuis Publishing & Groningen University Library, Groningen.
- Casalbore, D., Chiocci, F.L., Scarascia Mugnozza, G., Tommasi, P., Sposato, A., 2011. Flash flood hyperpycnal flows generating shallow water landslides at fiumara mouths in Western Messina Straits (Italy). *Mar. Geophys. Res.* 32, 257–271. <https://doi.org/10.1007/s11001-011-9128-y>.
- Cavazza, W., DeCelles, P.G., 1993. Geometry of a Miocene submarine canyon and associated sedimentary facies in southeastern Calabria, southern Italy. *Geol. Soc. Am. Bull.* 105, 1297–1309.
- Clapham, A.J., 1999. *The Characterisation of two mid-Holocene submerged forests*. Unpublished PhD thesis. Liverpool John Moores University, Liverpool.
- Di Rita, F., Fletcher, W.J., Aranbarri, J., Margaritelli, G., Lirer, F., Magri, D., 2018. Holocene forest dynamics in central and western Mediterranean: periodicity, spatio-temporal patterns and climate influence. *Nature Sci. Rep.* 8, 8929. <https://doi.org/10.1038/s41598-018-27056-2>.
- Donato, S.V., Reinhardt, E.G., Boyce, J.L., Pilarczyk, J.E., Jupp, B.P., 2009. Particle-size distribution of inferred tsunami deposits in Sur Lagoon Sultanate of Oman. *Mar. Geol.* 257, 54–64.
- Drake, D.E., 1999. Temporal and spatial variability of the sediment grain-size distribution on the Eel shelf: the flood layer of 1995. *Mar. Geol.* 154, 169–182. [https://doi.org/10.1016/S0025-3227\(98\)00111-X](https://doi.org/10.1016/S0025-3227(98)00111-X).
- Erlandson, J.M., 2001. The archaeology of aquatic adaptations: paradigms for a new millennium. *J. Archaeol. Res.* 9, 287–350. <https://doi.org/10.1023/A:1013062712695>.
- Evans, J.G., 2003. *Environmental Archaeology and the Social Order*. Routledge, London and New York.
- Farr, R.H., 2003. Analisi dell'industria litica proveniente dai siti di Umbro e Penitenzeria, Bova Marina, Calabria. *Atti, XXVII Calabria 2002. Riunione Scientifica dell'Istituto Italiano di Preistoria e Protostoria* 38, 757–760.
- Farr, R.H., 2006. Seafaring as social action. *J. Maritime Archaeol.* 1, 85–99. <https://doi.org/10.1007/s11457-005-9002-7>.
- Felix, M., Peakall, J., McCaffrey, W.D., 2006. Relative importance of processes that govern the generation of particulate hyperpycnal flows. *J. Sediment. Res.* 76, 382–387. <https://doi.org/10.2110/jsr.2006.022>.
- Flemming, N.C., Harff, J., Moura, D., Burgess, A., Bailey, G.N. (Eds.), 2017. *Submerged Landscapes of the European Continental Shelf: Quaternary Palaeoenvironments*. COST European Cooperation in Science and Technology. Wiley Blackwell, Oxford. <https://doi.org/10.1002/9781118927823>.
- Follieri, M., Giardini, M., Magri, D., Sadori, L., 1998. Palynostratigraphy of the last glacial period in the volcanic region of central Italy. *Quat. Int.* 47–48, 3–20.
- Fugazzola Delpino, M.A., D'Eugenio, G., Pessina, A., 1993. La Marmotta. *Bollettino di Paleontologia Italiana* 48, 181–342.
- Grove, A.T., Rackham, O., 2003. *The Nature of Mediterranean Europe: An Ecological History*. Yale University Press, Yale and London.
- Guerrieri, L., Triglia, A., Iadanza, C., 2008. The I. F. F. I. Project (Italian landslide inventory): Methodology and Results Experts meeting on guidelines for mapping areas at risk of landslides in Europe JRC, Ispra 23-24 October 2007: https://esdac.jrc.ec.europa.eu/Library/Themes/Landslides/Meeting102007/Landslides_IFFI.pdf (accessed on 15/07/2019).
- Heiri, O., Lotter, A.F., Lemcke, G., 2001. Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *J. Paleolimnol.* 25, 101–110. <https://doi.org/10.1023/A:1008119611481>.

- Ingold, T., 2000. *The Perception of the Environment, Essays on Livelihood, Dwelling and Skill*. Routledge, Abingdon and New York.
- Joannin, S., et al., 2012. Pollen-based reconstruction of Holocene vegetation and climate in southern Italy: the case of Lago Trifoglietti. *Clim. Past* 8, 1973–1996.
- Khan, S.M., Imran, J., Bradford, S., Syvitski, J., 2005. Numerical modelling of hyperpynal plumes. *Mar. Geol.* 222–223, 193–211. <https://doi.org/10.1016/j.margeo.2005.06.025>.
- Kyoji, S., Canuti, P. (Eds.), 2009. *Landslides: Disaster Risk Reduction*. Springer, Berlin and Heidelberg. https://doi.org/10.1007/978-3-540-69970-5_30.
- Lamb, M.P., Mohrig, D., 2009. Do hyperpynal-flow deposits record river-flood dynamics? *Geology* 37 (12), 1067–1070. <https://doi.org/10.1130/G30286A.1>.
- Lamb, M.P., Myrow, P.M., Lukens, C., Houck, K., Strauss, J., 2008. Deposits from wave-influenced turbidity currents: Pennsylvanian Minturn Formation Colorado, USA. *J. Sediment. Res.* 78, 480–498. <https://doi.org/10.2110/jsr.2008.052>.
- Lambeck, K., Antonioli, F., Purcell, A., Silenzi, S., 2004. Sea level change along the Italian coast for the past 10,000 years. *Quat. Sci. Rev.* 23, 1567–1598. <https://doi.org/10.1016/j.quascirev.2004.02.009>.
- Le Pera, E., Sorriso-Valvo, M., 2000. Weathering, erosion and sediment composition in a high-gradient river, Calabria, Italy. *Earth Surf. Processes Landforms* 25 (3), 277–292. [https://doi.org/10.1002/\(SICI\)1096-9837\(200003\)25:3<3C277::AID-ESP799>3E3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1096-9837(200003)25:3<3C277::AID-ESP799>3E3.0.CO;2-Z).
- Lear, E., 2003. *Diario di un viaggio a piedi, Reggio Calabria e la sua Provincia (25 Luglio-5 Settembre 1847)*. Laruffa Editore, Reggio Calabria.
- Magny, M., Vannière, B., Calo, C., Millet, L., Leroux, A., Peyron, O., Zanchetta, G., La Mantia, T., Tinner, W., 2011. Holocene hydrological changes in south western Mediterranean as recorded by lake-level fluctuations at Lago Preola, a coastal lake in southern Sicily, Italy. *Quat. Sci. Rev.* 30, 2459–2475. <https://doi.org/10.1016/j.quascirev.2011.05.018>.
- Michelaki, K., Hancock, R.G.B., Braun, G.V., 2012. Using provenance data to assess archaeological landscapes: an example from Calabria, Italy. *J. Archaeol. Sci.* 39, 234–246. <https://doi.org/10.1016/j.jas.2011.08.034>.
- Michelaki, K., Braun, G.V., Hancock, R.G.B., 2015. Local clay sources as histories of human-landscape interactions: a ceramic taskscape perspective. *J. Archaeol. Method Theory* 22, 783–827. <https://doi.org/10.1007/s10816-014-9204-0>.
- Mulder, T., Syvitski, J., 1995. Turbidity currents generated at river mouths during exceptional discharges to the world oceans. *J. Geol.* 103, 285–299. <https://doi.org/10.1086/629747>.
- Mulder, T., Syvitski, J., Migeon, S., Faugeres, J., Savoye, B., 2003. Marine hyperpynal flows: initiation, behavior and related deposits. A review. *Mar. Petroleum Geol.* 20, 861–882. <https://doi.org/10.1016/j.marpetgeo.2003.01.003>.
- Noti, R., van Leeuwen, J.F.N., Colombaroli, D., Vescovi, E., Pasta, S.C., La Mantia, T., Tinner, W., 2009. Mid- and Late-Holocene vegetation and fire history at Biviere di Gela, a coastal lake in southern Sicily, Italy. *Vegetation History Archaeobotany* 18, 371–387. <https://doi.org/10.1007/s00334-009-0211-0>.
- Patterson, R.T., Blenkinsop, J., Cavazza, W., 1995. Planktic foraminiferal biostratigraphy and $^{87}\text{Sr}/^{86}\text{Sr}$ isotopic stratigraphy of the Oligocene-to-Pleistocene sedimentary sequence in the southeastern Calabrian microplate, southern Italy. *J. Paleontol.* 69 (1), 7–20.
- Piper, D.J.W., Normark, W.R., 2009. Processes that initiate turbidity currents and their influence on turbidites: a marine geology perspective. *J. Sediment. Res.* 79, 347–362. <https://doi.org/10.2110/jsr.2009.046>.
- Reinhardt, E.G., Patterson, R.T., Cavazza, W., Blenkinsop, J., 2000. Use of bulk-rock samples for strontium isotope analysis: implications of differential diagenesis on high precision results. *Chem. Geol.* 164, 331–343.
- Ridente, D., Martorelli, E., Bosman, A., Chiocci, F.L., 2014. High-resolution morpho-bathymetric imaging of the Messina Strait (Southern Italy). new insights on the 1908 earthquake and tsunami. *Geomorphology* 208, 149–159.
- Robb, J.E., Farr, R.H., 2005. Substances in motion: Neolithic Mediterranean “Trade”. In: Blake, E., Knapp, A.B. (Eds.), *The Archaeology of Mediterranean Prehistory*. Blackwell Studies in Global Archaeology, 6. Blackwell Publishing Ltd., Chichester, pp. 24–46.
- Robb, J.E., Chesson, M., Farr, R.H., Lazrus, P., Michelaki, K., White, C., Wolff, C., Yoon, D., 2008. Bova Marina Archaeological Project: Preliminary Report, 2008 session. University of Cambridge.
- Robb, J.E., Michelaki, K., 2007. Agency, skill, and creativity: pottery decoration in Neolithic southern Italy. In: Madsen, N., Jensen, H., Jessen, M. (Eds.), *Excavating the Mind: Cross-sections Through Culture, Cognition And Materiality*. Aarhus University Press, Aarhus.
- Robb, J.E., van Hove, D., 2003. Gardening, foraging, and herding: Neolithic land use and social territories in southern Italy. *Antiquity* 77, 241–254.
- Robb, J.E., 1997. Bova Marina Field Survey: Preliminary Report, 1997 season. University of Cambridge.
- Robb, J.E., 2002. Scavi Neolitici e di Età del Bronzo ad Umbro, Bova Marina, Reggio Calabria. Atti, XXXV Lipari 2000, Riunione Scientifica dell'Istituto Italiano di Preistoria e Protostoria, 35, 955–958.
- Robb, J.E., 2004. Il Neolitico dell'Aspromonte meridionale. Atti, XXXVII Calabria 2002, Riunione Scientifica dell'Istituto Italiano di Preistoria e Protostoria 38, 175–188.
- Robb, J.E., 2007. Scavi Neolitici a Umbro e Penitenzeria (Bova Marina, Calabria): Rapporto Preliminare. In Laganà, C. and Tinè, V. (Eds.) *La Preistoria della Calabria: Atti del 2 Convegno, Pellarò: Gruppo Archeologico Pellarrese*.
- Sommerfield, C.K., Nittrouer, C.A., 1999. Modern accumulation rates and a sediment budget for the Eel shelf: a flood-dominated depositional environment. *Mar. Geol.* 154 (1–4), 227–241. [https://doi.org/10.1016/S0025-3227\(98\)00115-7](https://doi.org/10.1016/S0025-3227(98)00115-7).
- Sonnenburg, E.P., Boyce, J.I., 2008. Data-fused digital bathymetry and side-scan sonar as a base for archaeological inventory of submerged landscapes in the Rideau Canal, Ontario, Canada. *Geoarchaeology* 23, 654–674.
- Stace, C., 2010. *New Flora of the British Isles*, third ed. Cambridge University Press, Cambridge.
- Syvitski, J.P., Morehead, M.D., 1999. Estimating river-sediment discharge to the Eel margin, northern California. *Mar. Geol.* 154, 13–24.
- Thomas, J., 1996. *Time, Culture and Identity, an Interpretative Archaeology*. Routledge, London and New York.
- Tinner, W., van Leeuwen, J.F.N., Colombaroli, D., Vescovi, E., van der Knaap, W.O., Henne, P.D., Pasta, S., D'Angelo, S., La Mantia, T., 2009. Holocene environmental and climatic changes at Gorgo Basso, a coastal lake in southern Sicily, Italy. *Quat. Sci. Rev.* 28, 1498–1510. <https://doi.org/10.1016/j.quascirev.2009.02.001>.
- Traykovski, P., Geyer, W.R., Irish, J.D., Lynch, J.F., 2000. The role of wave-induced density-driven fluid mid flows for cross-shelf transport on the Eel River continental shelf. *Cont. Shelf Res.* 20, 2113–2140.
- Wheatcroft, R.A., Borgeld, J.C., 2000. Oceanic flood deposits on the northern California Shelf: large-scale distribution and small-scale physical properties. *Cont. Shelf Res.* 20, 2163.