



# Optics Letters

## 2.9 $\mu\text{m}$ lasing from a $\text{Ho}^{3+}/\text{Pr}^{3+}$ co-doped $\text{AlF}_3$ -based glass fiber pumped by a 1150 nm laser: publisher's note

SHUNBIN WANG,<sup>1</sup> JIQUAN ZHANG,<sup>1</sup> NIANNIAN XU,<sup>1</sup> SHIJIE JIA,<sup>1</sup> GILBERTO BRAMBILLA,<sup>2</sup> AND PENGFEI WANG<sup>1,3,\*</sup>

<sup>1</sup>Key Lab of In-Fiber Integrated Optics of Ministry of Education of China, Harbin Engineering University, Harbin 150001, China

<sup>2</sup>Optoelectronics Research Centre, University of Southampton, Southampton SO17 1BJ, UK

<sup>3</sup>Key Laboratory of Optoelectronic Devices and Systems of Ministry of Education and Guangdong Province, College of Optoelectronic Engineering, Shenzhen University, Shenzhen 518060, China

\*Corresponding author: pengfei.wang@tudublin.ie

Received 25 February 2020; posted 25 February 2020 (Doc. ID 391563); published 4 March 2020

---

**This publisher's note contains corrections to Opt. Lett. 45, 1216 (2020).** © 2020 Optical Society of America

<https://doi.org/10.1364/OL.391563>

---

On the first page and on the third page of [1], “157 mW” should have been “173 mW.” The article was corrected online on 24 February 2020.

### REFERENCE

1. S. Wang, J. Zhang, N. Xu, S. Jia, G. H. Brambilla, and P. Wang, *Opt. Lett.* **45**, 1216 (2020).