

ARTICLE

Variability of physical meteorology in urban areas at different scales: implications for air quality

Denise Hertwig^a, Sue Grimmond^{*a}, Simone Kotthaus^b, Christina Vanderwel^c, Hannah Gough^a, Martial Haeffelin^b, Alan Robins^d

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Air quality in cities is influenced not only by the emissions and chemical transformations but also by the physical state of the atmosphere which varies both temporally and spatially. Increasingly, tall buildings (TB) are common features of the urban landscape, yet their impact is not well understood on urban air flow and dispersion, and their effects are not appropriately captured in parameterisation schemes. Here, hardware models of areas within two global mega-cities (London, Beijing) are used to analyse the impact of TB on flow and transport in isolated and cluster settings. Results show that TB generate strong updrafts and downdrafts that affect street-level flow fields. Velocity differences do not decay monotonically with distance from the TB, especially in the near wake region where the flow is characterised by recirculating winds and jets. Lateral distance from an isolated TB centreline is crucial, with flow strongly impacted multiple TB heights away. Evaluation of a wake-flow scheme (ADMS-Build) in the isolated TB case indicates important characteristics are not captured. There is better agreement for a slender, shorter TB than a taller non-cuboidal TB. Better prediction of flow occurs horizontally further away and vertically further from the surface. TB clusters modify plume dispersion shape downwind. Strong updrafts generated by TB cluster overlapping wakes lift pollutants out of the canopy, causing a much deeper tracer plume in the lee of the cluster and an elevated plume centreline with maximum concentrations around the TB mean height. Enhanced vertical spread of the pollutants in the near-wake of the cluster results in overall lower maximum concentrations, but higher concentrations above mean TB height. These results have important implications for interpreting observations in areas with TB. Using real world ceilometer observations in two mega-cities (Beijing, Paris), we assess the urban boundary layer diurnal seasonal variability and evaluate a mixed layer height (MLH) model with parameters derived from a third mega-city (London). The MLH model works well in central Beijing but less well in suburban Paris. The variability of the physical meteorology across different vertical scales discussed in this paper provide additional contextual background for interpreting air quality observations.

1. Introduction

Air quality in cities is impacted not only by emissions and chemical transformations but also by the physical state of the atmosphere. In this paper we focus on the latter, particularly considering the impact of tall buildings (TB) and their implications. We consider both isolated tall buildings, which are common in Europe, and clusters of tall buildings, which occur more frequently in Asia and the Americas.

Critical in our consideration is the vertical spatial structure of the urban atmosphere. We identify three scales with the following horizontal and vertical dimensions^{1–3} (Fig. 1):

- (1) micro-scale – buildings to streets impact the urban canopy layer (UCL) and the roughness sub-layer (RSL)
- (2) local-scale – neighbourhoods (areas with common repeated characteristics, e.g. low density residential, high density commercial) influence the inertial sub-layer (ISL) or constant flux layer above the RSL
- (3) meso-scale – a city impacts the urban boundary layer (UBL), the depth of which changes with both time of day and season.

Flow, turbulence and dispersion vary rapidly in three dimensions based on the positioning of individual roughness elements (RE) (e.g. buildings, trees) in the UCL-RSL⁴. The UCL extends to the heights of the RE but the vertical extent of the RSL varies with both the RE heights (mean, maximum, variability) and the spacing between them^{5–9} (Fig. 1, purple equations). The UCL, RSL and ISL combine to make the surface layer (SL) or lowest ~10% of the UBL (Fig. 1).

Observations and modelling within cities need to consider this structure of the urban atmosphere. Basic micrometeorological theory is developed for the ISL. For example, the logarithmic wind profile and the Monin-Obukhov similarity theory both apply in the ISL but not within the UCL-RSL^{2,10}. Thus, our understanding and the ability to predict the transport close to the surface (UCL+RSL) is poor. As bluff body RE (e.g. buildings) are not porous, unlike trees, they can channel flow within the UCL and influence building ventilation^{11,12}. To enhance our theoretical understanding of processes and/or to undertake numerical predictions we typically simplify the urban form to capture what are thought to be the most relevant features^{4,13–15}. For example, buildings may be treated as having smooth walls¹⁶, rather than accounting for variations caused by the presence of windows, doors, balconies (etc.) or non-cuboid shapes¹⁷. Similarly, buildings may be treated as having flat roofs that are all of the same height^{13,14,18} rather than being sloped at different angles and/or of variable heights¹⁷.

Given issues of access, observations in the real world are often undertaken in the UCL. These micro-scale measurements are challenging to interpret and generalise because small changes in sensor location can result in large changes in the flow and radiative characteristics because of the three-dimensional scaling well below the typical RE height^{2,16,19,20}.

^a Department of Meteorology, University of Reading, Reading RG6 6BB, UK

^b Institut Pierre Simon Laplace, École Polytechnique, CNRS, Institut Polytechnique de Paris, Palaiseau, France

^c Aerodynamics and Flight Mechanics Research Group, University of Southampton, Southampton SO17 1BJ, UK

^d EnFlo, Department of Mechanical Engineering Sciences, University of Surrey, Guildford GU2 7XH, UK

* Corresponding Author: C.S.Grimmond@reading.ac.uk

Please do not adjust margins

The MLH model coefficients used were derived¹⁴ from observations taken in central London where surface roughness is high¹⁰. The sensor was operated at King's College London (KCL, 51° 30' 42.408'' N, 0° 7' 0.3066'' W, 1 January 2011–2 March 2011) and Marylebone Road (MR, 51° 31' 21.108'' N, 0° 9' 16.4376'' W, 9 March 2011–31 December 2017). The differences in heights above ground-level (agl) of 32.1 m (KCL) and 4 m (MR) are accounted for.

In Paris, only the larger scale MLH analyses are undertaken, using a sensor located at SIRTa (48.713° N and 2.208° E) in a suburban area 20 km south of Paris⁴⁰.




Figure 2: Study domain with model buildings (lengths and heights in full-scale) for (a) London/Southwark and (b) Beijing/Haidian with 3D views in the horizontal plane (upper) and 2D views for vertical cross-sections along the east-west axis (lower). View from the south. Circles indicate the extents of the main model domains. Note the model scales differ in (a) and (b).

2.2 Wind-tunnel experiments of isolated tall buildings (TB)

An open-return boundary-layer wind-tunnel (test-section length: 20 m; cross-section: 3.5 m x 1.5 m) was used for flow experiments in a hardware scale model of buildings for the area in south London (Fig. 2a, section 2.1). The ~700-m extent study domain has two isolated TB with heights of $H_{TB} = H_{81} = 81$ m (hereafter: T_{81}) and $H_{134} = 134.3$ m (T_{134}) surrounded by low-rise buildings (average H : ~13.4 m) in a mixed residential-commercial neighbourhood (Fig. 2a). The hardware model is built at a scale of 1:200. T_{134} with a triangular footprint is located at the south-eastern edge of the domain (Fig. 2a) and the hexagonal T_{81} is located close to the model centre. Hence, the flow at T_{134} is also affected by the roughness transition between approach flow and model⁴¹. The depth of the approach-flow boundary layer is $\sim H_{134}$.

Point-wise velocity measurements^{42,43} obtained using a two-component laser Doppler anemometry system^{e.g.44} are analysed. In each experiment neutral atmospheric stability conditions (i.e. thermal flow effects are negligible) are used. The wind-tunnel free-stream velocity U_{ref} is approximately constant at 2 m s⁻¹ for all cases. Experimental design, measurement techniques and uncertainties are available⁴¹. Here we analyse two configurations of the model: (i) *Core* and (ii) *No Tall*. In the *Core* case the hardware model includes both the tall buildings and the low-rise building canopy (Fig. 2a). Whereas in the *No Tall* case only the low-rise canopy buildings ($H < 32$ m) are present (i.e. there are no TB).

Analysis of the wind-tunnel observations for these two conditions allows the influence of the presence of the TB to be assessed. The buildings in this case have flat roofs, smooth walls but are close to 'true' general footprint shape (i.e. not rectangular in cross-section). The street orientations and building arrangements are realistic, but other roughness elements (e.g. trees, bus shelters, lamp posts) are missing. Thus, the model can be regarded as a realistic characterisation of the actual 'chaotic' building forms found in a small area of London. The combination of large building scale (1:200), the tallest building (T_{134}) and wind tunnel dimensions allow profiles throughout the UCL but it is not possible to undertake measurements beyond the RSL of the full model.

2.3 Water-flume experiments with a cluster of tall buildings

A closed-loop water flume (test-section length: 6.25 m; width: 1.2 m; constant water depth: 0.6 m) with particle image velocimetry (PIV^{e.g.45}) and planar laser-induced fluorescence (PLIF^{e.g.46}) systems allowed simultaneous velocity and scalar concentration measurements in the Beijing-Haidian hardware model (Fig. 2b, section 2.1). A two-dimensional measurement plane is illuminated by a 100 mJ Nd:YAG pulsed laser operating at 2 Hz. Two CMOS cameras (4 MP resolution) are used for the PIV flow measurements and one 16-bit sCMOS camera (5.4 MP) for PLIF, both with a shutter speed of 1200 μ s. The free-stream velocity U_{ref} is kept at 0.45 m s⁻¹, assuring a constant water flow rate through the flume. No thermal effects are modelled.

The dispersion of a passive scalar tracer is determined using an aqueous solution of fluorescent dye (Rhodamine 6G) released 5 mm upstream of the model domain from a 3 mm diameter ground source with a continuous source strength of 500 mm³ s⁻¹. The tracer concentration at the source, C_s , was 0.3 mg l⁻¹. The PLIF calibration procedure and general aspects of measurement uncertainty of the technique are discussed⁴⁷.

In the ~580-m extent model domain (Fig. 2b) there is a cluster of 14 tall buildings (maximum/minimum heights: $H_{72} = 72$ m / $H_{48} = 48$ m) surrounded by a low-rise canopy (typical building heights between 4.8 m and 14.4 m). The domain is a characterisation of a small part of the Haidian District, Beijing. The hardware model has a more detailed representation of the building cross-section than the London case (Fig. 2b cf. Fig. 2a), but still has flat roofs. Some simplification is required because of the small scale of the models. The two scales used (1:2400 and 1:4800⁴⁸) allow analysis of both the RSL and ISL characteristics, but limited detail within the UCL (Fig. 1). The depth of the approach-flow boundary layer is ~85 mm in model scale (~203 m at a scale of 1:2400; i.e. $\sim 3H_{72}$).

To investigate the influence of the Beijing TB cluster, multiple model configurations are used: (i) low-rise canopy without the tall-building cluster (*Low*), (ii) tall-building cluster with low-rise canopy (*Tall + Low*; Fig. 2b) and tall-building cluster in isolation without the low-rise buildings (*Tall*).

2.4 ADMS-Build wake model

The ADMS-Build wake model^{31,49} parameterises the velocity-deficit in the wake of an isolated, cuboidal-shaped building through a 3D extension of the 2D Counihan⁵⁰ constant eddy-viscosity wake theory. ADMS-Build provides an analytical expression for the flow field in the momentum deficit region of the main building wake (see Hertwig et al.⁴¹ eq. 3-5 and Fig. 12).

The beginning of the main wake is determined by modelling the extent of the recirculation (cavity) zone that develops on the lee of the building (see inset Fig. 1) as a function of height, crosswind width and along-wind building length. The inflow to the model (i.e. flow upwind of the building) is based on wind tunnel flow conditions for the case without tall buildings (*No Tall*). Implementation of the model and further evaluation results are given elsewhere⁴¹.

2.5 A simple urban boundary layer height model

The average MLH is estimated in Beijing and Paris using a simple parametrisation with coefficients derived in London³⁷. The modelled MLH varies both diurnally and seasonally with differences in cloud cover. The cloud conditions considered are (Table 1): clear nights followed by cumulus clouds during the day (clearCu); same conditions in both the day and night - cloudless (clear), cumulus (Cu) or stratus (St); or when the MLH is below the cloud base height (CBH). Cloud conditions cause variations in minimum MLH (z_{min}), the morning transition (MT) growth rate (GR_{MT}) and the nocturnal decay rate (DR_{noc} ; Table 1). The MLH above ground level (agl) is approximated by calculations at six key times through the day (Table 1). These times vary with sunrise (SR) and sunset (SS) to identify the timing of minimum and maximum depth of MLH, and the beginning and end of both the morning and the evening transitions periods.

Automatic lidar and ceilometer (ALC) measurements of attenuated backscatter allow the mixed layer height to be determined from observations. Here Vaisala CL31 observations are used. A background correction⁵¹ is undertaken prior to using the CABAM (Characterising the Atmospheric Boundary layer based on ALC Measurements) algorithm³⁶. The algorithm considers precipitation, cloud cover and cloud type, all obtained from the ALC measurements.

Instruments installed in Beijing³⁹, very close to the TB cluster (Fig. 2b), and in suburban Paris⁴⁰ (section 2.1; <https://www.lmd.polytechnique.fr/sirta/mlh/>) are used to obtain the attenuated backscatter data. The APHH-Beijing³⁹ data available for analysis are from November 2016 to June 2017. Although observations occur in all four seasons, both summer and autumn are incomplete. For Paris, data are analysed for a year (2018⁵²). During the analysis, the diurnal variability of the MLH is assessed using the median and the interquartile range (IQR). The four seasons are defined by month: winter (DJF, December, January, February), spring (MAM), summer (JJA), and autumn (SON).

	All	Clear	ClearCu	Cu	St	MLH<CBH
z_{min} (m agl)	338	218	281	397	340	277
GR_{MT} (m h ⁻¹)	187	182	229	186	91	140
DR_{noc} (m h ⁻¹)	1.28	1.66	2.16	1.17	0.59	1.19




Table 1: Simple mixed layer height (MLH) model equations for height and time (lower) with parameters derived for London³⁷ based on cloud cover state (upper). For the minimum height of the MLH (z_{min}) the value is the 75th percentile, whereas the morning transition (MT) growth rate (GR_{MT}) and the nocturnal decay rate (DR_{noc}) are median values. The cloud classes are: clear nights followed by cumulus clouds during the day (clearCu); day and night - cloudless (clear), cumulus (Cu) or stratus (St); MLH is below the cloud base height (CBH).

3 Results and Discussion

3.1 Tall-building wakes and model comparison in London

Prior analyses of tall-building wakes in the London domain⁴¹ (Fig. 2a) found wind fields above the low-rise canopy to be impacted for downwind distances of at least $5H_{TB}$ of the towers. As numerical weather prediction (NWP) models are used to force dispersion and air quality models, their horizontal resolution is important. With the next generation of NWP models expected to be of the order of 100 m^{e.g.53}, the tall-building wakes will span several grid cells.

Here we use a Cartesian coordinate system to present the results. Thus, the longitudinal axis (x direction; u velocity component) is aligned with the mean forcing wind direction (i.e. along-wind). The y axis (v velocity) is the lateral (transverse) direction. The z axis (w velocity) indicates the vertical component. Time-averaged velocities (U_i) are written in uppercase and are scaled by the free-stream velocity (U_{ref}).

The logarithmic law allows the wind profile to be modelled in the ISL^{e.g.54}. Although not directly applicable within the RSL¹⁰ it is often used above the UCL in air-quality models like ADMS-Urban, SIRANE or QUIC-Plume. Analysis of measurements across the *Core* configuration (Fig. 2a) provides the ensemble mean vertical profiles (z) of the longitudinal (along-wind) velocity (U/U_{ref}) at increasing radial distances (r_i) from the TB (Fig. 3). These results show the spatial variability of the flow in the presence of the tall buildings and the extent of wake effects with increasing distance and number of measurement points included in the ensemble average. Ensemble-averaged wind profiles of the *No Tall* case (i.e. when T_{81} and T_{134} are removed) serve as a reference, as the profiles follow the logarithmic law from above the height of the roughness length plus displacement height ($z_0 + z_d$). However, for both T_{81} (Fig. 3a,c) and T_{134} (Fig. 3b,d) the *Core* profiles show noticeable deviations from the logarithmic law for all radii.

The flow profiles averaged over different radial distances only converge back to the reference state well above roof-level ($z = H_{81}$) for the slender T_{81} tower near the model centre. The taller and wider T_{134} building has a much broader wake⁴¹, with lateral fanning enhanced by the shape and orientation of the tower relative to the south-easterly forcing wind direction (Fig. 3d). The differences from the *No Tall* reference are enhanced (cf. T_{81}) and remain large even when averaging over all available *Core* sites (note: no measurements

are available above a height of H_{134}).




Figure 3: Profiles of mean longitudinal (along-wind) velocity (U/U_{ref}) for the *Core* configuration of the London wind-tunnel model, which are ensemble averages for increasing distances from the tall building (radii r_1 to r_8 are 0.5 times tower height (H_{TB}) increments, colour-coded) plus the domain average across all sites for both *Core* and *No Tall* configurations for buildings (a) T_{81} (wind direction $\theta = 0^\circ$) and (b) T_{134} ($\theta = 101^\circ$). Logarithmic wind profile (log-law fit) using *No Tall* data with morphometrically^{8,41} derived roughness length (z_0) and displacement height (z_d) shown. Measurement locations (dots) and radii (arcs) for: (c) T_{81} and (d) T_{134} , with the direct downwind transect in the building wakes (dashed lines) indicated.

To explore further the impact of the T_{134} wake (Fig. 3d) velocity differences (*Core* - *No Tall*) are analysed for both the mean along-wind velocity ($\Delta U/U_{ref}$; Fig. 4a,c,e) and the mean vertical velocity ($\Delta W/U_{ref}$; Fig. 4b,d,f) without the spatial averaging. The velocity differences are shown as a function of their straight-line distance from the centre of T_{134} (d/H_{134}), with the lateral displacement of the sites ($|y - y_{134}|/H_{134}$) from the building centreline indicated, at three heights: just above the low-rise canopy ($z/H_{134} = 0.27$; Fig. 4a,b), at close to half T_{134} ($z/H_{134} = 0.48$; Fig. 4c,d) and near roof-level of T_{134} ($z/H_{134} = 0.91$; Fig. 4e,f).

The strongest impact on the flow is measured directly behind the building (\diamond), with a velocity deficit ΔU of up to ~ 1.2 times the reference velocity U_{ref} (Fig. 4c) and residual updrafts ($\Delta W/U_{ref} > 0$) of up to $\sim 0.28 U_{ref}$ (Fig. 4b) and downdrafts ($\Delta W/U_{ref} < 0$) of $\sim 0.22 U_{ref}$ (Fig. 4f). The changes in the mean vertical flow over large distances are crucial, as in dispersion modelling it is typically assumed that on average $W = 0 \text{ m s}^{-1}$, i.e. the vertical pollutant exchange is governed by turbulent motions and not by the mean flow e.g. ²⁸. This is clearly not the case in the presence of TB, which generate strong updrafts and downdrafts that affect street-level flow fields e.g. ³⁵. The velocity differences (Fig. 4) do not decay monotonically with distance from the TB, especially in the near wake region where the flow is characterised by recirculating winds. Instead, the lateral distance of the measurement points from the building centreline is crucial. Despite being several building heights

away (i.e. several 100 m), locations with small $|y - y_{134}|/H_{134}$ are strongly impacted by the TB.

Some urban air-quality models (e.g. ADMS–Urban) have flow modules to account for the impact of isolated buildings on pollutant dispersion. Wake models (e.g. ADMS–Build³) represent flow around cuboidal buildings of low height to length (or width) ratios, without accounting for the impacts of neighbouring low-rise buildings. We evaluate ADMS–Build (section 2.4) at the London site using our measurements in the main wake (i.e. behind the recirculation zone of the near wake, starting $\sim 51 \text{ m}$ downwind of T_{81} ; $\sim 161 \text{ m}$ downwind of T_{134} ⁴¹; inset Fig. 1). Data are analysed for all available measurement heights, stratified by straight-line distance (d/H_{TB}) between the tall building and the comparison sites (Fig. 5).




Figure 4: Difference between *Core* and *No Tall* cases of the London model for the mean (a,c,e) longitudinal velocity ($\Delta U/U_{ref}$) and (b,d,f) vertical velocity ($\Delta W/U_{ref}$) as a function of straight-line distance (d/H_{134}) from the centre of building T_{134} (south-east approach flow) at three heights z/H_{134} (a,b) 0.27, (c,d) 0.48 and (e,f) 0.91. Lateral displacement of the measurement points ($|y - y_{134}|/H_{134}$, colour) from centreline of T_{134} (dashed line, Fig. 2d) indicated: along the T_{134} centreline (\diamond), off the centreline (\circ) and affected by the wake of building T_{81} (Δ).

Overall, the agreement between ADMS–Build and wind tunnel is better for the slender, shorter T_{81} (Fig. 5a) than for the larger and wider T_{134} (Fig. 5b). The percentage of locations where the model and measurement agree within a tolerance threshold of $\pm 0.1 U_{ref}$ for either T_{81} or T_{134} clearly depends on the respective distance from the tower. The flow at sites closer to the towers is less well predicted (cf. sites at greater distances). Model performance improves with height above the urban canopy (i.e. larger U/U_{ref} conditions). This is partially explained by neglect of roughness effects from the low-rise buildings surrounding the towers within ADMS–Build. For some sites this causes an over-prediction of velocities closer to the ground. The wake-flow close to T_{134} (located at the edge of the wind-tunnel model; Fig. 2a) is also affected by the roughness transition between the upwind boundary-layer and the building model, which is not accounted for in ADMS–Build⁴¹. Such changes in roughness characteristics (e.g. induced by the local building morphology of a neighbourhood) occur frequently in mega-cities (e.g. central business districts, residential quarters, urban parks).




Figure 5: Mean longitudinal (along-wind) velocity (U/U_{ref}) in the London study area from the ADMS–Build wake model versus wind tunnel (WT) for profiles in the main wake behind (a) building T_{81} (wind direction $\theta = 0^\circ$) and (b) T_{134} ($\theta = 101^\circ$) with straight-line distance (d/H_{TB}) from the centre of the towers (colour). Inset maps show the (x, y) locations of the comparison sites in the main building wake; each site has data pairs at several heights. Ideal 1:1 (solid), $\pm 0.1 U/U_{ref}$ threshold (dotted) and linear regression fit (dashed). Percentage of the ADMS–Build data within $\pm 0.1 U/U_{ref}$ of the WT measurements are given for each distance range.

3.2 Beijing: Plume characteristics in a cluster of tall buildings

The effect of tall buildings on the dispersion of pollutants continuously emitted from an upwind ground source is analysed in the TB cluster in Beijing (Fig. 2b and 6). For reference, we compare the pollutant dispersion to an open area (i.e. without buildings, Fig. 6a) to identify changes in mean plume characteristics for dispersion through a low-rise building canopy (*Low* configuration; Fig. 6b) and through a high-rise building cluster surrounded by the low-rise buildings (*Tall + Low* (Fig. 2b); Fig. 6c). The time-averaged concentrations (C) are scaled by the concentration of the tracer solution at the release point (C_s). The elevation of the plume centreline is used to analyse the impact of the TB cluster. This is measured by the height of the maximum tracer concentration (C_{max}/C_s) along the longitudinal direction. The vertical plume half-width is the distance over which C_{max}/C_s has dropped to half its local value vertically.




Figure 6: Contours of scaled concentrations ($\log_{10}(C/C_s)$) in the (x, z) plane in the surface layer for areas (a) without buildings, (b) with the *Low* configuration without the tall-building cluster and (c) *Tall + Low* (i.e. tall building cluster with the surrounding low-rise buildings in the Beijing neighbourhood). For (b,c) the forcing wind direction is from the north ($\theta = 0^\circ$; arrows in insets of building geometries). The heights of the maximum concentration (C_{max}/C_s ; solid lines) and of $0.5 C_{max}/C_s$ (dashed lines) are indicated. In (c) grey lines indicate when values are less reliable as tall buildings block the camera view. All spatial dimensions are scaled by the height of the tallest building in the cluster ($H_{72} = 72$ m). The source is located at $x/H_{72} = 0$.

Compared to the open-reference, the elevation of the plume centreline is noticeably enhanced in the presence of the *low-rise* canopy (Fig. 6a,b). An initial upward momentum of pollutants when released (i.e. at the source) is observed (Fig. 6a) causing C_{max} to be initially away from the ground. However, in the presence of the building canopy, this is a secondary effect on the plume dispersion. The increased vertical transport and mixing of pollutants in the building canopy contribute to a faster decay of concentrations with increasing distance from the release point (cf. open reference). The along-wind decay is further enhanced by a stronger lateral plume spread from pollutant channelling and lateral plume branching in response to the local street topology (not measured in this experiment).

The vertical plume half-width is noticeably increased at the far end of the measurement domain (~ 8.5 building heights away from the source) in the *Low* configuration (cf. to the open case). Including the *Tall*-building cluster in the centre of the model domain leads to strong changes in the plume shape downwind of $x/H_{72} = 2.5$ (Fig. 6c). The strong updrafts generated in the overlapping wakes of the tall buildings lift pollutants out of the canopy, resulting in a much deeper tracer plume on the leeward side of the cluster (larger vertical plume half-width) and an elevated plume centreline (maximum concentrations just below $z = H_{72}$). Note that the first building impacted by the plume is located at $x/H_{72} = 3.25$ (i.e. the TB at $x/H_{72} = 2.75$ is in front of the measurement plane (camera perspective)).

The height of the plume envelope increases longitudinally (Fig. 6c; increasing from $z/H_{72} = 1$ to 2 over $x/H_{72} = 3.25$ to 8.5) and correlates with the region of maximum wind shear (and maximum velocity variances; not shown). The enhanced mixing and vertical spread of the pollutants in the wake of the cluster results in overall lower maximum concentrations. Furthermore, on the windward face of the TB below the region of strong upward flow at roof-level, there is a region of enhanced downward motion (downwash; Fig. 1) and outflow near the building's base resulting in a further lateral distribution of pollutants. A large portion of the pollutant plume behind the TB cluster is transported above the street canyons of the low-rise buildings. Hence, higher concentrations are observed above $z/H_{72} = 1$ compared to the open-terrain and *Low* cases at the same distances from the source.

To study the wake region behind the cluster over larger distances, measurements in a 1:4800 scale model of the *Tall* configuration (no low-rise buildings) are analysed. Vertical profiles of mean concentrations normalised by the local maximum concentration at various up- and down-wind distances to the tall building cluster show the change in plume shape (Fig. 7). While the region upwind of the buildings has maximum concentrations close to the ground and a rapid decay of concentrations with height, in the wake region the pollutants are distributed over a much larger vertical depth, with C_{max} 's elevation growing with distance from the buildings.

We compare the measured concentration profiles with fits for a reflected Gaussian plume model⁵⁵:

$$C(z) \propto \exp\left(-\frac{(z-z_s)^2}{2\sigma_z^2}\right) + \exp\left(-\frac{(z+z_s)^2}{2\sigma_z^2}\right),$$

where z_s is the height of the source and σ_z characterises the vertical plume spread (fitted).

As the pollutants are released from a ground source (at $x/H_{72} = 0$), upwind of the buildings (i.e. over open terrain) the pollutant dispersion is modelled by setting $z_s = 0$ m. While the measured upwind profiles decay less rapidly with height, overall their shape agrees with the Gaussian model (Fig. 7). After passing through the TB cluster, the plume shape in the wake is well approximated if a $z_s > 0$ m is used i.e. concentration profiles in the wake of the tall buildings resemble those obtained as if an elevated source release had occurred. Here, the "fictitious" source height z_s/H_{72} varies between 0.85 at $x/H_{72} = 11$ and 1.06 at $x/H_{72} = 16$ (Fig. 7).

Model development work supported by experimental evidence is needed for current (operational) air-quality and dispersion models to be able to capture such effects and make the models suitable for applications in cities with TB.




Figure 7: Vertical profiles of normalised concentrations (C/C_{max}) for the *Tall* case of the Beijing neighbourhood ($\theta = 0^\circ$; arrow in inset of building geometry; scale of 1:4800) showing measurements (circles) and Gaussian plume model fits (dashed lines) at two locations upwind of the building cluster (blue, cyan; modelled for a ground source release) and at three locations in the wake region (green, yellow, red; modelled for a release from an elevated source). The x/H_{72} locations of the profiles are indicated in the inset showing the concentration plume ($\log_{10}(C/C_s)$; contours as in Fig. 6). For clarity profiles show only every third measurement point and measurements closest to the ground (with larger uncertainties) are removed.

3.3 Mixed layer height (MLH) in Beijing and Paris

Here, the MLH is derived from observations of attenuated backscatter from aerosols and moisture. Hence, it is indicative of the depth of the atmospheric layer that the aerosols have mixed into, rather than the depth of the atmospheric layer that is undergoing active vertical motion (referred to as the mixing height)⁵⁶. Layers of aerosols create gradients in the observed attenuated backscatter (here 10 m vertical and 15 s temporal resolution) that are used to identify the MLH⁵⁶.

The observed diurnal maxima of MLH during the study years (section 2.5) in Beijing and Paris are for both cities larger in spring and summer than in the autumn and winter (black lines, Figure 8). With the warmer season peak daytime medians (thick lines) being similar to the 75th percentile (thin upper lines) in autumn in both cities.

The simple MLH model (Table 1) applied in the two mega-cities uses parameters derived from six years of observations undertaken in central London³⁷. Synoptic conditions generally differ between the three cities, with a clear oceanic influence in London, continental climates in Paris and a mixture of orographic and maritime effects dominating Beijing³⁹. MLH growth is driven by the strength of vertical motions and hence the turbulent sensible heat flux (Q_H) relative to the synoptic conditions modifying the. The model accounts for differences in daylength (section 2.5) as well as cloud cover and cloud type but does not consider energy forcing differences associated with geographic location, surface cover, or human activities. As it effectively only has linear variations between the six key times within a day (Table 1, Fig. 8), the model performance should focus on these periods rather than other times.

On average the model (with London coefficients, Table 1) works surprisingly well in Beijing (Figure 8, row 1) where parameterised

diurnal patterns mostly fall within the inter-quartile range of the observations. For the London – Paris comparison again (Figure 8, row 2), the curves reveal a MLH overprediction in winter and underprediction in summer. To evaluate the performance we compare the peak daytime values modelled with the corresponding observed MLH values (data point closest in time). Both the mean bias error (MBE) and the mean absolute error (MAE) are calculated. The overall MBE and MAE are greater for Paris (-230 m/431 m) than for Beijing (-136 m/306 m). The MBE by season indicates that peak values for suburban Paris are overestimated in winter by about 100 m and underestimated in summer by about 440 m. However, as the modelled MLH peak is dependent on the growth from the minimum it is important to assess this. The modelled z_{\min} (Table 1) in Beijing and in Paris are generally reasonable with the MBE and MAE below ± 40 m and 100 m for Beijing and below ± 80 m and 150 m for Paris across all seasons. However, for Paris in Winter, modelled values clearly overestimate the observations (MBE = 142 m)(Fig. 8). This suggests the growth rates derived from London characteristics are not large enough to capture MLH variability in Paris. Previous estimates of GR_{MT} at SIRTa from ALC data were stratified by season (cf. cloud, Table 1). These varied from (mean \pm standard deviation)⁵⁷: DJF: 149 ± 119 , SON: 196 ± 137 , MAM: 225 ± 138 , JJA: 247 ± 142 m h⁻¹. These values are consistent with Table 1 but support the expectation of a larger peak MLH in Paris.

Given the lower latitudes (section 2.1) of both cities (cf. London) the peak solar noon radiation on a clear day should be greater than in London. This could contribute to the MLH increment at the continental, suburban European site. The simple MLH parameterisation accounts for differences in cloud cover and cloud type, so that the greater probability of clear-sky days at the continental Paris site⁵⁸ does not explain the contrasts to the London conditions. It is likely the continental synoptic setting of Paris (e.g. absence of sea breeze effects) may explain some of the larger growth rates and resulting greater MLH at this site during spring and summer. A strong relation between peak MLH and peak incoming solar radiation has been found in Paris⁵⁷. Given clear skies in London more frequently in winter and clear skies in summer are not well sampled the synoptic differences are not fully captured by cloud-type.

Potential increase in solar radiation might have less impact in the Asian mega-city given increased aerosols will reduce the shortwave receipt at the surface^{39,59,60}. Further, energy partitioning may restrict the growth of the mixed layer. In central London, the surface energy balance is dominated by Q_H ^{61–63}. In Beijing, Q_H is much greater in the summer at the IAP site than it is in suburban Meiyun⁶⁴. These two Beijing sites have very similar net all wave radiation (Q^*) but there are differences in outgoing longwave radiation characteristics, with the urban area having greater nocturnal emissions⁶⁴. The suburban site has larger latent heat fluxes (Q_E) making the Bowen ratio ($\beta = Q_H/Q_E$) smaller. However given the land cover characteristics, the urban IAP site has smaller summertime β than expected⁶⁴. This is thought to be associated with extensive external water use for irrigation of vegetation and for wetting roads to reduce airborne particles and cleaning⁶⁴. Thus, the summertime β are much greater in central London than in urban Beijing which could mean even if more energy were available from radiation, the MLH growth is still restricted.

In both Beijing and London, the MLH observations were taken near the centre of the mega-cities, whereas in Paris the site is in a suburban area some distance from the centre (section 2.1). Urban density greatly affects anthropogenic heat flux (Q_F) emissions. This additional source of energy (i.e. beyond Q^*) is typically larger in dense central business districts than lower density residential areas

e.g. 65–68. In colder climates the Q_F emissions are often greater in the winter than the summer^{66,69–71} because of building heating needs. Thus, winter Q_F emissions provide a proportionally larger additional source of energy relative to Q^* than in summer, given the shortwave radiation differences e.g. 40. As Q_F is always a source of energy, unlike nocturnal Q^* , it can have a large impact on the wintertime MLH. At night, this additional source of energy combined with storage heat flux (ΔQ_S) release and outgoing longwave radiation can help to maintain a source of Q_H , hence maintaining the MLH at a greater depth than it would do otherwise. Obviously, this is important as both winter and nocturnal periods are times when the MLH is lower (Fig. 8), hence any increase in boundary-layer volume will have proportionally larger impacts on potential dilution/concentration of air pollutants and hence on local air quality. From machine learning analyses of the meteorological drivers of PM₁ variability at the Paris site, it is found the impact of MLH up to about 900 m is particularly pronounced⁷².

During winter, empirical coefficients (Table 1) derived from the central London observations appear to capture the MLH evolution at the central Beijing site well, while suburban Paris (Fig. 7) conditions are overestimated. This is likely explained by the lower Q_F and ΔQ_S at the suburban location providing less energy to maintain Q_H and thereby a considerable MLH. MLH at SIRTa has been observed to be lower than in central Paris during winter and at night⁷³.

Overall, these results suggest the form of the model is reasonable and further analyses of both ALC MLH data and surface energy balance partitioning with consideration of synoptic would allow a simple, extremely fast, first order model to be developed with broader applicability.




Figure 8: Diurnal variations of observed (obs) and modelled mixed layer height (MLH) by season in (row 1) Beijing and (row 2) suburban Paris showing median (thick line) and interquartile range (thin lines). Modelled MLH (colour) are stratified by cloud conditions (colour, Table 1). The number of days (N) with data in each season varies. Observations (black) are Vaisala CL31 ceilometer attenuated backscatter data analysed with the CABAM algorithm (section 2.5). Seasons are winter: DJF, spring: MAM, summer: JJA, autumn: SON. note: y-axes differ between rows)

4 Conclusions

Physical meteorology as well as chemical emissions and transformations need to be understood to interpret air quality changes and variations across mega-cities. Given physical meteorological processes vary at micro, local and city scales (Fig. 1), it is important to be aware of these scales and their impact on atmospheric transport. Depending on the height of observations, the physical and chemical characteristics will be influenced more/less by individual roughness elements. Given

the complexity of the urban canopy layer (UCL, Fig. 1), observations taken within this region are very dependent on the exact measurement site. This creates a challenge given that outdoor human exposure is most frequently at an order of 1–2 m agl. In mega-cities, there are increasing numbers of tall buildings (TB) in isolated and cluster settings. These expand the depth of both the UCL and the roughness sub-layer (RSL, Fig. 1), where atmospheric motions are more chaotic; and increase the inhomogeneity of the urban atmosphere complicating the situation further.

Models that are rapid enough to provide air quality forecasts parameterise the urban physical meteorology. To date these parameterisations have not considered TB adequately. Using hardware scale models that characterize real areas in two global mega-cities (London, Beijing) we explore the impact of isolated TB and a cluster of TB on flow and dispersion. The studies consider the UCL, RSL and ISL. Using real world observations in two global mega-cities (Beijing, Paris) we assess the UBL height modelled with parameters derived from a third mega-city (London).

Isolated TB wakes impact the flow above the background low-rise buildings for at least $5H_{TB}$, with stronger impacts in regions close to the centre-line of the wake. Wind profiles clearly still deviate from the logarithmic law at $4H_{TB}$ radial distances from the TB. Flow profiles only converge back to the reference state well above roof-level over the horizontal distances analysed in this study. A taller and wider TB, or a group of such buildings, has a much broader wake, varying with building shape and wind direction.

Directly behind the TB the strongest impact on the mean flow is evident, based on the magnitude of the velocity deficit and residual up- and down-drafts. As dispersion models typically assume $W = 0$ m s⁻¹ with vertical pollutant exchange governed by turbulent motions and not by the mean flow, changes in mean vertical flow over distances of interest are crucial.

Velocity differences do not decay monotonically with distance from the TB, especially in the near wake region where the flow is characterised by recirculation. Lateral distance from the TB centreline is crucial as strong effects persist close to the TB centreline even several H_{TB} away (i.e. several 100 m).

Evaluation of ADMS-Build in the main TB wake considering straight-line distance (d/H_{TB}) found better agreement for a slender, shorter TB (cf. larger TB). Prediction of flow horizontally, is better further away (cf. nearer); whereas the prediction is better higher above the background low-rise buildings while overprediction occurs closer to them given they are unaccounted for by the model.

A **TB cluster** causes strong changes in the plume dispersion shape downwind of the source. Strong updrafts generated in the overlapping wakes of the TB lift pollutants out of the canopy, resulting in a much deeper tracer plume in the lee of the cluster (larger vertical plume half-width) and an elevated plume centreline (maximum concentrations just below H_{TB}). Enhanced mixing and vertical spread of the pollutants in the wake of the cluster results in overall lower maximum concentrations for a surface source, but higher concentrations above H_{TB} (cf. *open terrain* and *low-rise*) at the same distance from the source.

A *low-rise* canopy (cf. *open terrain*) increases the vertical transport and mixing of pollutants in the building canopy. This contributes to a faster decay of concentrations with increasing distance from the release point. The along-wind decay is further enhanced by a stronger lateral plume spread from pollutant channelling and lateral plume branching in response to the local street topology.

The building cluster causes vertical both displacement of the maximum concentration and wider spread of pollutants. This vertical distribution of pollutants in the wake of the TB can be approximated using a reflected Gaussian plume model for a release from an elevated source (“fictitious” source height is $\sim H_{TB}$ but growing with distance from the buildings).

Observed diurnal maxima of **mixed layer height** (MLH) are greater in spring and summer (cf. autumn and winter), with warm seasons medians daytime peaks being similar to the 75th percentiles of the cooler seasons.

A simple MLH model (Table 1) with central London empirical coefficients works surprisingly well in central Beijing, but less well in suburban Paris where values are often over/under-predicted in winter/summer. Synoptic conditions, land cover characteristics, and human activities not accounted for by this simple parameterisation likely explain the discrepancies. Still, the results suggest the form of the model is reasonable offering a simple, extremely fast, first order MLH calculation method.

Our results indicate a need for more observations at each of the scales. As both the wind tunnel and water flume results are for neutral stability conditions, the influence of TB (isolated and clusters) under non-neutral condition needs to be studied. At the microscale, large differences in surface temperatures occur^{16,17}. The variability of the urban form (e.g. TB height, number, low-rise building characteristics) and a wider range of upwind fetch conditions need to be considered.

Using the hardware model results, model development work can be used to improve current (operational) air-quality and dispersion models to enable them to capture the effects found here, thus making the models more suitable for applications in cities with TB. Future work should combine thermal influences with TB observations (hardware, real world) for dispersion models so that there is improved capability to model vertical variability (wind, temperature, humidity) between buildings e.g.⁷⁴.

The simple MLH model performance suggests further analyses of observations of both MLH data and surface energy balance partitioning across cities, with different synoptic and regional settings, would be beneficial. Beyond improving our understanding of the variability of the MLH within cities it would increase the range of data for establishing empirical coefficients. Thus, it is useful to consider these results in light of surface energy balance characteristics. Comparison of observed Bowen ratio (β) between central London^{61,62} and suburban residential areas^{75,76}, and similarly in Beijing⁶⁴, indicates the latter is smaller. There is observational evidence that urban-rural contrasts in partitioning explain spatial boundary layer depth variations^{57,77–79} when experiencing the same synoptic forcing. The timing of synoptic events, notably clear sky conditions, clearly need to be considered to ensure the information within the empirical coefficients are appropriately disentangled.

Automatic lidar and ceilometer (ALC) observations provide much more information than MLH. For example, the high resolution attenuated backscatter gives vertical profiles of aerosol distribution⁸⁰ which when modelled is improved with enhanced atmospheric chemistry details⁸¹. Thus, there is clear benefit from conducting physical and chemical based urban meteorology research together to enhance our ability to predict the air quality exposure in mega-cities, which also benefit the delivery of other integrated urban services (IUS)^{82–84}.

Conflicts of interest: There are no conflicts to declare.

Acknowledgements SG thanks the RSC and the conference organizers for the invitation to contribute this paper.

The research reported here has been funded by NERC NE/N00700X/1, Newton Fund/Met Office CSSP-China (SG, DH, SK), EPSRC UK Fluids Network visiting scientist award, ERC urbisphere (855005), and ACTRIS-2 (SK). Facilities at University of Southampton are supported by UKRI and University of Surrey are supported by NERC. The EPSRC MAGIC project allowed use of the London hardware model. For the Beijing cluster model work we acknowledge the help of George Meachim (UR), Matthew Coburn (Southampton), Christoph Kent (UR) and Shiguang Miao (IUM, CMA). For the Paris MLH processing we thank Marc-Antoine Drouin for ongoing support. ALC measurements at SIRTa performed by Météo-France. SIRTa also is supported by CNES, Ecole Polytechnique, INSU-CNRS, and IPSL and in London are funded with grants from NERC, EPSRC, Met Office, EU7, H2020 and ERC with support from KCL and University of Reading.

References

- H. A. Cleugh and C. S. B. Grimmond, in *The Future of the World's Climate*, eds. A. Henderson – Sellers and K. Mc Guffie, Elsevier, 2012.
- T. R. Oke, G. Mills, A. Christen and J. A. Voogt, *Urban Climates*, Cambridge University Press, Cambridge, 2017.
- M. Piringer, C. S. B. Grimmond, S. M. Joffre, P. Mestayer, D. R. Middleton, M. W. Rotach, A. Baklanov, K. De Ridder, J. Ferreira, E. Guilloteau, A. Karppinen, A. Martilli, V. Masson and M. Tombrou, *Water, Air, & Soil Pollution: Focus*, 2002, **2**, 1–16.
- P. Kastner-Klein and M. W. Rotach, *Boundary-Layer Meteorology*, 2004, **111**, 55–84.
- C. S. B. Grimmond and T. R. Oke, *Journal of Applied Meteorology*, 1999, **38**, 1262–1292.
- M. R. Raupach, *Boundary-Layer Meteorology*, 1992, **60**, 375–395.
- M. Roth, *Quarterly Journal of the Royal Meteorological Society*, 2000, **126**, 941–990.
- M. Kanda, A. Inagaki, T. Miyamoto, M. Gryschka and S. Raasch, *Boundary-Layer Meteorology*, 2013, **148**, 357–377.
- J. T. Millward-Hopkins, A. S. Tomlin, L. Ma, D. Ingham and M. Pourkashanian, *Boundary-Layer Meteorology*, 2011, **141**, 443–465.
- C. W. Kent, S. Grimmond, J. Barlow, D. Gatey, S. Kotthaus, F. Lindberg and C. H. Halios, *Boundary-Layer Meteorol*, 2017, **164**, 183–213.
- H. Gough, T. Sato, C. Halios, C. S. B. Grimmond, Z. Luo, J. F. Barlow, A. Robertson, R. Hoxey and A. Quinn, *Journal of Wind Engineering and Industrial Aerodynamics*, , DOI:10.1016/j.jweia.2018.02.010.
- H. L. Gough, Z. Luo, C. H. Halios, M.-F. King, C. J. Noakes, C. S. B. Grimmond, J. F. Barlow, R. Hoxey and A. D. Quinn, *Building and Environment*, 2018, **137**, 246–256.
- V. Masson, *Boundary-Layer Meteorology*, 2000, **94**, 357–397.
- H. Kusaka, H. Kondo, Y. Kikegawa and F. Kimura, *Boundary-Layer Meteorology*, 2001, **101**, 329–358.
- M. Kanda, *Theoretical and Applied Climatology*, 2006, **84**, 23–33.
- W. Morrison, S. Kotthaus, C. S. B. Grimmond, A. Inagaki, T. Yin, J.-P. Gastellu-Etchegorry, M. Kanda and C. J. Merchant, *Remote Sensing of Environment*, 2018, **215**, 268–283.
- W. Morrison, T. Yin, N. Lauret, J. Guilleux, S. Kotthaus, J.-P. Gastellu-Etchegorry, L. Norford and S. Grimmond, *Remote Sensing of Environment*, 2020, **237**, 111524.
- I. P. Castro, Z.-T. Xie, V. Fuka, A. G. Robins, M. Carpentieri, P. Hayden, D. Hertwig and O. Coceal, *Boundary-Layer Meteorol*, 2017, **162**, 207–230.
- C. S. B. Grimmond, *Theoretical and Applied Climatology*, 2006, **84**, 3–22.
- H. P. Schmid, *Agricultural and Forest Meteorology*, 1997, **87**, 179–200.
- V. Masson, C. S. B. Grimmond and T. R. Oke, *Journal of Applied Meteorology*, 2002, **41**, 1011–1026.
- C. S. B. Grimmond, M. Blackett, M. J. Best, J.-J. Baik, S. E. Belcher, J. Beringer, S. I. Bohnenstengel, I. Calmet, F. Chen, A. Coutts, A. Dandou, K. Fortuniak, M. L. Gouvea, R. Hamdi, M. Hendry, M. Kanda, T. Kawai, Y. Kawamoto, H. Kondo, E. S. Krayenhoff, Lee, T. Loridan, A. Martilli, V. Masson, S. Miao, K. Oleson, R. Ooka, G. Pigeon, A. Porson, Y.-H. Ryu, F. Salamanca, G. J. Steeneveld, M. Tombrou, J. A. Voogt, D. T. Young and N. Zhang, *International Journal of Climatology*, 2011, **31**, 244–272.
- R. E. Britter and S. R. Hanna, *Annual Review of Fluid Mechanics*, 2003, **35**, 469–496.
- S. E. Belcher, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 2005, **363**, 2947–2968.
- C. A. McHugh, D. J. Carruthers and H. A. Edmunds, *International Journal of Environment and Pollution*, 1997, **8**, 666–674.
- R. Berkowicz, *Environmental Monitoring and Assessment*, 2000, **65**, 323–331.
- M. D. Williams, M. J. Brown, B. Singh and D. Boswell, *QUIC-PLUME Theory Guide*, Los Alamos National Laboratory, 2004.
- L. Souhac, P. Salizzoni, F.-X. Cierco and R. Perkins, *Atmospheric Environment*, 2011, **45**, 7379–7395.
- A. J. Cimorelli, S. G. Perry, A. Venkatram, J. C. Weil, R. J. Paine, R. B. Wilson, R. F. Lee, W. D. Peters and R. W. Brode, *Journal of Applied Meteorology*, 2005, **44**, 682–693.
- A. Jones, D. Thomson, M. Hort and B. Devenish, in *Air Pollution Modeling and Its Application XVII*, eds. C. Borrego and A.-L. Norman, Springer US, Boston, MA, 2007, pp. 580–589.
- A. Robins and C. McHugh, *IJEP*, 2001, **16**, 161.
- Petersen, Guerra and Bova, *air waste manag assoc*, 2017, **67**, 826–835.
- L. A. Brixey, D. K. Heist, J. Richmond-Bryant, G. E. Bowker, S. G. Perry and R. W. Wiener, *Journal of Environmental Monitoring*, 2009, **11**, 2171.
- E. Aristodemou, L. M. Boganegra, L. Mottet, D. Pavlidis, A. Constantinou, C. Pain, A. Robins and H. ApSimon, *Environmental Pollution*, 2018, **233**, 782–796.
- V. Fuka, Z.-T. Xie, I. P. Castro, P. Hayden, M. Carpentieri and A. G. Robins, *Boundary-Layer Meteorol*, 2018, **167**, 53–76.
- S. Kotthaus and C. S. B. Grimmond, *Quarterly Journal of the Royal Meteorological Society*, 2018, **144**, 1511–1524.
- S. Kotthaus and C. S. B. Grimmond, *Quarterly Journal of the Royal Meteorological Society*, 2018, **144**, 1525–1538.
- A. G. Robins and R. W. Macdonald, *Review of Flow and Dispersion in the Vicinity of Groups of Buildings*, Atmospheric Dispersion Modelling Liaison Committee (ADMLC), UK, 2001.
- Z. Shi, T. Vu, S. Kotthaus, R. M. Harrison, S. Grimmond, S. Yue, T. Zhu, J. Lee, Y. Han, M. Demuzere, R. E. Dunmore, L. Ren, D. Liu, Y. Wang, O. Wild, J. Allan, W. J. Acton, J. Barlow, B. Barratt, D. Beddows, W. J. Bloss, G. Calzolari, D. Carruthers, D. C. Carslaw, Q. Chan, L. Chatzidiakou, Y. Chen, L. Crilley, H. Coe, T. Dai, R. Doherty, F. Duan, P. Fu, B. Ge, M. Ge, D. Guan, J. F. Hamilton, K. He, M. Heal, D. Heard, C. N. Hewitt, M. Hollaway, M. Hu, D. Ji, X. Jiang, R. Jones, M. Kalberer, F. J. Kelly, L. Kramer, B. Langford, C. Lin, A. C. Lewis, J. Li, W. Li, H. Liu, J. Liu, M. Loh, K. Lu, F. Lucarelli, G. Mann, G. McFiggans, M. R. Miller, G. Mills, P. Monk, E. Nemitz, F. O'Connor, B. Ouyang, P. I. Palmer, C. Percival, O. Popoola, C.

- Reeves, A. R. Rickard, L. Shao, G. Shi, D. Spracklen, D. Stevenson, Y. Sun, Z. Sun, S. Tao, S. Tong, Q. Wang, W. Wang, X. Wang, X. Wang, Z. Wang, L. Wei, L. Whalley, X. Wu, Z. Wu, P. Xie, F. Yang, Q. Zhang, Y. Zhang, Y. Zhang and M. Zheng, *Atmos. Chem. Phys.*, 2019, **19**, 7519–7546.
- 40 M. Haefelin, L. Barthès, O. Bock, C. Boitel, S. Bony, D. Bouniol, H. Chepfer, M. Chiriaco, J. Cuesta, J. Delanoë, P. Drobinski, J.-L. Dufresne, C. Flamant, M. Grall, A. Hodzic, F. Hourdin, F. Lapouge, Y. Lemaître, A. Mathieu, Y. Morille, C. Naud, V. Noël, W. O'Hirok, J. Pelon, C. Pietras, A. Protat, B. Romand, G. Scialom and R. Vautard, *Ann. Geophys.*, 2005, **23**, 253–275.
- 41 D. Hertwig, H. L. Gough, S. Grimmond, J. F. Barlow, C. W. Kent, W. E. Lin, A. G. Robins and P. Hayden, *Boundary-Layer Meteorol.*, 2019, **172**, 239–270.
- 42 D. Hertwig, H. L. Gough, S. Grimmond, J. F. Barlow and A. G. Robins, <https://zenodo.org/record/2563700>, 2019.
- 43 D. Hertwig, H. L. Gough, S. Grimmond, J. F. Barlow and A. G. Robins, [https://zenodo.org/ to update](https://zenodo.org/record/2563700), 2020.
- 44 C. Tropea, *Meas. Sci. Technol.*, 1995, **6**, 605–619.
- 45 J. Westerweel, G. E. Elsinga and R. J. Adrian, *Annu. Rev. Fluid Mech.*, 2013, **45**, 409–436.
- 46 J. P. Crimaldi, *Exp Fluids*, 2008, **44**, 851–863.
- 47 C. Vanderwel and S. Tavoularis, *Exp Fluids*, 2014, **55**, 1801.
- 48 D. Hertwig, H. L. Gough, S. Grimmond and C. Vanderwel, <https://zenodo.org/record/3768763>, 2020.
- 49 A. G. Robins, D. D. Apsely and CERC, *Modelling building effects in ADMS*, CERC, 2020.
- 50 J. Counihan, J. C. R. Hunt and P. S. Jackson, *J. Fluid Mech.*, 1974, **64**, 529–564.
- 51 S. Kotthaus, E. O'Connor, C. Munkel, C. Charlton-Perez, M. Haefelin, A. M. Gabey and C. S. B. Grimmond, *Atmospheric Measurement Techniques*, 2016, **9**, 3769–3791.
- 52 S. Kotthaus, M. Haefelin, M.-A. Drouin, J.-C. Dupont, C. S. B. Grimmond, A. Haefele, M. Hervo, Y. Poltera and M. Wiegner, *in preparation*.
- 53 H. W. Lean, J. F. Barlow and C. H. Halios, *Q.J.R. Meteorol. Soc.*, 2019, **145**, 1674–1689.
- 54 C. W. Kent, C. S. B. Grimmond, D. Gatey and J. F. Barlow, *Journal of Wind Engineering and Industrial Aerodynamics*, 2018, **173**, 100–111.
- 55 S. R. Hanna, R. Britter and P. Franzese, *Atmospheric Environment*, 2003, **37**, 5069–5082.
- 56 S. Kotthaus, C. H. Halios, J. F. Barlow and C. S. B. Grimmond, *Atmospheric Environment*, 2018, **190**, 401–414.
- 57 S. Pal and M. Haefelin, *J. Geophys. Res. Atmos.*, , DOI:10.1002/2015JD023268.
- 58 S. Kotthaus, S. Grimmond, J.-A. Bravo-Aranda, M. Haefelin and A. Haefele, *EMS Annual Meeting Abstracts*, 2017, **14**, EMS2017-560-1.
- 59 T. Petäjä, L. Järvi, V.-M. Kerminen, A. J. Ding, J. N. Sun, W. Nie, J. Kujansuu, A. Virkkula, X. Yang, C. B. Fu, S. Zilitinkevich and M. Kulmala, *Sci Rep.*, 2016, **6**, 18998.
- 60 T. V. Kokkonen, S. Grimmond, S. Murto, H. Liu, A.-M. Sundström and L. Järvi, *Atmos. Chem. Phys.*, 2019, **19**, 7001–7017.
- 61 S. Kotthaus and C. S. B. Grimmond, *Urban Climate*, 2014, **10**, 261–280.
- 62 S. Kotthaus and C. S. B. Grimmond, *Urban Climate*, 2014, **10**, 281–307.
- 63 B. Crawford, C. S. B. Grimmond, H. C. Ward, W. Morrison and S. Kotthaus, *Quarterly Journal of the Royal Meteorological Society*, 2017, **143**, 817–833.
- 64 J. Dou, S. Grimmond, Z. Cheng, S. Miao, D. Feng and M. Liao, *International Journal of Climatology*, 2019, **39**, 2793–2810.
- 65 M. Iamarino, S. Beevers and C. S. B. Grimmond, *International Journal of Climatology*, 2012, **32**, 1754–1767.
- 66 D. J. Sailor, *International Journal of Climatology*, 2011, **31**, 189–199.
- 67 A. M. Gabey, C. S. B. Grimmond and I. Capel-Timms, *Theoretical and Applied Climatology*, , DOI:10.1007/s00704-018-2367-y.
- 68 I. Capel-Timms, S. T. Smith, T. Sun and S. Grimmond, *Dynamic Anthropogenic activities impacting Heat emissions (DASHv1.0): Development and evaluation*, Atmospheric Sciences, 2020.
- 69 I. G. Hamilton, M. Davies, P. Steadman, A. Stone, I. Ridley and S. Evans, *Building and Environment*, 2009, **44**, 807–817.
- 70 L. Allen, F. Lindberg and C. S. B. Grimmond, *International Journal of Climatology*, 2011, **31**, 1990–2005.
- 71 F. Lindberg, C. S. B. Grimmond, N. Yogeswaran, S. Kotthaus and L. Allen, *Urban Climate*, 2013, **4**, 1–15.
- 72 R. Stirnberg, J. Cermak, S. Kotthaus, M. Haefelin, H. Andersen, J. Fuchs, M. Kim, J.-E. Petit and O. Favez, *Meteorology-driven variability of air pollution (PM1) revealed with explainable machine learning*, Aerosols/Field Measurements/Troposphere/Physics (physical properties and processes), 2020.
- 73 D. Cimini, M. Haefelin, S. Kotthaus, U. Löhnert, P. Martinet, E. O'Connor, C. Walden, M. C. Coen and J. Preissler, *Bull. of Atmos. Sci. & Technol.*, 2020, **1**, 23–42.
- 74 N. E. Theeuwes, R. J. Ronda, I. N. Harman, A. Christen and C. S. B. Grimmond, *Boundary-Layer Meteorol.*, 2019, **173**, 321–348.
- 75 H. C. Ward, J. G. Evans and C. S. B. Grimmond, *Boundary-Layer Meteorology*, 2014, **152**, 65–89.
- 76 H. C. Ward, S. Kotthaus, C. S. B. Grimmond, A. Björkegren, M. Wilkinson, W. T. J. Morrison, J. G. Evans, J. I. L. Morison and M. Iamarino, *Environmental Pollution*, , DOI:10.1016/j.envpol.2014.12.031.
- 77 J. F. Barlow, *Urban Climate*, 2014, **10**, 216–240.
- 78 N. E. Theeuwes, J. F. Barlow, A. J. Teuling, C. S. B. Grimmond and S. Kotthaus, *npj Climate and Atmospheric Science*, 2019, **2**, 15.
- 79 S. Pal, I. Xueref-Remy, L. Ammoura, P. Chazette, F. Gibert, P. Royer, E. Dieudonné, J.-C. Dupont, M. Haefelin, C. Lac, M. Lopez, Y. Morille and F. Ravetta, *Atmospheric Environment*, 2012, **63**, 261–275.
- 80 E. Warren, C. Charlton-Perez, S. Kotthaus, H. Lean, S. Ballard, E. Hopkin and S. Grimmond, *Atmospheric Environment*, 2018, **191**, 532–547.
- 81 E. Warren, C. Charlton-Perez, S. Kotthaus, F. Marengo, C. Ryder, B. Johnson, D. Green, H. Lean, S. Ballard and S. Grimmond, *Atmospheric Environment*, 2019, 117177.
- 82 A. Baklanov, C. S. B. Grimmond, D. Carlson, D. Terblanche, X. Tang, V. Bouchet, B. Lee, G. Langendijk, R. K. Kolli and A. Hovsepian, *Urban Climate*, 2018, **23**, 330–341.
- 83 S. Grimmond, V. Bouchet, L. T. Molina, A. Baklanov, J. Tan, K. H. Schlünzen, G. Mills, B. Golding, V. Masson, C. Ren, J. Voogt, S. Miao, H. Lean, B. Heusinkveld, A. Hovsepian, G. Teruggi, P. Parrish and P. Joe, *Urban Climate*, 2020, **33**, 100623.
- 84 WMO, *Guidance on Integrated Urban Hydrometeorological, Climate and Environmental Services - Volume I: Concept and Methodology*, WMO-No: 1234, World Meteorological Organization, Geneva, 2019.