

Accepted Manuscript

BJGP OPEN

Alternative approaches to managing respiratory tract infections: A survey of public perceptions

Alex Moore, Rebecca Cannings-John, Christopher C Butler, Clodna McNulty, Nick Francis

DOI: <https://doi.org/10.3399/BJGPO.2020.0124>

To access the most recent version of this article, please click the DOI URL in the line above.

Received 14 July 2020

Revised 23 October 2020

Accepted 02 November 2020

© 2020 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>). Published by BJGP Open. For editorial process and policies, see: <https://bjgpopen.org/authors/bjgp-open-editorial-process-and-policies>

When citing this article please include the DOI provided above.

Author Accepted Manuscript

This is an 'author accepted manuscript': a manuscript that has been accepted for publication in BJGP Open, but which has not yet undergone subediting, typesetting, or correction. Errors discovered and corrected during this process may materially alter the content of this manuscript, and the latest published version (the Version of Record) should be used in preference to any preceding versions

Alternative approaches to managing respiratory tract infections: A survey of public perceptions

Moore A, Cannings-John R, Butler CC, McNulty C, Francis NA

Name	Address	Qualifications
Alex Moore	Greenways, The Coombe, Betchworth, Surrey, RH3 7BT	MBBCh, BSc
Rebecca Cannings-John	Centre for Trials Research Cardiff University Neuadd Meirionnydd Heath Park Cardiff CF14 4YS	BSc, MSc, PhD
Christopher C Butler	Nuffield Department of Primary Care Health Sciences University of Oxford Radcliffe Primary Care Building Oxford OX2 6GG United Kingdom	FMedSci, MD, MBChB, BA, FRCGP
Cliona AM McNulty	Public Health England, Primary Care Unit, 4th Floor, Twyver House, Bruton Way, Gloucester, GL1 1DQ	MB, BS; FRCPathology
Nick A Francis	School of Primary Care, Population Sciences and Medical Education University of Southampton Southampton SO16 5ST	PhD, MD, PGD (Epidemiology), BA, MRCP

Abstract

Background

Respiratory tract infections (RTIs) are a common reason for people to consult in primary care and contribute to antibiotic overuse and antimicrobial resistance. Alternative approaches to supporting patients with RTIs may help, but it is important to understand public perceptions about these approaches before they are widely implemented.

Aim

To describe public perceptions regarding finger-prick testing, back-up antibiotic prescriptions (BUP), and alternatives to traditional consultations for RTIs, and identify factors associated with favouring these approaches.

Design and setting

Online national survey (HealthWise Wales) with linked primary care health record data.

Methods

Survey item response distributions were described. Associations between responses about consultation alternatives, BUP and finger-prick point of care testing, and potential explanatory variables, were explored using logistic regression.

Results

8,752 participants completed the survey between 2016 and 2018. 76.7% (3,807/4,966) and 71.2% (3,529/4,953) of respondents with valid responses were in favour of being able to consult with a pharmacist or nurse in their GP surgery, or with a community pharmacist, respectively. 92.8% (8034/8659) of respondents indicated they would be happy to have a finger-prick test to guide antibiotic prescribing, and 31.8% (2746/8646) indicated they would like to be given a BUP if their clinician thought immediate antibiotics were not required. 47.4% (2342/4944) and 42.3% (2095/4949) were in favour of having video and email consultations respectively. Characteristics associated with different response options were identified.

Conclusion

Consulting with pharmacists, using electronic communication tools, and finger-prick testing are widely acceptable approaches. BUP was described as acceptable less often and is likely to require greater information and support when used.

Word counts

Abstract: 258

Main body: 2,660

Keywords

Primary healthcare, Respiratory tract infections, Antimicrobial resistance

How this fits in

Primary care services are struggling to keep up with the demand for respiratory tract infection consultations and the pressure to prescribe unnecessary antibiotics is driving antimicrobial resistance. This study explored public views about alternative approaches to traditional consultations through an online survey. There is scope for incorporating such interventions in current practice, to reduce the strain on GPs, although it is important to ensure adequate information and support are provided.

Introduction

Respiratory tract infections (RTIs) are one of the most frequent reasons for consulting in primary care, with 27% of symptomatic people visiting a general practitioner (GP) over a 12-month period.(1) This contributes significantly to GP workload in the United Kingdom (UK), which increased by 16% from 2007 to 2014.(2)

Antimicrobial resistance (AMR) poses a threat to public health, worsens patient outcomes and increases healthcare costs.(3, 4) Antibiotic consumption drives AMR at both individual and societal levels,(5) and few new antibiotics are being developed.(6) Primary care is responsible for 72% of healthcare-related antibiotic prescriptions (7) and is therefore a key priority for tackling AMR.(8) RTIs accounted for about half of all antibiotic prescriptions in UK primary care between 2013 and 2015.(9) UK national guidance recommends that GPs do not prescribe antibiotics for most RTIs,(10) but prescribing continues to be excessive.(11, 12)

Alternatives to face-to-face GP consultations may help reduce the pressure on primary care services from RTIs, and use of antibiotics. Non-medical healthcare practitioners, such as pharmacists and nurses, are increasingly managing common infections. Some community pharmacists also provide services such as point-of-care testing (POCT) to help determine the need for GP consultations or antibiotics.(13) Telephone consultations have been used for years, video and email consultations are becoming more common,(14) especially during the COVID-19 pandemic.

Back-up prescriptions (BUPs), also known as delayed antibiotic prescriptions, reduce antibiotic prescribing and consumption for RTIs, without increasing symptom burden, in clinical trials.(15) BUPs aim to de-medicalise minor infections and cut antibiotic use by empowering patients to share control over their treatment plan.(16, 17) Point of care testing (POCT) can help reduce antibiotic prescribing for RTIs in primary care.(16, 18) Rapid tests using biomarkers such as C-reactive protein (CRP) can help reduce uncertainty about the need for antibiotics.(19, 20) GPs report that use of POCT can reduce diagnostic uncertainty, but that costs and effect on workflow are potential barriers.(21)

In this study we aimed to explore public perceptions regarding various approaches to consulting for RTIs, and perceptions about BUPs and POCT to target high levels of antibiotic use. Understanding public perceptions may help inform the implementation of different approaches to this challenging problem.

Methods

We administered a questionnaire as part of a national online cohort study in Wales, called HealthWise Wales (HWW).(22) All adults (age 16 years or older) who are usually resident or receive their healthcare in Wales were eligible for inclusion in HWW. Participants were recruited through television, radio and social media advertising campaigns, as well as promotion through the NHS (hospitals and GP surgeries), pharmacies, large employers, and at cultural events. HWW includes multiple cross-sectional surveys, encompassing general questionnaires and more focused areas of research. Our sample consisted of HWW participants who completed a module (questionnaire) called, 'Caring for Coughs and Colds' (CCC). The module included 31 questions covering five domains: perceived consulting frequency, drivers of consulting, perceptions of serious illness, alternative sources of information and views about alternative approaches to consulting for RTIs (Supplementary Table S1). Response options used five-point Likert scales, ranging from "strongly agree" to "strongly disagree" and "very much in favour" to "completely against". Questionnaire development was based on previously published studies (23) used an iterative approach to develop items and a pilot questionnaire to refine items.

Analysis

Questionnaire responses were summarised using numbers and percentages for categorical variables and mean and standard deviation for continuous data.

Drivers of consulting

After examining response distributions, the five-point Likert scales for the drivers of consulting and components of a consultation domains were collapsed into binary outcomes of “agree” (strongly agree/agree) and “do not agree” (neither agree nor disagree/disagree/strongly disagree) for ease of understanding and to allow for a logistic regression analysis. We conducted multivariable logistic regression analyses to explore predictors of being in favour of: consulting with a community pharmacist; consulting with a pharmacist or nurse in a general practice; having a video consultation; having an email consultation; receiving a BUP; confidence in knowing when to take a BUP; and having a POCT.

Perceived features of serious illness

Response options for the domain on perceived features of serious illness were left as the original wider five-point scale ranging from “very serious” to “not serious at all”.

Alternative sources of advice

The first 3,775 participants were mistakenly been asked to respond to questions on sources of advice using either “acceptable” or “not acceptable”. Unfortunately, it is difficult to interpret these responses, and therefore they were not analysed leading to a high level of missing data for these questions. Questions on alternative sources of advice from the remaining responders were recorded using a five-point Likert scale ranging from “very much in favour” to “completely against”. After examining the response distributions these were collapsed into a three-point scale (“in favour”, “neither in favour nor against” and “against”) and then dichotomised into “in favour” versus “not in favour” (“neither in favour nor against” and “against”) and a logistic regression model was conducted to examine characteristics associated with favouring alternative sources of advice.

Data on personal characteristics were obtained from other core HWW modules. Levels of relative deprivation were calculated from home postcode using the Welsh Index of Multiple Deprivation (WIMD), a measure utilised by the Welsh Government.(24) Consultation and antibiotic prescribing data were obtained through anonymous data linkage with primary care health record data held within the Secure Anonymised Information Linkage (SAIL) Databank. SAIL is an archive for personal, anonymised data, which can be linked on an individual level,(25) and holds anonymised data for around 80% of Welsh GP surgeries.(26) Read codes in the primary care record were used to identify consultations for the symptoms of an RTI and antibiotic prescriptions across a 3-year period from January 2015 to December 2017 (Supplementary Tables S2 and S3). SAIL data was also used to identify the presence of Read codes for four key comorbidities (asthma, diabetes mellitus (DM), cardiovascular disease (CVD) and chronic obstructive pulmonary disease (COPD)) (Supplementary Table S4).

For the regression models, the following pre-hypothesised explanatory variables were used: gender, age, deprivation quintile, rurality, current mental health problem, history of mental health problem, marital status, having children, smoking status, reported physical exercise, the four comorbid conditions (asthma, DM, CVD, COPD), consulting and prescribing frequency, and responses to three of the questionnaire items on symptoms reflecting infection seriousness (perception of a cough lasting one week or more, perception of a cough with

green phlegm lasting two days or more, perception of a noisy and wheezy chest). For each analysis, the univariable association between outcome and each independent variable was calculated, and variables associated with outcome at $p=0.1$ were entered into a multivariable analysis. Results from the models are reported as odds ratios (ORs) with 95% confidence intervals (CIs).

Results

8,752 participants completed the HWW CCC module between October 2016 and April 2018. Study participant characteristics are presented in Table 1. Just under three-quarters were female, and the mean (standard deviation) age was 49(17) years. 4,387 (51.6%) came from the two most affluent quintiles, with only 12% coming from the most deprived. Less than 10% were current smokers, and just over a third reported that they had not engaged in any exercise in the past week. Responses to questions about alternative sources of health information from the first 3,775 (43%) participants were discarded because of an error in the response options for these participants. There were no marked differences observed between responders retained in the main analysis and those excluded due to response options (Supplementary Table S5). 1,662 (19%) participants could not be linked to GP data within SAIL.

The most common reason for deciding to consult about an RTI was worry about an infection being serious (90.5% of participants). However, perceived need for an antibiotic prescription was also an important driver (Figure 1). The illness features most associated with a perception of serious illness were fever, prolonged dry cough (for three weeks) and green phlegm (Figure 2).

<Figure 1: Public perceptions of reasons to consult a healthcare professional for a respiratory tract infection>

<Figure 2: Public perceptions of seriousness of features associated with respiratory tract infections>

Of the 4,966 participants who completed a question about consulting for an RTI with a pharmacist or nurse in their usual GP surgery, 76.7% (3,807) were in favour (figure 3). Of the 4,953 participants who completed the question about consulting with a community pharmacist with training in assessing RTIs, 71.2% (3,529/4953) were in favour (figure 3). 47.4% (2342/4944) and 42.3% (2095/4949) were in favour of having video and email consultations for RTIs respectively. 31.8% (2746/8646) and 92.8 (8034/8659) were in favour of having back up prescription and finger-prick blood tests respectively. These proportions did not vary significantly by socio-economic status (Table 2).

<Figure 3: Public views about possible sources of advice/information for respiratory tract infections>

Female gender was associated with being in favour of consulting with a local pharmacist or pharmacist or nurse in their own surgery (Table 3). Adults without children were more likely to favour being seen by a local pharmacist than parents, and individuals who thought that a one-week cough was serious, a marker of increased concern about RTI symptoms, were more likely to accept a pharmacist or nurse consultation (Table 3). Younger (<65 years) and sedentary (no weekly exercise) participants, and those not reporting a mental health problem,

were more in favour of video consultations (Table 3). Similarly, younger participants, and those that believe having a wheezy chest is an indicator of serious infection were more likely to favour email consultations.

Back-up prescriptions and finger-prick blood tests

2,746 (31.8%) agreed they would like to be given a BUP if their clinician thought they did not need immediate antibiotics. 2,922 (33.8%) respondents indicated that they would not feel comfortable in knowing when to take a BUP. There was an association between socio-economic deprivation and wanting to be given a BUP, with those from more deprived backgrounds being more in favour of this approach (Table 2). However, in a multivariable analysis socio-economic status was not significant. Those who reported that they would be happy to receive a BUP being more likely to be male, young, have a current mental health problem, be a parent and be sedentary (report not engaging in any exercise). A history of COPD was associated with twice the odds of wanting to be given a BUP. Consulting less frequently, receiving antibiotics more frequently, and having greater concern about symptoms (believing that a one-week cough and a noisy, wheezy chest are indicators of serious infection) were all associated with a greater odds of wanting to be offered a BUP. Women, younger adults (under 30), those who more sedentary and those who consulted less frequently, were all more likely to indicate that they would not feel comfortable in knowing when to take a BUP if they were given one (Table 3).

8,034 (92.8%) participants agreed that they would generally be happy to be offered a finger-prick blood test to help guide antibiotic treatment during a consultation. Women, younger people, smokers (current and previous) and frequent antibiotic recipients were less likely to report finger-prick blood testing acceptable (Table 3).

Discussion

Summary

Key findings in this survey of public perceptions about consulting for RTIs are that more than 90% of respondents indicated that they would be happy to have a finger-prick blood test in primary care to help inform the need for antibiotics, and around three-quarters would be happy to consult with a community pharmacist or nurse or pharmacist in their surgery. Women, adults without children, and people with greater concern about the seriousness of cough, were more in favour of having the option of consulting with non-medical practitioners. Less than half of respondents thought electronic consultations (video/email) were acceptable, with younger people being more in favour of this approach. Less than a third of participants indicated they would want to be given a BUP, and a third said they would not feel comfortable deciding when to use a BUP.

Strengths and limitations

Our study benefited from being able to link perceptions reported in an online survey to primary care health record data. We obtained data from over 8,700 people, however we had over-representation of females (73.9% female compared to 50.7% of the UK population), (27) and middle-aged (30-64 year-old age group) participants. Our survey was only available to those with internet access between October 2016 and April 2018, which was 85% of Welsh households in 2018. (28) Our respondents were on average less socio-economically deprived than the general population, which is likely why we had a lower proportion of current

smokers than in the general population. Nevertheless, we were able to include large numbers from all socio-economic groups and control for socioeconomic status in our analyses. We found an association between socio-economic status and views about use of BUP, but this was not significant in a multivariable analysis, and we found no other significant associations with socio-economic status. We lost 3775 (43%) of our sample due to a mistake with response outcomes; however, the samples that we excluded and those retained were broadly similar and appears to exhibit little bias. Finally, views expressed in an online survey may differ from those expressed in other settings or following greater provision of information. For example, questions about back-up prescribing may have differed if respondents had a more detailed understanding of the reasons for using this approach and how it works.

Interpretation and comparison with existing literature

How people appraise their symptoms and perceive the severity of their illness was identified as a key driver of consulting in a previous qualitative study and survey of adults across England.(23) Women were more likely to consult, but there were no differences by age or region in this study. Worry about cough has previously been shown as a driver of consulting in general practice.(29) We found that those who perceived a one-week cough as serious were more likely to indicate that consulting with a pharmacist would be acceptable. This suggests that those with greater concern about RTI symptoms want different consultation options, but this needs confirming.

A 2014 public survey found a similar proportion (just over a third favouring or strongly favouring) back-up/delayed prescriptions, as in our study (31.8%).(30) Their findings are consistent with our finding that women and older people are less likely to favour delayed/back-up antibiotics, but they also found that women and parents were more aware of BUPs.(31) We also explored perceptions about deciding when to use BUP antibiotics and found that a third were not comfortable making this decision. A previous qualitative interview study found that most patients were happy to be given BUP, but some found it confusing to be issued a prescription after being told that it was viral.(32) A similar study in Australia identified concerns about the use of BUP, and in particular knowing when to take them.(33) GPs have expressed mixed views about BUPs.(34, 35)

Implications for research and practice

Our findings support the view that several approaches to managing RTIs are likely to be acceptable to the public. Pharmacists, nurses and allied health professionals already play an important role in managing RTIs, and our findings suggest that many value these options, with men and parents reporting less acceptance. These findings may help those designing services, and further research could explore the reasons for these differences. We found less enthusiasm for digital (video and email) consultations for RTIs. Previous qualitative research has found that this approach is generally very acceptable to patients, although technical issues can be a barrier to use.(36, 37) In the context of the COVID-19 pandemic remote consultations have become the norm, and it would be interesting to see how public perceptions on use of these technologies have changed.

It is encouraging to see that more than 90% were happy to have a finger-prick blood test to help guide antibiotic prescribing. POCT can improve antibiotic prescribing for RTIs,(16) including acute exacerbations of COPD.(38) Our findings suggest that members of the public have mixed views about the role of BUPs. There is clearly a need for more public information about BUPs, including a rationale for use of BUP and clear instructions on when to initiate

them. These findings can be used to help develop strategies to improve the management of RTI and reduce the use of antibiotics in primary care.

Additional information

Funding:

This study received no direct external funding, but was supported by Healthwise Wales, which is funded by the Welsh Assembly Government through Health and Care Research Wales.

Ethical approval:

Healthwise Wales received ethical approval from Wales Research Ethics Committee (REC) 3 on 16th March 2015 (reference 15/WA/0076). A substantial amendment was submitted when the Caring for Coughs and Colds module was added and this received approval from Wales REC 3 on 13/10/2016.

Competing interests

The authors declare no competing interests.

Acknowledgements

This study was facilitated by HealthWise Wales, the Health and Care Research Wales initiative, which is led by Cardiff University in collaboration with SAIL, Swansea University. We would like to acknowledge the Healthwise Wales team who supported the study (Shantini Paranjothy, Pauline Ashfield-Watt, Ameeta Richardson, and Mark Atkinson), Fiona Lugg-Widger, who helped develop the questionnaire, and the Healthwise Wales study participants.

References

XUncategorized References

1. McNulty CAM, Collin SM, Cooper E, et al. Public understanding and use of antibiotics in England: findings from a household survey in 2017. *BMJ Open*. 2019;9(10).

2. Hobbs FDR, Bankhead C, Mukhtar T, et al. Clinical workload in UK primary care: a retrospective analysis of 100 million consultations in England, 2007– 14. *The Lancet*. 2016;387(10035):2323-30.
3. Aryee A, Price N. Antimicrobial stewardship - can we afford to do without it? *Br J Clin Pharmacol*. 2015;79(2):173-81.
4. Cosgrove SE, Carmeli Y. The impact of antimicrobial resistance on health and economic outcomes. *Clin Infect Dis*. 2003;36(11):1433-7.
5. Bell BG, Schellevis F, Stobberingh E, et al. A systematic review and meta-analysis of the effects of antibiotic consumption on antibiotic resistance. *BMC Infect Dis*. 2014;14:13.
6. Cars O, Hogberg LD, Murray M, et al. Meeting the challenge of antibiotic resistance. *BMJ*. 2008;337:a1438.
7. Public Health England. English Surveillance Programme for Antimicrobial Utilisation and Resistance (ESPAUR) Report 2018-2019.
8. Costelloe C, Metcalfe C, Lovering A, et al. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: systematic review and meta-analysis. *BMJ*. 2010;340(7756).
9. Dolk FCK, Pouwels KB, Smith DRM, et al. Antibiotics in primary care in England: which antibiotics are prescribed and for which conditions? *J Antimicrob Chemother*. 2018;73(suppl_2):ii2-ii10.
10. National Institute for Health and Clinical Excellence, Public Health England. Summary of antimicrobial prescribing guidance - managing common infections 2020 [Available from: <https://www.nice.org.uk/Media/Default/About/what-we-do/NICE-guidance/antimicrobial%20guidance/summary-antimicrobial-prescribing-guidance.pdf>].
11. Gulliford MC, Dregan A, Moore MV, et al. Continued high rates of antibiotic prescribing to adults with respiratory tract infection: survey of 568 UK general practices. *BMJ Open*. 2014;4(10).
12. Smieszek T, Pouwels KB, Dolk FCK, et al. Potential for reducing inappropriate antibiotic prescribing in English primary care. *J Antimicrob Chemother*. 2018;73(suppl_2):ii36-ii43.
13. Wakeman M, Cork T, Watwood D. A Pilot Study investigating Point of Care C-Reactive Protein (POC CRP) Testing in Community Pharmacy to deliver appropriate Interventions in Respiratory Tract Infections (RTIs). *J Pharm Pharmacol*. 2019;71(s1):11-2.
14. Brant H, Atherton H, Ziebland S, et al. Using alternatives to face-to-face consultations: a survey of prevalence and attitudes in general practice. *Br J Gen Pract*. 2016;66(648):e460-6.
15. Little P, Moore M, Kelly J, et al. Delayed antibiotic prescribing strategies for respiratory tract infections in primary care: pragmatic, factorial, randomised controlled trial. *BMJ*. 2014;348(mar05 4).
16. Spurling GK, Del Mar CB, Dooley L, et al. Delayed antibiotic prescriptions for respiratory infections. *Cochrane Database of Systematic Reviews*. 2017.
17. Little P. Delayed prescribing of antibiotics for upper respiratory tract infection. *BMJ : BMJ*. 2005;331(7512):301.
18. Aabenhus R, Jensen J-US, Jørgensen KJ, et al. Biomarkers as point-of-care tests to guide prescription of antibiotics in patients with acute

respiratory infections in primary care. Cochrane Database of Systematic Reviews. 2014;2014(11).

19. Stanton N, Francis N, Butler C. Reducing uncertainty in managing respiratory tract infections in primary care. *Br J Gen Pract*. 2010;60(581).

20. Huang Y, Chen R, Wu T, et al. Association between point-of-care CRP testing and antibiotic prescribing in respiratory tract infections: a systematic review and meta-analysis of primary care studies. *Br J Gen Pract*. 2013;63(616):e787.

21. Eley CV, Sharma A, Lecky DM, et al. Qualitative study to explore the views of general practice staff on the use of point-of-care C reactive protein testing for the management of lower respiratory tract infections in routine general practice in England. *BMJ Open*. 2018;8(10).

22. Hurt L, Ashfield-Watt P, Townson J, et al. Cohort profile: HealthWise Wales. A research register and population health data platform with linkage to National Health Service data sets in Wales. *BMJ Open*. 2019;9(12):e031705.

23. McNulty CAM, Nichols T, French DP, et al. Expectations for consultations and antibiotics for respiratory tract infection in primary care: the RTI clinical iceberg. *Br J Gen Pract*. 2013;63(612):e429.

24. Welsh Government. Welsh Index of Multiple Deprivation 2014 [Available from: <https://gov.wales/statistics-and-research/welsh-index-multiple-deprivation/what-is-wimd/?lang=en>].

25. Welsh Government. SAIL Databank 2019 [Available from: <https://saildatabank.com>].

26. Jones K, Ford D, Lyons R. The SAIL Databank: 10 years of spearheading data privacy and research utility, 2007-2017. 2017.

27. Coates S. Overview of the UK population: November 2018. 2018.

28. Statistics for Wales. National Survey for Wales, 2017-18: Internet use and digital skills. 2018.

29. Cornford CS, Morgan M, Ridsdale L. Why do mothers consult when their children cough? *Fam Pract* 1993;10(2):193-6.

30. McNulty CAM, Lecky DM, Hawking MKD, et al. Delayed/back up antibiotic prescriptions: what do the public think? *BMJ Open*. 2015;5(11).

31. McNulty C. Personal communication. ed2020.

32. McDermott L, Leydon G, Halls A, et al. Qualitative interview study of antibiotics and self-management strategies for respiratory infections in primary care. *BMJ Open*. 2017;7(11).

33. Lum EPM, Page K, Nissen L, et al. Australian consumer perspectives, attitudes and behaviours on antibiotic use and antibiotic resistance: a qualitative study with implications for public health policy and practice. *BMC public health*. 2017;17(1):799.

34. Peters S, Rowbotham S, Chisholm A, et al. Managing self-limiting respiratory tract infections: a qualitative study of the usefulness of the delayed prescribing strategy. *Br J Gen Pract*. 2011;61(590):e579.

35. Ryves R, Eyles C, Moore M, et al. Understanding the delayed prescribing of antibiotics for respiratory tract infection in primary care: a qualitative analysis. *BMJ Open*. 2016;6(11).

36. Donaghy E, Atherton H, Hammersley V, et al. Acceptability, benefits, and challenges of video consulting: a qualitative study in primary care. *Br J Gen Pract*. 2019;69(686):e586.
37. Atherton H. Digitally enabled primary care: the emperor's new clothes? *Br J Gen Pract*. 2019;69(686):420-1.
38. Butler CC, Gillespie D, White P, et al. C-Reactive Protein Testing to Guide Antibiotic Prescribing for COPD Exacerbations. *New Engl J Med*. 2019;381(2):111-20.

Accepted Manuscript – BJGPO – BJGPO.2020.0224

Table 1. Study participant characteristics (N=8,752)

	N (%)
Gender	
Male	2273 (26.1)
Female	6452 (73.9)
Missing	27
Age	
Under 30 years	1522 (17.4)
30-64 years	5512 (63.0)
65 years or older	1718 (19.6)
Missing	0
Deprivation quintile	
1 (most deprived)	975 (11.5)
2	1427 (16.8)
3	1710 (20.1)
4	2279 (26.8)
5 (least deprived)	2108 (24.8)
Missing	253
Rurality	
Urban settlement of over 10,000	5130 (60.4)
Town or fringe settlement	1507 (17.7)
Village, hamlet and isolated dwellings	1862 (21.9)
Missing	253
Current mental health problem	
No	6010 (71.8)
Yes	2362 (28.2)
Missing	380
History of mental illness	
No	5664 (66.9)
Yes	2800 (33.1)
Missing	288
Marital status	
Not married	1940 (30.4)
Married	4445 (69.6)
Missing	2,367
Have children	
No	2998 (35.8)
Yes	5375 (64.2)
Missing	379
Smoking status	
Never smoked	4756 (56.2)
Previous smoker	2890 (34.2)
Current smoker	810 (9.6)
Missing	296
Physical exercise in the past week	
None	3107 (36.7)
Some but less than 1 hour	1197 (14.1)
1 hour or more, but less than 3 hours	1994 (23.6)
3 hours or more	2166 (25.6)
Missing	288
Ever diagnosed with asthma	
No	5926 (83.6)
Yes	1164 (16.4)
Missing*	1662
Ever diagnosed with DM	
No	6657 (93.9)
Yes	433 (6.1)
Missing*	1662

Ever diagnosed with CVD	
No	6854 (96.7)
Yes	236 (3.3)
<i>Missing*</i>	1662
Ever diagnosed with COPD	
No	7032 (99.2)
Yes	58 (0.8)
<i>Missing*</i>	1662
Consulting frequency	
Never	4866 (71.4)
Less than once a year	1076 (15.8)
About once a year	812 (11.9)
Twice or more per year	65 (1.0)
<i>Missing*</i>	1933
Antibiotic prescribing frequency	
Less than once a year	5791 (81.7)
Once or more per year	1299 (18.3)
<i>Missing*</i>	1662

All results reported as n(%)

*Due to non-linkage

Accepted Manuscript – BJGPO – BJGPO.2020.014

Table 2 – Variation in responses by socio-economic status

Variable	Deprivation quintile					Missing	Total
	1 most deprived	2	3	4	5 least deprived		
In favour of:							
Consulting with Pharmacist or Nurse in practice	376/498 (75.5)	588/783 (75.1)	714/920 (77.6)	1088/1425 (76.4)	918/1187 (77.6)	123/153 (80.4)	3807/4966 (76.7)
Consulting with community pharmacist	347/494 (70.2)	556/783 (71.0)	650/918 (70.8)	1006/1422 (70.7)	859/1182 (72.7)	111/154 (72.1)	3529/4953 (71.2)
Video consultation	245/494 (49.6)	357/779 (45.8)	434/916 (47.4)	632/1420 (44.5)	607/1184 (51.3)	67/151 (44.4)	2342/4944 (47.4)
Email consultation	220/492 (44.7)	325/780 (41.7)	377/915 (41.2)	613/1421 (43.1)	493/1187 (41.5)	67/154 (43.5)	2095/4949 (42.3)
Back up prescription	351/963 (36.4)	469/1411 (33.2)	542/1685 (30.1)	678/2251 (29.9)	624/2085 (27.0)	82/251 (32.7)	2746/8646 (31.8)
Finger-prick blood test	875/958 (91.3)	1276/1416 (90.1)	1572/1690 (93.0)	2110/2257 (93.5)	1966/2087 (94.2)	235/251 (93.6)	8034/8659 (92.8)

All results reported as n(%)

Accepted Manuscript – BJCP 2020.0124

Table 3. Summarised results for the multivariable analysis to identify associations between explanatory and outcome variables

	Local pharmacist ¹	Pharmacist or nurse ¹	Video consultation ¹	Email ¹	Receive BUP ²	Take BUP ²	POCT ²
Adjusted OR ³ (95% CI)							
Gender							
Male (ref.)	1.00	1.00			1.00	1.00	1.00
Female	1.48 (1.29, 1.71)	1.46 (1.26, 1.70)			0.80 (0.70, 0.91)	1.18 (1.02, 1.37)	0.64 (0.49, 0.83)
Age							
Under 30 years (ref.)	1.00		1.00	1.00	1.00	1.00	1.00
30-64 years	1.19 (0.97, 1.46)		1.03 (0.85, 1.24)	0.73 (0.55, 0.97) 0.62 (0.44, 0.88)	0.79 (0.66, 0.94) 0.96 (0.77, 1.20)	0.49 (0.40, 0.61)	1.43 (1.12, 1.82)
65 years or older	1.01 (0.80, 1.29)		0.75 (0.59, 0.96)			0.51 (0.39, 0.68)	1.65 (1.16, 2.35)
Deprivation quintile							
1 (most deprived) (ref.)			1.00		1.00		1.00
2			0.84 (0.65, 1.08)		0.86 (0.70, 1.05)		0.80 (0.58, 1.10)
3			0.88 (0.68, 1.13)		0.83 (0.68, 1.01)		1.28 (0.91, 1.80)
4			0.82 (0.64, 1.05)		0.87 (0.72, 1.05)		1.11 (0.80, 1.54)
5 (least deprived)			0.99 (0.78, 1.26)		0.86 (0.71, 1.04)		1.36 (0.97, 1.90)
Rurality							
Urban settlement >10,000 (ref.)			1.00	1.00		1.00	
Town or fringe settlement			0.92 (0.76, 1.12)	0.96 (0.76, 1.21)		1.02 (0.85, 1.21)	
Village, hamlet and isolated dwellings			1.03 (0.85, 1.24)	0.90 (0.73, 1.10)		0.98 (0.83, 1.16)	
Current mental health problem							
No (ref.)			1.00		1.00	1.00	1.00
Yes			0.83 (0.71, 0.97)		1.24 (1.09, 1.41)	1.03 (0.88, 1.19)	0.94 (0.76, 1.17)
History of mental illness							
No (ref.)		1.00					
Yes		1.12 (0.97, 1.30)					
Marital status							
Not married (ref.)				1.00		1.00	
Married				0.93 (0.76, 1.15)		1.00 (0.85, 1.18)	
Have children							
No (ref.)	1.00			1.00	1.00	1.00	
Yes	0.82 (0.70, 0.95)			1.04 (0.85, 1.29)	1.17 (1.02, 1.34)	1.00 (0.85, 1.18)	
Smoking status							
Never smoked (ref.)			1.00		1.00	1.00	1.00
Previous smoker			1.09 (0.94, 1.27)		1.08 (0.95, 1.22)	0.92 (0.80, 1.06)	0.99 (0.79, 1.23)
Current smoker			0.83 (0.65, 1.06)		1.17 (0.96, 1.42)	0.99 (0.78, 1.25)	0.72 (0.54, 0.98)
Physical exercise in the past week							
None (ref.)			1.00	1.00	1.00	1.00	1.00
Some but <1 hour			1.26 (1.02, 1.56)	0.79 (0.61, 1.02)	0.87 (0.73, 1.03)	1.18 (0.97, 1.43)	1.01 (0.76, 1.36)
1 hour or more but <3 hours			1.20 (1.00, 1.43)	1.13 (0.92, 1.39)	0.83 (0.72, 0.97)	0.93 (0.79, 1.10)	1.01 (0.78, 1.30)
3 hours or more			1.21 (1.01, 1.44)	1.15 (0.94, 1.41)	0.81 (0.70, 0.94)	0.79 (0.66, 0.93)	1.13 (0.86, 1.47)
Ever diagnosed with asthma							
No (ref.)							
Yes							
Ever diagnosed with DM							
No (ref.)					1.00		
Yes					1.20 (0.95, 1.50)		
Ever diagnosed with CVD							
No (ref.)			1.00	1.00	1.00		
Yes			0.84 (0.56, 1.26)	0.67 (0.41, 1.09)	1.19 (0.87, 1.62)		
Ever diagnosed with COPD							
No (ref.)					1.00		
Yes					2.27 (1.21, 4.27)		
Consulting frequency							
Never (ref.)				1.00	1.00	1.00	1.00
Less than once a year				1.01 (0.81, 1.26)	1.25 (1.07, 1.46)	0.93 (0.78, 1.11)	1.36 (1.02, 1.82)
About once a year				0.86 (0.66, 1.12)	1.27 (1.05, 1.54)	0.71 (0.57, 0.88)	1.20 (0.87, 1.66)
Twice or more per year				0.88 (0.37, 2.08)	1.32 (0.75, 2.33)	1.36 (0.71, 2.63)	1.34 (0.56, 3.17)
Antibiotic prescribing frequency							
Less than once a year (ref.)			1.00		1.00		1.00
Once or more per year			0.87 (0.69, 1.10)		1.51 (1.23, 1.85)		0.58 (0.42, 0.80)
Perception of 1-week cough							
Not serious (ref.)		1.00			1.00	1.00	
Serious		1.33 (1.08, 1.65)			1.61 (1.37, 1.89)	0.85 (0.70, 1.02)	
Perception of green phlegm for 2 days							
Not serious (ref.)					1.00		1.00
Serious					1.08 (0.95, 1.23)		0.89 (0.71, 1.11)
Perception of noisy and wheezy chest							
Not serious (ref.)				1.00	1.00		1.00
Serious				0.85 (0.72, 1.00)	1.24 (1.09, 1.41)		0.86 (0.69, 1.07)

1 In favour of consultation
2 Agree with use in primary care consultation
3 Adjusted for all other factors in the model




Figure 1. Public perceptions of reasons to consult a healthcare professional for a respiratory tract infection

299x176mm (72 x 72 DPI)




Figure 2: Public perceptions of seriousness of features associated with respiratory tract infections




Figure 3. Public views about possible sources of advice/information for respiratory tract infections

299x176mm (72 x 72 DPI)