

Taking the Pulse of Earth's Tropical Forests using Networks of Highly Distributed Plots

Author: "ForestPlots.net", with individual contributors as listed in table (alphabetical by country and name)

Abstract

Tropical forests are the most diverse and productive ecosystems on Earth. While better understanding of these forests is critical for our collective future, until quite recently efforts to measure and monitor them have been largely disconnected. Networking is essential to discover the answers to questions that transcend borders and the horizons of funding agencies. Here we show how a global community has responded to the challenges of tropical ecosystem research with diverse teams measuring forests tree-by-tree in hundreds of long-term plots. We review the major scientific discoveries of this work and show how this process is changing tropical forest science. Our core approach involves linking long-term grassroots initiatives with standardized protocols and data management to generate robust scaled-up results. By connecting tropical researchers and elevating their status, this 'social research network' model recognises the key role of the data originator in scientific discovery. Conceived in 1999 with RAINFOR (South America), our permanent plot networks have been adapted to Africa (AfriTRON) and Southeast Asia (T-FORCES) and widely emulated worldwide. Multiple initiatives are integrated via ForestPlots.net cyber-infrastructure, linking colleagues from 53 countries. The combined networks are transforming understanding of tropical forests and their biospheric role. Together we have discovered how, where and why forest carbon and biodiversity are responding to climate change, and how they feedback on it. Long-term pan-tropical collaboration has revealed a large long-term carbon sink and its trends, as well as making clear which drivers are most important, which forest processes are affected, where they are changing, what the lags are, and the likely future responses of tropical forests as the climate continues to change. By leveraging a remarkably old technology, plot networks have already sparked a modern revolution in tropical forest science. In the future, humanity will benefit greatly by nurturing the grassroots communities now collectively capable of generating unique, long-term understanding of Earth's most precious forests.

Key Words: Amazonia, Africa, Southeast Asia, rainforest, RAINFOR, AfriTRON, T-FORCES, forest plots, permanent sample plots, monitoring, dynamics, carbon sink, global change, biodiversity.

Introduction

As the most diverse and productive ecosystems on Earth, tropical forests play essential roles in the carbon and water cycles and maintenance of global biodiversity. Tropical forest lands are also home to more than a billion people and thousands of cultures. Having first provided the environments and germplasm that sustained foragers and farmers since the earliest days of humanity, today they underpin a large fraction of our globalized diet and intense demand for water, food and clean air. They also affect our health in multiple ways, providing rich pharmacopeias to traditional and modern societies, and capable of changing the course of history when pandemic zoonotic pathogens emerge as forests and wildlife are exploited. Tropical forests are also critical to determining the degree and impact of anthropogenic climate change. Because of their extent, carbon density and productivity, they may both slow global heating by absorbing carbon into their biomass and soils, or accelerate it as deforestation and high temperatures damage forests and release carbon to the atmosphere.

Tropical carbon and biodiversity are therefore critical targets for environmental measurement and monitoring. While vital to our past and future, efforts to measure and monitor them have until recently been localised and largely disconnected. Although aspects of their ecology can be sensed remotely on-the-ground, tree-by-tree measurement is essential. Indeed ground measurements are irreplaceable – whether to address a plethora of ecological questions (e.g., Wright, this volume), inform and validate ecosystem models (e.g., Malhi et al., this volume), or assist with interpreting remotely acquired data (e.g., Chave et al. 2019, Duncanson et al. 2019, Phillips et al. 2019). Yet the very features that enhance tropical forests' ecological value, such as remoteness, diversity and high rainfall, make fieldwork challenging. Tropical forest science and scientists from forest-rich countries are often under-resourced and academically marginalised. Often colonized from afar and distant from economic centres, tropical nature and many who explore it remain peripheral to national and global academic and political priorities.

The focus of this paper is specifically about the power of new collaborative networks to transform tropical forest science – *what* we do, *how* we do it, and eventually *who* does it - to understand tropical forest functioning and dynamics over large temporal and spatial scales. Conceived and funded starting in South America in 1999 (RAINFOR, Malhi et al. 2002) and later adapted to Africa (AfriTRON, Lewis et al. 2009) and Southeast Asia (T-FORCES, Qie et al. 2017) our approach encourages international grassroots initiatives and links them with standardized field methods and data management. Now, with ForestPlots.net (Lopez-González et al. 2011, 2015) we support multiple networks with cyber-infrastructure that enables tropical scientists to do together what was previously impossible alone. Providing tools to ensure tropical scientists can manage, share and analyse their data themselves, ForestPlots.net is a global platform where data originators are in control and free to collaborate, support, or lead as much as they like. However, while much has been accomplished the wider challenges still run deep. Our aim of supporting the best possible science within a model of equitable access to data and other resources, remains as much an aspiration as a claim of achievements already made.

Here, we review how the continental networks and ForestPlots.net emerged, in terms of collaborators, institutions, people and plots. Next, we focus on key scientific achievements of the combined networks, including a comprehensive understanding of the variation in biomass carbon stock, growth rates, and carbon residence time among continents. Then, we review multiple discoveries concerning large-scale changes over time, with insights emerging from hundreds of permanent plots that have transformed our understanding of the role that tropical forests play in the biosphere. Finally, we return to the challenges

of building and sustaining long-term science networks in the tropics and outline key priorities for the future.

1. Network Development

Tropical research plots that tag, measure, identify and follow forests tree-by-tree have existed for decades. They long precede any continental or global network, but no plot survives since before 1939 and few predate 1970. The earliest efforts were closely connected to the imperial- and post-imperial projects of European nations. As such, these were largely motivated by questions of timber inventory and wood production, and only later diversity and wider ecological questions. The very first permanent sample plots we are aware of in the tropics were installed in 1857 by the German forester Brandis, who worked for the British in Burma (now Myanmar) and later other parts of India (Dawkins and Philip, 1998). **In India** a few extant Forest Department plots date to 1939 (Pomeroy et al. 2003). Important early work in **Southeast Asia** included plots installed by Don Nicholson and J.E.D. Fox in the 1950s through to the 1970s, as well as Peter Ashton since the 1960s and John Proctor since the 1970s. **In Africa**, early permanent plots include those installed by William Eggeling in Uganda in the 1930s. Among plots surviving today are one in Mpanga Forest, Uganda, set up by Alan Hamilton in 1968, and those established by Mike Swaine in Ghana and Hans Woell in Liberia in the 1970s, and later plots by Jan Reistma and Lee White (Gabon), Bonaventure Sonké (Cameroon), Kofi Affum Baffoe (Ghana), and Henri-Félix Maître and colleagues (Gabon, Congo, C.A.R.). **In Australia**, North Queensland saw the first plot sampling, for timber, in the 1930s, with many sites from the 1970s still maintained today by the national science agency (CSIRO). Separately Joe Connell, co-originator of the influential Janzen-Connell hypothesis, installed and expanded long-term ecological plots in 1963.

In the tropical Americas, T.A.W. Davis and Paul Richards installed ecological plots in Guyana in the 1930s (Davis and Richards 1933) but these do not survive, while Frank Wadsworth established long-term plots in Puerto Rico's subtropical forests starting in 1943 (e.g. Drew et al. 2009). In Suriname, Schulz and colleagues established silvicultural studies in the 1950s and 60s that were used to design the CELOS Management System (Werger 2011). Neotropical ecological plots that persist today include many in Venezuela by Jean-Pierre Veillon in the 1950s, 60s and 70s (Vilanova et al. 2018) and Rafael Herrera, Ernesto Medina and colleagues in the 1970s, as well as a few in Brazilian Amazonia by João Murça Pires, H. Dobzhansky and G.A. Black and later Ghilleen Prance, and several in Costa Rica since 1969 by Diana and Milton Lieberman. Elsewhere, Alwyn Gentry, John Terborgh, Terry Erwin, Gary Hartshorn, David Neill and Rodolfo Vásquez set up the first long-term plots in western Amazon in the late 1970s and 80s (Gentry 1988a, Monteagudo et al. 2020). Eastern and central Amazon plots survive established by Ima Vieira and Rafael Salomão in Pará (Salomão 1991, Pires and Salomão 2000), Tom Lovejoy, Niro Higuchi and colleagues near Manaus, and Henri-Félix Maître in French Guiana. The earliest extant plots in southern Amazon originate with Tim Killeen, Luzmila Arroyo, Beatriz Marimon and José Roberto Rodrigues. The first long-term tropical large plot was established in Costa Rica (Hubbell 1979), which represented a separate innovation that permitted plot-level analysis of multi-species demography, followed soon after by the first 50-ha plot in Panama (Hubbell and Foster 1983, Wright this volume) and later developments by the Smithsonian Institution and the ForestGEO network (e.g. Anderson-Teixeira et al. 2015).

RAINFOR (Red Amazónica de Inventarios Forestales) is the first international tropical forest network encompassing hundreds of long-term plots. At root, RAINFOR was inspired by Alwyn Gentry, a virtuoso tropical botanist who established the first globally standardized floristic inventories. In the 1970s Gentry developed a 0.1-ha sampling design to rapidly inventory diversity in species-rich tropical forests, capturing all stems ≥ 2.5 cm diameter. He and his colleagues applied it with throughout the

tropical Americas as well as parts of Africa, India, Southeast Asia, Australasia, and some northern and southern temperate forests. By the time of his untimely death at the age of 48 in 1993, Gentry had completed 226 of these samples, comprising an inventory of thousands of tree and liana species including many new to science. His legacy lives on in multiple ways. After studying with Walter Lewis and recruited by Peter Raven in the early 1970's, Gentry was a key figure in the Missouri Botanical Garden's golden age of tropical botany. He collected more than 80,000 plant specimens, approximately half of which are tropical trees and lianas. He pioneered a new approach to the challenge of identifying plants in the world's most diverse forests (Gentry and Vásquez 1993) which has inspired generations of botanists throughout Latin America. Perhaps most importantly, it was Gentry who embodied the ambition of combining efficient ecological sampling with high-quality identifications and replicating these to create highly distributed measurements of the world's forests (e.g. Gentry 1988b, Clinebell et al. 1995, Phillips and Miller 2002, Phillips and Raven 1997). He also established permanent plots (Gentry 1988a) that feature in the first continental and pan-tropical analyses of forest carbon and dynamics (Phillips and Gentry 1994, Phillips et al. 1994, Phillips et al. 1998), which in turn led to the creation of RAINFOR (Malhi et al. 2002, López-Gonzalez and Phillips 2012) and its protocols (e.g. Phillips et al. 2002). Originating in 1999 from a small nucleus of researchers and plots and supported by EU funding to Brazil's LBA initiative, RAINFOR grew to tackle the challenge of analysing Amazonian forests and climate responses tree-by-tree from the ground-up. By bringing different groups together RAINFOR facilitated the development of long-term international collaborations to measure and understand not only forest dynamics and diversity but also biogeochemistry and carbon fluxes.

While RAINFOR has grown steadily, other plot networks later emerged with complementary foci in South America. Some are daughter initiatives to RAINFOR, others were formed separately, but most share a similar ethos and strongly overlapping protocols. To the extent that they can be combined together these networks represent an impressive 'Observatory for Neotropical Forests'. Below (Table 1) we report key information about many vibrant networks worldwide that specifically contribute to ForestPlots.net, while here we briefly enumerate national and international neotropical networks, most but not all of which we support. These include (with dates when fieldwork used these network labels) Tropical Ecology Assessment and Monitoring (TEAM, 2002), Amazon Tree Diversity Network (ATDN, 2003), Programa de Pesquisa em Biodiversidade (PPBio, 2004, Brazil), Global Ecosystems Monitoring (GEM, 2010; Malhi et al. this volume), Latin American Seasonally Dry Tropical Forest Network (DryFlor, 2012), Red Colombiana de Monitoreo de los Bosques (ColTree, 2012), Secondary Forest Network (2ndFOR, 2015), Peru Monitoring Network (MonANPerú, 2017), sANDES (Tree Diversity, Composition and Carbon in Andean Montane Forests, 2019), and Red de Bosques Andinos (RBA, 2020), as well as global networks and meta-networks including ForestGEO (Anderson-Teixeira et al. 2015), sPlot (Bruehlheide et al. 2019), FOS (Schepaschenko et al. 2019) and TmFO in logged forests (Sist et al. 2015). Each of these has notable achievements of their own and at the time of writing this in 2020 almost all have active research programmes.

In Africa, our early networking focussed on assessing whether there were similar patterns of changes in carbon stocks as observed in South American forests and the causes of such changes. Efforts began in 2001 to recensus many of the earlier plots installed in post-independence Africa (UK funding to Phillips, Malhi and Lewis), which were later formalised as the African Tropical Rainforest Observation Network (AfriTRON; Lewis et al. 2009). These plots span 12 African countries with moist forests from Sierra Leone in the west to Tanzania in the east. Like RAINFOR in Amazonia, AfriTRON pools expertise and data to tackle long-term, large-scale questions relating to the ecology and biogeochemistry of tropical forests. Networks sharing a similar ethos with programmes in Africa now include TEAM, DynAfFor (Gourlet-Fleury et al. 2013), TmFO and ForestGEO. Recently, the SEOSAW (Ryan et al. in

review) and AfriMont networks have been established, extending long-term plots into the extensive southern woodlands and savannas and Africa's distinctive montane forests.

Our work in Southeast Asia began in 2001 to assess forest carbon balance and later developed into a network once Lan Qie undertook considerable fieldwork and networking. European Research Council investment (T-FORCES 2012 grant to Phillips, Malhi and Lewis) enabled intensive campaigns to develop long-term plot networking in Borneo (Qie et al. 2017), and also supported African censuses. While smaller than its Amazonian and African counterparts the Asian network builds on plots installed by a number of foresters and botanists as long as 60 years ago. Critically, RAINFOR, AfriTRON, T-FORCES and TmFO use the same field and analytical protocols.

Insert Table 1 here: **Table 1. Networks contributing to ForestPlots.net (August 2020).**

How can we combine the different strengths of these and other initiatives to maximise their impact on science and society? To achieve this requires shared data management tools and horizontal organisational structures that foster leadership by tropical scientists. Our plot data management scheme was originally conceived in 2000 as a desktop database to support RAINFOR analyses of spatial variation in wood density, biomass, productivity, and changes in biomass over time (Baker et al. 2004a, b; Malhi et al. 2004). This was expanded to draw together inventory data from more than 100 sites in Amazonia and subsequently African forest plots to include some of the longest running monitoring sites worldwide (Peacock et al. 2007).

Since 2009 we have developed a global, on-line Structured Query Language resource with sophisticated programming that supports multiple applications (López-Gonzalez et al. 2011). Now, ForestPlots.net provides ecological informatics to colleagues in scientist-led networks from 53 countries working across 44 tropical nations (Fig. 1). Key advances in this platform include the ability to manage complex time-series data, to track species linked to high-quality botanical records, and to analyse records with common BiomasaFP R protocols (López-Gonzalez et al. 2015). While focussed on species identity, tree growth, mortality and carbon dynamics, ForestPlots.net encompasses many related forest attributes including lianas, soils, and plant traits.

At their heart long-term plots are an intensely human enterprise and so we document too the personal contributions to plot establishment and continued monitoring. By tracking who did what, and when, we also honour the inter-generational aspect of plots that allows modern analysts to stand on the shoulders of giants. With ForestPlots.net data contributors retain control and are able to manage, share and analyse their records using a common toolset. If new projects requesting to use their data are proposed they can agree to collaborate, or not, as they wish. Contributors often propose their own multi-site projects. ForestPlots.net can provide DOIs to datasets, further ensuring that contributors are properly acknowledged. Developing this functionality has supported a surge in multi-site and multi-national analyses that are increasingly initiated by scientists from the tropics, gradually supplanting the traditional model where researchers from the Global North lead. In sum, ForestPlots.net enables the level of control and collaboration that individual researchers wish for while underpinning network and multi-network integration. In turn, this is empowering data owners and networks and helping to transform the face of tropical ecology.




Figure 1. Current extent of ForestPlots.net

Top: Pantropical plot sampling density per 2.5 degree square with the 3,653 multiple- and single-inventory plots hosted at ForestPlots.net. These plots contribute to networks including RAINFOR, AfriTRON, T-FORCES, ATDN, BIOTA, ColTree, FATE, GEM, Nordeste, PELD, PPBio, RAS, RBA and SEOSAW. Forest cover based on the Global Land Cover 2000 database (JRC, 2003) with tree cover categories: broad-leaved evergreen; mixed leaf type; and regularly flooded. Our plots also extend into neotropical and African savannas; **Bottom:** The same plot sampling but displayed at higher-resolution (1-degree grid cells) for each focal continent, South America, Africa, and Southeast Asia and Australia.

The networks and ForestPlots share a 20-year history, but as already seen the history of plot monitoring is much longer. The first recorded census in ForestPlots.net dates from 1939 in Budongo, Uganda. Forty years later, 676 censuses had been completed from 90 plots, but since 1979 fieldwork has accelerated greatly with 9,000 censuses completed across 3,500 plots by 2019 (Fig. 2a). This acceleration is reflected by the growing community of contributors, which by 2019 had surpassed 1,800 individuals (Fig. 2b). ForestPlots.net itself has grown steadily both in terms of censuses uploaded and in outputs (Fig. 3). The neotropics dominate much of this inventory and monitoring effort as well as the growth of ForestPlots.net in particular, but contributions from Africa and other continents are increasing (Figs. 2, 3). Scientific outputs emerging from this collective effort span local to global scales but have an increasingly pan-tropical theme (Fig. 3b).





Figure 2. Growth of pan-tropical forest monitoring since the mid-twentieth-century.

Top: Plot-censuses curated at ForestPlots.net by date of census;

Bottom: Cumulative number of contributors to ForestPlots.net by date of first recorded fieldwork. Growth was slow following the first census in 1939, only reaching 100 censuses by 1969. For early censuses, records of field team personnel and leaders are often sparse or absent. Note that ‘contributors’ is defined inclusively to reflect members of indigenous communities, protected area guards, parataxonomists, students, and technicians, as well as principal investigators, botanists, and other specialists.




Figure 3. Growth of ForestPlots.net and contributing networks since 2000.

Top: Cumulative upload of unique plot censuses to ForestPlots.net by date of upload, with pre-2009 uploads to pre-internet versions allocated evenly back to network beginnings;


Bottom: Cumulative peer-reviewed scientific articles based on network plots, excluding research based on single-plot studies.

2. Environmental Representation




While it is not possible to intensively sample the whole tropical forest extent, in practice RAINFOR, AfriTRON and T-FORCES have managed to cover almost the entire climatic and geographic space across the humid tropics with permanent plots (Fig. 4a) as well as extensively sample the biome space of the terrestrial tropics except for semi-arid biomes (Fig. 4b). Within each continent coverage has been focused on the moist tropical lowlands with sampling extending into montane and drier forest systems most effectively in South America (Fig 4c). Plots also cover the complex edaphic variation present in Amazonia (Quesada et al. 2012) where they encompass landscape-level variability within old-growth forests (Anderson et al. 2009, 2010). This effective representation of structurally intact moist forests provides good support for large-scale inferences from what is, inevitably, a limited sample of the domain. It is important to note that many tropical countries lack statistical inventories of forests, let alone long-term monitoring, so that research plots fill critical gaps in global and national observations.

Yet significant work remains to be done to increase representativeness, better understand impacts of geological and edaphic variation, and expand sampling in remote areas especially in parts of Amazonia, the central Congo Basin, and New Guinea (c.f. Brearley et al. 2019). Fuller environmental coverage can help networks address challenges such as monitoring of protected area effectiveness (Baker et al. 2020) and providing calibration-validation of Earth Observation space-borne sensors (Chave et al. 2019). Beyond the lowland humid tropics, special effort is also needed for long-term, ground-based monitoring in particular environments. Expansion is especially required for: (i) secondary forests and those impacted by disturbance events such as logging, fragmentation, and wildfires (e.g. Chazdon et al. 2016, Elias et al. 2020); (ii) montane forests, which harbour exceptional concentrations of endemism and are at great risk of biodiversity loss due to deforestation and climate change, and therefore represent urgent conservation opportunities (e.g. Malizia et al. 2020); (iii) Asian dry forests, and (iv) the wider extent of tropical dry forest and savanna biomes, which are home to distinctive biotas and significant carbon stocks of their own (DRYFLOR 2016, Pennington et al. 2018). ForestPlots.net partner groups are expanding research and monitoring in such critical areas beyond the structurally intact lowland forests that have been the main focus of RAINFOR and AfriTRON.

1. Geographic distance (km)



2. Environmental distance (Euclidean distance [SD])



[Previous Page] Figure 4. Network coverage of geographical and climate space

Analyses include >1500 ForestPlots.net permanent plots from 15 networks (RAINFOR, AfriTrON, T-FORCES, Andes, ColTree, ECOFOR, FATE, GEM, I-For, Nordeste, PELD, PPBio, RAS, RBA and TROBIT). **(a) Top panels:** (1) Geographic distance between multi-census plots across the humid tropical forest biome; and (2) Minimum climate dissimilarity (Euclidean distance on variables scaled by their standard deviation, accounting for mean annual temperature, temperature seasonality, mean annual precipitation and precipitation seasonality), where for each cell environmental distance represents how dissimilar a location is to the most climatically similar plot in the network. Note that some poorly sampled areas are mostly deforested, such as Central America, Madagascar, and much of tropical South and Southeast Asia. The baseline map depicts WWF terrestrial ecoregions (Olson et al. 2001). **(b) Middle panel:** Tropical plots displayed in global biome space (Whittaker diagram), showing the main concentration of plots from lowland wet through to moist forests and savanna, with some samples in cooler montane climates. **(c) Lower panels:** Plots displayed within tropical humid and sub-humid climate space (represented by light grey pixels), with plots displayed colour-coded by continent and symbol size corresponding to total census effort. Note the important differences in baseline climatic conditions between continents.

3. Discovery: Forest Ecology across the Tropical Continents

RAINFOR, AfriTRON and T-FORCES plots have generated ecological and biogeographical insights that have only been achievable via large-scale collaboration. RAINFOR has revealed that Amazonian forests differ substantially from one another, even those that share essentially identical climates. For example, basal-area weighted wood density of northeastern forests is 50% greater than that of southern and western forests. This reflects floristic differences (Baker et al. 2004, ter Steege et al. 2006, Honorio Coronado et al. 2009, Patiño et al. 2009), which, in turn, are associated with large differences in forest dynamics. Stem turnover is twice as fast in the west and south as the east (Phillips et al. 2004) due to younger soils with poorer structure providing less rooting support (Quesada et al. 2012, Schiatti et al. 2016) and in spite of only modest productivity differences (Malhi et al. 2004, 2014). In contrast, biomass in north-eastern Amazonia is higher than elsewhere due to the reduced mortality risk and hence bigger trees and denser wood (Baker et al. 2004, Malhi et al. 2006, Marimon et al. 2014, Pallqui et al. 2014, Johnson et al. 2016, Alvarez et al. 2018, Phillips et al. 2019).

In Africa, AfriTRON plots also show that species-driven differences in wood density prevail at large scales. In mature forests, soil-related compositional differences cause significant variation in basal-area weighted wood density. Forests on younger and more fertile acrisols and cambisols have 10 and 20% lighter wood than those on arenosols and histosols, respectively (Lewis et al. 2013). Similarly to Amazonia, African forests growing on older, less fertile soils have higher biomass (Lewis et al. 2013). Local and regional variation in soils and forest attributes are important within both continents but the key difference is that only Amazonia has clear continental-scale gradients in wood density, due to the powerful influence of Andean orogeny in the west. This leads to young, geologically dynamic landscapes with fertile, less-developed soils, influencing speciation, immigration and extinction, and contrasts with the ancient, stable Brazilian and Guyanan Shields of the east. The influence of climatic differences on Amazon floristics is of second-order importance.

Large-scale analysis thus reveals how soils and species help control the carbon that tropical forests store. This has implications for monitoring carbon stocks using remotely-sensed data. In tropical forests neither soil nor tree composition is easily perceived from space. For example, RAINFOR plots show that LiDAR-derived biomass estimates of Amazonian forests are compromised because they do not perceive the critical large-scale floristic gradients (Mitchard et al. 2014). Accounting for such


limitations by relating plot-derived woody density and allometry to LiDAR sampling shows that plots greatly improve biomass maps (Mitchard et al. 2014, Avitabile et al. 2016). Thus the role of soils and species composition in affecting biomass carbon is a key reason why ground data are essential for mapping forests (Chave et al. 2019). While Earth Observation has great benefits in terms of spatial coverage and ability to update frequently, the incorporation of plot-derived compositional data greatly improves our understanding of carbon storage patterns over large scales.

When networks using the same protocols are combined it is also possible to discover and explore variation between continents too. Common protocols have revealed major pan-tropical variation in vertical structure, including tree height and height-diameter allometry (Feldpausch et al 2011) which have impacts on biomass (Feldpausch et al 2012; Sullivan et al 2018). African forests average one-third higher biomass per unit area than Amazon forests (Lewis et al. 2013), yet have roughly one-third *fewer* stems >10 cm diameter per unit area. This may be driven by systematically lower tree mortality in these forests (Hubau et al. 2020, Sullivan et al. 2020). Similarly, comparing climatically and edaphically similar forests in parts of Borneo with northwest Amazonia reveals that Bornean forests produce much more wood, with trees growing up to 50% more rapidly than those of Amazonia. This suggests that differences in phylogenetic composition of tree communities, especially the dominance of the dipterocarp family in tropical Asia (Corlett and Primack 2011), determine the efficiency with which atmospheric carbon is converted to woody carbon (Banin et al. 2014).

Tree species composition and dominance strongly control forest function within continents too. For example, a recent RAINFOR study discovered that Amazon woody productivity is enhanced in more phylogenetically diverse forests (Coelho de Sousa et al. 2019). Yet while Amazonian forests are very diverse, research led by the ATDN network shows that remarkably few species dominate in terms of stems (ter Steege et al. 2013), while RAINFOR plots show that different species dominate biomass stocks and woody productivity (Fauset et al. 2015). Evidence also suggests that a number of such ‘hyperdominants’ may have been long favoured by indigenous people as part of wider human influences on old-growth Amazon forests (Levis et al. 2017, Oliveira et al. 2020). These and other studies show that identity matters. Dominant species and their evolutionary history thus affect forest ecology and forest values, whether in terms of storing carbon, converting solar energy into wood or sustaining whole cultures.

These insights show that two of the defining challenges of the twenty-first century, climate change and biodiversity loss, are closely linked. How then do we best devise conservation strategies to achieve the targets of biodiversity protection and climate mitigation and adaptation? Can we rely for example on carbon conservation via schemes like REDD+ to protect tropical diversity too? The answers to these questions depend on the relationship between diversity and carbon storage but assessing this has been challenging due to the scarcity of inventories in which both carbon stocks and species identifications have been reliably quantified. By combining RAINFOR, AfriTRON and T-FORCES plots we found that for tropical trees diversity-carbon storage relationships barely exist at all (Sullivan et al. 2017, Fig. 5). For example, South America, the continent with the richest forests, actually stores the least carbon per hectare, while within continents there is no association. Independent data from the RAS network data support this, showing that strong carbon-biodiversity relationships are only found in disturbed and secondary forests but not old-growth (Ferreira et al. 2018). As mature forests exhibit all possible combinations of tree diversity and carbon stocks it is clear that both need to be explicitly considered to protect the climate and biodiversity. In addition, long-term carbon storage is threatened by defaunation of large-bodied frugivores, often essential for dispersing large-seeded heavy-wooded tree species (Peres et al. 2016). *We cannot simply focus on carbon and achieve biodiversity conservation, and vice versa.*

404



405

406

407 **Figure 5. Pantropical forest carbon storage and species richness.**

408

409 There is a lack of within or across continent relationships between carbon stocks and tree species richness per
410 hectare in structurally intact old-growth tropical forests. Figure adapted from Sullivan et al. 2017.

411

412




Figure 6. Tropical continental macroecology

Remarkable continental differences in species richness, stem density and carbon stocks emerge among lowland tropical moist forests when densely-sampled plot networks are combined. Graphics depict probability densities such that the whole area for each continent sums to 1. Note that the y-axis scale for each variable thus *itself varies depending on the range of the x-axis*: for continents with larger variation in x, the probability density at any point along the y axis is correspondingly smaller. Analysis adapted from Sullivan et al. 2017 and 2020.

When network data are combined surprisingly large and coherent continental-level differences emerge (Fig. 6). African forests are remarkably species-poor at the 1-ha scale whereas South American and Asian forests are more than twice as rich on average, but also vary much more in species-richness and diversity. The very richest forests in the world are located in parts of Western Amazonia, vindicating a claim by Gentry (Gentry 1988a, b) from more than three decades ago. African forests have many fewer stems than their Asian and South American counterparts, but South American forests have considerably less biomass. In terms of carbon gains Borneo's forests are outliers, being up to twice as productive as other forests. Yet it is in South America where woody carbon turns over fastest. Fully half the carbon in neotropical trees has been replaced since 1970.

Overall these comparisons reveal remarkable continental differences in tropical forests that are not strongly driven by rainfall, temperature or soil (Sullivan et al. 2020). The implication is that other factors related to the evolutionary and historical happenstance of each continent matter. We draw three higher level conclusions from this. *First, global-scale ecological modelling ignores biological composition at its peril. Second, if there was ever any doubt, each continent clearly needs its own robust research and monitoring programme. And third, each region likely responds to climate change in its own, idiosyncratic way.*

4. Discovery: Tropical Forest Change

The single most significant scientific impact of these multiple permanent plot networks has been to transform our understanding of how tropical forests function in the Earth system.

As the most diverse and carbon-rich tropical biome, the fate of humid tropical forests will impact the future of all life on Earth. Until quite recently it was axiomatic that old-growth tropical forests are at equilibrium when considered over sufficiently large scales, and that any changes observed at smaller scales are driven by natural disturbance-recovery processes. However, large-scale imbalances observed in the global carbon balance have cast doubt on this assumption (e.g. Taylor and Lloyd 1992). Over time, network analyses have helped to recast our understanding of contemporary old-growth tropical forests as being non-stationary systems. Their carbon, biodiversity and ecosystem processes are widely recognised as dynamic and continually responsive to multiple anthropogenic drivers (e.g. Lewis et al. 2004b, Pan et al. 2011, Malhi et al. 2014, Levis et al. 2017, McDowell et al. 2018, Reis et al. 2018). Key discoveries at this intersection between global change science and forest ecology and biodiversity include:

(1) A pantropical increase in tree turnover rates, representing the first evidence for a widespread impact of global anthropogenic change on old-growth tropical forests (Phillips and Gentry 1994).

The finding that these forests were changing was controversial at the time - let alone the inference that global drivers were responsible - and contradicted established ecological orthodoxy. The debate that ensued helped generate new questions and analyses (e.g. Sheil 1996, Phillips and Sheil 1997) and address potential biases (e.g. Sheil 1995, Condit 1997, Lewis et al. 2004a, Gloor et al. 2009, Espirito-Santo 2014, Kohyama et al. 2019). A quarter of a century of research since then has repudiated the notion that 'intact' tropical forests are unaffected by atmospheric changes and reinforced the central concept that all tropical forests are being influenced by a suite of large-scale anthropogenic drivers.

(2) Biomass dynamics have also accelerated in Amazonia. In parallel with the increases in stem dynamics, as RAINFOR grew it became clear that carbon fluxes via biomass growth and mortality were

also increasing. Moreover, the increased gains in stems (recruitment) and biomass (woody productivity) clearly preceded increases in stem and biomass losses (mortality) (Lewis et al. 2004b, Phillips et al. 2004, 2008, Brien et al. 2015, Nogueira et al. 2018). The mechanism underlying this acceleration of forest dynamics must therefore involve stimulated productivity via increased resources for plant growth, rather than direct stimulation of tree mortality such as by drought (Lewis, Phillips and Malhi 2004).

(3) **The Amazon forest carbon sink.** In conjunction with faster growth and turnover, the biomass density of Amazonian forests has increased (Phillips et al. 1998, Baker et al. 2004, Pan et al. 2011). Old-growth Amazonian forests have absorbed (net) atmospheric carbon for at least three decades now (Brien et al. 2015), providing a true “subsidy from nature” with flux magnitude matching or exceeding net losses from neotropical deforestation (Aragão et al. 2014, Gatti et al. 2014). Thus, monitoring networks have shown that most Amazonian nations are on balance *not* net emitters of carbon (Espírito-Santo et al. 2014, Phillips and Brien et al. 2017). The location, magnitude and persistence of this old-growth carbon sink has important implications for guiding approaches to meeting nationally differentiated targets for controlling climate change (Vicuña Miñano et al. 2018).

(4) **The African forest carbon sink.** The AfriTRON network discovered a long-term net biomass increase similar in magnitude to that of the Amazon in the 1990s and early 2000s (Lewis et al. 2009). The consistency of these results on a second continent supports the idea that global drivers of change can affect even the most remote forests. The fact that biomass is increasing across the entire wood density spectrum of tree species implies that forests are responding to increasing CO₂ (Lewis et al. 2009). The long-term increase in carbon stocks of African forests was recently updated and confirmed, with three times as many plots showing continued sink strength (Hubau et al. 2020).

(5) **The Pan-Tropical forest carbon sink.** Once the T-FORCES network allowed sufficient plot coverage across remaining Bornean forest a similar increase in aboveground biomass over recent decades was revealed (Qie et al. 2017). Thus the three continental networks discovered that old-growth tropical forests as a whole have been a long-term sink. Our ground measurements imply this totalled more than one billion tonnes of carbon each year over the 1990s and early 2000s, i.e. half the terrestrial global carbon sink (Pan et al. 2011), which is sufficient to significantly slow climate change. The fact that the main blocs of remaining tropical forests are *en masse* out-of-equilibrium and undergoing biomass increases of similar magnitude implies a common global driver of growth. Increasing atmospheric CO₂ is the most parsimonious candidate and is consistent with predictions from first principles (e.g., Huntingford et al. 2013), inference from CO₂ fertilization experiments (Terrer et al. 2019), analyses of the global carbon budget (Ballantyne et al. 2012, Gaubert et al. 2019), observed greening of forests unaffected by land-use change (Piao et al. 2019), and recent plot analyses showing a significant role of CO₂ (Hubau et al. 2020).

(6) **The Amazon sink is slowing.** After 30 years of monitoring Amazonian forests, the RAINFOR plots show that the rate of increase in forest growth is declining. Tree mortality rates have increased in some regions, leading to a slow decline in the magnitude of the net biomass accumulation (Brien et al. 2015). The subsidy from nature provided by tropical forests may be time-limited.

(7) **Recent droughts in Amazonia have had large impacts.** Long-term plots also monitored immediately before and soon after droughts reveal that these forests can switch rapidly from being a major sink to a source of carbon. Both the 2005 and 2010 Amazon droughts had a net impact on the order of 1 Pg of carbon, driven primarily by drought-induced mortality (Phillips et al. 2009, Lewis et

al. 2011; Doughty et al. 2015, Feldpausch et al. 2016). RAINFOR and GEM have quantified the drought sensitivity of the world's biggest rainforest and identified the key process affected (mostly tree mortality rather than growth, and not photosynthesis). The impact on the biomass carbon sink of the 2010 drought and non-drought years match independent inferences from measurements of atmospheric [CO₂] using aircraft (Gatti et al. 2014).

(8) The African and Amazon sinks have diverged. Thirty years of monitoring AfriTRON plots show that African forests have continued to function as a carbon sink, although the most intensively monitored plots suggest that the sink may be declining (Hubau et al. 2020). When analysed together with RAINFOR data, within-plot changes over time reveal a common set of drivers that suggest the sinks will decline, with African forests lagging Amazonian forests by 15-20 years (Hubau et al. 2020). Changes across both continents are best explained by a combination of the positive effects of increasing CO₂ enhancing productivity and negative effects of higher temperatures and droughts in suppressing growth and accelerating mortality, combined with the intrinsic properties of forests themselves. The time-lag of the African sink saturation is due to longer carbon residence times in African forests, so that mortality catches-up slower than in faster turnover forests. Amazonian forests are often harder hit because they are hotter and can be drought-prone (Hubau et al. 2020). Together, the pan-tropical plot networks have revealed long-term trends in carbon storage and determined which drivers matter, which processes are affected, where they are impacting, and what the lags are.

(9) The future of the tropical forest carbon sink. Monitoring the present and the recent past of forest behaviour can also reveal likely future scenarios as the climate continues to change. Our plot networks provide two powerful and independent lines of evidence. First, the *long-term sensitivity to climate* emerges from a space-for-time analysis based on 813 plots across the Earth's tropical forests. This shows how maximum temperature and dry season intensity combine to determine the equilibrium climate controls on forest carbon, acting on productivity and mortality to limit forest carbon storage in *the long-term* (Sullivan et al. 2020). Forests exhibit remarkable thermal resilience under low amounts of warming, but in the hottest forests (>32.2°C max. temp.) biomass carbon drops off rapidly. Most of the biome will exceed this value with one further degree of warming (approximately equivalent to a 2°C increase above pre-industrial levels). Second, analysing recent changes in productivity and mortality as a function of recent climates, and coupling them with future climate scenarios, confirms that the carbon sink is likely to decline (Hubau et al. 2020). A key uncertainty with these latter projections is the extent to which local resilience due to shallow water-tables (Sousa et al. 2020) may mitigate effects, and whether more compositional changes will extend the carbon sink further if species better-adapted to the new conditions compensate for others' losses. The first analysis by Sullivan et al. (2020) confirms that lagged species-related resilience is likely as long as forests do not experience substantial warming.

(10) Tropical forest biodiversity is changing. RAINFOR data show that an entire group of plants, lianas (woody vines), are increasing in dominance across Amazonia (Phillips et al. 2002). Large lianas in turn contribute to higher tree mortality (Phillips et al. 2005). Tree community composition is changing too. In the Andes, plots of ABERG, RBA, and RedSPP collectively show 'thermophilization' – communities becoming more warm-adapted (e.g Fadrique et al. 2018). Climate change is inducing large-scale change in tropical lowland trees too, as wet-adapted taxa in Amazonia face greater mortality risks from drought (Esquivel Muelbert et al. 2017, 2019) while a shift towards drought-deciduous tree species is observed in west African plots experiencing a multi-decadal drought (Fauset et al. 2012, Aguirre-Gutiérrez et al. 2019). In both continents these community responses to drought coincided with biomass gains. Nonetheless, because of the long generation times of tropical trees the compositional

change has not kept pace with the drying of Amazonia (Esquivel-Muelbert et al. 2019). This suggests that further community change is inevitable, even before accounting for losses driven by deforestation and disturbance of remaining forests (Barlow et al. 2016). Current models lack the capacity to account for variation in tropical woody plant biodiversity and demographic processes and their lagged responses to global change drivers.

In sum, highly distributed, long-term monitoring of the world's richest forests has profoundly increased our understanding of nature's sensitivity to climate change. Looking forward, many of the key uncertainties that remain concern the responses of tropical biodiversity itself. This includes the extent to which the great biocomplexity of tropical forests themselves will provide an effective and timely insurance policy in the face of rapidly changing climates. To understand this, we must continue to monitor.

5. Challenges and the Future of Tropical Forest Monitoring

Large-scale plot networks have not only made a series of crucial scientific discoveries and advances, but the 'social research network model' pioneered by RAINFOR since 2000 has influenced how tropical ecology itself is being done. Tropical ecology has undergone a remarkable shift from a small cadre of researchers working in one or two sites to a more globalised and decentralised process with greatly increased contributions from tropical scientists. This has been made possible by supporting highly-distributed researchers and field sites, establishing mechanisms for shared data management, fostering an equitable concept of data ownership, and embracing groups who are often marginalised in research. Importantly, the network model is nurtured by researchers placing trust in the sharing of hard-won data to answer big questions and recognition of the value of developing trusting relationships over time. Finally, the growth of interactive multi-site, multi-cultural science has benefited hugely from standardized field and analytical methods that have been agreed upon, formalised and promoted. The ForestPlots.net experience demonstrates that collaborative, multi-polar structures help ensure breadth and resilience, while encouraging the leaders of the future.

The transformative power of this approach has now led to the establishment of multiple plot-centred networks that are reshaping our understanding of tropical ecosystems. However, these networks face a number of key challenges to sustain the achievements made and enact even deeper transformational change, which we set out here.

1. How can networks support leadership in the Global South? No one project can reverse the impact of centuries of global inequality, but this is no reason to refrain from tackling the structural impediments to attaining a more equitable world. Ecology and conservation science remain biased towards temperate ecosystems in terms of funding, researchers, and topical focus (Di Marco et al. 2017, Reboledo et al. 2020), and tropical ecology sometimes detached from policy-making processes and with most high-impact papers still led from the North. Together with open data-sharing and long-term collaboration, more leadership of forest science from tropical countries helps address these disparities and achieve more impact on forest and carbon management (e.g., Vargas et al. 2017, Baker et al. 2020). Supporting tropical students at different levels up to PhD and mentoring beyond the doctoral degree is also important. To help, ForestPlots.net has made shared tools widely available, and especially data management and analytic tools that support data contributors as much as users. To ensure fieldwork is valued and leadership in tropical researchers is fostered, we have developed a Code of Conduct to encourage contributions, support scientists in tropical countries, and promote mentoring of junior

scientists. To oversee this we created a diverse steering committee that currently supports more than 30 projects each year (<http://www.forestplots.net/en/join-forestplots/research-projects>). As a result, the proportion of ForestPlots.net research projects and products being led by tropical nationals has greatly increased, from less than 10% at the start of RAINFOR (2000-2004), to 35% in 2009 and 55% by 2019. In spite of such gains diversifying leadership is a long-term process. Ultimately, sustained funding in and by tropical countries themselves will ensure they not only have strong training programmes to develop the core field and analytical skills scientists need, but equal opportunities for career development.

2. How should we value and recognise collaboration and leadership? Most of the obvious reward structures in science - job security, income, grant success, peer reputation and public acclaim – can favour a ‘me first’ approach. Credit accrues to individuals, but true collaboration involves trust, sharing and encouraging others. Collaboration is gratifying, but letting go of our egos can be challenging, and in larger groups there is greater risk that individuals feel their contributions go unnoticed. Likewise, the essential major effort needed ‘backstage’ in ForestPlots.net to check data, update and develop data management, and support requests to utilize data, goes unseen. A partial developmental solution to this involves providing network contributors the opportunity to lead analyses with the expectation that *these new leaders then support others* with their analyses. Another approach is to reflect the diversity of contributions that underpin the success of networks by using a group author that shares credit amongst all (as in the current paper). These steps can promote the recognition of multiple contributions and development of tomorrow’s leaders.

3. How do we properly value the long-term? Project and thesis time-scales last from one to five years, but the lifespans of trees are measured in decades and centuries. What can seem vitally important in a hypothesis-driven research grant or a PhD may, in fact, have little relevance to the longer natural rhythms of nature. What if the dominant processes governing climate responses of forests turn out to involve lifetime accumulated ecophysiological stress, tree demography and species migration? Clearly very long-term research is essential to decode these processes. Meanwhile, maintaining permanent plots is as much an expression of hope in the future as a stake in an immediate scientific outcome, as rewards may accrue to others distant in time and space. Indeed, we have all benefited from researchers installing plots from the 1930s onwards. These pioneers never dreamt that their careful tree measurements and botanical identifications would help reveal the impacts of climate change on tropical forests. Long-term research programmes are simply irreplaceable, enabling us to discover, quantify, identify the causes of, and ultimately tackle environmental change.

4. Can we ensure fieldwork and human skills are valued for what they are? Technology provides many benefits to the scientific endeavor, but there are risks too, particularly in a field where long-term measurements may be perceived as old-fashioned (Ríos-Saldaña et al. 2018). A serious risk is that the tail wags the dog: when technological advance is an end in itself, it is unlikely that scientific and human progress will follow. Another risk is that our community and funders forget the basic truth that human skills are essential to measure and identify tropical trees. It is notable that those measuring, climbing and collecting tropical trees in permanent plots are among the least well-paid of all actors in the global scientific endeavour. Yet these true ‘key workers’ are irreplaceable as tree measurement in many locations is completely dependent on human labour and skill (Fig. 7) and, more broadly, *combinations* of people and technology provide the best results (next section). Moreover, because tropical tree floras usually run into the thousands of species (e.g., >4,700 tree species in Peru, Vásquez et al. 2018), identification depends on the work of well-trained botanists to collect material from canopies, make vouchers, and identify and permanently store them in herbaria. Without physical collections and the

immense knowledge and skills that produce them, identifications are untestable hypotheses whose quality cannot be evaluated. But with vouchers, we have the names that are essential to test questions about diversity, composition, functional traits, and wood density and biomass.

5. How should we fund proven networks long-term? As perhaps the most pressing immediate concern, this questions intersects closely with all of the above. *Few organisations have the vision to support long-term endeavours where leadership and credit is shared diffusely, many benefits accrue after decades, and where the most exciting discoveries may be unforeseeable.* We recommend the following, potentially transformational changes to address the challenges and unlock the benefits of ambitious, long-term forest monitoring:

(i) *Science Agencies have the foresight to build long-term research capacity*, and consciously adopt the challenge of international ecosystem monitoring and tropical career development;

(ii) *Space Agencies recognise that tropical fieldwork can measure the things they can not and validate the attributes that they can*, and contribute to the labour and unique skills of tropical field scientists and help overcome the challenges they face;

(iii) *Development and Conservation Agencies who depend on a robust understanding of the long-term health of forests*, recognise that high quality, long-term, on-the-ground monitoring of trees, and supporting these skills, is vital for their agenda;

(iv) *National and international climate adaptation and mitigation funders focus on what long-term, accurate forest monitoring can do to assess carbon fluxes and determine land-based emissions* which will impact forest management, nature-based solutions, and nationally determined contributions (NDCs) to reducing greenhouse gases for decades to come.

Every one of these user groups requires successful networks with long-term, research-grade tropical forest plots to discern the status and change of biodiversity and to assess the stocks and flows of carbon.

691



692

693

694 **Figure 7. Accurately measuring and identifying trees in remote tropical forests requires**
695 **dedication, skill and courage.**

696

697 To measure the diameter of this giant *Ceiba* (Malvaceae) tree in Colombia's Chocó, three people each
698 needed to climb more than 10 meters. Such techniques can be the most practical and accurate options
699 for measuring large trees. Here, like many of our sites, there is no electric power, let alone a field station,
700 and chronic insecurity due to political and social conflicts and narcotrafficking means that aircraft and
701 laser-scanners are not deployable. Images: Pauline Kindler.

702

703

704

705 **6. Achievements, Impact and Potential**

706

707 Despite the challenges, tropical forest science has come a very long way. Until recently, tropical ecology
708 suffered from a massive data deficit. We had plenty of theory and conjecture, but few comparable
709 observations over time and space to (deductively) put these ideas to the test or (inductively) generate
710 new ones. ForestGEO, RAINFOR, AfriTRON, and the wider ForestPlots community have contributed
711 much to resolving this. *By leveraging a remarkably old technology, forest plot networks have sparked*
712 *a modern revolution in tropical forest science.* They provide the means by which we have quantified
713 the trajectory of tropical forest carbon balance, including its climate sensitivity, and now provide a Pan-
714 Tropical Observatory for tracking these vital indicators of Earth's health going forward.

715

716 Permanent plots are now the prism through which ecologists address a rich suite of ecological questions,
717 but they have also changed the way *others* see forests. For example, well-identified permanent plots
718 have proved fertile ground for botanists to discover new tree species and genera (e.g. Reitsma et al.

1988, Baker et al. 2017, Wurdack and Farfan Rios 2017, Vásquez et al. 2018, Gosline et al. 2019, Vásquez & Soto, 2020), ethnoecologists to quantify forest people's values (Phillips and Gentry 1993, Lawrence et al. 2005), atmospheric scientists to explore organic volatiles production (Harley et al. 2004), ecophysiologicalists to assess why trees die (Rowland et al. 2015, McDowell et al. 2018), modelers to verify ecosystem simulations (Johnson et al. 2016), and foresters to predict and manage wood production and its impacts (Berry et al. 2008, Gourlet-Fleury et al. 2013). They provide critical infrastructure for whole-biodiversity and cross-taxa inventory, including exploration of cryptic canopy and soil faunal and microbial biodiversity. The impacts of these networks on policy are also growing. In Peru, for example, permanent plots are being used to validate national contributions to the Paris Climate Accord (Vicuña et al 2018, Baker et al. 2020). In Ghana, plots were needed to quantify historical and current carbon stocks, helping to establish baseline forest reference levels for the flagship Cocoa Forest REDD+ Programme (FCPF, 2017). In Gabon stratified-random sampling of high-quality AfriTRON plots is being used for the National Forest Inventory (Poulsen et al. in press. Ecol. Apps). Internationally, RAINFOR, AfriTRON, T-FORCES and 2ndFor provide the new IPCC default values on old-growth and secondary forest carbon sequestration to assist countries develop their own nationally determined contributions as part of the UNFCCC process (Requena Suarez et al. 2019).

What of the future? As new technologies for probing forests become available, the hundreds of standardised long-term plots and networks of skilled tropical researchers represent critical infrastructure to enhance and calibrate new insights as they arise. The benefits of working within established plots go beyond simply having confidence in species identifications and hence biomass. By leveraging their labour and insights, we can increase the scientific value of new technology. For example, the ability to match individual trees from laser-scanning surveys to tagged, censused individuals provides critical information on growth and identity (Disney et al. 2018). Integrating long-term botanical and ecological records of plots with terrestrial and airborne laser-scanning in designated 'super-sites' (Chave et al. 2019) can help overcome limitations of different approaches, providing greater certainty to biomass estimates (e.g., Schepaschenko et al. 2019). Hence forest networks can help unlock the value of space-based efforts to monitor forests. Just as the constellation of Earth-observing environmental satellites is a public good, the plot constellation is highly complementary, critical global infrastructure. Last but not least, as intact tropical ecosystems continue to shrink, burn and fray at the edges, permanent plots provide an indispensable baseline for understanding biodiversity and ecosystem processes. They should be our North Star for guiding sorely needed restoration efforts throughout this century.

So far this effort has relied on the goodwill of hundreds of colleagues and dozens of grants from many different sources (see Acknowledgments), but only long-term funding will ensure that the vital public benefits of plot networks continue to flow. Such support is surprisingly difficult to obtain (see Box 1). Twenty years of hard-won scientific results show that reliable and highly distributed monitoring is irreplaceable. They underscore the importance of welcoming all contributors to this effort, and of valuing the diverse skills needed to understand tropical biodiversity and its dynamics. Ultimately, we will understand the nature of tropical forests best when the science is global, local skills are fairly valued, and when the development of tropical scientists is at its heart. Indeed, we know of no other model capable of achieving this.

[Main Body of Text Ends]

[Insert Box]

Box 1. What Does It Take?

Clearly long-term ground-based monitoring of tropical forests requires a sustained global team effort. But just how much does it take to deliver tropical forest plot data in practice? It requires both skilled people and their labour, and funds. So here we address this question in terms of *the human effort made thus far* and the *financial investment needed to monitor across continents*.

(a) The Human Contribution: Network efforts include not only in-country field campaigns but much besides. To deliver from conception to product, high-quality data collected over many years and in dozens of countries requires multiple teams that are well-led and consistently trained in the proper protocols, quality control, and data management. In RAINFOR and AfriTRON this includes national or local field-team members to establish and remeasure plots, others to collect and identify plants and collect and analyse soils, colleagues to organize and manage the data, and others to sustain and lead the process nationally and globally – not to mention those who support these processes with essential administration, herbarium assistance, database development, analytical packages, IT support, technical training and so on. Naturally some individuals contribute in several ways and roles change over time as lives change. All local, national and global efforts ultimately depend on funding.

In sum, the average effort *in the field, herbarium, and lab* to install a typically remote and diverse 1-ha tropical forest plot and analyse its species and soil comes to 98 person-days, with an additional effort of 38 person-days *to support and sustain these teams and data management*. Together a total of *136 person-days is needed on average to deliver high-quality data from a new plot*.

Recensusing a plot is less demanding (for example soil collection is not repeated and there are fewer plants to identify) but still considerable: 45 person-days in the field and herbarium, and 31 person-days to support and sustain. Therefore, *76 person-days are required to deliver high quality data from a recensused plot*. These represent long-term averages. Naturally circumstances vary somewhat from site-to-site and country-to-country.

Thus far our teams have established 3,653 plots in tropical forests of which 1,626 are recensused, from as little as once up to as many as 40 times each. Of the 3,653 plots the modal size is between 0.9 and 1.1 ha but there are smaller plots too (1,728 are ≥ 0.9 ha, and 1,925 are < 0.9 ha). Of the 1,626 recensused plots, 67% are ≥ 0.9 ha (1,088) and 33% are < 0.9 ha (538).

If we conservatively assume that plots ≥ 0.9 ha (average size = 1.2 ha) require 136 days to install and 76 days to recensus, and those < 0.9 ha require half this effort (also likely to be conservative due to fixed costs for even the smallest plots), then the total effort to install these plots has been 176,044 person-days, and recensusing them has taken 341,204 person-days. In total this comes to 1,417 years.

As if one remarkably talented and tireless individual had been working continuously since AD 603.

(b) Cost of Sustained Continental Monitoring: *How much does it cost to monitor Earth's remaining old-growth tropical forests with ground networks?* This is a critical question given the exceptional

ecological value of these systems, the threats they are under, and the role they have and can play in modifying the rate of global climate change.

At first sight this question appears difficult to answer, or to even agree upon the terms of reference. Scientists would ask and likely argue: Monitoring what? For whom? With what precision, level of confidence, or spatial and temporal resolution? Recognising such difficulties we take a pragmatic approach and reframe the question. Instead we posit, *How much will it cost to monitor tropical forests using all the permanent plots that we have already remeasured?*

This question is tractable *practically* (these plots represent a known quantity: we know exactly *where* they are, *what* most of the species are, and to a large extent *who* can actually do the work – *each of which is critical*), it makes sense *scientifically* (the plots already have a baseline monitoring period against which we can assess any change, *which is essential*), and it is justifiable *quantitatively* (using somewhat smaller datasets than this we have already detected long-term changes in carbon balance, productivity and tree mortality on each continent, reported short-term changes in response to El Niño droughts and other climate anomalies, and attributed changes in carbon and biodiversity to climate drivers, *all of which establish proof-of-concept*). So here goes:

* There are 915 remeasured ForestPlots.net plots in tropical forest South America ($290 < 0.9\text{ha} + 640 > 0.9\text{ha}$), 462 in tropical forest Africa ($119 + 353$), 192 in tropical forest Asia ($106 + 86$) and 32 in tropical forest Australasia ($22 + 10$). With all 1,626 plots monitored on a four-year cycle this requires revisiting 406 plots annually, of which 272 are ≥ 0.9 ha.

* Recensus costs can vary from site-to-site. Botanical identification is especially challenging in most of South America due to the extraordinary diversity, while some African forests are exceptionally remote. Employment, social security and health costs vary but are rising almost everywhere. On average, considering all *the direct and indirect human effort required* (above) and *additional direct costs* (including consumables, equipment, travel, subsistence, insurance, visas, permits, shipping, training, IT), the current cost to deliver a high-quality tropical recensus is $\approx 18,000$ USD for plots ≥ 0.9 ha, and at least half this for plots that are < 0.9 ha.

[Similarly to install plots is a significant operation, and requires more expert time to collect and identify hundreds of trees. The total cost to properly install a high-quality tropical forest plot is $\approx 27,000$ USD for a 1 ha plot. When forests are recensused this start-up investment is leveraged as a contribution: this enables the subsequent monitoring of forest dynamics but it's not new spend.]

Thus, the annual delivery cost for a pantropical, practical ground-based *recensus programme* capable of tracking and attributing forest change to published standards is estimated as:

$(272 * 18,000 + 134 * 9,000) \approx 6.1$ million US dollars.

This *annual investment is sufficient to ensure that ground-measurements track the biome-wide and continent-specific biomass carbon balance of the remaining intact tropical moist forests, as well as their climate sensitivity*. It also provides ground calibration and validation for remote estimates of biomass. It also enables us to detect whether the tropical sink is now disappearing as predicted, and where and why, and what the consequences for biodiversity are, and to determine how much intact ecosystems can contribute to countries' nationally determined contributions (NDCs) to climate mitigation.

859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874

While \$6 million is a significant sum it is instructive to compare it to funding required for other large-scale science initiatives. The United States alone spends \$80 million annually (i.e., *thirteen times as much*) on its national forest inventory (Castillo and Alvarez, 2020). Space Agencies invest one to two orders of magnitude as much in each satellite mission designed to estimate biomass wall-to-wall for a few years (from ca. \$100 million to 500 million Euros for a single mission). And as we have seen, ground networks ultimately not only transcend the short-term time windows of such missions but add huge value to them.

In conclusion, the ongoing cost of monitoring Earth’s remaining tropical forests on the ground is extraordinarily small compared to the great scientific and practical benefits it provides. Now that the capacity to monitor tropical forests is established and proven it would be prudent to ensure this collective effort continues.

Acknowledgments

This paper is a product of the RAINFOR, AfriTRON and T-FORCES networks and the many other partner networks in ForestPlots.net which support long-term forest science and monitoring across tropical countries. These initiatives have been supported by numerous people and grants since their inception. We are particularly indebted to more than one thousand four hundred field assistants for their essential help in establishing and maintaining the plots, as well hundreds of rural communities and institutions. For additional assistance we thank Michel Baisie, Wemo Betian, Vincent Bezard, Mireille Breuer-Ndoundou Hockemba, Ezequiel Chavez, Douglas Daly, Armandu Daniels, Eduardo Hase, Muhammad Idhamsyah, Phillipe Jeanmart, Cisquet Keibou Opepa, Jeanette Kemp, Antonio Lima, Jon Lloyd, Mpanya Lukasu, Sam Moore, Klaus Scipal and Rodrigo Sierra. We thank Mark Burkitt for help developing the ForestPlots.net database. We acknowledge the long-term help provided by national and local government offices in all countries where colleagues work in facilitating the permission and documentation for fieldwork, as well as help provided by protected area and other authorities.

The networks have been supported by multiple grants, most notably the European Research Council (ERC Advanced Grant 291585 – ‘T-FORCES’), the Gordon and Betty Moore Foundation (#1656 ‘RAINFOR’, and MonANPe), the David and Lucile Packard Foundation, the European Union’s Fifth, Sixth, and Seventh Framework Programme (EVK2-CT-1999-00023 – ‘CARBONSINK-LBA’, 283080 – ‘GEOCARBON’, 282664 – ‘AMAZALERT’), the Natural Environment Research Council (NE/D005590/1 – ‘TROBIT’, NE/F005806/1 – ‘AMAZONICA’, ‘PPFOR’ E/M0022021/1, serval NERC Urgency and New Investigators Grants, the NERC/ State of São Paulo Research Foundation (FAPESP) consortium grants ‘BIO-RED’ (NE/N012542/1, 2012/51872-5), ‘ECOFOR’ (NE/K016431/1, 2012/51509-8), ‘ARBOLES’ (NE/S011811/1), the Royal Society (University Research Fellowships and Global Challenges Awards (‘FORAMA’, ICA/R1/180100), the National Geographic Society, the Centre for International Forestry (CIFOR), Gabon’s National Parks Agency (ANPN), US National Science Foundation (DEB 1754647), and Colombia’s Colciencias. We thank the National Council for Science and Technology Development of Brazil (CNPq) for support to the Cerrado/Amazonia Transition Long-Term Ecology Project (PELD/441244/2016-5), the PPBio Phytogeography of Amazonia/Cerrado Transition project (CNPq/PPBio/457602/2012-0), the Goiás Research Foundation (FAPEG/PELD: 2017/10267000329), and several PVE and Productivity Grants. Funding for plots in the Udzungwa Mountains (Tanzania) was obtained from the Leverhulme Trust under the Valuing the Arc project. We acknowledge grant CEBA (ref. ANR-10-LABX-25-01) and the support of the Forestry Development Authority of Liberia. We also acknowledge the support of the European Space Agency. Data from RAINFOR, AfriTRON and T-FORCES are stored and curated at ForestPlots.net, a cyber-infrastructure initiative developed at the University of Leeds that unites permanent plot records and supports scientists from the world’s tropical forests. The development of ForestPlots.net and curation of data has been funded by several grants including NE/B503384/1, NE/N012542/1 BIO-RED, ERC AdG 291585 T-FORCES’, NE/F005806/1 ‘AMAZONICA’, NERC New Investigators Awards, NE/N004655/1, ‘TREMOR’, the Gordon and Betty Moore Foundation (‘RAINFOR’, ‘MonANPe’), EU Framework 6, a Royal Society University Research Fellowship, and a Leverhulme Trust Research Fellowship.

The manuscript has been developed with the encouragement of Richard Primack and Reinmar Seidler and has benefited from the constructive comments from three anonymous reviewers; we thank them all. Finally we thank our late, great colleagues for their unique contributions that made possible all that the networks and ForestPlots.net have achieved together since the beginning: Samuel Almeida, Elisban

922 Armas, José Armas, Sandra Brown, Kwaku Duah, Gloria Galeano, Alwyn Gentry, Max Gunther, Moïse
923 Mikame, Norman Myers, Sandra Patiño, John Proctor, David Smith and Jean-Pierre Veillon.
924
925

Author Contributions

All authors have contributed to ForestPlots.net-associated networks by leading, collecting or supporting the collection of field data, or implementing and funding network data management, analyses and outputs. O.L.P. wrote the manuscript with contributions from most co-authors, M.J.S. contributed new analyses and M.J.S., G.L.P. and A.L.L. helped prepare the figures. O.L.P., T.R.B., G.L.-G. and S.L.L. conceived ForestPlots.net. R.B., T.F., D.G., E.G., E.H., W.H., A.E.-M., A.L., K.M., Y.M., G.C.P., B.S.-M., L.Q., and M.J.P.S have additionally contributed tools, funding or management to its development since.

Literature Cited

Aguirre-Gutiérrez, J., Oliveras, I., Rifai, S., Fauset, S., Adu-Bredu, S., Affum-Baffoe, K., Baker, T.R., Feldpausch, T.R., et al. 2019. Drier tropical forests are susceptible to functional changes in response to a long-term drought. *Ecology Letters*, 22, 855-865.

Alvarez-Davila, E., Cayuela, L., González-Caro, S., Aldana, A.M., Stevenson, P.R., Phillips, O., Cogollo, Á., Penuela, M.C., von Hildebrand, P., Jiménez, E. and Melo, O., 2017. Forest biomass density across large climate gradients in northern South America is related to water availability but not with temperature. *PloS One*, 12(3).

Anderson, L.O., Malhi, Y., Ladle, R.J., Aragao, L.E.O., Shimabukuro, Y., Phillips, O.L., Baker, T., Costa, A.C.L., Espejo, J.S., Higuchi, N. and Laurance, W.F., 2009. Influence of landscape heterogeneity on spatial patterns of wood productivity, wood specific density and above ground biomass in Amazonia. *Biogeosciences*, 6, 1883-1902.

Anderson, L. O., Malhi, Y., Aragão, L. E., Ladle, R., Arai, E., Barbier, N., & Phillips, O. 2010. Remote sensing detection of droughts in Amazonian forest canopies. *New Phytologist*, 187, 733-750.

Anderson-Teixeira, K.J., Davies, S.J., Bennett, A.C., Gonzalez-Akre, E.B., Muller-Landau, H.C., Joseph Wright, S., et al. 2015. CTFs-Forest GEO: a worldwide network monitoring forests in an era of global change. *Global Change Biology* 21, 528-549.

Aragao, L.E., Poulter, B., Barlow, J.B., Anderson, L.O., Malhi, Y., Saatchi, S., Phillips, O.L. and Gloor, E., 2014. Environmental change and the carbon balance of Amazonian forests. *Biological Reviews*, 89, 913-931.

Avitabile, V., Herold, M., Heuvelink, G.B., Lewis, S.L., Phillips, O.L., Asner, G.P., Armston, J., Ashton, P.S., Banin, L., Bayol, N et al., 2016. An integrated pan-tropical biomass map using multiple reference datasets. *Global Change Biology*, 22, 1406-20.

Baker, T.R., Phillips, O.L., Malhi, Y., Almeida, S., Arroyo, L., Di Fiore, A., Erwin, T., Killeen, T.J., Laurance, S.G., Laurance, W.F. et al., 2004. Variation in wood density determines spatial patterns in Amazonian forest biomass. *Global Change Biology*, 10, 545-62.

Baker, T.R., Phillips, O.L., Malhi, Y., Almeida, S., Arroyo, L., Di Fiore, A., Erwin, T., Higuchi, N., Killeen, T.J., Laurance, S.G. and Laurance, W.F., 2004. Increasing biomass in Amazonian forest plots. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 359, 353-365.

- Baker, T.R., Phillips, O.L., Laurance, W.F., Pitman, N.C., Almeida, S., Arroyo, L., DiFiore, A., Erwin, T. et al., 2009. Do species traits determine patterns of wood production in Amazonian forests? *Biogeosciences* 6, 297-307.
- Baker, T.R., Pennington, R.T., Magallon, S., Gloor, E., Laurance, W.F., Alexiades, M., Alvarez, E., Araujo, A., Arets, E.J., Aymard, G. and De Oliveira, A.A., 2014. Fast demographic traits promote high diversification rates of Amazonian trees. *Ecology Letters*, 17, 527-536.
- Baker, T.R., Vicuña Minano, E., Banda, K., Honorio Coronado, E., Lawson, I., Roucoux, K., del Castillo, D., Lojas, E., Monteagudo, A., Phillips, O., Vásquez, R., 2020. From plots to policy: how to ensure long-term forest plot data supports environmental management in intact tropical forest landscapes. *Plants, People and the Planet*
- Ballantyne, A.P., Alden, C.B., Miller, J.B., Tans, P.P. and White, J.W.C., 2012. Increase in observed net carbon dioxide uptake by land and oceans during the past 50 years. *Nature*, 488, 70-72.
- Banin, L., Lewis, S.L., Lopez-Gonzalez, G., Baker, T.R., Quesada, C.A., Chao, K.J., Burslem, D.F., Nilus, R., Abu Salim, K., Keeling, H.C. and Tan, S., 2014. Tropical forest wood production: a cross-continental comparison. *Journal of Ecology*, 102, 1025-1037.
- Barlow, J., Lennox, G.D., Ferreira, J., Berenguer, E., Lees, A.C., Mac Nally, R., Thomson, J.R., de Barros Ferraz, S.F., Louzada, J., Oliveira, V.H.F. and Parry, L., 2016. Anthropogenic disturbance in tropical forests can double biodiversity loss from deforestation. *Nature*, 535, 144-147.
- Berry, N.J., Phillips, O.L., Ong, R.C. and Hamer, K.C., 2008. Impacts of selective logging on tree diversity across a rainforest landscape: the importance of spatial scale. *Landscape Ecology*, 23, 915-929.
- Brienen, R.J., Phillips, O.L., Feldpausch, T.R., Gloor, E., Baker, T.R., Lloyd, J., Lopez-Gonzalez, G., Monteagudo-Mendoza, A., Malhi, Y., Lewis, S.L. and Martinez, R.V., 2015. Long-term decline of the Amazon carbon sink. *Nature*, 519, 344-348.
- Brearely, F.Q., Adinugroho, W.C., Cámara-Leret, R., Krisnawati, H., Ledo, A., Qie, L., Smith, T.E., Aini, F., Garnier, F., Lestari, N.S. and Mansur, M., 2019. Opportunities and challenges for an Indonesian forest monitoring network. *Annals of Forest Science*, 76, 54.
- Bruelheide, H., Dengler, J., Jiménez-Alfaro, B., Purschke, O., Hennekens, S.M., Chytrý, M., Pillar, V.D., Jansen, F., Kattge, J., Sandel, B. and Aubin, I., 2019. sPlot—A new tool for global vegetation analyses. *Journal of Vegetation Science*, 30, 161-186.
- Castillo, P.S.B. and M. Alvarez, 2020. *Forest Inventory and Analysis Fiscal Year 2018 Business Report*. United States Department of Agriculture, 78 pp.
- Chave, J., Davies, S.J., Phillips, O.L., Lewis, S.L., Sist, P., Schepaschenko, D., Armiston, J., Baker, T.R., Coomes, D. et al., 2019. Ground data are essential for biomass remote sensing missions. *Surveys in Geophysics* 40, 863-880.
- Chazdon, R.L., Broadbent, E.N., Rozendaal, D.M., Bongers, F., Zambrano, A.M.A., Aide, T.M., Balvanera, P., Becknell, J.M., Boukili, V., Brancalion, P.H. et al., 2016. Carbon sequestration potential of second-growth forest regeneration in the Latin American tropics. *Science Advances*, 2(5), p.e1501639.
- Clinebell, R., Phillips, O.L., Gentry, A.H., Stark, N., and Zuuring, H., 1995. Prediction of neotropical woody plant diversity from soil and climatic data. *Biodiversity and Conservation* 4, 56-90.

- Coelho de Souza, F., Dexter, K.G., Phillips, O.L., Pennington, R.T., Neves, D., Sullivan, M.J., Alvarez-Davila, E., Alves, Á., Amaral, I., Andrade, A. and Aragao, L.E., 2019. Evolutionary diversity is associated with wood productivity in Amazonian forests. *Nature Ecology & Evolution*, 3, 1754-1761.
- Condit, R.S., 1997, Forest turnover, diversity, and CO₂. *Trends in Ecology & Evolution* 12, 249-250.
- Corlett, R. T. and R.B. Primack. 2011. Tropical Rain Forests: An Ecological and Biogeographical Comparison, 2nd Edition. Wiley-Blackwell.
- Davis, T.A.W. and Richards, P.W., 1933. The vegetation of Moraballi Creek, British Guiana: an ecological study of a limited area of tropical rain forest. Part I. *Journal of Ecology*, 21, 350-384
- Dawkins, H.C. and Philip, M.S., 1998. Tropical moist forest silviculture and management: a history of success and failure. CAB international.
- Di Marco, M., Chapman, S., Althor, G., Kearney, S., Besancon, C., Butt, N., Maina, J.M., Possingham, H.P., von Bieberstein, K.R., Venter, O. and Watson, J.E., 2017. Changing trends and persisting biases in three decades of conservation science. *Global Ecology and Conservation*, 10, 32-42.
- Disney, M.I., Boni Vicari, M., Burt, A., Calders, K., Lewis, S.L., Raunonen, P. and Wilkes, P., 2018. Weighing trees with lasers: advances, challenges and opportunities. *Interface Focus*, 8(2), p.20170048.
- Doughty, C.E., Metcalfe, D.B., Girardin, C.A.J., Amézquita, F.F., Cabrera, D.G., Huasco, W.H., Silva-Espejo, J.E., Araujo-Murakami, A., Da Costa, M.C., Rocha, W., Feldpausch, T.R. et al., 2015. Drought impact on forest carbon dynamics and fluxes in Amazonia. *Nature*, 519, 78-82.
- Drew, A.P., Boley, J.D., Zhao, Y., Wadsworth, F.H., 2009. Sixty-two years of change in subtropical wet forest structure and composition at El Verde, Puerto Rico. *Interciencia*, 34, 34-40.
- DRYFLOR, 2016. Plant diversity patterns and their conservation implications in neotropical dry forests. *Science*, 353, 1383-1387.
- Duncanson, L., Armston, J., Disney, M., Avitabile, V., Barbier, N., Calders, K., Carter, S., Chave, J., Herold, M., Crowther, T.W., Falkowski, M. et al., 2019. The importance of consistent global forest aboveground biomass product validation. *Surveys in Geophysics*, 40, 979-999.
- Elias, F., Ferreira, J., Lennox, G.D., Berenguer, E., Ferreira, S., Schwartz, G., Melo, L.D.O., Reis Junior, D.N., Nascimento, R.O., Ferreira, F.N. and Espirito-Santo, F., 2020. Assessing the growth and climate sensitivity of secondary forests in highly deforested Amazonian landscapes. *Ecology*, 101(3), p.e02954.
- Espírito-Santo F.D., Gloor M., Keller M., Malhi Y., Saatchi S., Nelson B., Junior R.C., Pereira C., Lloyd J. et al. 2014. Size and frequency of natural forest disturbances and the Amazon forest carbon balance. *Nature Communications*, 5:3434.
- Esquivel-Muelbert, A., Baker, T.R., Dexter, K.G., Lewis, S.L., ter Steege, H., Lopez-Gonzalez, G., Monteagudo Mendoza, A., Brien, R., Feldpausch, T.R., Pitman, N., Alonso, A. et al. 2017. Seasonal drought limits tree species across the Neotropics. *Ecography*, 40, 618-629.
- Esquivel-Muelbert, A., Baker, T.R., Dexter, K.G., Lewis, S.L., Brien, R.J., Feldpausch, T.R., Lloyd, J., Monteagudo-Mendoza, A., Arroyo, L., Álvarez-Dávila, E., Higuchi, N. et al. 2019. Compositional response of Amazon forests to climate change. *Global Change Biology*, 25, 39-56.

- Fadrique, B., Báez, S., Duque, Á., Malizia, A., Blundo, C., Carilla, J., Osinaga-Acosta, O., Malizia, L., Silman, M., Farfán-Ríos, W. et al., 2018. Widespread but heterogeneous responses of Andean forests to climate change. *Nature* 564, 207-212.
- Fauset, S., Johnson, M.O., Gloor, M., Baker, T.R., Monteagudo, A., Brien, R.J., Feldpausch, T.R. et al., 2015. Hyperdominance in Amazonian forest carbon cycling. *Nature Communications* 6:6857.
- Feldpausch, T.R., Banin, L., Phillips, O.L., Baker, T.R., Lewis, S.L., Quesada, C.A., Affum-Baffoe, K., Arets, E.J., Berry, N.J., Bird, M. et al. 2011. Height-diameter allometry of tropical forest trees. *Biogeosciences*, 8, 1081-1106.
- Feldpausch, T.R., Lloyd, J., Lewis, S.L., Brien, R.J., Gloor, M., Monteagudo Mendoza, A., Lopez-Gonzalez, G., Banin, L., et al. 2012. Tree height integrated into pantropical forest biomass estimates. *Biogeosciences*, 9, 3381-3403.
- Feldpausch, T.R., Phillips, O.L., Brien, R.J.W., Gloor, E., Lloyd, J., Lopez-Gonzalez, G., Monteagudo-Mendoza, A., Malhi, Y., Alarcón, A., Dávila, E.Á. and Alvarez-Loayza, P., 2016. Amazon forest response to repeated droughts. *Global Biogeochemical Cycles*, 30, 964-982.
- Ferreira, J., Lennox, G.D., et al. 2018. Carbon-focused conservation may fail to protect the most biodiverse tropical forests. *Nature Climate Change*. DOI 10.1038/s41558-018-0225-7.
- Forest Carbon Partnership Facility (FCPF) Carbon Fund, 2017. *Emission Reductions Programme Document (ER-PD)*, *Ghana Cocoa Forest REDD+ Programme (GCFRP)*. World Bank, 339 pp.
- Fyllas, N.M., Patiño, S., Baker, T.R., Bielefeld Nardoto, G., Martinelli, L.A., Quesada, C.A., Paiva, R., Schwarz, M. et al., 2009. Basin-wide variations in foliar properties of Amazonian forest: phylogeny, soils and climate. *Biogeosciences* 6, 2677-708.
- Gatti, L.V., Gloor, M., Miller, J.B., Doughty, C.E., Malhi, Y., Domingues, L.G., Basso, L.S. et al. 2014. Drought sensitivity of Amazonian carbon balance revealed by atmospheric measurements. *Nature*, 506, 76-80.
- Gaubert, B., Stephens, B.B., Basu, S., Chevallier, F., Deng, F., Kort, E.A., Patra, P.K., Peters, W. et al., 2019. Global atmospheric CO₂ inverse models converging on neutral tropical land exchange, but disagreeing on fossil fuel and atmospheric growth rate. *Biogeosciences*, 16, 117-134.
- Gentry, A.H., 1988a, Tree species richness of upper Amazonian forests. *Proceedings of the National Academy of Sciences*, 85, 156-159.
- Gentry, A.H., 1988b, Changes in plant community diversity and floristic composition on environmental and geographical gradients. *Annals of the Missouri Botanical Garden*, 75, 1-34.
- Gentry, A.H. and Vasquez, R., 1993. *A field guide to the families and genera of woody plants of northwest South America (Colombia, Ecuador, Peru): with supplementary notes on herbaceous taxa*.
- Gloor, M., Phillips, O.L., Lloyd, J.J., Lewis, S.L., Malhi, Y., Baker, T.R., Lopez-Gonzalez, G., et al. 2009. Does the disturbance hypothesis explain the biomass increase in basin-wide Amazon forest plot data? *Global Change Biology*, 15, 2418-2430.
- Gosline, G., Marshall, A.R., Larridon, I., 2019. Revision and new species of the African genus *Mischogyne* (Annonaceae). *Kew Bulletin*, 74, 28.

- Gourlet-Fleury, S., F. Mortier, A. Fayolle, F. Baya, D. Ouédraogo, F. Bénédet, and N. Picard, 2013. Tropical forest recovery from logging: a 24 year silvicultural experiment from Central Africa. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368, no. 1625: 20120302.
- Harley, P., Vasconcellos, P., Vierling, L., Pinheiro, C.C.D.S., Greenberg, J., Guenther, A., Klinger, L., Almeida, S.S.D., Neill, D., Baker, T., Phillips, O., Malhi, Y. 2004. Variation in potential for isoprene emissions among Neotropical forest sites. *Global Change Biology*, 10, 630-650.
- Honorio Coronado E.N., Baker T.R., Phillips O.L., Pitman N.C., Pennington R.T., Vásquez Martinez R., Monteagudo, A. et al., 2009. Multi-scale comparisons of tree composition in Amazonian terra firme forests. *Biogeosciences* 30, 2719-31.
- Hubau, W., Lewis, S.L., Phillips, O.L., Affum-Baffoe K., Beeckman H., Cuní-Sánchez A., Daniels A.K., Ewango C.E.N. et al. 2020. Asynchronous carbon sink saturation in the world's largest tropical forests. *Nature* 579, 80-87.
- Hubbell, S.P., 1979. Tree dispersion, abundance, and diversity in a tropical dry forest. *Science*, 203, 1299-1309.
- Hubbell, S.P., Foster, R.B., 1983. Diversity of canopy trees in a neotropical forest and implications for the conservation of tropical trees, In *Tropical Rainforest Ecology and Management*. Eds. S.J. Sutton, T.C. Whitmore, A.C. Chadwick, pp. 25-41. Blackwell, Oxford.
- Johnson, M.O., Galbraith, D., Gloor, M., De Deurwaerder, H., Guimberteau, M., Rammig, A., Thonicke, K. et al., 2016, Variation in stem mortality rates determines patterns of above-ground biomass in Amazonian forests: implications for dynamic global vegetation models. *Global Change Biology* 22, 3996-4013.
- Joint Research Centre (2003) *Global Land Cover 2000 database*. European Commission, Joint Research Centre. https://forobs.jrc.ec.europa.eu/products/glc2000/data_access.php
- Jucker, T., Bongalov, B., Burslem, D.F., Nilus, R., Dalponte, M., Lewis, S.L., Phillips, O.L., Qie, L., Coomes, D.A., 2018. Topography shapes the structure, composition and function of tropical forest landscapes. *Ecology Letters* 21, 989-1000.
- Kohyama, T.S., Kohyama, T.I. and Sheil, D., 2019. Estimating net biomass production and loss from repeated measurements of trees in forests and woodlands: Formulae, biases and recommendations. *Forest Ecology and Management*, 433, 729-740.
- Lawrence, A., Phillips, O.L., Ismodes, A.R., Lopez, M., Rose, S., Wood, D. and Farfan, A.J., 2005. Local values for harvested forest plants in Madre de Dios, Peru: towards a more contextualised interpretation of quantitative ethnobotanical data. *Biodiversity & Conservation*, 14, 45-79.
- Levis, C., Costa, F.R., Bongers, F., Peña-Claros, M., Clement, C.R., Junqueira, A.B., Neves, E.G., Tamanaha, E.K., Figueiredo, F.O., Salomão, R.P. and Castilho, C.V., 2017. Persistent effects of pre-Columbian plant domestication on Amazonian forest composition. *Science*, 355, 925-931.
- Lewis, S.L., Phillips, O.L., Sheil, D., Vinceti, B., Baker, T.R., Brown, S., Graham, A.W., Higuchi, N., Hilbert, D.W., et al., 2004a. Tropical forest tree mortality, recruitment and turnover rates: calculation, interpretation and comparison when census intervals vary. *Journal of Ecology*, 92, 929-944.
- Lewis, S.L., Phillips, O.L., Baker, T.R., Lloyd, J., Malhi, Y., Almeida, S., Higuchi, N., Laurance, W.F., Neill, D.A., Silva, J.N.M. and Terborgh, J., 2004b. Concerted changes in tropical forest structure and dynamics: evidence from 50 South American long-term plots. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 359, 421-436.

- Lewis, S.L., Malhi, Y., Phillips, O.L., 2004. Fingerprinting the impacts of global change on tropical forests. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 359, 437-462.
- Lewis, S.L., Lopez-Gonzalez, G., Sonké, B., Affum-Baffoe, K., Baker, T.R., Ojo, L.O., Phillips, O.L., Reitsma, J.M., White, L., Comiskey, J.A., Ewango, C.E., et al. 2009. Increasing carbon storage in intact African tropical forests. *Nature*, 457, 1003-1006.
- Lewis S.L., Sonké, B., Sunderland, T., Begne, S.K., Lopez-Gonzalez, G., van der Heijden, G.M.F., Phillips, O.L. et al., 2013. Above-ground biomass and structure of 260 African tropical forests. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368, 20120295.
- López-Gonzalez G., Lewis, S.L., Burkitt, M., Phillips, O.L., 2011. ForestPlots.net: a web application and research tool to manage and analyse tropical forest plot data. *Journal of Vegetation Science* 22, 610-613.
- López-González, G. and Phillips, O.L., 2012. Estudiando el Amazonas: la experiencia de la Red Amazónica de Inventarios Forestales. *Revista Ecosistemas*, 21, 1-2.
- López-Gonzalez G., Sullivan, M.J.P., Baker, T.R., 2015. BiomasaFP: Tools for analysing data downloaded from ForestPlots.net. R package version 1.1.
- Malhi, Y., Baker, T.R., Phillips, O.L., Almeida, S., Alvarez, E., Arroyo, L., Chave, J., Czimczik, C.I., Fiore, A.D., Higuchi, N. and Killeen, T.J., 2004. The above-ground coarse wood productivity of 104 Neotropical forest plots. *Global Change Biology*, 10, 563-591.
- Malhi, Y., Girardin, C., Metcalfe, D., Doughty, C., Aragão, L.E.O., Rifai, S., Shenkin, A., Aguirre Gutierrez, J., Dahlsjö, C. et al. The Global Ecosystems Monitoring network: monitoring ecosystem productivity and carbon cycling across the tropics. *Biological Conservation* (this volume)
- Malhi, Y., Phillips, O.L., Lloyd, J., Baker, T., Wright, J., Almeida, S., Arroyo, L., Frederiksen, T., Grace, J., Higuchi, N. et al., 2002. An international network to monitor the structure, composition and dynamics of Amazonian forests (RAINFOR). *Journal of Vegetation Science* 13, 439-50.
- Malhi Y, Wood D, Baker TR, Wright J, Phillips OL, Cochrane T, Meir P, Chave J, Almeida S, Arroyo L et al., 2006. The regional variation of aboveground live biomass in old-growth Amazonian forests. *Global Change Biology* 12, 1107-1138.
- Malhi, Y., Farfán Amézquita, F., Doughty, C.E., Silva-Espejo, J.E., Girardin, C.A., Metcalfe, D.B., Aragão, L.E., Huaraca-Quispe, L.P., et al. 2014. The productivity, metabolism and carbon cycle of two lowland tropical forest plots in south-western Amazonia, Peru. *Plant Ecology & Diversity* 7, 85-105.
- Malhi, Y., Gardner, T.A., Goldsmith, G.R., Silman, M.R. and Zelazowski, P., 2014. Tropical forests in the Anthropocene. *Annual Review of Environment and Resources*, 39.
- Malizia, A, Blundo, C., Carilla, J., Osinaga-Acosta, O., Cuesta, F., Duque, A., Aguirre, N., Aguirre, Z., Ataroff, M., Baez, S. et al. (2020). Elevation and latitude drives structure and tree species composition in Andean forests: results from a large-scale plot Network. *PLoS ONE* (in press)
- Marimon B.S., Marimon-Junior B.H., Feldpausch T.R., Oliveira-Santos C., Mews H.A. et al., 2014. Disequilibrium and hyperdynamic tree turnover at the forest–cerrado transition zone in southern Amazonia. *Plant Ecology & Diversity* 7, 281-92.

- McDowell, N., Allen, C.D., Anderson-Teixeira, K., Brando, P., Brien, R., Chambers, J., Christoffersen, B., Davies, S., Doughty, C., Duque, A. and Espirito-Santo, F., 2018. Drivers and mechanisms of tree mortality in moist tropical forests. *New Phytologist*, 219, 851-869.
- Mitchard, E.T.A., Feldpausch, T.R., Brien, R.J.W., Lopez-Gonzalez, G., Monteagudo, A., Baker, T.R. et al., 2014. Markedly divergent estimates of Amazon forest carbon density from ground plots and satellites. *Global Ecology and Biogeography*. doi:10.1111/geb.12168.
- Monteagudo Mendoza A, Vásquez Martínez R., Rojas Gonzales R., Phillips O.L., Baker T.R., Dueñas Linares H., Pickavance G.C., Núñez Vargas P et al., 2020. Primer Catálogo de los Árboles de la Amazonía de Madre De Dios, Perú. Universidad Andina del Cusco, Cusco, Peru, 240pp.
- Nemani, R. R. et al., 2003. Climate-driven increases in global terrestrial net primary production from 1982 to 1999. *Science* 300, 1560–1563.
- Nogueira, D.S., Marimon, B.S., Marimon-Junior, B.H., Oliveira, E.A., Morandi, P., Reis, S.M., Elias, F., Neves, E.C., Feldpausch, T.R., Lloyd, J. and Phillips, O.L., 2019. Impacts of fire on forest biomass dynamics at the southern Amazon edge. *Environmental Conservation*, 46, 285-292.
- Norden, N., González-M, R., Avella-M, A., Salgado-Negret, B., Alcázar, C., Rodríguez-Buriticá, S., Aguilar-Cano, J., et al., 2020, Building a socio-ecological monitoring platform for the comprehensive management of tropical dry forests. *Plants, People, Planet*. <https://doi.org/10.1002/ppp3.10113>.
- Oliveira, E.A., Marimon-Junior, B.H., Iriarte, J., Morandi, P.S., Maezumi, S.Y., Nogueira, D.S., Aragão, L.E.O.C. and Feldpausch, T.R. 2020. Legacy of Amazonia Dark Earth soil on forest structure and species composition. *Global Ecology and Biogeography*, in press.
- Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V. Underwood, E.C., D'Amico, J.A., et al., 2001. Terrestrial ecoregions of the world: a new map of life on Earth. *Bioscience*, 51, 933-938.
- Pallqui, N.C., Monteagudo, A., Phillips, O.L., Lopez-Gonzalez, G., Cruz, L., Galiano, W., Chavez, W., Vasquez, R., 2014. Dinámica, biomasa aérea y composición florística en parcelas permanentes Reserva Nacional Tambopata, Madre de Dios, Perú. *Revista Peruana de Biología* 21, 235-42.
- Pan, Y., Birdsey, R.A., Fang, J., Houghton, R., Kauppi, P.E. et al., 2011. A large and persistent carbon sink in the world's forests. *Science*, 333, 988–993. <https://doi.org/10.1126/science.1201609>.
- Pan, Y., Birdsey, R.A., Phillips, O.L. and Jackson, R.B., 2013. The structure, distribution, and biomass of the world's forests. *Annual Review of Ecology, Evolution, and Systematics*, 44, 593-622.
- Patiño, S., Lloyd, J., Paiva, R., Baker, T.R., Quesada, C.A., Mercado, L.M. et al., 2009. Branch xylem density variations across the Amazon Basin. *Biogeosciences*, 6, 545-568.
- Peacock, J., Baker, T.R., Lewis, S.L., Lopez-Gonzalez, G., Phillips, O.L., 2007. The RAINFOR database: monitoring forest biomass and dynamics. *Journal of Vegetation Science*, 18, 535-42.
- Pennington, T., Lehmann, C.E., Rowland, L. 2018. Tropical savannas and dry forests. *Current Biology* 28, 541-45.
- Peres, C.A., Emilio, T., Schietti, J., Desmoulière, S.J. and Levi, T., 2016. Dispersal limitation induces long-term biomass collapse in overhunted Amazonian forests. *Proceedings of the National Academy of Sciences*, 113, 892-897.

- Phillips, O.L., 1996. Long-term environmental change in tropical forests: increasing tree turnover. *Environmental Conservation* 23, 235-248.
- Phillips, O. and Gentry, A.H., 1993. The useful plants of Tambopata, Peru: I. Statistical hypotheses tests with a new quantitative technique. *Economic Botany*, 47, 15-32.
- Phillips, O.L. and Gentry, A.H., 1994. Increasing turnover through time in tropical forests. *Science*, 263, 954-958.
- Phillips, O. L., P. Hall, A. H. Gentry, S. A. Sawyer and R. Vasquez, 1994. Dynamics and species richness of tropical rain forests. *Proceedings of the National Academy of Sciences of the United States of America* 91, 2805-2809.
- Phillips, O.L. and Raven, P.H., 1997. A strategy for sampling neotropical forests. *Neotropical Biodiversity and Conservation*, 1, 141-165.
- Phillips, O. and Sheil, D. 1997. Forest turnover, diversity and CO₂. *Trends in Ecology & Evolution* 12, 404-404.
- Phillips, O.L., Malhi, Y., Higuchi, N., Laurance, W.F., Núñez, P.V., Vásquez, R.M., Laurance, S.G., Ferreira, L.V., Stern, M., Brown, S. and Grace, J., 1998. Changes in the carbon balance of tropical forests: evidence from long-term plots. *Science*, 282, 439-442.
- Phillips, O.L., Baker, T., Feldspach, T., & Brien, R. J. W., 2002. Field manual for plot establishment and remeasurement (RAINFOR). Amazon Forest Inventory Network, Sixth Framework Programme (2002–2006). URL: <http://www.geog.leeds.ac.uk/projects/rainfor>.
- Phillips, O. and Miller, J.S., 2002, Global patterns of plant diversity: Alwyn H. Gentry's forest transect data set. *Monographs in Systematic Botany* 89:1-319. Missouri Botanical Press.
- Phillips, O.L., Baker, T.R., Arroyo, L., Higuchi, N., Killeen, T.J., Laurance, W.F., Lewis, S.L., Lloyd, J., Malhi, Y., Monteagudo, A. and Neill, D.A., 2004. Pattern and process in Amazon tree turnover, 1976–2001. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 359, 381-407.
- Phillips, O.L., Aragão, L.E., Lewis, S.L., Fisher, J.B., Lloyd, J., López-González, G., Malhi, Y., Monteagudo, A., Peacock, J., Quesada, C.A. and Van Der Heijden, G., 2009. Drought sensitivity of the Amazon rainforest. *Science*, 323, 1344-1347.
- Phillips, O.L. and Brien, R.J., 2017. Carbon uptake by mature Amazon forests has mitigated Amazon nations' carbon emissions. *Carbon Balance and Management*, 12, 1.
- Phillips, O.L., Sullivan, M.J., Baker, T.R., Mendoza, A.M., Vargas, P.N. and Vásquez, R. 2019. Species matter: wood density influences tropical forest biomass at multiple scales. *Surveys in Geophysics*, 404, 913-935.
- Piao, S., Wang, X., Park, T., Chen, C., Lian, X., He, Y., Bjerke, J.W., Chen, A., Ciais, P., Tømmervik, H. and Nemani, R.R., 2019. Characteristics, drivers and feedbacks of global greening. *Nature Reviews Earth & Environment*, pp.1-14.
- Pires, J.M. and Salomão, R.P. 2000. Dinâmica da diversidade arbórea de um fragmento de floresta tropical primária na Amazônia oriental - 1. Período: 1956 a 1992. *Bol. Mus. Para. Emílio goeldi, Sér. Botânica*, 16, 63-110.

- Pomeroy, M., R. B. Primack, and S. N. Rai. 2003. Changes in four rain forest plots of the Western Ghats, India, 1939–93. *Conservation and Society* 1, 113–135.
- Qie L, Lewis SL, Sullivan MJ, Lopez-Gonzalez G, Pickavance GC, Sunderland T, Ashton P, Hubau W et al., 2017, Long-term carbon sink in Borneo’s forests halted by drought and vulnerable to edge effects. *Nature Communications* 8:1966.
- Quesada CA, Phillips OL, Schwarz M, Czimczik CI, Baker TR, Patiño S, Fyllas NM, Hodnett MG, Herrera R et al., 2012. Basin-wide variations in Amazon forest structure and function are mediated by both soils and climate. *Biogeosciences* 9, 2203-2246. doi:10.5194/bg-9-2203-2012.
- Reboredo Segovia, A.L., Romano, D., Armsworth, P.R., 2020. Who studies where? Boosting tropical conservation research where it is most needed. *Frontiers in Ecology and the Environment*, 18, 159-166.
- Requena Suarez, D., Rozendaal, D.M., De Sy, V., Phillips, O.L., Alvarez-Dávila, E., Anderson-Teixeira, K., Araujo-Murakami, A., Arroyo, L., Baker, T.R., Bongers, F., et al., 2019. Estimating aboveground net biomass change for tropical and subtropical forests: Refinement of IPCC default rates using forest plot data. *Global Change Biology*, 25, 3609-3624.
- Reis, S.M., Marimon, B.S., Marimon-Junior, B.H., Morandi, P.S., Oliveira, E.A., Eias, F., Neves, E.C., Oliveira, B., Nogueira, D.S., Umetsu, R.K., Feldpausch, T.R. and Phillips, O.L., 2018. Climate and fragmentation affect forest structure at the Southern border of Amazonia. *Plant Ecology & Diversity*, 11, 13-25.
- Reitsma, J.M. 1988. *Forest vegetation of Gabon*. TROPENBOS Technical Series, ISBN: 90-5113-007-4.
- Ríos-Saldaña, C.A., Delibes-Mateos, M. and Ferreira, C.C., 2018. Are fieldwork studies being relegated to second place in conservation science? *Global Ecology and Conservation*, 14, p.e00389.
- Rowland, L., da Costa, A.C.L., Galbraith, D.R., Oliveira, R.S., Binks, O.J., Oliveira, A.A.R., Pullen, A.M., Doughty, C.E., Metcalfe, D.B., Vasconcelos, S.S. and Ferreira, L.V., 2015. Death from drought in tropical forests is triggered by hydraulics not carbon starvation. *Nature*, 528,119-122.
- Salomão, R.D., 1991. Uso de parcelas permanentes para estudos da vegetação da floresta tropical úmida. I. Município de Marabá, Pará. Boletim do Museu Paraense Emílio Goeldi. *Serie Botânica*, 7, 543-604.
- Schepaschenko, D., Chave, J., Phillips, O.L. et al., 2019. The Forest Observation System, building a global reference dataset for remote sensing of forest biomass. *Sci Data* 6, 198 (2019). <https://doi.org/10.1038/s41597-019-0196-1>
- Schietti, J., Martins, D., Emilio, T., Souza, P.F., Levis, C., Baccaro, F.B., et al. 2016. Forest structure along a 600 km transect of natural disturbances and seasonality gradients in central-southern Amazonia. *Journal of Ecology*, 104, 1335-1346.
- Sheil, D., 1995. Evaluating turnover in tropical forests. *Science*, 268, 894-896.
- Sheil, D., 1996. Species richness, tropical forest dynamics and sampling: questioning cause and effect. *Oikos*, 76, 587-590.
- Sist, P., Rutishauser, E., Peña-Claros, M., Shenkin, A., Hérault, B., Blanc, L., Baraloto, C., Baya, F., Benedet, F., da Silva, K.E., Descroix, L., et al. 2015. The Tropical managed Forests Observatory: a research network addressing the future of tropical logged forests. *Applied Vegetation Science*, 18, 171-174.

- 1409
1410 Sousa, T.R., Schietti, J., Coelho de Souza, F., Esquivel-Muelbert, A., Ribeiro, I.O., Emílio, T. et al.,
1411 2020. Palms and trees resist extreme drought in Amazon forests with shallow water tables. *Journal of*
1412 *Ecology*.
1413
1414 Sullivan, M.J., Talbot, J., Lewis, S.L., Phillips, O.L., Qie, L., Begne, S.K., Chave, J., Cuni-Sanchez, A.,
1415 Hubau, W., Lopez-Gonzalez, G. and Miles, L., 2017. Diversity and carbon storage across the tropical
1416 forest biome. *Scientific Reports*, 7, 1-12.
1417
1418 Sullivan, M.J., Lewis, S.L., Hubau, W., Qie, L., Baker, T.R., Banin, L.F., Chave, J., Cuni-Sanchez, A.,
1419 Feldpausch, T.R., et al. 2018. Field methods for sampling tree height for tropical forest biomass
1420 estimation. *Methods in Ecology and Evolution*, 9, 1179-1189.
1421
1422 Sullivan, M.J.P., Lewis, S.L., Affum-Baffoe, K., Castilho, C., Costa, F., Cuni Sanchez, A., Ewango,
1423 C.E.N., Hubau, W., Marimon, B., Monteagudo-Mendoza, A., et al. 2020. Long-term thermal sensitivity
1424 of Earth's tropical forests. *Science* 368, 869-874. DOI: 10.1126/science.aaw7578.
1425
1426 Taylor, J.A. and Lloyd, J., 1992. Sources and sinks of atmospheric CO₂. *Australian Journal of Botany*
1427 40, 407-418.
1428
1429 ter Steege, H., Pitman, N., Sabatier, D., Castellanos, H., Van Der Hout, P., Daly, D.C., Silveira, M., et
1430 al., 2003. A spatial model of tree α -diversity and tree density for the Amazon. *Biodiversity &*
1431 *Conservation*, 12, 2255-2277.
1432
1433 ter Steege, H., Pitman, N.C., Phillips, O.L., Chave, J., Sabatier, D., Duque, A., Molino, J.F., Prévost,
1434 M.F., Spichiger, R., Castellanos, H., Von Hildebrand, P., 2006. Continental-scale patterns of canopy
1435 tree composition and function across Amazonia. *Nature* 443, 444-447.
1436
1437 ter Steege, H., Pitman, N.C., Sabatier, D., Baraloto, C., Salomão, R.P., Guevara, J.E., Phillips, O.L.,
1438 Castilho CV et al., 2013. Hyperdominance in the Amazonian tree flora. *Science* 342, 1243092.
1439
1440 Terrer, C. et al., 2019. Nitrogen and phosphorus constrain the CO₂ fertilization of global plant biomass.
1441 *Nature Climate Change* 9, 684–689.
1442
1443 Vargas, R., Alcaraz-Segura, D., Birdsey, R., Brunsell, N.A., Cruz-Gaistardo, C.O., de Jong, B.,
1444 Etchevers, J., Guevara, M., Hayes, D.J., Johnson, K. and Loescher, H.W., 2017. Enhancing
1445 interoperability to facilitate implementation of REDD+: Case study of Mexico. *Carbon Management*,
1446 8, 57-65.
1447
1448 Vásquez, R., Rojas, R., Monteagudo, A.M., Valenzuela, L.G., Huamantupa, I., 2018. *Catalogo de los*
1449 *Arboles del Perú*. Q'ueña Revista de la Sociedad Botánica del Cusco 9 (1), número especial.
1450
1451 Vásquez, M. R. & Soto Shareva, Y. 2020. *Virola pseudosebifera* (Myristicaceae), una nueva especie
1452 para la Selva Alta del Perú. *Revista Q'ueña*, 10, 07-12.
1453
1454 Vicuña Miñano, E., Baker, T.R., Banda, K., Coronado, E.H., Monteagudo, A., Phillips, O.L., Torres,
1455 D.D.C., Rios, W.F., Flores, G., Huaman, D. and Huaman, K.T., 2018. El sumidero de carbono en los
1456 bosques primarios Amazónicos es una oportunidad para lograr la sostenibilidad de su conservación.
1457 *Folia Amazónica*, 27, 101-109.
1458
1459 Vilanova, E., Ramírez-Angulo, H., Torres-Lezama, A., Aymard, G., Gámez, L., Durán, C., Hernández,
1460 L., Herrera, R., van der Heijden, G., Phillips, O.L. and Ettl, G.J., 2018. Environmental drivers of forest
1461 structure and stem turnover across Venezuelan tropical forests. *PloS one*, 13(6).
1462

- 1463 Werger, M.J.A. (ed.). 2011. *Sustainable Management of Tropical Rainforests: the CELOS Management*
1464 *System*. Tropenbos International, Paramaribo, Suriname.
- 1465
- 1466 Wright, S.J. (this volume). The Smithsonian Tropical Research Institute: ecological and applied
1467 research. *Biological Conservation*.
- 1468
- 1469 Wurdack, K.J. and Farfan-Rios, W., 2017. *Incadendron*: a new genus of Euphorbiaceae tribe
1470 Hippomaneae from the sub-Andean cordilleras of Ecuador and Peru. *PhytoKeys*, 85, 69-86.
- 1471

1472 Taking the Pulse of Earth's Tropical Forests using Networks of Highly Distributed Plots

1473 Author: ForestPlots.net (Cecilia Blundo, Julieta Carilla, Ricardo Grau, Hieu

1474 Dang Tran)

1475

1476 Table of Contributing Authors:

Country	First Name	Last Name	Institution
Argentina	Cecilia	Blundo	Instituto de Ecología Regional (IER); Universidad Nacional de Tucumán (UNT); Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)
Argentina	Julieta	Carilla	Instituto de Ecología Regional (IER); Universidad Nacional de Tucumán (UNT); Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)
Argentina	Ricardo	Grau	Instituto de Ecología Regional (IER); Universidad Nacional de Tucumán (UNT); Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)
Argentina	Agustina	Malizia	Instituto de Ecología Regional (IER); Universidad Nacional de Tucumán (UNT); Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)
Argentina	Lucio	Malizia	Facultad de Ciencias Agrarias, Universidad Nacional de Jujuy. Jujuy, Argentina.
Argentina	Oriana	Osinaga-Acosta	Instituto de Ecología Regional (IER); Universidad Nacional de Tucumán (UNT); Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)
Australia	Michael	Bird	James Cook University (JCU)
Australia	Matt	Bradford	CSIRO (Commonwealth Scientific and Industrial Research Organisation)
Australia	Damien	Catchpole	School of Land & Food University of Tasmania
Australia	Andrew	Ford	CSIRO (Commonwealth Scientific and Industrial Research Organisation)
Australia	Andrew	Graham	CSIRO Tropical Forest Research Centre
Australia	David	Hilbert	
Australia	Jeanette	Kemp	Environmental Protection Agency (EPA)
Australia	Susan	Laurance	Centre for Tropical Environmental and Sustainability Science (TESS) and College of Marine and Environmental Sciences, James Cook University
Australia	William	Laurance	Centre for Tropical Environmental and Sustainability Science (TESS) and College of Marine and Environmental Sciences, James Cook University
Australia	Francoise	Yoko Ishida	Centre for Tropical Environmental and Sustainability Science, College of Science and Engineering, James Cook University, Cairns, Australia
Australia	Andrew	Marshall	University of the Sunshine Coast; University of York; Flamingo Land Ltd.
Austria	Hannsjoerg	Woell	Sommersbergseestrasse
Belgium	Jean-Francois	Bastin	Ghent University
Belgium	Hans	Beeckman	Royal Museum for Central Africa - Service of Wood Biology
Belgium	Pascal	Boeckx	Isotope Bioscience Laboratory-ISOFYS Ghent University
Belgium	Jan	Bogaert	Gembloux Agro-Bio Tech, Université de Liege
Belgium	Charles	De Canniere	Landscape Ecology and Vegetal Production Systems Unit
Belgium	Thales	de Haulleville	CAVElab Computational & Applied Vegetation Ecology, Ghent University
Belgium	Jean-Louis	Doucet	Tropical Forestry, Forest Ressources Management, Gembloux Agro-Bio Tech, University of Liege
Belgium	Olivier	Hardy	Université Libre de Bruxelles (ULB) Evolutionary Biology and Ecology
Belgium	Wannes	Hubau	Royal Museum for Central Africa, Belgium
Belgium	Elizabeth	Kearsley	Royal Museum for Central Africa / Ghent University
Belgium	Hans	Verbeeck	Department of Environment Ghent University
Belgium	Jason	Vleminckx	Service Evolution Biologique et Ecologie
Belize	Steven W.	Brewer	Belize Foundation for Research and Environmental Education
Bolivia	Alfredo	Alarcón	IBIF
Bolivia	Alejandro	Araujo-Murakami	Museo de Historia Natural Noel Kempff Mercado, Universidad Autonoma Gabriel Rene Moreno

Bolivia	Eric	Arets	PROMAB
Bolivia	Luzmila	Arroyo	Museo de Historia Natural Noel Kempff Mercado, Universidad Autonoma Gabriel Rene Moreno
Bolivia	Ezequiel	Chavez	Museo Noel Kempff
Bolivia	Todd	Fredericksen	IBIF
Bolivia	René	Guillén Villaroel	Consultor independiente
Bolivia	Gloria	Gutierrez Sibauty	Jardin Botanico Municipal de Santa Cruz
Bolivia	Timothy	Killeen	Museo de Historia Natural Noel Kempff Mercado
Bolivia	Juan Carlos	Licona	IBIF
Bolivia	John	Lleigue	PROMAB
Bolivia	Casimiro	Mendoza	Forest Management in Bolivia
Bolivia	Samaria	Murakami	PROMAB
Bolivia	Alexander	Parada Gutierrez	Museo de Historia Natural Noel Kempff Mercado, Universidad Autónoma Gabriel Rene Moreno
Bolivia	Guido	Pardo	Universidad Autónoma del Beni Riberalta
Bolivia	Marielos	Peña-Claros	PROMAB
Bolivia	Lourens	Poorter	PROMAB
Bolivia	Marisol	Toledo	Museo de Historia Natural Noel Kempff
Bolivia	Jeanneth	Villalobos Cayo	Herbario del Sur de Bolivia
Bolivia	Laura Jessica	Viscarra	Museo de Historia Natural Noel Kempff Mercado
Bolivia	Vincent	Vos	Universidad Autónoma del Beni
Brazil	Jorge	Ahumada	Conservation International
Brazil	Everton	Almeida	Instituto de Biodiversidade e Floresta, Universidade Federal do Oeste do Pará
Brazil	Jarcilene	Almeida	Universidade Federal de Pernambuco
Brazil	Edmar	Almeida de Oliveira	Universidade do Estado de Mato Grosso
Brazil	Wesley	Alves da Cruz	Universidade do Estado de Mato Grosso (UNEMAT)
Brazil	Atila	Alves de Oliveira	Projeto TEAM – Manaus
Brazil	Fabício	Alvim Carvalho	Universidade Federal de Juiz de Fora (UFJF)
Brazil	Flávio	Amorim Obermuller	Universidade Federal do Rio de Janeiro
Brazil	Ana	Andrade	Instituto Nacional de Pesquisas da Amazônia, Projeto Dinâmica Biológica de Fragmentos Florestais
Brazil	Fernanda	Antunes Carvalho	Departamento de Genética, Ecologia e Evolução. Universidade Federal de Minas Gerais
Brazil	Simone	Aparecida Vieira	Universidade Estadual de Campinas
Brazil	Ana Carla	Aquino	Laboratório de Ecologia de Comunidades e Funcionamento de Ecossistemas- ECoFERP, Departamento de Biologia, Faculdade de Filosofia, Ciências e Letras - USP- Ribeirão Preto-SP
Brazil	Luiz	Aragão	National Institute for Space Research (INPE)
Brazil	Ana Claudia	Araújo	Universidade Federal de Roraima (UFRR)
Brazil	Marco Antonio	Assis	UNESP - São Paulo State University
Brazil	Jose	Ataliba Mantelli Aboin Gomes	Carbonozero Consultoria Ambiental
Brazil	Fabício	Baccaro	Departamento de Biologia, Universidade Federal do Amazonas (UFAM)
Brazil	Plínio	Barbosa de Camargo	Centro de Energia Nuclear na Agricultura, Universidade de São Paulo
Brazil	Paulo	Barni	UERR - Campus Rorainópolis
Brazil	Jorcely	Barroso	Universidade Federal do Acre
Brazil	Luis Carlos	Bernacci	Instituto Agronômico de Campinas
Brazil	Kauane	Bordin	Universidade Federal do Rio Grande do Sul

Brazil	Marcelo	Brilhante de Medeiros	Embrapa
Brazil	Igor	Broggio	Universidade Estadual do Norte Fluminense (UENF)
Brazil	José Luís	Camargo	Instituto Nacional de Pesquisas da Amazônia, Projeto Dinâmica Biológica de Fragmentos Florestais
Brazil	Domingos	Cardoso	Universidade Federal da Bahia (UFBA)
Brazil	Maria	Carmozina Araújo	TBC
Brazil	Maria	Carniello	Universidade do Estado de Mato Grosso (UNEMAT)
Brazil	Andre Luis	Casarin Rochelle	Instituto de Biologia, Universidade Estadual de Campinas
Brazil	Carolina	Castilho	Embrapa Roraima
Brazil	Antonio Alberto Jorge Farias	Castro	Universidade Federal do Piauí, Teresina (UFPI)
Brazil	Wendeson	Castro	Botany and Plant Ecology Laboratory, Federal University of Acre
Brazil	Sabina	Cerruto Ribeiro	Universidade Federal do Acre
Brazil	Flávia	Costa	INPA- National Institute of Amazonian Research
Brazil	Rodrigo	Costa de Oliveira	UERR - Campus Boa Vista
Brazil	Italo	Coutinho	Universidade Federal do Ceará
Brazil	Sandra	Cristina Müller	Universidade Federal do Rio Grande do Sul
Brazil	John	Cunha	Universidade Federal de Campina Grande
Brazil	Lola	da Costa	Universidade Federal do Para
Brazil	Lucia	da Costa Ferreira	Núcleo de Estudos e Pesquisas Ambientais, Universidade Estadual de Campinas
Brazil	Richarly	da Costa Silva	Instituto Federal de Educação, Ciência e Tecnologia do Acre
Brazil	Marta	da Graça Zacarias Simbine	Universidade Estadual de Campinas
Brazil	Vitor	de Andrade Kamimura	UNESP - São Paulo State University
Brazil	Haroldo Cavalcante	de Lima	Instituto de Pesquisas Jardim Botânico do Rio de Janeiro
Brazil	Lia	de Oliveira Melo	Universidade Federal do Oeste do Pará
Brazil	Luciano	de Queiroz	UEFS - Depto. de Ciências Biológicas
Brazil	Jano	de Rutte Corzo	TBC
Brazil	José Romualdo	de Sousa Lima	Universidade Federal do Agreste de Pernambuco (UFAPE)
Brazil	Mário	do Espírito Santo	Universidade Estadual de Montes Claros
Brazil	Tomas	Domingues	FFCLRP-USP/Br
Brazil	Nayane Cristina	dos Santos Prestes	UNEMAT
Brazil	Steffan	Eduardo Silva Carneiro	Universidade Federal de Jataí
Brazil	Fernando	Elias	Universidade Federal do Pará, Instituto de Ciências Biológicas
Brazil	Gabriel	Eliseu	Universidade Federal de Jataí
Brazil	Thaise	Emilio	Universidade de Campinas
Brazil	Camila Laís	Farrapo	Universidade Federal de Lavras (UFLA)
Brazil	Letícia	Fernandes	Universidade Federal do Acre
Brazil	Gustavo	Ferreira	Universidade Federal de Jataí
Brazil	Joice	Ferreira	Embrapa
Brazil	Leandro	Ferreira	Museu Goeldi
Brazil	Socorro	Ferreira	Embrapa Amazônia Oriental
Brazil	Marcelo	Fragomeni Simon	EMBRAPA
Brazil	Maria Aparecida	Freitas	Instituto Nacional de Pesquisas da Amazônia
Brazil	Queila S	García	UFMG - Universidade Federal de Minas Gerais

Brazil	Angelo	Gilberto Manzatto	Fundação Universidade Fedral de Rondônia - UNIR
Brazil	Paulo	Graça	INPA- Instituto Nacional de Pesquisas Amazônicas
Brazil	Frederico	Guilherme	Universidade Federal de Jatai
Brazil	Luis	Gustavo Canesi Ferreira	TBC
Brazil	Eduardo	Hase	Instituto Nacional de Pesquisas da Amazônia
Brazil	Niro	Higuchi	Instituto Nacional de Pesquisas da Amazônia - Coordenação de Pesquisas em Silvicultura Tropical
Brazil	Mariana	Iguatemy	Jardim Botânico do Rio de Janeiro
Brazil	Reinaldo	Imbrozio Barbosa	National Institute for Research in Amazonia
Brazil	Margarita	Jaramillo	Universidade Federal de Roraima (UFRR/PRONAT)
Brazil	Carlos	Joly	Universidade Estadual de Campinas/UNICAMP
Brazil	Joice	Klipel	Universidade Federal do Rio Grande do Sul (UFRGS)
Brazil	Iêda	Leão do Amaral	Instituto Nacional de Pesquisas da Amazônia/CPBO
Brazil	Carolina	Levis	Universidade Federal de Santa Catarina (UFSC)
Brazil	Antonio S	Lima	Museu Goeldi
Brazil	Maurício	Lima Dan	INCAPER- Instituto Capixaba de Pesquisa, Assistência Técnica e Extensão Rural
Brazil	Aline	Lopes	INPE- Instituto Nacional de Pesquisas Espaciais
Brazil	Herison	Madeiros	Universidade de São Paulo
Brazil	William E.	Magnusson	Instituto Nacional de Pesquisas da Amazônia (INPA)
Brazil	Rubens	Manoel dos Santos	Universidade Federal de Lavras (UFLA)
Brazil	Beatriz	Marimon	Universidade do Estado de Mato Grosso
Brazil	Ben Hur	Marimon Junior	Universidade do Estado de Mato Grosso
Brazil	Roberta	Marotti Martelletti Grillo	Instituto de Biociências, Universidade Estadual Paulista
Brazil	Luiz	Martinelli	Centro de Energia Nuclear na Agricultura, Universidade de São Paulo
Brazil	Simone	Matias Reis	Universidade do Estado de Mato Grosso
Brazil	Salomão	Medeiros	Semiarid National Institute (INSA)
Brazil	Milton	Meira-Junior	Universidade de Brasília, Departamento de Engenharia Florestal
Brazil	Thiago	Metzker	IBAM - Instituto Bem Ambiental
Brazil	Paulo	Morandi	Universidade do Estado de Mato Grosso, Campus de Nova Xavantina
Brazil	Natanael	Moreira do Nascimento	Universidade Federal de Jatai
Brazil	Magna	Moura	EMBRAPA
Brazil	Sandra	Müller	Universidade Federal do Rio Grande do Sul
Brazil	Laszlo	Nagy	University in Campinas
Brazil	Henrique	Nascimento	INPA
Brazil	Marcelo	Nascimento	Universidade Estadual do Norte Fluminense Darcy Ribeiro (UENF)
Brazil	Adriano	Nogueira Lima	LMF/INPA
Brazil	Raimunda	Oliveira de Araújo	INPA
Brazil	Jhonathan	Oliveira Silva	Universidade Federal do Vale do São Francisco
Brazil	Marcelo	Pansonato	USP- University of São Paulo
Brazil	Gabriel	Pavan Sabino	UNESP - São Paulo State University
Brazil	Karla Maria	Pedra de Abreu	Instituto Federal do Espírito Santo (Ifes)
Brazil	Pablo José Francisco	Pena Rodrigues	Instituto de Pesquisas Jardim Botânico do Rio de Janeiro
Brazil	Maria	Piedade	Instituto Nacional de Pesquisas da Amazônia - INPA - Grupo MAUA
Brazil	José	Roberto Rodrigues Pinto	Universidade de Brasília, Departamento de Engenharia Florestal
Brazil	Carlos	Quesada	Instituto Nacional de Pesquisas da Amazônia
Brazil	Eliana	Ramos	Instituto Nacional da Mata Atlântica

Brazil	Rafael	Ramos	Universidade Estadual de Campinas
Brazil	Priscyla	Rodrigues	Universidade Federal do Vale do São Francisco
Brazil	Thaiane	Rodrigues de Sousa	RAINFOR-PPBIO
Brazil	Rafael	Salomão	Universidade Federal Rural da Amazônia - UFRA/CAPES
Brazil	Flávia	Santana	INPA- National Institute of Amazonian Research
Brazil	Marcos	Scaranello	Instituto de Biologia, Universidade Estadual de Campinas
Brazil	Rodrigo	Scarton Bergamin	Universidade Federal do Rio Grande do Sul
Brazil	Juliana	Schietti	Universidade Federal do Amazonas (UFAM)
Brazil	Jochen	Schöngart	INPA/ Max-Planck Project
Brazil	Gustavo	Schwartz	EMBRAPA- Empresa Brasileira de Pesquisa Agropecuária (Amazônia Oriental)
Brazil	Natalino	Silva	Serviço Florestal Brasileiro
Brazil	Marcos	Silveira	Museu Universitário, Universidade Federal do Acre
Brazil	Cristiana	Simão Seixas	Universidade Estadual de Campinas
Brazil	Marta	Simbine	Instituto de Biologia, Universidade Estadual de Campinas
Brazil	Ana Claudia	Souza	UNESP - São Paulo State University
Brazil	Priscila	Souza	Instituto Nacional de Pesquisas da Amazônia (INPA)
Brazil	Rodolfo	Souza	Universidade Federal Rural de Pernambuco
Brazil	Tereza	Sposito	IBAM - Instituto Bem Ambiental
Brazil	Edson	Stefani Junior	Instituto de Biologia, Universidade Estadual de Campinas
Brazil	Julio Daniel do	Vale	PUCPR - Pontifícia Universidade Católica do Paraná
Brazil	Alberto	Vicentini	INPA
Brazil	Ima Célia Guimarães	Vieira	Museu Paraense Emilio Goeldi
Brazil	Dora	Villela	Universidade Estadual do Norte Fluminense Darcy Ribeiro (UENF)
Brazil	Marcos	Vital	Universidade Federal de Roraima
Brazil	Haron	Xaud	Embrapa Roraima
Brazil	Katia	Zanini	Universidade Federal do Rio Grande do Sul
Brazil	Charles Eugene	Zartman	Instituto Nacional de Pesquisas Amazônicas
Brunei	Faizah binti Hj	Metali	Universiti Brunei Darussalam
Brunei	Kamariah	Salim	Environmental and Life Sciences, Faculty of Science, Universiti Brunei Darussalam
Brunei	Muhd Shahruneey	Saparudin	Universiti Brunei Darussalam
Brunei	Rafizah	Serudin	Universiti Brunei Darussalam
Brunei	Rahayu	Sukri	Universiti Brunei Darussalam
Cameroon	Serge	Begne	Plant Systematic and Ecology Laboratory, Department of Biology, Higher Teachers' Training College, University of Yaounde I
Cameroon	George	Chuyong	Faculty of Science, Department of Botany and Plant Physiology, University of Buea, Buea, Cameroon
Cameroon	Marie Noel	Djuikouo	Faculty of Science, Department of Plant Science, University of Buea.
Cameroon	Christelle	Gonmadje	National Herbarium, Yaounde
Cameroon	Murielle	Simo-Droissart	Plant Systematics and Ecology Laboratory, Higher Teachers' Training College, University of Yaoundé I
Cameroon	Bonaventure	Sonké	Plant Systematics and Ecology Laboratory, Higher Teachers' Training College, University of Yaoundé I
Cameroon	Hermann	Taedomg	Department of Plant Biology, Faculty of Sciences, University of Yaounde 1, Cameroon // Bioversity international, Yaoundé-Cameroon
Cameroon	Lise	Zemagho	Plant Systematics and Ecology Laboratory, Higher Teachers' Training College, University of Yaoundé I
Canada	Sean	Thomas	Faculty of Forestry, University of Toronto

Central African Republic	Fidèle	Baya	Ministère des Eaux, Forêts, Chasse et Pêche (MEFCP), Bangui, Central African Republic
Chile	Gustavo	Saiz	Universidad Católica de la Santísima Concepción
Chile	Javier	Silva Espejo	Universidad de La Serena
China	Dexiang	Chen	Research Institute of Tropical Forestry, Chinese Academy of Forestry
China	Alan	Hamilton	Kunming Institute of Botany
China	Yide	Li	Research Institute of Tropical Forestry, Chinese Academy of Forestry
China	Tushou	Luo	Research Institute of Tropical Forestry, Chinese Academy of Forestry
China	Han	Xu	Research Institute of Tropical Forestry, Chinese Academy of Forestry
China	Zhang	Zhou	Research Institute of Tropical Forestry, Chinese Academy of Forestry
Colombia	Esteban	Álvarez Dávila	UNAD, Coltree
Colombia	Juan Carlos	Andrés Escobar	ColTree
Colombia	Henry	Arellano-Peña	Nuevo Estándar Biotropical NEBIOT SAS
Colombia	Jaime	Cabezas Duarte	Universidad del Tolima
Colombia	Jhon	Calderón	Asociación GAICA, Universidad de Nariño
Colombia	Lina Maria	Corrales Bravo	Universidad del Tolima
Colombia	Borish	Cuadrado	Parques Nacionales Naturales, Territorial Caribe
Colombia	Hermes	Cuadros	Universidad del Atlantico
Colombia	Alvaro	Duque	Departamento de Ciencias Forestales, Universidad Nacional de Colombia - Sede Medellín
Colombia	Luisa Fernanda	Duque	Socioecosistemas y clima sostenible, Fundacion con Vida
Colombia	Sandra Milena	Espinosa	Corporación COL-TREE
Colombia	Rebeca	Franke-Ante	Parques Nacionales Naturales de Colombia
Colombia	Hernando	García	Instituto de Investigación de Recursos Biológicos Alexander von Humboldt
Colombia	Alejandro	Gómez	UNAL, Colombia
Colombia	Roy	González-M.	Instituto de Investigación Recursos Biologicos Alexander von Humboldt
Colombia	Álvaro	Idárraga-Piedrahíta	Fundación Jardín Botánico de Medellín, Herbario “Joaquín Antonio Uribe” (JAUM)
Colombia	Eliana	Jimenez	Universidad Nacional de Colombia sede Amazonia
Colombia	Rubén	Jurado	Asociación GAICA, Universidad de Nariño
Colombia	Wilmar	López Oviedo	Coltree
Colombia	René	López-Camacho	Facultad del Medio Ambiente y Recursos Naturales, Universidad Distrital Francisco José de Caldas
Colombia	Omar Aurelio	Melo Cruz	Universidad de Tolima
Colombia	Irina	Mendoza Polo	Socioecosistemas y clima sostenible, Fundacion con Vida
Colombia	Edwin	Paky	ColTree
Colombia	Karen	Pérez	Fundación Orinoquia Biodiversa
Colombia	Angel	Pijachi	ColTree
Colombia	Camila	Pizano	Departamento de Biología, Facultad de Ciencias Naturales, Universidad Icesi
Colombia	Adriana	Prieto	Instituto de Ciencias Naturales, Universidad Nacional de Colombia
Colombia	Laura	Ramos	Universidad de los Llanos
Colombia	Zorayda	Restrepo Correa	Servicios Ecosistemicos y Cambio Climatico (SECC) Fundación Con Vida & Corporación COL-TREE
Colombia	James	Richardson	Universidad del Rosario
Colombia	Elkin	Rodríguez	Parques Nacionales Naturales, Territorial Caribe
Colombia	Gina M.	Rodriguez M.	Fundacion Ecosistemas Secos de Colombia
Colombia	Agustín	Rudas	Instituto de Ciencias Naturales, Universidad Nacional de Colombia
Colombia	Pablo	Stevenson	Universidad de los Andes - ANDES herbarium
Czech Republic	Radim	Hédl	Institute of Botany, Czech Academy of Sciences
Czech Republic	Stanislav	Lhota	Czech University of Life Sciences Prague

Democratic Republic of Congo	Jacques	Mukinzi	World Wide Fund for Nature
Democratic Republic of Congo	Corneille	Ewango	Wildlife Conservation Society-DR Congo
Democratic Republic of Congo	Terese	Hart	Lukuru Wildlife Research Foundation
Democratic Republic of Congo	Emmanuel	Kasongo Yakusu	Université de Kisangani
Democratic Republic of Congo	Janvier	Lisingo	Faculté des Sciences, Laboratoire d'écologie et aménagement forestier, Université de Kisangani, Kisangani, Democratic Republic of Congo
Democratic Republic of Congo	Jean-Remy	Makana	Wildlife Conservation Society-DR Congo
Democratic Republic of Congo	Faustin	Mbayu	Université de Kisangani Faculté des Sciences Agronomiques République Démocratique du Congo
Democratic Republic of Congo	Vincent	Medjibe	TBC
Democratic Republic of Congo	Donatien	Nzala	TBC
Democratic Republic of Congo	Benjamin	Toirambe	Ministère de l'Environnement et Développement Durable, Kinshasa, Democratic Republic of Congo
Democratic Republic of Congo	John	Tshibamba Mukendi	Université de Kisangani Faculté des Sciences Agronomiques République Démocratique du Congo
Denmark	Lars	Kvist	Aarhus University
Denmark	Gustav	Nebel	University of Copenhagen
Ecuador	Selene	Báez	Escuela Politécnica Nacional del Ecuador
Ecuador	Carlos	Céron	Herbario Alfredo Paredes (QAP), Universidad Central del Ecuador
Ecuador	Daniel M.	Griffith	Universidad Técnica Particular de Loja
Ecuador	Juan Ernesto	Guevara Andino	Grupo de Investigación en Biodiversidad, Medio Ambiente y Salud-BIOMAS- Universidad de las Américas, Campus Queri, Quito Ecuador; Keller Science Action Center, The Field Museum, 1400 South Lake Shore Dr., Chicago, IL
Ecuador	David	Neill	Universidad Estatal Amazónica, Facultad de Ingeniería Ambiental
Ecuador	Walter	Palacios	Universidad Técnica del Norte, Herbario Nacional del Ecuador
Ecuador	Maria Cristina	Peñuela-Mora	Grupo de Ecosistemas Tropicales y Cambio Global, Universidad Regional Amazónica ikiam
Ecuador	Gonzalo	Rivas-Torres	Universidad San Francisco de Quito-USFQ, Colegio de Ciencias Biológicas y Ambientales-COCIBA & Galapagos Institute for the Arts and Sciences-GAIAS; Herbario QUSF; University of Florida, Gainesville-US.
Ecuador	Gorky	Villa	FindingSpecies
Ethiopia	Sheleme	Demissie	Mekelle University
Ethiopia	Tadesse	Gole	independent researcher
Ethiopia	Techane	Gonfa	Environment, Climate Change and Coffee Forest Forum (ECCCCFF)
Finland	Kalle	Ruokolainen	University of Turku
France	Michel	Baisie	CIRAD
France	Fabrice	Bénédet	Centre de coopération International en Recherche Agronomique pour le Développement (CIRAD)
France	Wemo	Betian	CNRS
France	Vincent	Bezard	ONF
France	Damien	Bonal	INRAE
France	Jerôme	Chave	Centre National de la Recherche Scientifique
France	Vincent	Droissart	AMAP, Univ Montpellier, IRD, CNRS, CIRAD, INRA, Montpellier, France.
France	Sylvie	Gourlet-Fleury	Forêts et Sociétés (F&S), Centre de coopération International en Recherche Agronomique pour le Développement (CIRAD), Montpellier, France
France	Annette	Hladik	Département Hommes Natures Sociétés Museum national d'histoire naturelle
France	Nicolas	Labrière	Centre National de la Recherche Scientifique
France	Pétrus	Naisso	Cirad

France	Maxime	Réjou-Méchain	AMAP, Univ Montpellier, IRD, CNRS, CIRAD, INRA, Montpellier, France.
France	Plinio	Sist	Cirad
French Guiana	Lilian	Blanc	CIRAD
French Guiana	Benoît	Burban	INRA Kourou
French Guiana	Géraldine	Derroire	Cirad, UMR Ecologie des Forêts de Guyane (AgroparisTech, CNRS, INRAE, Université des Antilles, Université de la Guyane)
French Guiana	Aurélié	Dourdain	Cirad, UMR Ecologie des Forêts de Guyane (AgroparisTech, CNRS, INRAE, Université des Antilles, Université de la Guyane)
French Guiana	Clement	Stahl	INRAE
Gabon	Natacha Nssi	Bengone	Ministry of Forests, Seas, Environment and Climate
Gabon	Eric	Chezeaux	Rougier-Gabon
Gabon	Fidèle	Evouna Ondo	Agence Nationale des Parcs Nationaux Gabon
Gabon	Vianet	Mihindou	Agence Nationale des Parcs Nationaux / Ministère des Forêts, des Eaux, de la Mer, de l'Environnement, Chargé du Plan Climat, des Objectifs de Développement Durable et du Plan d'Affectation des Terres
Gabon	Lee	White	Institut de Recherche en Ecologie Tropicale (CENAREST) Gabon/Agence Nationale des Parcs Nationaux
Germany	Heike	Culmsee	Georg-August-University Göttingen
Germany	Cristabel	Durán Rangel	University of Freiburg
Germany	Viviana	Horna	TBC
Germany	Florian	Wittmann	Max Planck Institute for Chemistry
Ghana	Stephen	Adu-Bredu	Forestry Research Institute of Ghana (FORIG)
Ghana	Kofi	Affum-Baffoe	Mensuration Unit, Forestry Commission of Ghana
Ghana	Ernest	Foli	Forestry Research Institute of Ghana (FORIG)
Guinea	Michael	Balinga	Center for International Forestry Research
Guyana	Anand	Roopsind	Iwokrama International Centre for Rainforest Conservation and Development
Guyana	James	Singh	Guyana Forestry Commission
Guyana	Raquel	Thomas	Iwokrama International Centre for Rainforest Conservation and Development
Guyana	Roderick	Zagt	Utrecht University
India	Indu	K Murthy	Centre for Sustainable Technologies, Indian Institute of Science
Indonesia	Kuswata	Kartawinata	CIFOR
Indonesia	Kuswata	Kartawinata	Herbarium Borgoriense, Indonesian Institute of Sciences (LIPI)
Indonesia	Edi	Mirmanto	Indonesian Institute of Science, Bogor, Indonesia
Indonesia	Hari	Priyadi	Centre for International Forestry Research (CIFOR)
Indonesia	Ismayadi	Samsedin	Forest Research and Development Agency (FORDA)
Indonesia	Terry	Sunderland	Center for International Forestry Research
Indonesia	Ishak	Yassir	Balitek-KSDA Samboja
Italy	Francesco	Rovero	University of Florence and MUSE - Museo delle Scienze
Italy	Barbara	Vinceti	Bioversity International
Ivory Coast	Bruno	Hérault	Cirad
Japan	Shin-Ichiro	Aiba	Hokkaido University
Japan	Kanehiro	Kitayama	Graduate School of Agriculture, Kyoto University
Liberia	Armandu	Daniels	Forestry Development Authority of the Government of Liberia (FDA)
Liberia	Darlington	Tuagben	Forestry Development Authority of the Government of Liberia (FDA)
Liberia	John T.	Woods	University of Liberia
Malaysia	Muhammad	Fitriadi	Sungai Wain Protection Forest
Malaysia	Kho Lip	Khoon	Malaysian Palm Oil Board
Malaysia	Colin	Maycock	Universiti Malaysia Sabah
Malaysia	Reuben	Nilus	Sabah Forestry Department
Malaysia	Sylvester	Tan	Sarawak Forestry Corporation
Mozambique	Almeida	Sitoe	Eduardo Mondlane University

Nicaragua	Indiana	Coronado G.	Herbarium UNAN-Leon, Universidad Nacional Autónoma de Nicaragua
Nigeria	Lucas	Ojo	University of Abeokuta
Norway	Rafael	de Assis	Natural History Museum of Norway
Norway	Axel Dalberg	Poulsen	University of Oslo
Norway	Douglas	Sheil	Norwegian University of Life Sciences
Peru	Karen	Arévalo Pezo	Universidad Nacional de la Amazonía Peruana
Peru	Hans	Buttgenbach Verde	Universidad Nacional Agraria La Molina (UNALM), Peru
Peru	Victor	Chama Moscoso	Jardin Botanico de Missouri
Peru	Jimmy Cesar	Cordova Oroche	Universidad Nacional de la Amazonia Peruana (UNAP)
Peru	Fernando	Cornejo Valverde	Andes to Amazon Biodiversity Program
Peru	Massiel	Corrales Medina	Universidad Nacional de San Agustín de Arequipa
Peru	Nallaret	Davila Cardozo	Facultad de Ciencias Biológicas, Universidad Nacional de la Amazonía Peruana
Peru	Jhon	del Aguila Pasquel	Instituto de Investigaciones de la Amazonia Peruana
Peru	Gerardo	Flores Llampazo	Universidad Nacional Jorge Basadre de Grohmann (UNJBG)
Peru	Luis	Freitas	Instituto de Investigaciones de la Amazonia Peruana (IIAP)
Peru	Darcy	Galiano Cabrera	Universidad Nacional de San Antonio Abad del Cusco
Peru	Roosevelt	García	TBC
Peru	Karina	Garcia Cabreara	Universidad Nacional de San Antonio Abad del Cusco
Peru	Diego	García Soria	Instituto de Investigaciones de la Amazonía Peruana
Peru	Leticia	Gatica Saboya	UNAP
Peru	Julio Miguel	Grandez Rios	Universidad Nacional de la Amazonia Peruana (UNAP)
Peru	Gabriel	Hidalgo Pizango	Instituto de Investigaciones de la Amazonia Peruana
Peru	Eurídice	Honorio Coronado	Instituto de Investigaciones de la Amazonía Peruana
Peru	Isau	Huamantupa-Chuquimaco	Universidad Nacional de San Antonio Abad del Cusco
Peru	Walter	Huaraca Huasco	Universidad Nacional de San Antonio Abad del Cusco
Peru	Yuri Tomas	Huilca Aedo	Universidad San Antonio Abad del Cusco
Peru	Jose Luis	Marcelo Peña	TBC
Peru	Abel	Monteagudo Mendoza	Universidad Nacional de San Antonio Abad del Cusco
Peru	Vanesa	Moreano Rodriguez	UNALM
Peru	Percy	Núñez Vargas	Universidad Nacional de San Antonio Abad del Cusco
Peru	Sonia Cesarina	Palacios Ramos	Universidad Nacional Agraria La Molina (UNALM), Peru
Peru	Nadir	Pallqui Camacho	Universidad Nacional de San Antonio Abad del Cusco
Peru	Antonio	Peña Cruz	Jardin Botanico de Missouri
Peru	Freddy	Ramirez Arevalo	Universidad Nacional de la Amazonía Peruana
Peru	José	Reyna Huaymacari	Universidad Nacional de la Amazonía Peruana (UNAP)
Peru	Carlos	Reynel Rodriguez	Universidad Nacional Agraria La Molina (UNALM), Peru
Peru	Marcos Antonio	Ríos Paredes	Universidad Nacional de la Amazonia Peruana
Peru	Lily	Rodriguez	Centro de Conservación, Investigación y Manejo, CIMA
Peru	Lily	Rodriguez Bayona	CIMA
Peru	Rocio	Rojas	Jardín Botánico de Missouri
Peru	Maria Elena	Rojas Peña	Universidad Nacional de la Amazonía Peruana (UNAP)
Peru	Norma	Salinas Revilla	Pontificia Universidad Catolica del Peru
Peru	Yahn Carlos	Soto Shareva	Jardin Botanico de Missouri, Oxapampa

Peru	Raul	Tupayachi	Asociacion Bosques Perú
Peru	Luis	Valenzuela Gamarra	Jardín Botánico de Missouri
Peru	Rodolfo	Vasquez Martinez	Jardín Botánico de Missouri
Peru	Jim	Vega Arenas	Universidad Nacional de la Amazonia Peruana
Republic of Congo	Christian	Amani	Université Officielle de Bukavu, Bukavu, Democratic Republic of Congo
Republic of Congo	Suspense	Averti Ifo	Université Marien N'Gouabi, Brazzaville
Republic of Congo	Yannick	Bocko	Université Marien Ngouabi, Brazzaville
Republic of Congo	Patrick	Boundja	Wildlife Conservation Society
Republic of Congo	Romeo	Ekoungoulou	Ecole Nationale Supérieure d'Agronomie et de Foresterie, Université Marien Ngouabi
Republic of Congo	Mireille	Hockemba	Wildlife Conservation Society
Republic of Congo	Shukui	Niu	TBC
Sierra Leone	Alusine	Fofanah	The Gola Rainforest National Park, Kenema, Sierra Leone
Singapore	David	Taylor	Department of Geography, National University of Singapore
Spain	Luis	Cayuela Delgado	Departamento de Biología, Geología, Física y Química Inorgánica, Universidad Rey Juan Carlos
Spain	Íñigo	Granzow-de la Cerda	Real Jardín Botánico - CSIC
Spain	Manuel	Macía	Departamento de Biología, Área de Botánica, Universidad Autónoma de Madrid
Spain	Juliana	Stropp	Museo Nacional de Ciencias Naturales (MNCN-CSIC)
Suriname	Maureen	Playfair	Centre for Agricultural Research in Suriname (CELOS)
Suriname	Verginia	Wortel	Centre for Agricultural Research in Suriname (CELOS)
Sweden	Toby	Gardner	Stockholm Environment Institute
Sweden	Robert	Muscarella	Department of Plant Ecology and Evolution, Uppsala University
Sweden	Hari	Priyadi	Southern Swedish Forest Research Centre
Switzerland	Ervan	Rutishauser	InfoFlora, Conservatoire et Jardin Botanique Geneve
Taiwan	Kuo-Jung	Chao	National Chung Hsing University
Tanzania	Pantaleo	Munishi	Sokoine University of Agriculture
The Netherlands	Olaf	Bánki	Naturalis Biodiversity Center
The Netherlands	Frans	Bongers	Wageningen University, Forest Ecology and Forest Management Group
The Netherlands	Rene	Boot	Tropenbos International
The Netherlands	Gabriella	Fredriksson	Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam
The Netherlands	Jan	Reitsma	Bureau Waardenburg BV
The Netherlands	Hans	ter Steege	Naturalis Biodiversity Center
The Netherlands	Tinde	van An del	Naturalis Biodiversity Center
The Netherlands	Peter	van de Meer	Van Hall Larenstein University of Applied Sciences
The Netherlands	Peter	van der Hout	Van der Hout Forestry Consulting
The Netherlands	Mark	van Nieuwstadt	TBC
The Netherlands	Bert	van Ulft	PROMAB
The Netherlands	Elmar	Veenendaal	Wageningen University, Plant Ecology and Nature Conservation Group
The Netherlands	Ronald	Vernimmen	Data for Sustainability
The Netherlands	Pieter	Zuidema	Wageningen University, Plant Ecology and Nature Conservation Group
The Netherlands	Joeri	Zwerts	Utrecht University
Uganda	Robert	Bitariho	The Institute of Tropical Forest Conservation (ITFC) , Mbarara University of Science and Technology (MUST), Mbarara, Uganda
Uganda	Colin	Chapman	George Washington University/ Makerere University
Uganda	Miguel	Leal	Wildlife Conservation Society
United Kingdom	Katharine	Abernethy	University of Stirling
United Kingdom	Miguel	Alexiades	University of Kent
United Kingdom	Timothy	Baker	School of Geography, University of Leeds
United Kingdom	Karina	Banda	School of Geography, University of Leeds

United Kingdom	Lindsay	Banin	UK Centre of Ecology & Hydrology
United Kingdom	Jos	Barlow	Lancaster University
United Kingdom	Belen	Fadrique	School of Geography, University of Leeds
United Kingdom	Amy	Bennett	School of Geography, University of Leeds
United Kingdom	Erika	Berenguer	Lancaster University, University of Oxford
United Kingdom	Nicholas	Berry	The Landscapes and Livelihoods Group (TLLG)
United Kingdom	Neil M	Bird	Overseas Development Institute
United Kingdom	George A.	Blackburn	Lancaster University
United Kingdom	Francis	Brearley	Manchester Metropolitan University
United Kingdom	Roel	Brienen	School of Geography, University of Leeds
United Kingdom	David	Burslem	University of Aberdeen
United Kingdom	Lidiany	Carvalho	University of Exeter
United Kingdom	Percival	Cho	Lancaster University
United Kingdom	Fernanda	Coelho	University of Leeds
United Kingdom	Murray	Collins	School of GeoSciences, University of Edinburgh
United Kingdom	David	Coomes	University of Cambridge
United Kingdom	Aida	Cuni-Sanchez	Department of Geography and Environmental Science, University of Dundee
United Kingdom	Greta	Dargie	School of Geography, University of Leeds
United Kingdom	Kyle	Dexter	School of GeoSciences, University of Edinburgh
United Kingdom	Mat	Disney	Department of Geography University College London
United Kingdom	Freddie	Draper	School of Geography, University of Leeds
United Kingdom	Muying	Duan	Imperial College, London
United Kingdom	Adriane	Esquivel-Muelbert	School of Geography, Earth & Environmental Sciences and Birmingham Institute of Forest Research, University of Birmingham
United Kingdom	Robert	Ewers	Imperial College London
United Kingdom	Sophie	Fauset	University of Plymouth
United Kingdom	Ted	Feldpausch	Geography, College of Life and Environmental Sciences, University of Exeter
United Kingdom	David	Galbraith	School of Geography, University of Leeds
United Kingdom	Martin	Gilpin	School of Geography, University of Leeds
United Kingdom	Emanuel	Gloor	School of Geography, University of Leeds
United Kingdom	John	Grace	University of Edinburgh
United Kingdom	Keith	Hamer	School of Biology, University of Leeds
United Kingdom	David	Harris	Royal Botanic Garden Edinburgh
United Kingdom	Kath	Jeffery	CENAREST & ANPN & Stirling University
United Kingdom	Tommaso	Jucker	University of Bristol, School of Biological Sciences
United Kingdom	Michelle	Kalamandeen	School of Geography, University of Leeds; Department of Plant Sciences, University of Cambridge and Living with Lake Centre, Laurentian University
United Kingdom	Bente	Klitgaard	Royal Botanic Gardens Kew
United Kingdom	Aurora	Levesley	University of Leeds
United Kingdom	Simon	Lewis	School of Geography, University of Leeds
United Kingdom	Jeremy	Lindsell	The Royal Society for the Protection of Birds, Centre for Conservation Science, Sandy, UK
United Kingdom	Gabriela	Lopez-Gonzalez	School of Geography, University of Leeds
United Kingdom	Jon	Lovett	School of Geography, University of Leeds & Royal Botanic Gardens, Kew
United Kingdom	Yadvinder	Malhi	Environmental Change Institute, School of Geography and the Environment, University of Oxford
United Kingdom	Toby	Marthews	UK Centre for Ecology & Hydrology
United Kingdom	Emma	McIntosh	School of Geography and the Environment, University of Oxford
United Kingdom	Karina	Melgaço	School of Geography, University of Leeds
United Kingdom	William	Milliken	The Royal Botanic Gardens
United Kingdom	Edward	Mitchard	School of GeoSciences, University of Edinburgh
United Kingdom	Peter	Moonlight	Royal Botanic Garden Edinburgh

United Kingdom	Sam	Moore	University of Oxford School of Geography
United Kingdom	Alexandra	Morel	Department of Geography and Environmental Science, University of Dundee
United Kingdom	Julie	Peacock	University of Leeds
United Kingdom	Kelvin	Peh	School of Biological Sciences, University of Southampton
United Kingdom	Colin	Pendry	Royal Botanical Garden Edinburgh
United Kingdom	R. Toby	Pennington	University of Exeter; Royal Botanic Garden Edinburgh
United Kingdom	Luciana de Oliveira	Pereira	University of Exeter
United Kingdom	Carlos	Peres	University of East Anglia
United Kingdom	Oliver	Phillips	School of Geography, University of Leeds
United Kingdom	Georgia	Pickavance	School of Geography, University of Leeds
United Kingdom	Thomas	Pugh	School of Geography, Earth & Environmental Sciences and Birmingham Institute of Forest Research, University of Birmingham
United Kingdom	Lan	Qie	School of Geography, University of Leeds
United Kingdom	Terhi	Riutta	University of Oxford
United Kingdom	Katherine	Roucoux	University of Leeds
United Kingdom	Casey	Ryan	University of Edinburgh
United Kingdom	Tiina	Sarkinen	Royal Botanical Gardens Edinburgh
United Kingdom	Camila	Silva Valeria	Lancaster University
United Kingdom	Dominick	Spracklen	School of Earth and Environment, University of Leeds
United Kingdom	Suzanne	Stas	School of Earth and Environment, University of Leeds
United Kingdom	Martin	Sullivan	University of Leeds
United Kingdom	Michael	Swaine	Department of Plant & Soil Science, Cruickshank Building, School of Biological Sciences, University of Aberdeen
United Kingdom	Joey	Talbot	School of Geography, University of Leeds and Institute for Transport Studies, University of Leeds
United Kingdom	James	Taplin	UK Research & Innovation
United Kingdom	Geertje	van der Heijden	University of Nottingham
United Kingdom	Laura	Vedovato	University of Exeter
United Kingdom	Simon	Willcock	University of Bangor
United Kingdom	Mathew	Williams	University of Edinburgh
USA	Luciana	Alves	Center for Tropical Research, Institute of the Environment and Sustainability, University of California, Los Angeles
USA	Patricia	Alvarez Loayza	Center for Tropical Conservation, Nicholas School of the Environment, Duke University
USA	Gabriel	Arellano	Ecology and Evolutionary Biology, University of Michigan
USA	Cheryl	Asa	Saint Louis Zoo
USA	Peter	Ashton	Department of Organismic and Evolutionary Biology, Harvard University
USA	Gregory	Asner	Center for Global Discovery and Conservation Science, Arizona State University
USA	Terry	Brcic	Wildlife Conservation Society – Programme Congo
USA	Foster	Brown	Woods Hole Research Center
USA	Robyn	Burnham	The University of Michigan Herbarium
USA	Connie	Clark	Nicholas School of the Environment
USA	James	Comiskey	National Park Service
USA	Gabriel	Damasco	University of California
USA	Stuart	Davies	Center for Tropical Forest Science, Smithsonian Tropical Research Institute
USA	Tony	Di Fiore	University of Texas at Austin
USA	Terry	Erwin	Smithsonian Institute
USA	William	Farfan-Rios	Washington University in Saint Louis; Center for Conservation and Sustainable Development at the Missouri Botanical Garden
USA	Jefferson	Hall	Smithsonian Tropical Research Institute, Smithsonian Institution Forest Global Earth Observatory (ForestGEO)

USA	David	Kenfack	Forest Global Earth Observatory (ForestGEO), Smithsonian Tropical Research Institute, Washington, DC, USA
USA	Thomas	Lovejoy	George Mason University, Virginia
USA	Roberta	Martin	Center for Global Discovery and Conservation Science, Arizona State University
USA	Olga Martha	Montiel	Missouri Botanical Garden
USA	John	Pipoly	Broward County Parks and Recreation
USA	John	Pipoly	Nova Southeastern University
USA	Nigel	Pitman	Science and Education, The Field Museum
USA	John	Poulsen	Nicholas School of the Environment
USA	Richard	Primack	Department of Biology, Boston University
USA	Miles	Silman	Wake Forest University
USA	Marc	Steininger	TBC
USA	Varun	Swamy	San Diego Zoo Institute for Conservation Research
USA	John	Terborgh	Center for Tropical Conservation, Nicholas School of the Environment, Duke University
USA	Duncan	Thomas	Biology Department, Washington State University, Vancouver, WA, USA
USA	Mathias	Tobler	TBC
USA	Peter	Umunay	Yale School of Forestry & Environmental Studies
USA	Maria	Uriarte	Columbia University
USA	Emilio	Vilanova Torre	Department of Environmental Science, Policy, and Management, University of California, Berkeley.
USA	Ophelia	Wang	School of Earth Sciences and Environmental Sustainability, Northern Arizona University
USA	Kenneth	Young	Department of Geography and the Environment University of Texas at Austin
Venezuela	Gerardo A.	Aymard C.	UNELLEZ-Guanare, Programa de Ciencias del Agro y el Mar, Herbario Universitario (PORT); Ci Progress GreenLife,
Venezuela	Lionel	Hernández	Universidad Nacional Experimental de Guayana
Venezuela	Rafael	Herrera Fernández	Instituto Venezolano de Investigaciones Científicas (IVIC)
Venezuela	Hirma	Ramírez-Angulo	Universidad de los Andes
Venezuela	Pedro	Salcedo	TBC
Venezuela	Elio	Sanoja	Universidad Nacional Experimental de Guayana
Venezuela	Julio	Serrano	Universidad de Los Andes
Venezuela	Armando	Torres-Lezama	Universidad de los Andes Merida
Viet Nam	Tinh Cong	Le	Viet Nature Conservation Centre
Viet Nam	Trai Trong	Le	Viet Nature Conservation Centre
Viet Nam	Hieu Dang	Tran	Viet Nature Conservation Centre

1477

1478