

Table 4. Subgroup of 229 patients with biopsy-proven NAFLD with available serum cytokine measurements, stratified by the rs2070673 *CYP2E1* genotype.

rs2070673	TT+TA	TA	<i>P</i> -value
N	108	121	
Cytokine(s) (pg/ml)			
IL-1 β	3.62 (1.92-6.22)	3.76 (1.89-6.64)	0.86
IL-1RA	378.05 (271.91-525.51)	370.03 (286.42-571.24)	0.51
IL-6	4.22 (2.82-6.74)	4.50 (2.67-6.42)	0.94
IL-7	13.53 (9.73-19.02)	12.31 (8.74-22.84)	0.30
IL-8	28.60 (16.14-59.16)	23.35 (14.94-41.21)	0.11
IL-9	105.36 (92.10-132.50)	103.82 (83.67-133.26)	0.34
IL-13	7.54 (4.58-16.15)	7.12 (4.32-14.62)	0.28
IL-16	119.43 (84.98-172.97)	108.01 (73.06-158.91)	0.14
IL-18	67.67 (47.72-84.77)	72.52 (57.44-97.09)	0.13
IP-10	914.38 (650.83-1251.43)	1107.09 (737.61-1521.95)	<i>0.01</i>

Data are expressed as medians (Q1-Q3). Significant p-values were highlighted in bold italics.