

10. Santos RAS, Oudit GY, Verano-Braga T, Canta G, Steckelings UM, Bader M. The renin-angiotensin system: going beyond the classical paradigms. *Am J Physiol Heart Circ Physiol*. 2019;316(5):H958-H970.
11. Jackson DJ, Busse WW, Bacharier LB, et al. Association of respiratory allergy, asthma, and expression of the SARS-CoV-2 receptor ACE2. *J Allergy Clin Immunol*. 2020;146(1):203-206.
12. Dhawale VS, Amara VR, Karpe PA, Malek V, Patel D, Tikoo K. Activation of angiotensin-converting enzyme 2 (ACE2) attenuates allergic airway inflammation in rat asthma model. *Toxicol Appl Pharmacol*. 2016;306:17-26.
13. Busnadiego I, Fernbach S, Pohl MO, et al. Antiviral activity of Type I, II, and III interferons counterbalances ACE2 inducibility and restricts SARS-CoV-2. *MBio*. 2020;11(5):e01928-20.
14. Qaradakh T, Gadanec LK, McSweeney KR, et al. The potential actions of angiotensin-converting enzyme II (ACE2) activator diminazene aceturate (DIZE) in various diseases. *Clin Exp Pharmacol Physiol*. 2020;47(5):751-758.

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

DOI: 10.1111/all.14731

## Association of endopeptidases, involved in SARS-CoV-2 infection, with microbial aggravation in sputum of severe asthma

To the Editor,

COVID-19 can be a serious multisystem disease caused by the SARS-CoV-2 coronavirus, and the current pandemic has affected more than 80 million people and caused nearly two million deaths worldwide. The SARS-CoV-2 virus attaches to angiotensin-converting enzyme 2 (ACE2) receptors on the host cell membrane, with the help of dipeptidyl peptidase 4 (DPP4), both exopeptidases.<sup>1</sup> Cleavage of the virus spike protein (S-protein) by endopeptidases, such as transmembrane protease, serine 2 (TMPRSS2) and furin, occurs following which the virus enters the host cell leading to virus replication.<sup>1</sup> Other enzymes, such as the sialyltransferases, ST6GAL1 and ST3GAL4, play a role for the synthesis of influenza A virus entry receptors<sup>2</sup>; however, their role in SARS-CoV-2 infection has not been elucidated.

Asthma is a chronic inflammatory airway disease affecting 350 million people worldwide. It has not been linked to serious outcomes when presenting with COVID-19 infection, although a higher risk of death has been reported in severe asthma populations.<sup>3</sup> The heterogeneous inflammatory nature of asthma raises the possibility that the type of asthmatic inflammation might determine the outcome of SARS-CoV-2 infection in asthma. Type 2 (T2) inflammatory markers have been associated with decreased ACE2 expression in asthma<sup>4,5</sup> that could underlie the reduced risk of SARS-CoV-2 infection in asthmatics. In contrast, non-T2 asthma, particularly neutrophilic asthma, has been associated with higher ACE2 and endopeptidases (TMPRSS2 and furin) expression as compared with the T2-high phenotype<sup>4,5</sup> that might imply a worse outcome with COVID-19 infection.


Airway microbial imbalances have been reported in asthma, particularly in severe non-T2 asthma, and are characterized by decreased microbial  $\alpha$ -diversity with increased pathogenic bacterial abundances in association with neutrophilia.<sup>6</sup> Endopeptidases involved in S-protein cleavage such as furin may also play a role in the cleavage of pathogenic bacteria such *Streptococcus pneumoniae* and *Pseudomonas aeruginosa* or bacterial toxins.<sup>7,8</sup> High expression of such endopeptidases may be associated not only with a higher risk of SARS-CoV-2 infection but also with microbial imbalances in severe asthma. Therefore, the aim of this study was to investigate associations of sputum endopeptidases gene expression with metagenomics composition and whether they could be used to stratify asthma patients according to risk of SARS-CoV-2 infection.

We examined the relation of SARS-CoV-2-associated endopeptidases with the airway bacterial composition, SARS-CoV-2-associated exopeptidases and sialyltransferases, and inflammatory profile (cells and proteins) in 120 sputum samples collected from severe nonsmoking asthmatics, severe smoking asthmatics, mild-moderate asthmatics, and healthy controls of the Unbiased BIOMarkers in PREDiction of respiratory disease outcomes (U-BIOPRED) adult cohort.<sup>9</sup> Definition of asthma severity within the U-BIOPRED cohort has been presented in details elsewhere.<sup>9</sup> Sputum transcriptomics, SomaScan<sup>®</sup> proteomics, and metagenomics were assayed as described previously.<sup>5,6</sup> Gene set variation analysis (GSVA) was performed to obtain the enrichment score (ES) of the endopeptidase genes (TMPRSS2 and furin). Spearman correlation coefficients were computed between endopeptidases ES and sputum inflammatory markers and metagenomics  $\alpha$ -diversity measures. The median ES


is equal to zero. Subsequently, subjects were subdivided into two groups according to their ESs, that is, endopeptidase-high (ES >0,  $n = 60$ ) and endopeptidase-low expression group (ES <0,  $n = 60$ ). These were compared according to sputum inflammatory markers, metagenomics  $\alpha$ -diversity measures, and gene expression of the exopeptidases, ACE2, DPP4, and sialyltransferases (ST3GAL4 and ST6GAL1). The two groups were also compared with respect to current intake of antibiotics, oral corticosteroid (OCS), OCS normalized dosage (in mg), and history of hypertension and diabetes diagnoses. The differential bacterial abundance between endopeptidase groups was computed using edgeR after relative log expression normalization, while proteomics differential abundance was computed using limma. Pathway enrichment analysis of differentially abundant

proteins in the endopeptidase-high group was performed using the Reactome database in g: Profiler (<https://biit.cs.ut.ee/gprofiler/gost>).

Severe nonsmoking ( $n = 61$ ) and smoking ( $n = 23$ ) asthmatics showed the highest median expression ES of endopeptidase as compared to mild-moderate asthmatics ( $n = 20$ ) and healthy controls ( $n = 16$ ) (Figure 1A), consistent with previous findings.<sup>5</sup> The endopeptidases ESs were significantly correlated with sputum neutrophil absolute counts ( $r_s = 0.55$ ,  $p = 7.7 \times 10^{-11}$ ) and percentages ( $r_s = 0.58$ ,  $p = 2.4 \times 10^{-12}$ ), which suggests that the endopeptidases were neutrophil-derived. The endopeptidases ESs were inversely associated with bacterial  $\alpha$ -diversity measures ( $r_s$  for observed species =  $-0.44$ , Shannon =  $-0.38$ , Chao1 =  $-0.46$ , Simpson =  $-0.35$ , all  $ps < 1 \times 10^{-4}$ ).



**FIGURE 1** (A) Protease (endopeptidases) genes enrichment scores (ES) in induced sputum were compared between the 4 U-BIOPRED adult subcohorts. (B) Sputum neutrophils (in absolute counts) were compared between endopeptidase-high and endopeptidase-low groups. (C) Sputum eosinophils (in absolute counts) were compared between endopeptidase-high and endopeptidase-low groups. (D) Different metagenomics  $\alpha$ -diversity measures (observed, Shannon, Chao1, and Simpson) were compared between endopeptidase-high and protease-low groups. (E) ACE2 and DPP4 expression in induced sputum was compared between endopeptidase-high and endopeptidase-low groups. (F) ST3GAL4 and ST6GAL1 gene expression in induced sputum were compared between endopeptidase-high and endopeptidase-low groups. Analysis was performed using two-tailed Mann-Whitney  $U$  and Kruskal-Wallis  $H$  tests as appropriate



**FIGURE 2** (A) Bacterial species differential abundance in induced sputum between endopeptidase-high and endopeptidase-low groups. Values in positive  $\log_2$  fold change demonstrate higher abundance of bacterial species in endopeptidase-high group relative to endopeptidase-low group. Only statistically significant differentially abundant bacterial species with false discovery rate (FDR)  $\alpha < 0.05$  are depicted. (B) SomaScan® proteomics differential abundance in induced sputum between endopeptidase-high and endopeptidase-low groups. Values in positive  $\log_2$  fold change demonstrate higher abundances of proteins in endopeptidase-high group relative to endopeptidase-low group. Only labels of sputum proteins with at least twofold change are depicted on the figure. (C) Pathway enrichment analysis of differentially abundant proteins (DAPs) in the endopeptidase-high group using the Reactome pathways database. Only the top 15 significant pathways are depicted

The endopeptidase-high group (mean age =  $50.9 \pm 13.2$ , 53.3% females) had higher sputum neutrophils (Figure 1B), with no differences in sputum eosinophils (Figure 1C), and exhibited reduced bacterial  $\alpha$ -diversity measures as compared with the endopeptidase-low group (mean age =  $48.2 \pm 14.6$  years, 53.3% females) (Figure 1D). In addition, the endopeptidase-high group had a higher abundance of

pathogenic bacteria, such as *Moraxella catarrhalis* and *Haemophilus influenzae*, displaying a pattern of pathogenic bacterial aggravation compared with endopeptidase-low group (Figure 2A), while the latter had a higher abundance of commensal bacteria, such as *Rothia* and *Prevotella* species. The endopeptidase-high group showed higher sputum expression of the exopeptidases, ACE2 and DPP4 (Figure 1E), and

sialyltransferase ST3GAL4 (but not ST6GAL1) (Figure 1F) compared with the endopeptidase-low group, which might indicate higher risk of SARS-CoV-2 infection and possible associated COVID-19 morbidity. No significant differences were found between both groups considering current antibiotic and normalized OCS dose (data not shown). 250 proteins were differentially abundant between the high and low endopeptidase groups particularly with a higher abundance of inflammatory markers, such as interleukin-6 (IL-6), tumor necrosis factor (TNF) superfamily member 4 (LIGHT), tissue inhibitor of metalloproteinases 2 (TIMP2), macrophage migration inhibitor factor (MIF), TNF-stimulated gene 6 protein (TSG-6), and IL-8 proteins in endopeptidase-high group. Enrichment analysis in the endopeptidase-high group showed up-regulation of several pathways including innate immunity, neutrophil degranulation, cytokines signaling, Toll-like receptor, and platelet activation (Figure 2C). In serum, there was a higher levels of IL-6, IL-18, and C-reactive protein in the endopeptidase-high group ( $p < 0.05$ , data not shown).

These findings suggest that appropriate stratification of asthma patients is necessary to adequately estimate risk and/or morbidity of SARS-CoV-2 infection. The neutrophilia observed in the endopeptidase-high group might be directly associated with pathogenic bacteria aggravation in this group. This may suggest that these pathogenic bacteria presence or "bloating" is aggravating the immune system and changing the overall microbial population. In addition, we speculate that the presence of airway bacterial imbalances might be a consequence of the disturbed immune system in severe asthma such as inadequate phagocytic capacity of macrophages,<sup>10</sup> which might lead to higher risk of infections. In this cohort, clusters of severe asthma patients that exhibited bacterial aggravation were relatively stable after 12–18 months,<sup>6</sup> which suggest impairment of immune system over relatively long periods of time. Second, this bacterial aggravation might be associated with comorbid conditions, such as hypertension and diabetes, which are known risk factors for more severe COVID-19. In our study, the endopeptidase-high group showed higher gene expression of exopeptidases ACE2 (associated with hypertension) and DPP4, and the sialyltransferase ST3GAL4 (associated with diabetes) compared with the endopeptidase-low group, which might indicate the pathophysiologic involvement of both diseases in the endopeptidase-high group. However, there were no significant associations between endopeptidases high/low groups and reported history of diabetes and hypertension diagnosis in the included subjects (data not shown). Therefore, future studies are needed to explore whether both diseases may influence the airway microbiome composition in asthmatics.

The present findings suggest that personalized therapies, such as those targeting neutrophils (eg, anti-IL-17), endopeptidase inhibitors (eg, neprilysin inhibitors), and/or antimicrobial compounds, might be tailored to asthma patients with high risk of SARS-CoV-2 infection.

In conclusion, these findings in sputum highlight that it is important to assess overall microbial profile in relation to SARS-CoV-2-associated proteases in order to adequately assess risk of infection in patients with severe neutrophilic asthma.

## ACKNOWLEDGEMENTS

We would like to thank all the patients who gave written and signed consent to take part in the U-BIOPRED study. The study is registered on ClinicalTrials.gov (identifier: NCT01976767). IMA is supported by the EPSRC (EP/T003189/1), the UK MRC (MR/T010371/1) and by the Wellcome Trust (208340/Z/17/Z).

## FUNDING INFORMATION

Innovative Medicines Initiative, Grant/Award Number: 115010; European Federation of Pharmaceutical Industries and Associations; Seventh Framework Programme, Grant/Award Number: FP7/2007–2013

## CONFLICT OF INTEREST

SED reports personal fees from AZ, Cayman Chemicals, GSK, Merck, Novartis, Regeneron, Sanofi, Teva, outside the submitted work. RD reports receiving fees for lectures at symposia organized by Novartis, AstraZeneca and TEVA, consultation for TEVA and Novartis as member of advisory boards, and participation in a scientific discussion about asthma organized by GlaxoSmithKline. RD is a co-founder and current consultant, and has shares in Synairgen, a University of Southampton spin out company. PJS reports grants from Innovative Medicines Initiative (IMI) covered by the European Union and the European Federation of Pharmaceutical industries and Associations (EFPIA), during the conduct of the study. AHM has received research grants outside the submitted work from GSK, Boehringer Ingelheim, and Vertex, and she is the PI of a P4O2 (Precision Medicine for more Oxygen) public-private partnership sponsored by Health Holland involving many private partners that contribute in cash and/or in kind (Boehringer Ingelheim, Breathomix, Fluida, Ortec Logiqcare, Philips, Quantib-U, Smartfish, SODAQ, Thirona, TopMD, and Novartis), and she has served in advisory boards for AstraZeneca, GSK, and Boehringer Ingelheim with money paid to her institution. KFC has received honoraria for participating in Advisory Board meetings of GSK, AZ, Roche, Novartis, Merck, BI, and Shionogi regarding treatments for asthma, chronic obstructive pulmonary disease, and chronic cough and has also been remunerated for speaking engagements. All other co-authors have nothing to disclose.

## FUNDING INFORMATION

U-BIOPRED has received funding from the Innovative Medicines Initiative (IMI) Joint Undertaking under grant agreement no. 115010, resources of which are composed of financial contributions from the European Union's Seventh Framework Programme (FP7/2007–2013), and European Federation of Pharmaceutical Industries and Associations (EFPIA) companies' in-kind contributions ([www.imi.europa.eu](http://www.imi.europa.eu)).

Mahmoud I. Abdel-Aziz<sup>1,2</sup> 

Nazanin Zounemat Kermani<sup>3</sup>

Anne H. Neerincx<sup>1</sup>

Susanne J. H. Vijverberg<sup>1</sup>

Yike Guo<sup>3</sup>

Peter Howarth<sup>4</sup>

Sven-Erik Dahlen<sup>5</sup>Ratko Djukanovic<sup>4</sup>Peter J. Sterk<sup>1</sup>Aletta D. Kraneveld<sup>6,7</sup>Anke H. Maitland-van der Zee<sup>1,8</sup>Kian Fan Chung<sup>9</sup> Ian M. Adcock<sup>10</sup> 

On behalf the U-BIOPRED Consortium

<sup>1</sup>Department of Respiratory Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands<sup>2</sup>Department of Clinical Pharmacy, Faculty of Pharmacy, Assiut University, Assiut, Egypt<sup>3</sup>Data Science Institute, Imperial College London, London, UK<sup>4</sup>NIHR Southampton Respiratory Biomedical Research Unit, Clinical and Experimental Sciences and Human Development and Health, University of Southampton, Southampton, UK<sup>5</sup>Centre for Allergy Research, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden<sup>6</sup>Division of Pharmacology, Faculty of Science, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, Utrecht, The Netherlands<sup>7</sup>Faculty of Veterinary Medicine, Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands<sup>8</sup>Department of Pediatric Respiratory Medicine, Emma Children's Hospital, Amsterdam UMC, Amsterdam, The Netherlands<sup>9</sup>National Heart and Lung Institute, Imperial College London, and Royal Brompton and Harefield NHS Trust, London, UK<sup>10</sup>National Heart and Lung Institute, Imperial College London and the NIHR Imperial Biomedical Research Centre, London, UK**Correspondence**

Ian M. Adcock, National Heart and Lung Institute, Imperial College London and the NIHR Imperial Biomedical Research Centre, London, UK.  
 Email: ian.adcock@imperial.ac.uk

Mahmoud I. Abdel-Aziz and Nazanin Zounemat Kermani: Shared co-authorship

**ORCID**Mahmoud I. Abdel-Aziz  <https://orcid.org/0000-0001-9263-7137>Kian Fan Chung  <https://orcid.org/0000-0001-7101-1426>Ian M. Adcock  <https://orcid.org/0000-0003-2101-8843>**REFERENCES**

1. Sallenave JM, Guillot L. Innate immune signaling and proteolytic pathways in the resolution or exacerbation of SARS-CoV-2 in Covid-19: key therapeutic targets? *Front Immunol.* 2020;11:1229.
2. Broszeit F, Tzarum N, Zhu X, et al. N-Glycolylneuraminic acid as a receptor for influenza A viruses. *Cell Rep.* 2019;27(11):3284-3294.e6.
3. Williamson EJ, Walker AJ, Bhaskaran K, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature.* 2020;584:430-436.
4. Camiolo M, Gauthier M, Kaminski N, Ray A, Wenzel SE. Expression of SARS-CoV-2 receptor ACE2 and coincident host response signature varies by asthma inflammatory phenotype. *J Allergy Clin Immunol.* 2020;146(2):315-324.e7.
5. Kermani NZ, Song WJ, Badi Y, et al. Airway expression of SARS-CoV-2 receptor, ACE2, and proteases, TMPRSS2 and furin, in severe asthma. *Respir Res.* 2021;22(1):10. <https://doi.org/10.1186/s12931-020-01605-8>
6. Abdel-Aziz MI, Brinkman P, Vijverberg SJH, et al. Sputum microbiome profiles identify severe asthma phenotypes of relative stability at 12-18 months. *J Allergy Clin Immunol.* 2020.
7. Schulte T, Mikaelsson C, Beaussart A, et al. The BR domain of PspA interacts with extracellular DNA to promote bacterial aggregation; structural insights into pneumococcal biofilm formation. *Sci Rep.* 2016;6:e32371.
8. Braun E, Sauter D. Furin-mediated protein processing in infectious diseases and cancer. *Clin Transl Immunol.* 2019;8:e1073.
9. Shaw DE, Sousa AR, Fowler SJ, et al. Clinical and inflammatory characteristics of the European U-BIOPRED adult severe asthma cohort. *Eur Respir J.* 2015;46:1308-1321.
10. Liang Z, Zhang Q, Thomas CM, et al. Impaired macrophage phagocytosis of bacteria in severe asthma. *Respir Res.* 2014;15:72.

**SUPPORTING INFORMATION**

Additional supporting information may be found online in the Supporting Information section.