

Diagnostic accuracy of ultrasonography for the detection of hepatic steatosis: an updated meta-analysis of observational studies

Stefano Ballestri^{1*}, Alessandro Mantovani^{2*}, Christopher D. Byrne^{3,4}, Amedeo Lonardo⁵, Giovanni Targher²

*These two authors contributed equally to the manuscript

¹Internal Medicine Unit, Pavullo Hospital, Azienda USL, Modena, Italy

²Section of Endocrinology, Diabetes and Metabolism, Department of Medicine, University and Azienda Ospedaliera Universitaria Integrata of Verona, Verona, Italy

³Nutrition and Metabolism, Faculty of Medicine, University of Southampton, Southampton, UK;

⁴Southampton National Institute for Health Research Biomedical Research Centre, University Hospital Southampton, UK

⁵Metabolic Syndrome Unit, Ospedale Civile di Baggiovara, Azienda Ospedaliero-Universitaria, Modena, Italy

Word count: abstract 217; text 2,843 (*excluding* title page, abstract, key points, references, figure legends and tables); n. 4 **Figures** *plus* online-only supplementary Material (n. 4 **supplementary Tables** + n. 3 **supplementary Figures**). **References=49**

FUNDING STATEMENT: GT is supported in part by grants from the University School of Medicine of Verona, Verona, Italy. CDB is supported in part by the Southampton NIHR Biomedical Research Centre (IS-BRC-20004), UK.

CONFLICTS OF INTEREST STATEMENT: Nothing to declare.

Address for correspondence:

Prof. Giovanni Targher, MD
Section of Endocrinology, Diabetes and Metabolism
Department of Medicine
University and Azienda Ospedaliera Universitaria Integrata
Piazzale Stefani, 1
37126 Verona, Italy
Phone: +39-045-8123110
E-mail: giovanni.targher@univr.it

or alternatively to

Dr. Alessandro Mantovani, MD, PhD

Section of Endocrinology, Diabetes and Metabolism
Department of Medicine
University and Azienda Ospedaliera Universitaria Integrata
Piazzale Stefani, 1
37126 Verona, Italy
Phone: +39-045-8123748
E-mail: alessandro.mantovani@univr.it

ABSTRACT

Background: To examine the diagnostic accuracy of ultrasonography to detect any HS (defined as steatotic hepatocytes $\geq 5\%$ on histology) and moderate-severe HS (defined as steatotic hepatocytes $\geq 30\%$ on histology) by performing a systematic review and meta-analysis.

Methods: We systematically searched PubMed, Web of Science and Scopus databases, from January 2011 to February 2021, to identify studies conducted in adults investigating the diagnostic accuracy of ultrasonography versus histology for detecting either $\geq 5\%$ histologically-defined HS or moderate-severe HS ($\geq 30\%$). Meta-analysis was performed using random-effects modelling.

Results: Twelve studies were included involving a total of 2,921 individuals, 1,710 (58.5%) of whom had HS $\geq 5\%$ by histology. The overall sensitivity, specificity, positive likelihood ratio and negative likelihood ratio of ultrasonography for the detection of $\geq 5\%$ histologically-defined HS, compared to histology, were 82% (95% confidence interval 76-86%), 80% (72-86%), 4.0 (2.90-5.70), and 0.23 (0.18-0.30), respectively. Based on the pooled analysis of seven studies, the overall sensitivity, specificity, positive likelihood ratio and negative likelihood ratio of ultrasonography for the detection of $\geq 30\%$ histologically-defined HS were 85% (72-92%), 85% (73-93%), 5.72 (3.06-10.7), and 0.18 (0.10-0.33). Funnel plots did not reveal any significant publication bias.

Conclusions: Conventional ultrasonography allows for reliable and accurate detection of $\geq 5\%$ histologically-defined HS compared to histology. These findings call for an extensive use of conventional ultrasonography in the clinical arena.

KEY-WORDS: ultrasound; fatty liver; nonalcoholic fatty liver disease

INTRODUCTION

Hepatic steatosis (HS) is histologically defined by accumulation of triglycerides in >5% of hepatocytes [1], and it is now realized that lipid accumulation represents a disease state [2]. HS is associated with adverse hepatic and extra-hepatic clinical outcomes, spanning progressive fibrosing liver disease [3] to increased risks of incident type 2 diabetes mellitus (T2DM) [4], chronic kidney disease [5], cardiovascular disease [6, 7], and extra-hepatic malignancies [8]. HS results from an imbalance between hepatic *de novo* lipogenesis, and the capacity of hepatocytes to either oxidize or export excess lipid. HS is influenced by many exogenous and endogenous stimuli, such as excessive alcohol consumption, viruses, drugs, and hereditary or endocrine-metabolic diseases [1]. However, from a probabilistic point of view, HS will most commonly occur due to nonalcoholic fatty liver disease (NAFLD) [3]. In this specific setting, quantitative alterations of hepatic fat content play a key role in determining the development and progression of liver disease [9] and, therefore, it is important to be able to accurately detect HS to establish a diagnosis of NAFLD.

Currently, there are several diagnostic tests available to diagnose HS, including non-invasive biomarkers, imaging techniques and liver biopsy [10]. The utilization of such diagnostic methods largely depends on local availability of resources to be destined to the diagnosis and management of this common and frequently asymptomatic liver disease. For example, scientific guidelines provide conflicting advice(s) as to whether specific populations at risk for this condition should be subjected to screening for asymptomatic NAFLD [11]. Nevertheless, scientific societies and authorities agree that because of its widespread availability and excellent precision, liver ultrasonography should be the first-line diagnostic tool for assessing suspected HS [10, 12-14]. In addition, ultrasonography allows prompt identification of focal liver diseases, metastases, hepatocellular carcinoma and gallstones as causes of abnormal liver function tests. Compared to more expensive imaging methods (such as magnetic resonance-based techniques),

ultrasonography is also more widely available [15]. Ultrasonography has a poor sensitivity for detecting mild HS (<30% of steatotic hepatocytes on histology) [16, 17], whereas more recent investigations suggested that semi-quantitative indices, such as the ultrasonographic fatty liver indicator (US-FLI) applied to conventional ultrasonography, may facilitate accurate detection of mild HS, i.e., corresponding to a minimum amount of 10% HS, as assessed histologically [18].

Despite this background of evidence attesting to the superiority of ultrasonography in the initial assessment of HS, more contemporary comparative data are required that assess the diagnostic accuracy of ultrasonography with different levels of HS. Therefore, we undertook a systematic review and meta-analysis in order to compare the diagnostic accuracy of ultrasonography to detect: (i) any HS (defined as steatotic hepatocytes $\geq 5\%$ on histology); and (ii) moderate-severe HS (defined as steatotic hepatocytes $\geq 30\%$ on histology). We identified all observational studies, which were published over the last 10 years after the pioneering meta-analysis undertaken by Hernaez *et al.* in 2011 [16].

METHODS

Review Protocol Registration

The protocol of the meta-analysis was registered in advance in PROSPERO database (International Prospective Register of Systematic Reviews) with the following code registration number: CRD#42020183739.

Data Sources and Searches

We systematically searched in PubMed, Scopus and Web of Science databases to identify all observational studies, published from 1st January 2011 (in order to identify all studies published after the aforementioned meta-analysis by Hernaez *et al.* [16]) to 28th February 2021, assessing

the diagnostic accuracy of conventional ultrasonography for the detection of either $\geq 5\%$ histologically-defined HS or moderate-severe HS in relation to liver biopsy, which is the 'gold standard' method for diagnosing and staging HS [12]. Only studies conducted in adult individuals (>18 years) were included. No restrictions in terms of sex, race or ethnicity were adopted. Exclusion criteria were: (i) abstracts, reviews, editorials, commentaries and guidelines, and (ii) non-English-language articles. Studies using ultrasound techniques other than conventional ultrasonography (e.g., Doppler, transient elastography, contrast-enhanced ultrasound, or quantitative ultrasound fat estimation) were also excluded. Overall, the eligible studies were identified using the free text terms: "fatty liver" OR "nonalcoholic fatty liver disease" OR "NAFLD" OR "chronic viral hepatitis" OR "chronic liver diseases" AND "ultrasonography" OR "ultrasound" IN "humans". We performed a systematic review in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (<http://www.prisma-statement.org>). Additionally, given that the included studies were observational in design, we followed the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) guidelines for the meta-analysis of these studies [19].

Data Extraction and Quality Assessment

Two investigators (SB and AM) independently evaluated titles and abstracts and subsequently obtained full texts of relevant papers. Working independently, these two investigators read the articles and judged whether they met the inclusion criteria. Discrepancies were resolved by discussion with a third author (GT). For all studies, we extracted information on sample size, population characteristics, study country, sensitivity, specificity, number of true positive, true negative, as well as number of false positive, and false negative. In the case of multiple publications, we included the most up-to-date or comprehensive information. We did not

contact any corresponding authors of the eligible studies in order to obtain additional information. Two authors (SB and AM) independently assessed the risk of bias. Since all the eligible studies had an observational design, the Newcastle-Ottawa Scale (NOS) was used to assess the quality of each study, as recommended by the Cochrane Collaboration [20]. The NOS assigns a maximum of four stars for selection (or five stars in the case of cross-sectional studies), two stars for comparability, and three stars for outcome/exposure. We judged studies that received a score of at least eight stars to be at low risk of bias (i.e., thus reflecting the highest study quality).

Data Synthesis and Analysis

The primary measures of the meta-analysis were the diagnostic accuracy of conventional ultrasonography for the detection of either $\geq 5\%$ histologically-defined HS or $\geq 30\%$ histologically-defined HS (i.e., moderate-severe HS) compared to liver biopsy, taken as the gold standard. The overall sensitivity, specificity, positive likelihood ratio and negative (i.e., LR+ and LR- that are estimated respectively by the ratio of the proportion of positive and negative tests in the diseased vs. non-diseased individuals) and the diagnostic odds ratio (DOR, calculated as the LR+ divided by the LR-) were obtained for each eligible study and subsequently combined in the meta-analysis [20]. The meta-analysis was carried out using random-effects modelling as this methodology takes into account any differences between studies even if there is no statistically significant heterogeneity. The Cochran Q chi-square test and the I^2 -statistics were used to assess the heterogeneity across the eligible studies [20]. Presence of high heterogeneity was defined when the result of the Q test was significant ($p < 0.05$) and I^2 -statistics $> 50\%$ [20]. Subgroup analyses by study country, causes of HS, publication year (2010-2014 vs. 2015-2020), population age (above the median) or levels of body mass index ([BMI] above the median) were planned to be used where the heterogeneity was high ($I^2 > 50\%$) and the number of available studies was more than

ten [21]. We also performed the summary receiver operating characteristics (ROC) curve analysis and the respective area under the curve was used as a global measure of test performance [20]. The possibility of publication bias was evaluated by visual inspection of the funnel plots and by the Deeks's test, which is the preferred statistical analysis to formally test for publication bias in studies assessing diagnostic test accuracy [22]. The Fagan's nomogram was used to estimate the clinical value of the index test [23]. This nomogram is a two-dimensional graphical tool for estimating how much a diagnostic test result changes the probability that a subject has the disease in question. All statistical tests were two sided and a significance level of $p < 0.05$ was considered. All analyses were performed with STATA® version 16.1 (Stata, College Station, TX, USA). Specifically, we used '*metandi*' and '*midas*' commands.

RESULTS

Literature search and study characteristics

Supplementary Figure S1 shows the results of the literature research and study selection. Based on the titles and abstracts of 2,467 selected citations (after excluding all duplicates), we initially identified 16 potentially relevant studies that were published between 1st January 2011 and February 28th, 2021. **Supplementary Table S1** describes the syntax used and the records identified through database searching. After examining the full text of these publications, we excluded four studies owing to unsatisfactory inclusion criteria or unsatisfactory outcome measures (**Supplementary Table S2**). As a consequence of this selection strategy, twelve observational studies [18, 24-34] comparing the diagnostic accuracy of ultrasonography to liver histology were included in the meta-analysis and were assessed for quality (as summarized in **Supplementary Table 3**). These studies included a total of 2,921 middle-aged participants (mean

age 45 ± 7 years; mean BMI 29 ± 9 kg/m²), 1,710 (58.5%) of whom had $\geq 5\%$ histologically-defined HS. As also shown in the table, about two thirds of these participants had mild HS (5%-30% of steatotic hepatocytes on histology), whereas one third of participants had moderate-severe HS ($\geq 30\%$ histologically-defined HS). Among the 12 eligible studies, four studies were carried out in the Europe (Italy, Netherlands and Germany), four studies in the United States and four studies in Asian countries (South Korea, China, Pakistan and Taiwan). In addition, six studies included patients with NAFLD, four studies involved patients with chronic viral hepatitis B or C, and two studies enrolled patients with various chronic liver diseases, including NAFLD and chronic viral hepatitis. Overall, four studies received eight stars on the NOS (indicating an overall low risk of bias) and the remaining eight studies received six or seven stars (indicating an overall medium risk of bias), thereby suggesting an overall medium-low risk of bias.

Diagnostic Accuracy of Ultrasonography Versus Histology for Hepatic Steatosis $\geq 5\%$

Supplementary Table 3 summarizes the main characteristics of the 12 eligible studies [18, 24-34], providing also data on the diagnostic accuracy of ultrasonography compared to histology (gold standard) for the detection of $\geq 5\%$ histologically-defined HS. The overall sensitivity of ultrasonography to detect $\geq 5\%$ histologically-defined HS was 82% (95% confidence interval [CI] 76-86%), the specificity was 80% (95% CI 72-86%), the LR+ was 4.0 (95% CI 2.90-5.70) and the LR- was 0.23 (95% CI 0.18-0.30). The diagnostic odds ratio was 17 (95% CI 11-28), and the summary area under the ROC curve was 0.87 (95% CI 0.84-0.90), respectively (**Figure 1**, panels A and B). As expected, the heterogeneity for the area under the summary ROC curve was high ($I^2=97\%$; chi-square: 57.1; df: 2.0, $p < 0.001$). To obtain the post-test probability, we used the Fagan's nomogram for which we performed a simulation with an observed prevalence of 58% of $\geq 5\%$ histologically-defined HS, based on the pooled prevalence of the 12 eligible studies. The probability in this

model of someone having $\geq 5\%$ histologically-defined HS and not being detected by ultrasonography was 24% (**Figure 2**).

As shown in **Supplementary Table S4**, the diagnostic performance of ultrasonography for detecting $\geq 5\%$ histologically-defined HS remained comparable even when the eligible studies were stratified by study country, causes of HS, publication year, age of participants or BMI levels. In addition, study country, age of participants and levels of BMI explained only a small proportion of the aforementioned between-study heterogeneity. Notably, the Deeks' funnel plot asymmetry test did not show any potential publication bias ($p=0.92$) (**Supplementary Figure S2**).

Diagnostic Accuracy of Ultrasonography Versus Histology for Hepatic Steatosis $\geq 30\%$

Combining together the results of the 7 eligible studies [18, 25, 27, 29, 31, 32, 34] involving a total of 2,062 middle-aged individuals, which tested the diagnostic accuracy of ultrasonography for the detection of $\geq 30\%$ histologically-defined HS, we found that the overall sensitivity of ultrasonography to detect HS was 85% (95% CI 72-92%), specificity was 85% (95% CI 73-93%), the LR+ was 5.72 (95% CI 3.06-10.7) and the LR- was 0.18 (95% CI 0.10-0.33). The diagnostic odds ratio was 32 (95% CI 13-77), and the summary area under the ROC curve was 0.92 (0.89-0.94), respectively (**Figure 3**, panels A and B). As expected, the heterogeneity for the area under the summary ROC curve was high ($I^2=98\%$; chi-square: 99.1; df: 2.0, $p<0.001$). To obtain the post-test probability, we used Fagan's nomogram for which we performed a simulation with an observed prevalence of 26% for moderate-severe HS, based on the pooled prevalence of moderate-severe HS in these 7 eligible studies. The probability in this model of someone having moderate-severe HS and not being detected by ultrasonography was 6% (**Figure 4**). In addition, the Deeks' funnel plot asymmetry test did not show any significant publication bias ($p=0.43$) (**Supplementary Figure S3**).

DISCUSSION

Our updated systematic review and meta-analysis of 12 observational studies published during the last decade (from January 2011 to February 2021) was aimed at examining the diagnostic accuracy of ultrasonography versus histology (gold standard) for the detection of either $\geq 5\%$ histologically-defined HS or moderate-severe HS ($\geq 30\%$ histologically-defined HS). The major findings of our meta-analysis are that conventional ultrasonography allows for reliable and accurate detection of $\geq 5\%$ histologically-defined HS (sensitivity 82%, specificity 80%), as well as moderate-severe HS (sensitivity 85%, specificity 85%), compared to liver histology. It should be noted that approximately two thirds of the subjects included in this meta-analysis had mild HS (i.e., less than 30% steatotic hepatocytes on histology).

At first glance, our finding on the diagnostic accuracy of ultrasonography for the detection of $\geq 5\%$ histologically-defined HS is at variance with previous studies published before 2010. Indeed, we found that overall sensitivity of conventional ultrasonography for the detection of $\geq 5\%$ histologically-defined HS was much better (sensitivity 82%) than that reported in a previous meta-analysis of 49 studies published between 1979 and 2010 by Hernaez *et al.* (sensitivity 65%). In contrast, in both meta-analyses overall specificity was very similar (80% in our study compared with 81% [16]). Conversely, our finding on the diagnostic accuracy of ultrasonography for the detection of $\geq 30\%$ histologically-defined HS (sensitivity 85%, specificity 85%) appears to be very similar to that previously reported in the meta-analysis of Hernaez *et al.*, who showed 84.8% sensitivity and 93.6% specificity for the ultrasound detection of moderate-severe HS compared to histology [16].

The reasons for the improved diagnostic accuracy of ultrasonography for the detection of $\geq 5\%$ histologically-defined HS in our meta-analysis, compared with that of Hernaez *et al.* published

over a decade ago, are not apparent. However, it is important to remember that we have included only studies published during the last decade, whereas Hernaez *et al.* included studies published between 1979 and 2010. Thus, we consider that this improved diagnostic accuracy of ultrasonography for the detection of $\geq 5\%$ histologically-defined HS could be, in large part, due to technical evolution/improvement in the ultrasonographic equipment, and increased operator awareness of HS over the last years.

By definition, the presence of HS is a prerequisite for the diagnosis of NAFLD [12]. In NAFLD, the presence of imaging-defined or biopsy-proven HS is associated with liver disease progression, predicts overall mortality and the risk of incident T2DM and other extra-hepatic diseases [4, 5, 8, 35, 36]. Consistently, improvement or resolution of HS on ultrasonography decreases the risk of incident T2DM in patients with NAFLD [37]. Furthermore, HS has recently been identified as a major diagnostic and therapeutic target in NAFLD [38-41]. Accumulating clinical evidence shows that lifestyle changes and various drug treatment options may improve HS [42-47]. Therefore, the ability to monitor HS over time using non-invasive imaging methods is crucial in routine clinical practice, in order to gauge treatment responses. Collectively, the findings of our meta-analysis support a wider clinical use of liver ultrasonography, which is a cheap, reproducible and globally available tool that is both sensitive and specific in identifying HS of varying severity. Additionally, assessment of HS by the most recent ultrasonography equipment may be combined with the non-invasive measurement of liver fibrosis to assess the severity of NAFLD and its inherent cardiovascular risk [48].

New ultrasound techniques, including controlled attenuation parameter (CAP) obtained by transient elastography (Fibroscan[®], Echosens, Paris), have been also proposed to quantitatively assess HS [15]. Fibroscan[®] equipped with CAP has shown a good accuracy for detecting HS and has also the advantage of simultaneously estimating liver fibrosis, being coupled with liver stiffness

measurement [14, 15]. Moreover, spleen size, as assessed by ultrasonography, might be also useful for staging NAFLD, since a larger spleen volume has been associated with the more advanced forms of NAFLD (cirrhosis) [49].

Our meta-analysis has some important limitations that should be mentioned. Firstly, we did not include other newer ultrasound techniques (e.g., sonographic hepato-renal index) that could further improve quantification of low levels of HS. Secondly, the eligible studies did not provide any specific data about the diagnostic accuracy of conventional ultrasonography for the differential detection of mild HS (defined by less than 30% steatotic hepatocytes on histology), and moderate-severe HS ($\geq 30\%$ steatotic hepatocytes). Thirdly, we did not have individual patient data and, therefore, we were unable to test the diagnostic performance of liver ultrasonography in specific patient subgroups.

In conclusion, by evaluating the diagnostic accuracy of conventional ultrasonography for the detection of either $\geq 5\%$ histologically-defined HS or moderate-severe HS, we found that ultrasonography is both sensitive and specific in identifying HS, both in its mild and severe form, as compared to liver histology. These findings call for a more extensive use of conventional ultrasonography both in clinics and in research.

AUTHORS CONTRIBUTIONS

Conceptualization, Methodology, Software, Resources and Supervision: AL, GT; Investigation, Validation and Data Curation: SB, AM, AL, GT; Formal Statistical Analysis: AM, GT; Writing- Original draft preparation: CDB, GT; Writing-Reviewing and Editing: all Authors.

Supplementary Table S1. Syntax used and records identified through database searching.**PubMed <from January 1, 2011 to February 28, 2021>**

#1	Search "fatty liver AND ultrasonography" IN "humans"	2,333
#2	Search "fatty liver AND ultrasound" IN "humans"	2,703
#3	Search "nonalcoholic fatty liver disease AND ultrasonography" IN "humans"	1,636
#4	Search "nonalcoholic fatty liver disease AND ultrasound" IN "humans"	1,940
#5	Search "NAFLD AND ultrasonography" IN "humans"	1,608
#6	Search "NAFLD AND ultrasound" IN "humans"	1,913
#7	Search " chronic viral hepatitis AND ultrasonography" IN "humans"	612
#8	Search " chronic viral hepatitis AND ultrasound" IN "humans"	688
#9	Search " chronic viral disease AND ultrasonography" IN "humans"	1,585
#10	Search " chronic viral disease AND ultrasound" IN "humans"	1,724
#11	Search "NAFLD OR chronic viral disease AND ultrasonography" IN "humans"	2,730

Scopus <from January 1, 2011 to February 28, 2021>

#1	Search "fatty liver AND ultrasonography" IN "humans"	3,125
#2	Search "fatty liver AND ultrasound" IN "humans"	3,248
#3	Search "nonalcoholic fatty liver disease AND ultrasonography" IN "humans"	2,230
#4	Search "nonalcoholic fatty liver disease AND ultrasound" IN "humans"	2,262
#5	Search "NAFLD AND ultrasonography" IN "humans"	1,489
#6	Search "NAFLD AND ultrasound" IN "humans"	2,733
#7	Search "chronic viral hepatitis AND ultrasonography" IN "humans"	1,313
#8	Search "chronic viral hepatitis AND ultrasound" IN "humans"	2,805
#9	Search "chronic viral disease AND ultrasonography" IN "humans"	1,513
#10	Search "chronic viral disease AND ultrasound" IN "humans"	3,220
#11	Search "NAFLD OR chronic viral disease AND ultrasonography" IN "humans"	1,196

Web of Science <from January 1, 2011 to February 28, 2021>

#1	Search "fatty liver AND ultrasonography" IN "humans"	1,242
#2	Search "fatty liver AND ultrasound" IN "humans"	1,785
#3	Search "nonalcoholic fatty liver disease AND ultrasonography" IN "humans"	580
#4	Search "nonalcoholic fatty liver disease AND ultrasound" IN "humans"	758
#5	Search "NAFLD AND ultrasonography" IN "humans"	837
#6	Search "NAFLD AND ultrasound" IN "humans"	1,086
#7	Search "chronic viral hepatitis AND ultrasonography" IN "humans"	103
#8	Search "chronic viral hepatitis AND ultrasound" IN "humans"	298
#9	Search "chronic viral disease AND ultrasonography" IN "humans"	87
#10	Search "chronic viral disease AND ultrasound" IN "humans"	255
#11	Search "NAFLD OR chronic viral disease AND ultrasonography" IN "humans"	3,117

Supplementary Table S2. Studies excluded at the eligibility step of PRISMA diagram.

Author, Year	PMID	Country	Sample size (n)	Exclusion criteria
Xu L et al. 2017	28433586	China	366	Unsatisfactory inclusion criteria
García-Monzón C et al. 2015	25708133	Spain	111	Unsatisfactory study design
Nelson SM et al. 2020	31647137	USA	208	Unsatisfactory inclusion criteria
Loy M et al. 2016	27298639	Italy	88	Unsatisfactory study design

References for table S2

1. Xu L, Lu W, Li P, Shen F, Mi YQ, Fan JG. A comparison of hepatic steatosis index, controlled attenuation parameter and ultrasound as noninvasive diagnostic tools for steatosis in chronic hepatitis B. *Dig Liver Dis* 2017;49:910-917.
2. Garcia-Monzon C, Vargas-Castrillon J, Porrero JL, Alonso MT, Bonachia O, Castillo MJ, Marcos A, et al. Prevalence and risk factors for biopsy-proven non-alcoholic fatty liver disease and non-alcoholic steatohepatitis in a prospective cohort of adult patients with gallstones. *Liver Int* 2015;35:1983-1991.
3. Nelson SM, Hoskins JD, Lisanti C, Chaudhuri J. Ultrasound fatty liver indicator: a simple tool for differentiating steatosis from nonalcoholic steatohepatitis: validity in the average obese population. *J Ultrasound Med* 2020;39:749-759.
4. Loy M, Serra G, Chessa L. The prevalence of bright liver echo pattern in patients with chronic hepatitis C: correlation with steatosis and fibrosis. *J Ultrasound* 2016;19:91-98.

Supplementary Table S3. Main characteristics of the eligible studies (n=12) evaluating the sensitivity and specificity of ultrasonography for identifying hepatic steatosis on histology as gold standard.

Author, Year	PMID	Study country	Sample size (n)	Study population	Causes of HS (Ultrasonographic criteria used for identifying HS)	Age (years)	BMI (kg/m ²)	Patients with HS (≥5%) on histology (n)	Patients without HS on histology (n)	Patients with mild HS (5%-30%) on histology (n)	Patients with moderate-severe HS (≥30%) on histology (n)	True positive (n)	True negative (n)	False positive (n)	False negative (n)	NOS
Van Werven JR et al. 2010	20574093	Netherlands	42*	Patients undergoing hepatic resection for neoplasia	NAFLD (Increased echogenicity compared with right kidney and decreased visualization of diaphragm and intrahepatic vessel borders)	59	27	20	22	11	9	13	18	5	7	6
Lee SS et al. 2010	20185194	South Korea	161	Consecutive potential living liver donors	NAFLD (Increased echogenicity compared with renal cortex and loss of detail of the portal vein)	32	23	60	101	49	11	37	82	23	19	7
Wu J et al. 2012	21901286	USA	410*	Bariatric surgery patients	NAFLD (Increased echogenicity compared with renal cortex and loss of detail of the portal veins)	46	45	291	119	NA	NA	251	81	38	40	8
Wang JH et al. 2013	23828144	Taiwan	175	Consecutive patients with chronic hepatitis and indication for liver biopsy	NAFLD (Increased echogenicity compared with renal cortex)	46	25	111	64	83	28	101	33	10	31	7
Sohail S et al. 2014	23930869	Pakistan	152	Patients with chronic HCV	HCV (Increased echogenicity compared with renal cortex and loss of detail of the portal vein)	39	23	79	73	45	34	72	69	4	7	7
Macaluso P et al. 2014	24112998	Italy	515	Patients with chronic HCV	HCV (increase in liver-kidney contrast and evidence of vascular blurring and deep attenuation signs)	53	26	251	264	170	81	158	239	93	25	8
Bril F et al. 2015	25847730	USA	146	Patients with overweight or obesity undergoing liver biopsy	NAFLD (Increased echogenicity compared with renal cortex and loss of detail of the portal vein)	50	34	96	50	NA	NA	60	44	36	6	7
Petrick A et al. 2015	26003548	USA	513	Bariatric surgery patients	NAFLD (fall in echo amplitude with penetration to deep portion of the liver, extent of discrepancy in echo amplitude between liver and kidney, loss of echoes from the portal vein)	44	47	348	165	197	151	279	112	70	53	8

Chen J et al. 2016	27241724	China	189	Patients with chronic HBV	HBV (Increased echogenicity compared with renal cortex and loss of detail of the portal vein)	37	23	98	91	61	37	87	78	11	14	7
Kelly EM et al 2017	28151547	USA	109	Patients with chronic HBV	HBV (Increased echogenicity compared with renal cortex and loss of detail of the portal vein)	46	25	48	61	39	9	29	56	5	19	7
Ballestri S et al 2017	28641784	Italy	352	Patients with chronic liver diseases	NAFLD/HBV/HCV (presence of liver-kidney contrast, graded as mild/moderate or severe)	48	27	226	126	86	140	187	116	10	39	8
Petzold G et al. 2020	32357147	Germany	157	Patients with chronic liver diseases	NAFLD/HBV/HCV (Increased echogenicity compared with renal cortex and loss of detail of the portal vein)	48	28	82	115	38	44	32	87	28	10	7

Abbreviations: HS, hepatic steatosis; BMI, body mass index; HBV, Hepatitis B virus; HCV, Hepatitis C virus; NAFLD, nonalcoholic fatty liver disease; NASH, nonalcoholic steatohepatitis; NOS, Newcastle-Ottawa scale.

Note: *number of patients in whom liver ultrasound examination was available.

Supplementary Table S4. Subgroup analyses - Diagnostic accuracy of conventional ultrasonography, compared to histology, for the detection of $\geq 5\%$ histologically-defined hepatic steatosis (HS) in eligible studies (n=12), which were stratified by study country, causes of HS, publication year, age of participants or levels of body mass index.

	Sensitivity (95% CI)	I ²	Specificity (95% CI)	I ²	LR+ (95% CI)	LR- (95% CI)
Study country						
Asia (n=4)	0.82 (0.71-0.89)	71%	0.86 (0.77-0.91)	80%	5.81 (3.14-10.8)	0.21 (0.12-0.38)
Europe (n=4)	0.83 (0.79-0.86)	59%	0.81 (0.70-0.89)	86%	4.36 (2.70-7.06)	0.21 (0.17-0.26)
United States (n=4)	0.82 (0.71-0.89)	72%	0.71 (0.52-0.85)	40%	2.84 (1.76-4.59)	0.25 (0.18-0.35)
Causes of HS*						
NAFLD (n=6)	0.82 (0.75-0.87)	70%	0.73 (0.64-0.80)	87%	3.02 (2.27-4.00)	0.25 (0.19-0.33)
Viral hepatitis B or C (n=5)	0.79 (0.61-0.88)	88%	0.85 (0.70-0.94)	93%	4.79 (1.81-12.5)	0.24 (0.14-0.44)
Publication year						
From 2010 to 2014 (n=6)	0.81 (0.73-0.88)	82%	0.79 (0.69-0.86)	76%	3.81 (2.49-5.84)	0.24 (0.15-0.36)
From 2015 to 2020 (n=6)	0.82 (0.75-0.87)	78%	0.81 (0.66-0.90)	93%	4.23 (2.40-7.43)	0.22 (0.17-0.30)
Population age (above the median)						
≤ 46 years (n=7)	0.81 (0.73-0.87)	86%	0.82 (0.71-0.89)	89%	4.44 (2.71-7.28)	0.24 (0.16-0.34)
> 46 years (n=5)	0.85 (0.81-0.88)	62%	0.77 (0.64-0.86)	89%	3.68 (2.27-5.97)	0.20 (0.16-0.25)
Body mass index (above the median)						
≤ 27 kg/m ² (n=8)	0.79 (0.71-0.85)	81%	0.85 (0.78-0.90)	85%	5.42 (3.50-8.39)	0.24 (0.17-0.35)
> 27 kg/m ² (n=4) [§]	0.85 (0.81-0.88)	40%	0.65 (0.57-0.72)	72%	2.44 (2.01-2.97)	0.23 (0.19-0.29)

Abbreviations: CI, confidence interval; LR+, positive likelihood ratio; LR-, negative likelihood ratio.

Note: *The study by Ballestri *et al.* [18] was included only in the subgroup analysis of studies involving patients with chronic viral hepatitis B or C, but not in the subgroup analysis of patients with NAFLD, because there were no false positive or true negative cases with $\geq 5\%$ histologically-defined HS in the group of NAFLD patients.

[§]This subgroup included three bariatric studies^{26,30,31} and the study by Petzold *et al.*³⁴ with an overall mean BMI of 38.5 kg/m² for these four studies.

FIGURE LEGENDS

Figure 1. Overall sensitivity and specificity of ultrasonography to detect $\geq 5\%$ histologically-defined hepatic steatosis (HS) (panel A), and the summary receiver-operating characteristic (ROC) curve plots showing the test accuracy of ultrasonography compared to histology to distinguish between presence of $\geq 5\%$ histologically-defined HS and the absence of HS (panel B) in 12 eligible studies.

Figure 2. Fagan's nomogram for ultrasonography to detect $\geq 5\%$ histologically-defined hepatic steatosis (HS) from the absence of HS.

Figure 3. Overall sensitivity and specificity of ultrasonography to detect $\geq 30\%$ (moderate-severe) histologically-defined hepatic steatosis (HS) (panel A), and the summary receiver-operating characteristic (ROC) curve plots showing the test accuracy of ultrasonography compared to histology to distinguish between presence of moderate-severe HS and the absence of HS (panel B) in 7 eligible studies.

Figure 4. Fagan's nomogram for ultrasonography to detect moderate-severe hepatic steatosis (HS) from the absence of HS.

Supplementary Figure S1. The PRISMA flow diagram of the meta-analysis.

Supplementary Figure S2. The Deeks' funnel plot asymmetry test for eligible studies assessing the overall sensitivity and specificity of ultrasonography to detect $\geq 5\%$ histologically defined hepatic steatosis (HS) from the absence of HS.

Supplementary Figure S3. The Deeks' funnel plot asymmetry test for eligible studies assessing the overall sensitivity and specificity of ultrasonography to detect $\geq 30\%$ (moderate-severe) histologically defined hepatic steatosis (HS) from the absence of HS.

References

1. Bedossa P. Pathology of non-alcoholic fatty liver disease. *Liver Int* 2017;37 Suppl 1:85-9.
2. Lonardo A. Fatty liver and nonalcoholic steatohepatitis. Where do we stand and where are we going? *Dig Dis* 1999;17:80-9.
3. AISF. AISF position paper on nonalcoholic fatty liver disease (NAFLD): Updates and future directions. *Dig Liver Dis* 2017;49:471-83.
4. Mantovani A, Petracca G, Beatrice G, et al. Non-alcoholic fatty liver disease and risk of incident diabetes mellitus: an updated meta-analysis of 501 022 adult individuals. *Gut* 2020.
5. Mantovani A, Petracca G, Beatrice G, et al. Non-alcoholic fatty liver disease and risk of incident chronic kidney disease: an updated meta-analysis. *Gut* 2020.
6. Anstee QM, Mantovani A, Tilg H, et al. Risk of cardiomyopathy and cardiac arrhythmias in patients with nonalcoholic fatty liver disease. *Nat Rev Gastroenterol Hepatol* 2018;15:425-39.
7. Lonardo A, Nascimbeni F, Mantovani A, et al. Hypertension, diabetes, atherosclerosis and NASH: Cause or consequence? *J Hepatol* 2018;68:335-52.
8. Mantovani A, Petracca G, Beatrice G, et al. Non-alcoholic fatty liver disease and increased risk of incident extrahepatic cancers: a meta-analysis of observational cohort studies. *Gut* 2021.
9. Valenti L, Pelusi S. Redefining fatty liver disease classification in 2020. *Liver Int* 2020;40:1016-7.
10. Bugianesi E, Rosso C, Cortez-Pinto H. How to diagnose NAFLD in 2016. *J Hepatol* 2016;65:643-4.
11. Leoni S, Tovoli F, Napoli L, et al. Current guidelines for the management of non-alcoholic fatty liver disease: A systematic review with comparative analysis. *World J Gastroenterol* 2018;24:3361-73.
12. European Association for the Study of the L, European Association for the Study of D, European Association for the Study of O. EASL-EASD-EASO Clinical Practice Guidelines for the management of non-alcoholic fatty liver disease. *J Hepatol* 2016;64:1388-402.
13. Wong VW, Chan WK, Chitturi S, et al. Asia-Pacific Working Party on Non-alcoholic Fatty Liver Disease guidelines 2017-Part 1: Definition, risk factors and assessment. *J Gastroenterol Hepatol* 2018;33:70-85.
14. Byrne CD, Patel J, Scorletti E, et al. Tests for diagnosing and monitoring non-alcoholic fatty liver disease in adults. *BMJ* 2018;362:k2734.
15. Ballestri S, Nascimbeni F, Lugari S, et al. A critical appraisal of the use of ultrasound in hepatic steatosis. *Expert Rev Gastroenterol Hepatol* 2019;13:667-81.
16. Hernaez R, Lazo M, Bonekamp S, et al. Diagnostic accuracy and reliability of ultrasonography for the detection of fatty liver: a meta-analysis. *Hepatology* 2011;54:1082-90.
17. Eslam M, Newsome PN, Sarin SK, et al. A new definition for metabolic dysfunction-associated fatty liver disease: An international expert consensus statement. *J Hepatol* 2020;73:202-9.
18. Ballestri S, Nascimbeni F, Baldelli E, et al. Ultrasonographic fatty liver indicator detects mild steatosis and correlates with metabolic/histological parameters in various liver diseases. *Metabolism* 2017;72:57-65.
19. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000;283:2008-12.
20. Cumpston M, Li T, Page MJ, et al. Updated guidance for trusted systematic reviews: a new edition of the Cochrane Handbook for Systematic Reviews of Interventions. *Cochrane Database Syst Rev* 2019;10:ED000142.
21. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539-58.

22. Deeks JJ, Macaskill P, Irwig L. The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *J Clin Epidemiol* 2005;58:882-93.
23. Fagan TJ. Letter: Nomogram for Bayes theorem. *N Engl J Med* 1975;293:257.
24. van Werven JR, Marsman HA, Nederveen AJ, et al. Assessment of hepatic steatosis in patients undergoing liver resection: comparison of US, CT, T1-weighted dual-echo MR imaging, and point-resolved 1H MR spectroscopy. *Radiology* 2010;256:159-68.
25. Lee SS, Park SH, Kim HJ, et al. Non-invasive assessment of hepatic steatosis: prospective comparison of the accuracy of imaging examinations. *J Hepatol* 2010;52:579-85.
26. Wu J, You J, Yerian L, et al. Prevalence of liver steatosis and fibrosis and the diagnostic accuracy of ultrasound in bariatric surgery patients. *Obes Surg* 2012;22:240-7.
27. Wang JH, Hung CH, Kuo FY, et al. Ultrasonographic quantification of hepatic-renal echogenicity difference in hepatic steatosis diagnosis. *Dig Dis Sci* 2013;58:2993-3000.
28. Sohail S, Aziz S, Mirza T. Ultrasound based evaluation of hepatic steatosis and fibrosis in hepatitis C non-responders. *J Coll Physicians Surg Pak* 2013;23:548-52.
29. Macaluso FS, Maida M, Camma C, et al. Body mass index and liver stiffness affect accuracy of ultrasonography in detecting steatosis in patients with chronic hepatitis C virus genotype 1 infection. *Clin Gastroenterol Hepatol* 2014;12:878-84 e1.
30. Bril F, Ortiz-Lopez C, Lomonaco R, et al. Clinical value of liver ultrasound for the diagnosis of nonalcoholic fatty liver disease in overweight and obese patients. *Liver Int* 2015;35:2139-46.
31. Petrick A, Benotti P, Wood GC, et al. Utility of Ultrasound, Transaminases, and Visual Inspection to Assess Nonalcoholic Fatty Liver Disease in Bariatric Surgery Patients. *Obes Surg* 2015;25:2368-75.
32. Chen J, Wu D, Wang M, et al. Controlled attenuation parameter for the detection of hepatic steatosis in patients with chronic hepatitis B. *Infect Dis (Lond)* 2016;48:670-5.
33. Kelly EM, Feldstein VA, Etheridge D, et al. Sonography Predicts Liver Steatosis in Patients With Chronic Hepatitis B. *J Ultrasound Med* 2017;36:925-32.
34. Petzold G, Lasser J, Ruhl J, et al. Diagnostic accuracy of B-Mode ultrasound and Hepatorenal Index for graduation of hepatic steatosis in patients with chronic liver disease. *PLoS One* 2020;15:e0231044.
35. Mantovani A, Scorletti E, Mosca A, et al. Complications, morbidity and mortality of nonalcoholic fatty liver disease. *Metabolism* 2020;111S:154170.
36. Simon TG, Roelstraete B, Khalili H, et al. Mortality in biopsy-confirmed nonalcoholic fatty liver disease: results from a nationwide cohort. *Gut* 2020.
37. Nasr P, Fredrikson M, Ekstedt M, et al. The amount of liver fat predicts mortality and development of type 2 diabetes in non-alcoholic fatty liver disease. *Liver Int* 2020;40:1069-78.
38. Byrne CD, Targher G. Time to Replace Assessment of Liver Histology With MR-Based Imaging Tests to Assess Efficacy of Interventions for Nonalcoholic Fatty Liver Disease. *Gastroenterology* 2016;150:7-10.
39. Ajmera V, Park CC, Caussy C, et al. Magnetic Resonance Imaging Proton Density Fat Fraction Associates With Progression of Fibrosis in Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology* 2018;155:307-10 e2.
40. Caussy C, Reeder SB, Sirlin CB, et al. Noninvasive, Quantitative Assessment of Liver Fat by MRI-PDFF as an Endpoint in NASH Trials. *Hepatology* 2018;68:763-72.
41. Pelusi S, Valenti L. Hepatic fat as clinical outcome and therapeutic target for nonalcoholic fatty liver disease. *Liver Int* 2019;39:250-6.
42. Keating SE, Hackett DA, Parker HM, et al. Effect of aerobic exercise training dose on liver fat and visceral adiposity. *J Hepatol* 2015;63:174-82.
43. Sung KC, Ryu S, Lee JY, et al. Effect of exercise on the development of new fatty liver and the resolution of existing fatty liver. *J Hepatol* 2016;65:791-7.

44. Kim W, Kim BG, Lee JS, et al. Randomised clinical trial: the efficacy and safety of oltipraz, a liver X receptor alpha-inhibitory dithiolethione in patients with non-alcoholic fatty liver disease. *Aliment Pharmacol Ther* 2017;45:1073-83.
45. Kotler DP, He Q, Engelson ES, et al. The effect of recombinant human growth hormone with or without rosiglitazone on hepatic fat content in HIV-1-infected individuals: a randomized clinical trial. *Antivir Ther* 2016;21:107-16.
46. Eriksson JW, Lundkvist P, Jansson PA, et al. Effects of dapagliflozin and n-3 carboxylic acids on non-alcoholic fatty liver disease in people with type 2 diabetes: a double-blind randomised placebo-controlled study. *Diabetologia* 2018;61:1923-34.
47. Harrison SA, Rinella ME, Abdelmalek MF, et al. NGM282 for treatment of non-alcoholic steatohepatitis: a multicentre, randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet* 2018;391:1174-85.
48. Ballestri S, Mantovani A, Baldelli E, et al. Liver Fibrosis Biomarkers Accurately Exclude Advanced Fibrosis and Are Associated with Higher Cardiovascular Risk Scores in Patients with NAFLD or Viral Chronic Liver Disease. *Diagnostics (Basel)* 2021;11.
49. Tarantino G, Citro V, Balsano C. Liver-spleen axis in nonalcoholic fatty liver disease. *Expert Rev Gastroenterol Hepatol* 2021;15:759-69.