

# The association between midwifery staffing levels and the experiences of mothers on postnatal wards: Cross sectional analysis of routine data

L. Turner<sup>a,\*</sup>, D. Culliford<sup>b</sup>, J. Ball<sup>c</sup>, E. Kitson-Reynolds<sup>c</sup>, P. Griffiths<sup>d</sup>

<sup>a</sup> University of Southampton, School of Health Sciences, Highfield Campus, Southampton SO17 1BJ, United Kingdom

<sup>b</sup> NIHR Applied Research Collaboration Wessex, School of Health Sciences, University of Southampton, United Kingdom

<sup>c</sup> School of Health Sciences, University of Southampton, United Kingdom

<sup>d</sup> School of Health Sciences, University of Southampton, National Institute for Health Research Applied Research Centre (Wessex), United Kingdom

## ARTICLE INFO

### Keywords:

Postnatal  
Workforce  
Midwife  
Staffing  
Patient experience  
Maternity

## ABSTRACT

**Background:** Women have consistently reported lower satisfaction with postnatal care compared with antenatal and labour care. The aim of this research was to examine whether women's experience of inpatient postnatal care in England is associated with variation in midwifery staffing levels.

**Methods:** Analysis of data from the National Maternity Survey in 2018 including 17,611 women from 129 organisations. This was linked to hospital midwifery staffing numbers from the National Health Service (NHS) Workforce Statistics and the number of births from Hospital Episode Statistics. A two-level logistic regression model was created to examine the association of midwifery staffing levels and experiences in post-natal care.

**Results:** The median Full Time Equivalent midwives per 100 births was 3.55 (interquartile range 3.26–3.78). Higher staffing levels were associated with less likelihood of women reporting delay in discharge (adjusted odds ratio [aOR] 0.849, 95% CI 0.753–0.959,  $p = 0.008$ ), increased chances of women reporting that staff always helped in a reasonable time aOR1.200 (95% CI 1.052, 1.369,  $p = 0.007$ ) and that they always had the information or explanations they needed aOR 1.150 (95% CI 1.040, 1.271,  $p = 0.006$ ). Women were more likely to report being treated with kindness and understanding with higher staffing, but the difference was small and not statistically significant aOR 1.059 (0.949, 1.181,  $p = 0.306$ ).

**Conclusions:** Negative experiences for women on postnatal wards were more likely to occur in trusts with fewer midwives. Low staffing could be contributing to discharge delays and lack of support and information, which may in turn have implications for longer term outcomes for maternal and infant wellbeing.

### Explanation

While we recognise that not all gestational parents identify as women; this term was chosen as it has been used in the data source which was accessed for this study and represents most people having maternities.

### Statement of significance (problem)

Women report negative experiences of postnatal care compared with antenatal care and birth. There is a recognised shortage of midwives in maternity services, and this may be impacting on the quality of postnatal care.

### What is already known?

There is evidence that midwifery staffing levels are associated with birth outcomes but little empirical evidence on the impact of midwifery staffing levels in postnatal care

### What this paper adds?

This analysis of survey data supports previous findings that increased midwifery staffing is associated with benefits. This is the first study to examine the effects of organisational staffing on women's experience of postnatal care.

## 1. Introduction

The State of the World's Midwifery report estimates that 900,000

\* Corresponding author.

E-mail addresses: [lyt1g19@soton.ac.uk](mailto:lyt1g19@soton.ac.uk) (L. Turner), [d.j.culliford@soton.ac.uk](mailto:d.j.culliford@soton.ac.uk) (D. Culliford), [jane.ball@soton.ac.uk](mailto:jane.ball@soton.ac.uk) (J. Ball), [e.l.kitson-reynolds@soton.ac.uk](mailto:e.l.kitson-reynolds@soton.ac.uk) (E. Kitson-Reynolds), [Peter.Griffiths@soton.ac.uk](mailto:Peter.Griffiths@soton.ac.uk) (P. Griffiths).

<https://doi.org/10.1016/j.wombi.2022.02.005>

Received 29 September 2021; Received in revised form 8 January 2022; Accepted 10 February 2022

Available online 17 February 2022

1871-5192/© 2022 The Authors. Published by Elsevier Ltd on behalf of Australian College of Midwives. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

more midwives are needed internationally to provide safe care and positive birth experiences [1]. Investment in midwives has been predicted to substantially reduce maternal and infant mortality [2] and better workforce planning has been at the forefront of recent policy initiatives [3–5].

This study focuses on inpatient postnatal care as women have consistently reported lower satisfaction with postnatal care compared with antenatal and labour care [6–8]. A systematic review of 53 studies on expectations and experiences of inpatient postnatal care concluded that whilst women were generally satisfied with their care, they were sometimes critical of communication, feeding support and a lack of explanations [9]. In a qualitative study of ten women following caesarean section in Norway, women reported that staff had not fully taken on board their need for rest and adequate pain relief, and were more concerned with aiding breastfeeding than with their post-surgical recovery [10]. Similar concerns have been raised by researchers in Sweden, who examined feedback from 150 women. They reported that women felt insufficient attention was given to their physical and emotional needs and some felt neglected [11]. In a survey of 1290 first time mothers relating to care in the first 24 h after birth, only 41% felt they had all the emotional support they needed, 45% had all the information and advice they needed, and 56% had all their physical care needs met [12]. Fawcett [13] conducted a thematic analysis of women's experience on the postnatal ward. Some women report that hospital staff seemed stressed and overworked, and this impacted on whether they felt able to ask for help [13].

An association between registered nurse staffing and outcomes for acute inpatients has been found in many cross sectional studies, large longitudinal studies and evaluations of staffing policies [14–16]. Higher skill mix (i.e. total proportion of hours provided by registered nurses) has also been associated with improved outcomes [17]. There is much less evidence of the impact of staffing in maternity services. This is especially notable in postnatal care, as a recent scoping review found a only a small number of studies, which examined outcomes such as breastfeeding and neonatal readmissions, and none looked at the woman's postnatal experience in relation to staffing levels [18]. There has been an increase in caesarean section rates worldwide [19] which impacts on the complexity of care on postnatal wards and length of stay [20]. The rise in maternal age and obesity also increases the risk profile for some women [21,22]. Women in the UK are now offered discharge from hospital 24 h after caesarean section if they are recovering well and do not have complications [20], which alters the acuity case mix of women remaining in postnatal areas. It is important to know whether in-patient services are staffed appropriately for the level of demand and patient acuity, as little evidence currently exists on which to base staffing decisions [23,24].

In England an annual national survey of maternity care is undertaken which asks all women about experiences of care, including postnatal care. Similarly, midwifery staffing levels are routinely reported for all National Health Service (NHS) care providers, which deliver the majority of maternity care in England. Our research aims to examine whether the quality of postnatal care reported by women is associated with variation in midwifery staffing levels.

The following research question was addressed: Is registered midwife staffing at organisational level associated with variation in women's experience of inpatient postnatal care, controlling for other factors?

## 2. Methods

This study is a cross sectional analysis of linked routinely collected datasets in English hospital Trusts. A hospital Trust is an organisation which provides health services in a geographical area in England and

manages one or more acute hospitals. Individual patient data on women's experiences of care has been obtained from the National Maternity Survey 2018 via the UK Data Service [8]. The Maternity Survey was anonymous, confidential, and approved by the NHS Health Research Authority ethics process. Staffing and workload data was obtained from the NHS Workforce Statistics dataset [25] and the Hospital Episode Statistics [26]. The size of the datasets enables detailed statistical analysis of patient and organisation level variables. The use of secondary data is a cost effective way of analysing new questions as data has already been subject to cleaning and quality checks [27]. Ethics permission was granted for this secondary data analysis by the corresponding authors university (ERGO 62570).

Women were eligible for the annual National Maternity Survey if they had a live birth under the care of a participating hospital Trust during February 2018 and were aged 16 years or over. In 2018, 97.2% of births occurred in an NHS establishment, 0.4% in a non NHS establishment and 2% occurred at home [28]. All eligible women were invited to participate unless they had registered to opt-out. The survey was administered by post and completed on paper but women could complete it over the phone if their first language was not English [8]. Fieldwork for the survey took place between April and August 2018. Participation was voluntary and return of the questionnaire was taken as consent. The Full Time Equivalent (FTE) headcount for midwives was extracted from the NHS Workforce Statistics dataset for February 2018, which corresponds with the same time as the National Maternity survey. Records were linked by the unique hospital Trust code. The number of Full Time Equivalent midwives per 100 births for each hospital Trust was calculated using annual births recorded in the Hospital Episode Statistics. These data give an indication of the midwifery staffing resource available to provide all maternity care including (but not limited to) post-natal care. Detailed data on deployment of staff to post-natal care were unavailable. Data from 129 Trusts and 17,611 women were included in this secondary analysis. The included Trusts represented 98% (129/132) of those offering maternity services. Three Trusts were not included in the survey as they had less than 300 births in the study period. The participating Trusts varied in size, with the mean number of births per Trust reported as 4844 births (range 1122 to 15, 500).

The overall response rate was 37% (range of response rates 21%–61% between Trusts). The majority of women who responded were over 30 years of age (71%). The age profile differed between Trusts e.g. women aged 35 or more ranged from 17% to 57% between Trusts. Data on ethnicity and parity are not available at the individual level but have been obtained at a Trust level from data published online. There is variability in ethnicity of respondents between Trusts, with the lowest proportion of white women in one Trust as 34% and the highest 99%. Overall, 86% per cent of respondents were from a White ethnic background, with 8% Asian/Asian British, 3% Black or Black British, 2% mixed ethnicity and 1% Arab or other ethnic group [8]. Younger women and women from non-white ethnicities were under-represented compared with those giving birth in the same time period [29].

In the survey, four closed questions asked specifically about the woman's experience of postnatal care. These relate to whether they experienced a delay in discharge, if they were able to have help within a reasonable time, if they were given the information or explanations they needed, and whether they had been treated with kindness and understanding (Table 1). These four survey questions were only answered by women who had given birth in hospital, and they specifically relate to postnatal care in that setting. The responses were dichotomised into 'Yes, Always' (coded 1) and the alternative which included both 'Yes, sometimes' and 'No' (coded 0). This grouping was decided in advance of the analysis based on the implied quality standard [30]. A sensitivity

**Table 1**  
Questions in maternity survey.

Questions in maternity survey	Response options
On the day you left hospital, was your discharge delayed for any reason?	Yes No Yes, always Yes, sometimes
If you needed attention while you were in hospital after the birth, were you able to get a member of staff to help you within a reasonable time?	No I did not want/need this Don't know/can't remember Yes, always Yes, sometimes
Thinking about the care you received in hospital after the birth of your baby, were you given the information or explanations you needed?	No Don't know/can't remember Yes, always Yes, sometimes
Thinking about the care you received in hospital after the birth of your baby, were you treated with kindness and understanding?	No Don't know/can't remember

analysis was performed by grouping all the ‘Yes’ responses together to examine the effects of this alternative grouping (see Supplementary material). Missing values, don’t know or not applicable responses were removed prior to the analysis, using pairwise deletion. Trust level midwifery staffing levels (as Full Time Equivalent per 100 births) were analysed both as a continuous variable and also divided into tertiles to explore potential non-linear relationships. Three categories were used to ensure sufficient numbers in each category, as the number of Trusts is limited and to aid interpretability.

A two-level multilevel logistic regression model was created using Level-1 (mothers) nested within Level-2 (Trusts). Regression coefficients and adjusted odds ratios (aOR) were calculated for individual predictors as a precursor to fitting a full model. The null model was a two-level random intercept model with no predictors to explore the extent of between-trust variation in the outcomes. Covariates were added to the multilevel models in 3 blocks: (i) staffing and number of births per year (Trust level data) (ii) age group and type of birth (Individual level data) (iii) ethnicity (percentage white) and percentage primiparous respondents (Trust level data).

Akaike’s Information criterion (AIC) and Bayesian information criterion (BIC) goodness of fit data were calculated in order to select models which did not to overfit or underfit the data. Models were selected based on minimising AIC and BIC scores, with lower scores indicating the best fit. Where a difference of less than 2 on the AIC scores was noted, then this was not acted upon as it is not considered to be discriminatory at this level [31]. If AIC and BIC scores disagreed, then priority was given to the model lowest on AIC, and the model lower on BIC was scrutinised and compared for a sensitivity check. Interaction variables of staffing with age, mode of birth and parity were explored to see if this improved model fit using the same method for model selection. To see if results were sensitive to variation in non-responses between Trusts, analyses were repeated including a variable for the Trust response rate and results were scrutinised. The assumptions of the multivariable logistic regression model were examined using guidance by Schreiber-Gregory [32]. We calculated the number of women who would need to be exposed to a higher staffing level to achieve on additional positive outcome using the reciprocal of the absolute risk difference between staffing tertiles [33]. All data was analysed in Stata 16.1 and coding is presented in the Supplementary material.

**3. Results**


For the 129 Trusts studied, the median Full Time Equivalent midwives per 100 births was 3.55 (interquartile range 3.26–3.78). This equates to one midwife per 28 births. The distribution of staffing levels shows variation in staffing between Trusts, with clustering on the left tail (Fig. 1).

The majority of respondents reported that they did not have a delay in discharge (55%) and that staff always helped within a reasonable time (60%), they always had the information or explanations they needed (65%), and were always treated with kindness and understanding (74%). Response categories and frequencies are shown in the Supplementary material. A small proportion of data was missing for each of the four questions, ranging from 466 to 603 respondents (2.6%–3.4%).

Responses varied by age, type of birth and staffing levels. All the unadjusted rates for categorical variables are presented in Table 2.

**3.1. Predictor variables of age, type of birth, parity, ethnicity, and size of Trust**

Women in the older age bands reported a delay in discharge less frequently (adjusted odds ratio [aOR] 0.808–0.892 across older subgroups) and reported more frequently that they were always helped within a reasonable time (aOR 1.206–1.473), they always had the information or explanations they needed (aOR 1.177–1.339), and had always been treated with kindness and understanding (aOR 1.284–1.598). Some variation was also noted in women who had undergone different types of birth. Compared to those having a spontaneous birth with no instrumental intervention, those having an instrumental vaginal birth were more likely to report delay in discharge (aOR 1.406) and less likely to report that staff always helped within a reasonable time (aOR 0.769), that they were always given the information or explanations they needed (aOR 0.613) or always treated with kindness and understanding so often (aOR 0.604). Similar findings were



**Fig. 1.** Distribution of Full Time Equivalent midwives per 100 births in the 129 Trusts.

**Table 2**  
Patient experience outcomes by age, type of birth and staffing levels (% yes always).

	Delay in discharge	Staff help reasonable time	Information/ explanations	Treated with kindness and understanding
Responses analysed per variable	17,050	15,690	17,027	16,962
Missing responses	561	498	466	603
Don't know or not applicable		1423	118	46
Overall rate – % yes always	44.9%	59.6%	65.0%	74.4%
Age of mother				
16–24	48.6%	54.8%	62.0%	69.1%
25–29	46.1%	58.8%	65.3%	73.8%
30–34	44.5%	59.7%	63.8%	74.2%
35+	43.7%	61.0%	66.8%	76.0%
Type of birth				
Spontaneous birth (no instrumental intervention)	43.7%	63.2%	69.4%	79.1%
Instrumental birth	51.9%	57.0%	58.0%	69.5%
Planned caesarean	42.1%	55.0%	63.6%	70.8%
Emergency caesarean	45.2%	54.4%	57.9%	65.8%
Trust staffing levels				
low fte/100 births	47.1%	57.3%	63.6%	73.5%
fte 2.543–3.395				
mid fte/100 births	45.6%	59.7%	65.0%	74.0%
fte 3.396–3.706				
high fte/100 births	41.6%	62.1%	66.8%	75.9%
fte 3.707–5.217				

fte = Full Time Equivalent.

reported by those who had a caesarean section (Table 3). Full models for each of the outcomes are presented in the Supplementary material. The case mix of each Trust suggested that there may be small differences in the experience of cohorts of women in Trusts with more primiparous women (aOR 0.983 to 0.986 for four outcomes) and those with fewer

**Table 3**  
Adjusted odds ratio and 95% CI from multilevel model for predictor variables of age, type of birth, parity, ethnicity and size of Trust.

Question	Delay in discharge	Staff help reasonable time	Information/explanations	Kindness and understanding
16–24 Reference age group				
25–29	0.892 (0.784, 1.015) p = 0.083	1.206 (1.053, 1.382) p = 0.007	1.177 (1.029, 1.354) p = 0.017	1.284 (1.114, 1.481) p = 0.001
30–34	0.829 (0.733, 0.938) p = 0.003	1.327 (1.166, 1.511) p < 0.001	1.139 (1.003, 1.293) p = 0.045	1.384 (1.209, 1.585) p < 0.001
35+	0.808 (0.713, 0.915) p = 0.001	1.473 (1.291, 1.680) p < 0.001	1.339 (1.177, 1.524) p < 0.001	1.598 (1.392, 1.835) p < 0.001
Spontaneous birth (no instrumental intervention) – reference group				
Instrumental birth	1.406 (1.286, 1.538) P < 0.001	0.769 (0.700, 0.844) p < 0.001	0.613 (0.560, 0.672) p < 0.001	0.604 (0.547, 0.668) p < 0.001
Planned caesarean	0.950 (0.863, 1.046) P = 0.298	0.673 (0.609, 0.744) p < 0.001	0.743 (0.673, 0.821) p < 0.001	0.603 (0.541, 0.671) p < 0.001
Emergency caesarean	1.054 (0.967, 1.149) P = 0.230	0.688 (0.629, 0.752) p < 0.001	0.605 (0.554, 0.661) p < 0.001	0.504 (0.549, 0.554) p < 0.001
Trust level characteristics				
% Primiparous	Not in model with best fit	0.978 (0.967, 0.990) p < 0.001	0.986 (0.978, 0.995) p = 0.003	0.983 (0.973, 0.992) p < 0.001
% White ethnic group	Not in model with best fit	1.006 (1.001, 1.011) p = 0.023	1.003 (0.999, 1.006) p = 0.177	1.006 (1.001, 1.009) p = 0.007
No births per Trust	1.000 (1.000, 1.000) p = 0.536	1.000 (1.000, 1.000) p = 0.959	1.000 (1.000, 1.000) p = 0.601	1.000 (1.000, 1.000) p = 0.296

white women (aOR 1.003 to aOR 1.006), however the effect sizes were very small (Table 3). The size of the Trust in terms of the annual number of births did not appear to be associated with variation in any of the reported experiences.

### 3.2. Effects of staffing variation

The association between midwifery staffing and patient experience is shown in Table 4. Full models can be found in the Supplementary material.

In the multi-level model we found that every additional Full Time Equivalent midwife per 100 births was associated with 15% reduction in odds of reporting delay in discharge (aOR 0.849, 95% CI 0.753, 0.959, p = 0.008), a 20% increased odds of women reporting that staff always helped in a reasonable time (aOR 1.200, 95% CI 1.052, 1.369, p = 0.007) and a 15% increased odds of women always having the information or explanations they needed (aOR 1.150, 95% CI 1.040, 1.271, p = 0.006). For women being treated with kindness and understanding, the point estimate is in the direction of improved experiences with more staffing, but the relationship was not statistically significant in the adjusted model (aOR 1.059, 95% CI 0.949, 1.181 p = 0.306) (Table 4).

### 3.3. Estimated differences related to staffing levels

Based on these models it is estimated that Trusts in the highest staffing tertile would have 5.7% fewer women reporting a delay in discharge compared to Trusts with staffing in the lowest tertile (95% CI 2.7%, 8.8%). For every 18 women who receive care in these Trusts one fewer would experience a discharge delay in the higher staffed Trust (number needed to be exposed). Trusts with the highest tertile of staffing would have 4.1% more women saying that staff always helped in a reasonable time (95% CI 0.9%, 7.3%) and 2.7% more reporting that they had always been given the information or explanations they needed (95% CI 0.4%, 5.1%). This equates to one improved outcome in a high staffed Trust for every 24 women or 37 women respectively (Table 4). There appears to be a dose response effect as Trusts with mid-tertile staffing had predicted effects in between the lowest and highest values (Table 4).

### 3.4. Sensitivity analyses

As Trust response rates varied (from 21% to 61%), a variable for the Trust response rate was added to the model. This led to slightly larger

**Table 4**  
Outcomes by staffing levels. Results from multilevel models, nested in Trusts. (aOR and 95% CI presented).

Question	Delay in discharge	Staff help reasonable time	Information/explanations	Treated with kindness and understanding
FTE midwives per 100 births unadjusted	0.855 (0.761, 0.961) p = 0.008	1.246 (1.087, 1.427) p = 0.002	1.171 (1.059, 1.295) p = 0.002	1.110 (0.988, 1.247) p = 0.080
FTE midwives per 100 births adjusted	0.849 (0.753, 0.959) p = 0.008	1.200 (1.052, 1.369) p = 0.007	1.150 (1.040, 1.271) p = 0.006	1.059 (0.949, 1.181) p = 0.306
Reference group low fte per 100 births 2.543–3.395				
Mid fte per 100 births 3.396–3.706	0.920 (0.815, 1.037) p = 0.173	1.141 (0.999, 1.303) p = 0.051	1.089 (0.985, 1.203) p = 0.096	1.067 (0.956, 1.189) p = 0.246
High fte per 100 births 3.707–5.217	0.789 (0.697, 0.894) p < 0.001	1.191 (1.037, 1.367) p = 0.013	1.130 (1.018, 1.255) P = 0.022	1.080 (0.963, 1.211) p = 0.187
Absolute risk difference compared to reference group of low fte per 100 births, predicted from model				
Mid fte per 100 births	−2.0% (−5.0%, −0.9%)	3.1% (0.0%, 6.2%)	1.9% (0.3%, 4.1%)	1.2% (−0.8%, 3.2%)
High fte per 100 births	−5.7% (−8.8%, −2.7%)	4.1% (0.9%, 7.3%)	2.7% (0.4%, 5.1%)	1.4% (−0.7%, 3.6%)
Number of women exposed to provide benefit for 1 woman (calculated as 1/absolute risk difference)				
Highest tertile of staffing vs lowest tertile	18 (11–37)	24 (14–111)	37 (20–250)	71 (non significant)

fte = Full Time Equivalent.

estimates for the effect of staffing but did not substantively alter results (see Supplementary material). The model with size of organisation, staffing, age group and parity for the outcome of information and explanations had a better fit by the BIC criterion only, however the staffing coefficients were similar to the full model with six predictor coefficients (aOR 1.162 vs aOR 1.150). When alternative dichotomisation was used ('yes sometimes' and 'yes always' grouped together versus 'no') substantive conclusions were generally unchanged, although effect sizes tended to be larger (see Supplementary material for all sensitivity analyses).

Interaction variables improved the model fit for kindness and understanding when staffing levels interacted with age group and the percentage of primiparous women. Some subgroups reported more positive experiences in Trusts with higher staffing, although women aged 25–29 years and Trusts with the smallest proportion of primiparous women had findings in the opposite direction. Introduction of interaction variables did not improve model fit for the other three measures (see Supplementary material).

#### 4. Discussion

This cross sectional secondary analysis is the first study to examine the effects of organisational staffing on women's experience of postnatal care. Midwifery staffing levels varied considerably between hospitals and were associated with variation in a number of patient reported experiences of postnatal care, after adjusting for other variables. Women in Trusts with more midwifery staff were less likely to report they had experienced a delay in discharge. They were more likely to report that staff always helped them in a reasonable time and were always given the information or explanations they needed.

Higher midwifery staffing levels have been associated with reductions in postpartum haemorrhage [34], reduced need for neonatal resuscitation [35], maternal readmission [36,37] and increased exclusive breastfeeding rates at discharge [38]. This analysis of survey data supports and expands upon the previous findings that increased midwifery staffing is associated with benefits, by demonstrating differences in important experiences in post-natal care. The effect sizes observed are relatively small, but large numbers of women are affected, and the adverse experiences may have economic consequences. In the lower staffed Trusts, an estimate of 5.7% more women (1 in 18) reported that their discharge had been delayed. The reason for a delay in discharge is multifactorial and may not always be related directly to staffing levels. In the 2018 Maternity Survey [8], 21% of women reported they were waiting for medicines and 10% waiting for test results. Delay in discharge contributes to bed pressures [39] and has negative consequences for the woman's experience [9]. There is room for improvement as overall 45% of women in the survey reported their discharge had been delayed. In a previous survey, student midwives and

postnatal women identified that the postnatal discharge process was rushed and this resulted in poor quality discharge advice [40]. Both a delay in discharge and rushed discharge are unsatisfactory outcomes and have been linked to staffing pressures in postnatal wards.

A higher proportion of staff responding in a timely way and providing information may contribute to a mother's wellbeing. Psychological health has been recognised as a major public health challenge [41], with up to one in five women developing mental health problems during pregnancy or in the first year after birth [41]. It is known that women value support, reassurance, and information from health professionals at this time [42]. Previous work has suggested that midwives do not have enough time to talk to women and support them on postnatal wards [9] and this study provides further evidence that this may sometimes be the case. There may be yet unrecognised consequences of lack of support, which may manifest in readmissions or a decline in breastfeeding rates if not addressed [43,44]. The move to a continuity of carer model in some countries may alter the pattern and experience of care for postnatal women. Such changes require staff to be available at the right time and place to provide this care, and overall staffing levels to facilitate this [45].

The increased drive to provide personalised, respectful and compassionate care is seen in many national and international initiatives [46–48]. In this study, three quarters of women felt they were always treated with kindness and understanding. Although the primary analysis did not find a significant association for staffing levels, our sensitivity analyses do provide evidence that that women's experience of kindness and understanding may be affected by staffing levels. Babaei and Taleghani [49] suggested that workload can be a barrier to providing compassionate care. While negative interactions with patients are relatively rare, Bridges et al. [50] found that negative interactions are more common with lower staffing levels.

The main limitations of this study are its cross-sectional design and the level at which staffing has been measured. Data from the Maternity Survey has been linked to staffing data at an organisational level, so there is not an accurate picture of how many midwives were deployed in the postnatal ward area. It may be proportional to the total number of midwives, although this is not certain, and there may be variations in how individual Trusts deploy midwives to meet local needs. There may also be registered nurses and non-registered staff in postnatal areas which have not been accounted for, or movement within the organisation during shifts as some midwives may be relocated away from the postnatal ward to meet needs in other areas [44]. Confirmation of these results is needed from studies with more direct measures of postnatal ward staffing. New sources of data from electronic rosters has created the potential to undertake longitudinal studies with exposure to staffing measured at a ward or individual patient level, mirroring studies now being undertaken in general nursing [51]. Recall bias and cognitive halo effects may have influenced responses to the maternity survey, as data

was collected some months after the postnatal stay.

We did not consider staffing by other professional groups in this analysis. It is conceivable that other staff groups, such as doctors, adult nurses, neonatal nurses and nursery nurses contribute to women's experiences in the postnatal period. Although this survey data records women's perceptions of delays in discharge actual delays have not been empirically demonstrated. This could be the focus of future research, along with midwives' autonomy in the discharge processes.

This was a large national study including 129 Trusts and over 17,000 women which aids the generalisability of findings. Although the response rate of 37% is fairly typical of similar surveys, this does raise questions about the experience and views of non-responders, especially in Trusts with a lower response rate. The study respondents differed from the target population, for example, there were fewer younger mothers and women from non-white ethnic groups [29]. The sensitivity analyses of response rate did not suggest a bias arising from variation in response rates between Trusts.

This analysis of the Maternity Survey in the UK adds to the body of evidence examining staffing and outcomes in maternity care. We found that that variation in midwifery staffing at organisational level is associated with variation in women's experiences of postnatal care. Low staffing levels were linked to higher levels of adverse experiences that could have important consequences in terms of hospital resource use and maternal wellbeing. While we cannot assume the relationship is causal, nonetheless it seems plausible, and this is worthy of further exploration.

#### CRediT authorship contribution statement

**L. Turner:** Conceptualization, Methodology, Data curation, Analysis, Writing – original draft. **D. Culliford:** Methodology, Statistical analysis, Critical review of draft paper. **J. Ball:** Analysis and interpretation of data, Critical review of draft paper, Supervision. **E. Kitson-Reynolds:** Interpretation of data, Critical review of draft paper, Supervision. **P. Griffiths:** Conceptualization, Funding acquisition, Methodology, Supervision, Analysis and interpretation of data, Writing – review & editing.

#### Ethical statement

This work was approved by the University of Southampton Ethics and Research Governance panel ERGO 62570, 19.3.2021.

#### Funding

Peter Griffiths receives support from a Senior Investigator award made by the National Institute for Health Research and the National Institute for Health Research Applied Research Centre (Wessex).

This research was part funded by the National Institute for Health Research's Health Services & Delivery Research programme (Award ID: NIHR128056).

#### Conflict of interest

None declared.

Full ICJME forms can be supplied by each author on request.

#### Acknowledgement

The views expressed are those of the author(s) and not necessarily those of the National Institute for Health Research, the Department of Health and Social Care, 'arms-length' bodies or other government departments.

#### Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.wombi.2022.02.005>.

#### References

- [1] UNFPA, The State of the World's Midwifery, 2021. <https://www.unfpa.org/sites/default/files/resource-pdf/21-038-UNFPA-SoWMY2021-FastFacts-ENv4302.pdf>.
- [2] A. Nove, I.K. Friberg, L. de Bernis, et al., Potential impact of midwives in preventing and reducing maternal and neonatal mortality and stillbirths: a Lives Saved Tool modelling study, *Lancet Glob. Health* 9 (1) (2021) e24–e32.
- [3] The Institute for Fiscal Studies, Under Pressure? NHS Maternity Services in England, 2017. <https://www.ifs.org.uk/uploads/publications/bns/BN215.pdf>.
- [4] J. Buchan, B. Gershlick, A. Charlesworth, I. Seccombe, Falling Short: The NHS Workforce Challenge, The Health Foundation, 2019. <https://www.health.org.uk/publications/reports/falling-short-the-nhs-workforce-challenge>.
- [5] Health Education England, Maternity Workforce Strategy — Transforming the Maternity Workforce, 2019. [https://www.hee.nhs.uk/sites/default/files/document/MWS\\_Report\\_Web.pdf](https://www.hee.nhs.uk/sites/default/files/document/MWS_Report_Web.pdf).
- [6] S.J. Brown, M.-A. Davey, F.J. Bruinsma, Women's views and experiences of postnatal hospital care in the Victorian Survey of Recent Mothers 2000, *Midwifery* 21 (2) (2005) 109–126.
- [7] Scottish Government, Maternity Care Survey 2018: National Results, 2019. <https://www.gov.scot/publications/maternity-care-survey-2018-national-results/>.
- [8] Care Quality Commission, Picker Institute Europe, 2018 Survey of Women's Experiences of Maternity Care, 2020.
- [9] R. Malouf, J. Henderson, F. Alderdice, Expectations and experiences of hospital postnatal care in the UK: a systematic review of quantitative and qualitative studies, *BMJ Open* 9 (7) (2019) e022212, 1–28.
- [10] M. Nilsen Mørch, E. Sommereth, B. Dahl, Perceptions of postnatal care after emergency caesarean sections, *Nor. J. Clin. Nurs.* 14 (76793) (2019), e-76793, <https://doi.org/10.4220/Sykepleienf.2019.76793>.
- [11] A. Rudman, U. Waldenstrom, Critical views on postpartum care expressed by new mothers, *BMC Health Serv. Res.* 7 (178) (2007).
- [12] V. Bhavnani, M. Newburn, Left to your Own Devices: The Postnatal Care Experiences of 1260 First-Time Mothers, National Childbirth Trust, 2010. [https://www.nct.org.uk/sites/default/files/related\\_documents/PostnatalCareSurveyReport5.pdf](https://www.nct.org.uk/sites/default/files/related_documents/PostnatalCareSurveyReport5.pdf).
- [13] J. Fawcett, The Impact of Postnatal Care on a Woman's Overall Maternity Experience: An Analysis of Stories from Patient Opinion, 2016. <https://www.caropinion.org.uk/resources/blog-resources/1-files/postnatal-care-march-2016.docx>.
- [14] P. Griffiths, J. Ball, J. Drennan, et al., Nurse staffing and patient outcomes: strengths and limitations of the evidence to inform policy and practice. A review and discussion paper based on evidence reviewed for the National Institute for Health and Care Excellence Safe Staffing guideline development, *Int. J. Nurs. Stud.* 63 (2016) 213–225.
- [15] R.L. Kane, T.A. Shamliyan, C. Mueller, S. Duval, T.J. Wilt, The association of registered nurse staffing levels and patient outcomes: systematic review and meta-analysis, *Med. Care* (2007) 1195–1204.
- [16] M. McHugh, L. Aiken, D. Sloane, C. Windsor, C. Douglas, P. Yates, Effects of nurse-to-patient ratio legislation on nurse staffing and patient mortality, readmissions, and length of stay: a prospective study in a panel of hospitals, *Lancet* 397 (2021), 1905–1911.
- [17] De Twigg, Y. Kutzer, E. Jacob, K. Seaman, A quantitative systematic review of the association between nurse skill mix and nursing-sensitive patient outcomes in the acute care setting, *J. Adv. Nurs.* 75 (12) (2019) 3404–3423.
- [18] L. Turner, P. Griffiths, E. Kitson-Reynolds, Midwifery and nurse staffing of inpatient maternity services — a systematic scoping review of associations with outcomes and quality of care, *Midwifery* 103 (2021), 103118.
- [19] A.P. Betrán, J. Ye, A.-B. Moller, J. Zhang, A.M. Gülmezoglu, M.R. Torloni, The increasing trend in caesarean section rates: global, regional and national estimates: 1990–2014, *PLoS One* 11 (2) (2016), e0148343.
- [20] National Institute for Health and Care Excellence, Caesarean Birth, 2021. <https://www.nice.org.uk/guidance/ng192>.
- [21] D. Vitner, K. Harris, C. Maxwell, D. Farine, Obesity in pregnancy: a comparison of four national guidelines, *J. Matern. Fetal Neonatal Med.* 32 (15) (2019) 2580–2590.
- [22] J. Sheen, J.D. Wright, D. Goffman, et al., Maternal age and risk for adverse outcomes, *Am. J. Obstet. Gynecol.* 219 (4) (2018) 390.e1–390.e15.
- [23] J. Baxter, A. Macfarlane, Postnatal caesarean care: evaluating the skill mix, *Br. J. Midwifery* 13 (6) (2005) 378–384.
- [24] D.A. Forster, H.L. McLachlan, J. Yelland, J. Rayner, J. Lumley, M.A. Davey, Staffing in postnatal units: is it adequate for the provision of quality care? Staff perspectives from a state-wide review of postnatal care in Victoria, Australia, *BMC Health Serv. Res.* 6 (2006) 83.
- [25] NHS Digital, NHS Workforce Statistics, 2018. <https://digital.N.H.S.uk/data-and-information/publications/statistical/nhs-workforce-statistics>.
- [26] NHS Digital, Hospital Episode Statistics (HES), 2018. <https://digital.N.H.S.uk/data-and-information/data-tools-and-services/data-services/hospital-episode-statistics>.
- [27] Uk Data Service, Data Catalogue, 2021. <https://ukdataservice.ac.uk/>.
- [28] Office for National Statistics, Provisional Births in England and Wales, 2020. <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeaths>

- sandmarriages/livebirths/articles/provisionalbirthsinenglandandwales/2020#live-births-by-place-of-occurrence.
- [29] NHS Digital, Maternity Services Data Set (MSDS), 2018. <https://digital.N.H.S.uk/data-and-information/data-collections-and-data-sets/data-sets/maternity-services-data-set>.
- [30] Care Quality Commission, Many Women Have a Positive Experience of Maternity Services, But for Some Care Continues to Fall Short, 2020. <https://www.cqc.org.uk/news/releases/many-women-have-positive-experience-maternity-services-some-care-continues-fall-short>.
- [31] K. Burnham, D. Anderson, Multimodel inference: understanding AIC and BIC in model selection, *Sociol. Methods Res.* 33 (2004) 261–304.
- [32] D. Schreiber-Gregory, Logistic and Linear Regression Assumptions: Violation Recognition and Control, 2018. [https://www.researchgate.net/publication/341354759\\_Logistic\\_and\\_Linear\\_Regression\\_Assumptions\\_Violation\\_Recognition\\_and\\_Control](https://www.researchgate.net/publication/341354759_Logistic_and_Linear_Regression_Assumptions_Violation_Recognition_and_Control).
- [33] R. Bender, M. Blettner, Calculating the “number needed to be exposed” with adjustment for confounding variables in epidemiological studies, *J. Clin. Epidemiol.* 55 (5) (2002) 525–530.
- [34] P. Prapawichar, A. Ratinthorn, K. Utriyaprasit, C. Viwatwongkasem, Maternal and health service predictors of postpartum hemorrhage across 14 district, general and regional hospitals in Thailand, *BMC Pregnancy Childbirth* 20 (172) (2020).
- [35] E.D. Hodnett, N.K. Lowe, M.E. Hannah, et al., Effectiveness of nurses as providers of birth labor support in North American hospitals: a randomized controlled trial, *JAMA* 288 (11) (2002) 1373–1381.
- [36] V. Gerova, P. Griffiths, S. Jones, D. Bick, The Association Between Midwifery Staffing and Outcomes in Maternity Units in England: Observational Study Using Routinely Collected Data, 2010. <https://eprints.soton.ac.uk/168511/>.
- [37] S.J. Kim, K.T. Han, H.J. Lee, J.A. Kwon, E.C. Park, Positive effects of medical staffing on readmission within 30 days after discharge: a retrospective analysis of obstetrics and gynecology data, *Eur. J. Public Health* 26 (6) (2016) 935–939.
- [38] C. Dani, S. Papini, L. Iannuzzi, S. Pratesi, Midwife-to-newborn ratio and neonatal outcome in healthy term infants, *Acta Paediatr.* 00 (2020) 1–4.
- [39] P. Hendy, J. Patel, T. Kordbacheh, N. Laskar, M. Harbord, In-depth analysis of delays to patient discharge: a metropolitan teaching hospital experience, *Clin. Med.* 12 (4) (2012) 320–323.
- [40] M. Haith-Cooper, T. Stacey, F. Bailey, Hospital postnatal discharge and sepsis advice: perspectives of women and midwifery students, *Br. J. Midwifery* 26 (4) (2018) 248–253.
- [41] Royal College of Obstetricians and Gynaecologists, Maternal Mental Health — Women’s Voices, 2017. <https://www.rcog.org.uk/globalassets/documents/patients/information/maternalmental-healthwomens-voices.pdf>.
- [42] K. Finlayson, N. Crossland, M. Bonet, S. Downe, What matters to women in the postnatal period: a meta-synthesis of qualitative studies, *PLoS One* 15 (4) (2020), e0231415.
- [43] T.L. Brown, Nurse Staffing, Satisfaction and Hospital Readmission Rates in Skilled Nursing Facilities: A Quantitative, Correlational Study, University of Phoenix, 2020. <https://www.proquest.com/docview/2395252910/previewPDF/8B927F8E10AD4319PQ/1?accountid=13963>.
- [44] F. Dykes, A critical ethnographic study of encounters between midwives and breast-feeding women in postnatal wards in England, *Midwifery* 21 (3) (2005) 241–252.
- [45] H.L. McLachlan, D.A. Forster, J. Yelland, J. Rayner, J. Lumley, Is the organisation and structure of hospital postnatal care a barrier to quality care? Findings from a state-wide review in Victoria, Australia, *Midwifery* 24 (3) (2008) 358–370.
- [46] S. Miller, E. Abalos, M. Chamillard, et al., Beyond too little, too late and too much, too soon: a pathway towards evidence-based, respectful maternity care worldwide, *Lancet* 388 (10056) (2016) 2176–2192.
- [47] NHS England, Better Births Four Years On: A Review of Progress, 2020. <http://www.england.N.H.S.uk/wp-content/uploads/2020/03/better-births-four-year-s-on-progress-report.pdf>.
- [48] Nursing and Midwifery Council, Standards of Proficiency for Midwives, 2019. <https://www.nmc.org.uk/standards/standards-for-midwives/standards-of-proficiency-for-midwives/>.
- [49] S. Babaei, F. Taleghani, Compassionate care challenges and barriers in clinical nurses: a qualitative study, *Iran. J. Nurs. Midwifery Res.* 24 (3) (2019) 213.
- [50] J. Bridges, P. Griffiths, E. Oliver, R.M. Pickering, Hospital nurse staffing and staff–patient interactions: an observational study, *BMJ Qual. Saf.* 28 (9) (2019) 706–713.
- [51] P. Griffiths, A. Maruotti, A. Saucedo, et al., Nurse staffing, nursing assistants and hospital mortality: retrospective longitudinal cohort study, *BMJ Qual. Saf.* 28 (8) (2019) 609–617.