

Micronutrients to support vaccine immunogenicity and efficacy

Philip C. Calder ¹, Mette M. Berger ², Adrian F. Gombart ³, Grace A. McComsey ⁴, Adrian R. Martineau ⁵ and Manfred Eggersdorfer ^{6*}

¹ Faculty of Medicine, University of Southampton and NIHR Southampton Biomedical Research Centre, University Hospital Southampton NHS Foundation Trust, Tremona Road, Southampton SO16 6YD, UK; P.C.Calder@soton.ac.uk

² Lausanne University Hospital (CHUV) & University of Lausanne, Lausanne, Switzerland; Mette.Berger@chuv.ch

³ Linus Pauling Institute, Department of Biochemistry and Biophysics, Oregon State University, 307 Linus Pauling Science Center, Corvallis, OR 97331, USA; adrian.gombart@oregonstate.edu

⁴ Case Western Reserve University and University Hospitals of Cleveland, Cleveland, Ohio, USA; grace.mccomsey@uhhospitals.org

⁵ Centre for Immunobiology, Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London E1 2AT, UK; a.martineau@qmul.ac.uk

⁶ Department of Internal Medicine, University Medical Center Groningen, 9713 GZ Groningen, The Netherlands; m.eggersdorfer@bluewin.ch

*Correspondence: m.eggersdorfer@bluewin.ch

Abstract:

The world has entered the third year of the coronavirus disease 2019 (COVID-19) pandemic. Vaccination is the primary public health strategy to protect against infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), in addition to other measures such as mask wearing and social distancing. Vaccination has reduced COVID-19 severity and mortality dramatically. Nevertheless, incidence globally remains high and certain populations are still at risk for severe outcomes. Additional strategies to support immunity, including potentially enhancing the response to vaccination are needed. Many vitamins and trace minerals have recognised immunomodulatory actions, and their status and/or supplementation have been reported to correspond to incidence and severity of infection. Furthermore, a variety of observational and some interventional studies report that adequate micronutrient status or micronutrient supplementation associate with enhanced vaccine responses, including to COVID-19 vaccination. Such data suggest that micronutrient supplementation may hold the potential to improve vaccine immunogenicity and effectiveness, although additional interventional studies to further strengthen the existing evidence are needed. Positive findings from such research could have important implications for global public health, since deficiencies in several micronutrients that support immune function are prevalent in numerous settings, and supplementation can be implemented safely and inexpensively.

Keywords: Micronutrients; COVID-19 pandemic; vaccine immunogenicity and efficacy.

Citation: Lastname, F.; Lastname, F.; Lastname, F. Title. *Vaccines* **2022**, *10*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Last-name

Received: date

Accepted: date

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors.

Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The world has entered the third year of the coronavirus disease 2019 (COVID-19) pandemic. The primary public health strategies to protect against infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and the disease it causes

(COVID-19) include vaccination, as well as other measures such as mask wearing and social distancing. Thirty COVID-19 vaccines with full or emergency use authorizations have been developed globally, with ten approved for use by the World Health Organization (WHO) [1,2]. The vaccines include those based on mRNA technology, non-replicating viral vectors, inactivated virus, or viral protein subunits [1]. While over 10.5 billion vaccine doses have already been administered, the virus has continued to spread rapidly across the world [3]. The omicron variant has proven to be less severe than previous variants, but infection rates globally have increased substantially. This has resulted in record numbers of hospital admissions in some areas, and a substantial average daily death rate [3,4]. Beyond COVID-19, other acute respiratory tract infections for which vaccines are available, such as seasonal influenza, also remain major causes of morbidity and mortality. Indeed, the WHO reports that seasonal influenza results in up to 5 million cases of severe illness and 650,000 deaths per year [5].

2. Vaccination and COVID-19

Vaccination has reduced the risk of COVID-19 severity and mortality dramatically. The US CDC reports that in December 2021 and January 2022, unvaccinated US adults were 14 times more likely to die from COVID-19 compared with those who were vaccinated, and 41 times more likely than those who also received a booster dose [6]. Nevertheless, those who are vaccinated, and especially certain vulnerable populations, are still at risk for SARS-CoV-2 infection and severe outcomes. In the U.S., incidence of death in those who are 65 – 79 years of age and fully vaccinated has been comparable to that of unvaccinated 30 – 49 year olds, and for those who are at least 80 years old and fully vaccinated, incidence of death has been comparable to those who are 50 - 64 years of age and unvaccinated. Similarly, those who are at least 65 years old and have received a booster have a higher mortality rate than those who are 18 – 49 years old and unvaccinated [6].

Current vaccines appear to show diminished efficacy against common SARS-CoV-2 variants compared to the original viral strain, due to a mismatch between vaccine and viral antigens [7,8]. This problem is expected to persist as additional variants emerge and vaccine antigens continue to differ from those expressed by the virus. In addition, the efficacy of vaccinations decreases over time as immunity is known to wane, requiring booster doses of the vaccine [9,10]. Finally, vaccination may not be as efficacious in certain vulnerable groups, such as the elderly, as described above. Indeed, older people have diminished antibody responses to vaccines, including the seasonal influenza vaccine and certain COVID-19 vaccines [9,11,12]. Additional strategies to support immunity, including possibly enhancing the response to vaccination to limit the impact of COVID-19, are needed.

3. Micronutrient Nutrition and Immunity

The relationship between adequate nutritional status and immune function has been well described [13,14]. Many vitamins and trace minerals are well known to help ensure an optimal immune response to infection. The key mechanistic and complementary roles that vitamins (e.g. vitamins A, B₆, B₁₂, C, D, E, K, folate) and trace elements (e.g. zinc, iron, selenium, copper) play in supporting the innate and adaptive immune responses have been comprehensively reviewed recently [13–16]. Briefly, preclinical and clinical data indicate roles for specific micronutrients in maintaining the physical barriers in the skin, gastrointestinal, and respiratory tracts (e.g. promoting collagen synthesis, promoting tight junction protein expression); supporting the cells and functions of the innate and inflammatory responses (e.g. oxidative burst, phagocytosis, production of complement proteins and proinflammatory and anti-inflammatory cytokines, activity of natural killer cells); and supporting the cells and functions of the adaptive immune response (e.g. antigen presentation; T cell differentiation, proliferation, and function; B cell differentiation and

antibody production). A mechanistic understanding of vitamin D and immunity is presented below. The reader is directed to the reviews cited above for more details, including related to functions of specific micronutrients. With the exception of vitamin E, each of these micronutrients has been granted a health claim in the European Union for their role in supporting immune function [17].

The active form of vitamin D, calcitriol (1,25-dihydroxyvitamin D [1,25(OH)₂D]), is a potent modulator of the immune system. Immune cells from the lymphoid and myeloid lineages can express the vitamin D receptor (VDR), as well as the enzyme 25-hydroxyvitamin D3-1 α -hydroxylase which allows these cells to convert intracellular calcidiol (25-hydroxyvitamin D [25(OH)D]) to 1,25(OH)₂D [16,18,19]. This form then binds to the intracellular vitamin D receptor (VDR), which translocates to the nucleus, binds to promoter elements in target genes, thus altering gene expression and profoundly impacting cellular activity [18,19]. Endocrine, paracrine and intracrine mechanisms of action are considered fundamental ways by which vitamin D impacts immune function [18–20]. Thus, when 25(OH)D levels in the blood are insufficient or deficient, immune responses can be limited, potentially leading to increased incidence and severity of disease. Overall, vitamin D positively impacts immunity by supporting barrier function; supporting the differentiation of monocytes to macrophages, as well as the phagocytic and killing capacities of these macrophages; supporting antigen presentation; modulating the inflammatory response, typically by reducing the expression of pro-inflammatory cytokines and increasing the expression of anti-inflammatory cytokines; and impacting antibody production and the activities of various T cell subsets [13,14,21,22].

Clinically, the most data related to reducing the risk of respiratory infections has been reported for vitamin D. While the results of individual studies are mixed, recent meta-analyses of double blind, randomized placebo-controlled trials indicate that vitamin D supplementation reduces incidence of acute respiratory tract infections, particularly when supplementation occurs daily [23–26]. Further, multiple lines of associational clinical evidence indicate that vitamin D inadequacy is associated with increased incidence, severity, and mortality from COVID-19 [27–29]. Population-based cohort studies investigating associations between regular use of vitamin D supplements and subsequent risk of COVID-19 have yielded mixed results [30–34], although meta-analyses report protective associations overall [28,35]. Deficiency in other micronutrients is also described to associate with increased incidence and/or severity of infectious disease [13,14].

4. Micronutrient Nutrition and Vaccine Responses

Vaccination primarily engages adaptive immune responses, although there is growing evidence that innate immunity may also be affected via induction of trained immunity [36]. Based on an extensive body of preclinical and clinical data, it is widely accepted that both innate and adaptive immune responses are supported by an adequate status of a number of vitamins and trace minerals, particularly those named earlier [13,14,37–41]. Furthermore, while not all data are consistent, results from some randomized controlled clinical trials support a cause-and-effect relationship between micronutrient status and the immune response to vaccination. A recent systematic review and meta-analysis of nine clinical studies found lower seroprotection rates in people who were vitamin D deficient compared to those who were adequate, when vaccinated with H3N2 and B strains of seasonal influenza, although there was no difference in seroprotection against the H1N1 strain [42]. In another study, selenium supplementation in healthy adults was associated with more robust T cell responses to a live, attenuated polio vaccine, as compared to the unsupplemented group. Supplementation was associated with a more rapid clearance of the attenuated poliovirus vaccine from the body, and virus recovered from the feces contained a lower incidence of genetic mutations [43]. A study in Kenyan infants found that anemia and iron deficiency at the time of vaccination were associated with reduced antibody responses to diphtheria, pneumococcal, and pertussis vaccination. Furthermore, in

a follow-up double blind randomized trial, supplementation with a multi-nutrient powder that included iron improved antibody responses to measles vaccination compared to the multi-nutrient powder lacking iron [44].

Consistent with these data, some recently available studies suggest that adequate vitamin D status or supplementation is associated with the response to COVID-19 vaccination. Chillon et al. explored the relationship between vitamin D status and the IgG response to vaccination with two doses of BNT162b2 in a cohort of 126 healthy health care workers (87% women) in Germany [45]. In this study, the authors found no effect of vitamin D status on the IgG response to vaccination through 21 weeks after the second dose [45]. In contrast, data from a longitudinal study of 712 subjects in Greece (mean age 51 years; 62% female) found that replete vitamin D levels were significantly associated with higher antibody titers 3 months post-vaccination with BNT162b2 [46]. Likewise, recently available preliminary data from Jolliffe et al. also demonstrate an independent association between vitamin D supplement use and enhanced humoral responses to COVID-19 vaccination [47]. This study assayed anti-spike antibodies (combined IgG, IgA, and IgM) before and after administration of two doses of ChAdOx1 nCoV-19 or BNT162b2 in 9,101 adults (mean age 64 years; 71% female) in a population-based longitudinal study in the UK and examined 66 potential determinants of the antibody response for their possible association with seronegativity. In a fully adjusted, multivariable analysis of the vaccine response, regular vitamin D supplementation was associated with significantly lower risk of post-vaccination seronegativity. While no information was given on dose, frequency of intake, or circulating levels of 25-hydroxyvitamin D, these association data are consistent with a role for vitamin D in supporting vaccine responses, including to COVID-19 vaccines.

The impact of micronutrient supplementation may be even more pronounced in the elderly, who undergo immunosenescence and can suffer from increased rates of infection and poorer response to vaccines [11,37,48]. In one study, healthy individuals 65 years or older who were supplemented with 200 mg/day vitamin E showed more robust cellular immune responses, as well as increased antibody titers to two of three vaccines (hepatitis B and tetanus but not diphtheria), when compared to those in the placebo group [49]. Consistent with these results, another study investigating this same dose of vitamin E supplementation in healthy elderly men and women showed a significant positive impact on several measures of immune cell function, including chemotaxis and measures of phagocytosis in neutrophils; chemotaxis in lymphocytes; and proliferation, IL-2 production, and cytotoxicity in NK cells [50]. Institutionalized elderly patients who received zinc and selenium supplements, either with or without beta carotene, vitamin C and vitamin E, exhibited higher antibody titers to the influenza vaccine, and a near-significant reduction in respiratory tract infections, whereas supplementation with vitamins alone was associated with lower titers [51]. Finally, a recent study investigated the impact of vitamin D supplementation in healthy vitamin D insufficient individuals 65 years or older, who were vaccinated with varicella zoster virus (VZV). Vitamin D supplementation significantly improved secondary, antigen-specific cutaneous immune response to VZV challenge, characterized by a reduction of inflammatory monocyte infiltration and an increase in T cell recruitment to the site of challenge [52].

Attenuation of systemic inflammation is one potential pathway by which supplementation may help improve vaccine effectiveness. Indeed, a chronic high inflammatory state is known to occur in ageing populations as well as in obesity, HIV infection, and autoimmune diseases. These populations are known to have a suboptimal response to immunizations [53]. It is well documented that vitamin K provision is associated with a reduced production of proinflammatory cytokines, such as interleukin-6 (IL-6), tumor necrosis factor alpha (TNF- α), and interleukin-1 (IL-1) [54,55], which are among the most important cytokines activated during COVID-19, contributing to the cytokine storm in severe COVID-19 patients. Similarly, vitamin D has been shown to decrease systemic inflammation in COVID-19 [56] and in HIV [57,58] and other hyper-inflammatory states.

Supplementation with vitamin D and/or vitamin K to decrease systemic inflammation could enhance vaccine effectiveness.

Despite these promising data, it is important to acknowledge that not all studies reveal an improvement in vaccine response as a function of micronutrient supplementation or status. In addition to negative results described above, an intervention study by Provinciale et al. reported no effect of supplemental zinc (400 mg/d for 60 days) on the antibody response to seasonal influenza vaccination in older participants [59]. In addition, while the meta-analysis cited above concluded lower seroprotection rates in people who were vitamin D deficient compared to those who were adequate, the effects are inconsistent with some studies reporting no effect of vitamin D [60,61]. In a study of elderly subjects, there was no association found between levels of vitamin A, vitamin E, or zinc and antibody responses to influenza vaccination. However, it is important to note that in this study no participants were deficient in vitamin A or vitamin E, and only 20% had low serum zinc levels [62]. Studies such as these highlight the need for additional intervention trials, and the importance of taking into account the baseline intakes and status of their participants.

5. Micronutrient Inadequacy and Supplementation

Unfortunately, micronutrient inadequacies are prevalent globally, including inadequacies of those micronutrients that are important for immune function [37,63–69]. For example, a systematic review of 195 studies including 168,000 individuals from 44 countries reported widespread vitamin D inadequacy, with 37% of the studies reporting mean serum 25-hydroxyvitamin D levels lower than the threshold of deficiency (<50 nmol/L) [67]. A systematic review including nearly 250,000 participants from 46 countries indicated that 27% of the American, 80% of the Middle East/African, 62% of the Asian, and 19% of the European adult population had a threshold concentration of α -tocopherol below 20 μ mol/L, which is recommended by experts. Furthermore, 13% of the individuals had concentrations below 12 μ mol/L, which is the threshold for deficiency [68]. The WHO and the Food and Agriculture Organization of the United Nations (FAO) have reported that iron and vitamin A deficiencies are widespread and of global concern [63,65,66]. Finally, a recent systematic review of elderly adults in 13 Western countries found widespread insufficiencies in several trace elements including selenium, zinc, iron, and copper [70].

Supplementation should be performed with the goal of reaching adequate status of the micronutrients in question, to achieve beneficial outcomes. The importance of assessing vitamin D status (serum 25(OH)D) after supplementation, and achieving adequate vitamin D status to achieve health outcomes has been discussed [71]. As mentioned above, multiple associational studies, systematic reviews, and meta-analyses have concluded that inadequate status of circulating 25(OH)D is associated with higher incidence and severity of COVID-19. These findings have led some authors to conclude that achieving specific thresholds of circulating 25(OH)D (e.g. at least 75 nmol/L, or 125 nmol/L) might reduce the burden of SARS-CoV-2 infection [28,72]. Indeed, in a large population-based cohort study using public health records, vitamin D supplementation prior to COVID-19 infection was linked to only a modest decrease in infection in the overall population, but substantial reductions in infection, severe outcomes, and mortality in the subjects that achieved serum 25(OH)D levels of at least 75 nmol/L [30].

6. Conclusions

Despite the availability of vaccines, both endemic and pandemic respiratory diseases such as influenza and COVID-19 lead to extensive morbidity and mortality worldwide. When available, vaccines are by far the most important and effective weapons in the arsenal against these and other infectious diseases. Nevertheless, they are not 100% effective, as described above for vaccines against SARS-CoV-2. Similarly, since the 2004 – 2005

influenza season, the US CDC estimates that influenza vaccination has ranged from 10% - 60% effective for preventing outpatient medical visits due to laboratory-confirmed influenza [73]. Therefore, there is interest in identifying modifiable risk factors for poor immunogenicity and vaccine failure. Given the data presented above, inadequate micronutrient status is worth considering as one of these factors. Unfortunately, nutritional gaps are prevalent in several micronutrients reported to support immune function, including vitamins A, D, and E, as well as iron, zinc and selenium. Supplementation combining the micronutrients at highest risk of deficit should be considered a safe and effective way to prevent or correct inadequacies, and existing data indicate such a strategy could support vaccine responses, thereby reducing the incidence and severity of respiratory diseases. On a population level, even an incremental improvement in vaccine immunogenicity could meaningfully reduce the impact of disease, and its associated social and economic costs.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: P.C.C. has research funding from Bayer Consumer Care; acts as an advisor/consultant to BASF AS, DSM, Cargill, Smartfish, Nutrileads, Bayer Consumer Care, and GSK Consumer Healthcare; has received reimbursement for travel and/or speaking from Danone/Nutricia, Fresenius Kabi, Baxter, GSK Consumer Healthcare, Abbott, Smartfish, Biogredia and the California Walnut Commission; and is Past President and member of the Board of Directors of the European Branch of the International Life Sciences Institute. M.M.B. receives honoraria for lectures from Baxter, B. Braun, DSM, Fresenius Kabi, Nestle, and Nutricia; and has research funding from ESPEN, Foundation Nutrition 2000, and Swiss Foundation for Research. GM has served as scientific consultant for Gilead, GSK/ViiV, Merck, Theratechnologies, Janssen; and has received funding support from Gilead, ViiV, Tetrphase, Roche, Genentech, Vanda, Astellas, Merck. M.E. acts as member of the Scientific Board of PM International, President of the Gesellschaft für angewandte Vitaminforschung and consults nutrition and supplement companies on request. A.F.G. has received research funding from Bayer Consumer Care; has acted as an advisor/consultant for GSK, DSM, Kellogg's and The Coca-Cola Company; and has received reimbursement for travel and/or speaking from Bayer Consumer Care. A.R.M. declares receipt of funding in the last 36 months to support vitamin D research from the following companies who manufacture or sell vitamin D supplements: Pharma Nord Ltd, DSM Nutritional Products Ltd, Thornton & Ross Ltd and Hyphens Pharma Ltd. A.R.M. also declares support for attending meetings from the following companies who manufacture or sell vitamin D supplements: Pharma Nord Ltd and Abiogen Pharma Ltd. A.R.M. also declares participation on the Data and Safety Monitoring Board for the VITALITY trial (Vitamin D for Adolescents with HIV to reduce musculoskeletal morbidity and immunopathology). A.R.M. also declares unpaid work as a Programme Committee member for the Vitamin D Workshop. A.R.M. also declares receipt of vitamin D capsules for clinical trial use from Pharma Nord Ltd, Synergy Biologics Ltd and Cytoplan Ltd.

References

1. World Health Organization COVID-19 Vaccine Tracker. Available online: <https://covid19.trackvaccines.org/agency/who/> (accessed on 10 February 2022).
2. Wikipedia List of COVID-19 Vaccine Authorizations. Available online: https://en.wikipedia.org/wiki/List_of_COVID-19_vaccine_authorizations (accessed on 20 January 2022).
3. COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU). Available online: <https://coronavirus.jhu.edu/map.html> (accessed on 28 February 2022).
4. Iuliano, A.D.; Brunkard, J.M.; Boehmer, T.K.; Peterson, E.; Adjei, S.; Binder, A.M.; Cobb, S.; Graff, P.; Hidalgo, P.; Panaggio, M.J.; et al. Trends in Disease Severity and Health Care Utilization During the Early Omicron Variant

- Period Compared with Previous SARS-CoV-2 High Transmission Periods — United States, December 2020–January 2022. *MMWR* **2022**, *71*, 146–152, doi:10.15585/mmwr.mm7104e4. 308
309
5. World Health Organization Influenza (Seasonal). Available online: [https://www.who.int/news-room/fact-sheets/detail/influenza-\(seasonal\)](https://www.who.int/news-room/fact-sheets/detail/influenza-(seasonal)) (accessed on 10 February 2022). 310
311
6. Centers for Disease Control and Prevention COVID Data Tracker: Rates of COVID-19 Cases and Deaths by Vaccination Status. Available online: <https://covid.cdc.gov/covid-data-tracker/#rates-by-vaccine-status> (accessed on 1 March 2022). 312
313
314
7. Collie, S.; Champion, J.; Moultrie, H.; Bekker, L.-G.; Gray, G. Effectiveness of BNT162b2 Vaccine against Omicron Variant in South Africa. *N Engl J Med* **2021**, NEJMc2119270, doi:10.1056/NEJMc2119270. 315
316
8. Planas, D.; Veyer, D.; Baidaliuk, A.; Staropoli, I.; Guivel-Benhassine, F.; Rajah, M.M.; Planchais, C.; Porrot, F.; Robillard, N.; Puech, J.; et al. Reduced Sensitivity of SARS-CoV-2 Variant Delta to Antibody Neutralization. *Nature* **2021**, *596*, 276–280, doi:10.1038/s41586-021-03777-9. 317
318
319
9. Tartof, S.Y.; Slezak, J.M.; Fischer, H.; Hong, V.; Ackerson, B.K.; Ranasinghe, O.N.; Frankland, T.B.; Ogun, O.A.; Zamparo, J.M.; Gray, S.; et al. Effectiveness of mRNA BNT162b2 COVID-19 Vaccine up to 6 Months in a Large Integrated Health System in the USA: A Retrospective Cohort Study. *Lancet* **2021**, *398*, 1407–1416, doi:10.1016/S0140-6736(21)02183-8. 320
321
322
323
10. Andrews, N.; Tessier, E.; Stowe, J.; Gower, C.; Kirsebom, F.; Simmons, R.; Gallagher, E.; Chand, M.; Brown, K.; Ladhani, S.N.; et al. Vaccine Effectiveness and Duration of Protection of Comirnaty, Vaxzevria and Spikevax against Mild and Severe COVID-19 in the UK. medRxiv preprint doi: <https://doi.org/10.1101/2021.09.15.21263583>, 2021; 324
325
326
327
11. Goodwin, K.; Viboud, C.; Simonsen, L. Antibody Response to Influenza Vaccination in the Elderly: A Quantitative Review. *Vaccine* **2006**, *24*, 1159–1169, doi:10.1016/j.vaccine.2005.08.105. 328
329
12. Müller, L.; Andrée, M.; Moskorz, W.; Drexler, I.; Walotka, L.; Grothmann, R.; Ptok, J.; Hillebrandt, J.; Ritchie, A.; Rabl, D.; et al. Age-Dependent Immune Response to the Biontech/Pfizer BNT162b2 Coronavirus Disease 2019 Vaccination. *Clin Infect Dis.* **2021**, *73*, 2065–2072, doi:10.1093/cid/ciab381. 330
331
332
13. Gombart, A.F.; Pierre, A.; Maggini, S. A Review of Micronutrients and the Immune System—Working in Harmony to Reduce the Risk of Infection. *Nutrients* **2020**, *12*, 236, doi:10.3390/nu12010236. 333
334
14. Calder, P.C. Foods to Deliver Immune-Supporting Nutrients. *Curr Opin Food Sci* **2022**, *43*, 136–145, doi:10.1016/j.cofs.2021.12.006. 335
336
15. Calder, P.C. Nutrition, Immunity and COVID-19. *BMJNPH* **2020**, *3*, 74–92, doi:10.1136/bmjnph-2020-000085. 337
16. Calder, P.C.; Carr, A.C.; Gombart, A.F.; Eggersdorfer, M. Optimal Nutritional Status for a Well-Functioning Immune System Is an Important Factor to Protect against Viral Infections. *Nutrients* **2020**, *12*, 1181, doi:10.3390/nu12041181. 338
339
340
17. European Commission of the European Union EU Register of Nutrition and Health Claims Made on Foods. Available online: https://ec.europa.eu/food/safety/labelling_nutrition/claims/register/public/?event=register.home (accessed on 10 February 2022). 341
342
343
344
18. Ao, T.; Kikuta, J.; Ishii, M. The Effects of Vitamin D on Immune System and Inflammatory Diseases. *Biomolecules* **2021**, *11*, 1624, doi:10.3390/biom11111624. 345
346
19. Bilezikian, J.P.; Bikle, D.; Hewison, M.; Lazaretti-Castro, M.; Formenti, A.M.; Gupta, A.; Madhavan, M.V.; Nair, N.; Babalyan, V.; Hutchings, N.; et al. MECHANISMS IN ENDOCRINOLOGY: Vitamin D and COVID-19. *Eur J Endocrinol* **2020**, *183*, R133–R147, doi:10.1530/EJE-20-0665. 347
348
349

20. Edfeldt, K.; Liu, P.T.; Chun, R.; Fabri, M.; Schenk, M.; Wheelwright, M.; Keegan, C.; Krutzik, S.R.; Adams, J.S.; Hewison, M.; et al. T-Cell Cytokines Differentially Control Human Monocyte Antimicrobial Responses by Regulating Vitamin D Metabolism. *Proc Natl Acad Sci* **2010**, *107*, 22593–22598, doi:10.1073/pnas.1011624108. 350–352
21. Charoenngam, N.; Holick, M.F. Immunologic Effects of Vitamin D on Human Health and Disease. *Nutrients* **2020**, *12*, 2097, doi:10.3390/nu12072097. 353–354
22. Cannell, J.J.; Vieth, R.; Umhau, J.C.; Holick, M.F.; Grant, W.B.; Madronich, S.; Garland, C.F.; Giovannucci, E. Epidemic Influenza and Vitamin D. *Epidemiol. Infect.* **2006**, *134*, 1129–1140, doi:10.1017/S0950268806007175. 355–356
23. Autier, P.; Mullie, P.; Macacu, A.; Dragomir, M.; Boniol, M.; Coppens, K.; Pizot, C.; Boniol, M. Effect of Vitamin D Supplementation on Non-Skeletal Disorders: A Systematic Review of Meta-Analyses and Randomised Trials. *Lancet Diabetes Endocrinol* **2017**, *5*, 986–1004, doi:10.1016/S2213-8587(17)30357-1. 357–359
24. Rejnmark, L.; Bislev, L.S.; Cashman, K.D.; Eiríksdóttir, G.; Gaksch, M.; Grübler, M.; Grimnes, G.; Gudnason, V.; Lips, P.; Pilz, S.; et al. Non-Skeletal Health Effects of Vitamin D Supplementation: A Systematic Review on Findings from Meta-Analyses Summarizing Trial Data. *PLoS ONE* **2017**, *12*, e0180512, doi:10.1371/journal.pone.0180512. 360–363
25. Martineau, A.R.; Jolliffe, D.A.; Hooper, R.L.; Greenberg, L.; Aloia, J.F.; Bergman, P.; Dubnov-Raz, G.; Esposito, S.; Ganmaa, D.; Ginde, A.A.; et al. Vitamin D Supplementation to Prevent Acute Respiratory Tract Infections: Systematic Review and Meta-Analysis of Individual Participant Data. *BMJ* **2017**, *356*, doi:10.1136/bmj.i6583. 364–366
26. Jolliffe, D.A.; Camargo, C.A.; Sluyter, J.D.; Aglipay, M.; Aloia, J.F.; Ganmaa, D.; Bergman, P.; Bischoff-Ferrari, H.A.; Borzutzky, A.; Damsgaard, C.T.; et al. Vitamin D Supplementation to Prevent Acute Respiratory Infections: A Systematic Review and Meta-Analysis of Aggregate Data from Randomised Controlled Trials. *Lancet Diabetes Endocrinol* **2021**, *9*, 276–292, doi:10.1016/S2213-8587(21)00051-6. 367–370
27. Kaufman, H.W.; Niles, J.K.; Kroll, M.H.; Bi, C.; Holick, M.F. SARS-CoV-2 Positivity Rates Associated with Circulating 25-Hydroxyvitamin D Levels. *PLoS ONE* **2020**, *15*, e0239252, doi:10.1371/journal.pone.0239252. 371–372
28. Chiodini, I.; Gatti, D.; Soranna, D.; Merlotti, D.; Mingiano, C.; Fassio, A.; Adami, G.; Falchetti, A.; Eller-Vainicher, C.; Rossini, M.; et al. Vitamin D Status and SARS-CoV-2 Infection and COVID-19 Clinical Outcomes. *Front. Public Health* **2021**, *9*, 736665, doi:10.3389/fpubh.2021.736665. 373–375
29. Desai, A.P.; Dirajlal-Fargo, S.; Durieux, J.C.; Tribout, H.; Labbato, D.; McComsey, G.A. Vitamin K & D Deficiencies Are Independently Associated With COVID-19 Disease Severity. *Open Forum Infect Dis* **2021**, *8*, ofab408, doi:10.1093/ofid/ofab408. 376–378
30. Oristrell, J.; Oliva, J.C.; Casado, E.; Subirana, I.; Domínguez, D.; Toloba, A.; Balado, A.; Grau, M. Vitamin D Supplementation and COVID-19 Risk: A Population-Based, Cohort Study. *J Endocrinol Invest* **2022**, *45*, 167–179, doi:10.1007/s40618-021-01639-9. 379–381
31. Loucera, C.; Peña-Chilet, M.; Esteban-Medina, M.; Muñozerro-Muñiz, D.; Villegas, R.; Lopez-Miranda, J.; Rodriguez-Baño, J.; Túnez, I.; Bouillon, R.; Dopazo, J.; et al. Real World Evidence of Calcifediol or Vitamin D Prescription and Mortality Rate of COVID-19 in a Retrospective Cohort of Hospitalized Andalusian Patients. *Sci Rep* **2021**, *11*, 23380, doi:10.1038/s41598-021-02701-5. 382–385
32. Ma, H.; Zhou, T.; Heianza, Y.; Qi, L. Habitual Use of Vitamin D Supplements and Risk of Coronavirus Disease 2019 (COVID-19) Infection: A Prospective Study in UK Biobank. *Am J Clin Nutr* **2021**, *113*, 1275–1281, doi:10.1093/ajcn/nqaa381. 386–388
33. Holt, H.; Talaei, M.; Greenig, M.; Zenner, D.; Symons, J.; Relton, C.; Young, K.S.; Davies, M.R.; Thompson, K.N.; Ashman, J.; et al. Risk Factors for Developing COVID-19: A Population-Based Longitudinal Study (COVIDENCE UK). *Thorax* **2021**, thoraxjnl-2021-217487, doi:10.1136/thoraxjnl-2021-217487. 389–391

34. Talaei, M.; Faustini, S.; Holt, H.; Jolliffe, D.A.; Vivaldi, G.; Greenig, M.; Perdek, N.; Maltby, S.; Bigogno, C.M.; Symons, J.; et al. Determinants of Pre-Vaccination Antibody Responses to SARS-CoV-2: A Population-Based Longitudinal Study (COVIDENCE UK). *BMC Med* **2022**, *20*, 87, doi:10.1186/s12916-022-02286-4.
35. Dissanayake, H.A.; de Silva, N.L.; Sumanatilleke, M.; de Silva, S.D.N.; Gamage, K.K.K.; Dematapitiya, C.; Kuruppu, D.C.; Ranasinghe, P.; Pathmanathan, S.; Katulanda, P. Prognostic and Therapeutic Role of Vitamin D in COVID-19: Systematic Review and Meta-Analysis. *J Clin Endocrinol Metab* **2021**, dgab892, doi:10.1210/clinem/dgab892.
36. Netea, M.G.; Giamarellos-Bourboulis, E.J.; Domínguez-Andrés, J.; Curtis, N.; van Crevel, R.; van de Veerdonk, F.L.; Bonten, M. Trained Immunity: A Tool for Reducing Susceptibility to and the Severity of SARS-CoV-2 Infection. *Cell* **2020**, *181*, 969–977, doi:10.1016/j.cell.2020.04.042.
37. Maggini, S.; Pierre, A.; Calder, P. Immune Function and Micronutrient Requirements Change over the Life Course. *Nutrients* **2018**, *10*, 1531, doi:10.3390/nu10101531.
38. Maggini, S.; Wintergerst, E.S.; Beveridge, S.; Hornig, D.H. Selected Vitamins and Trace Elements Support Immune Function by Strengthening Epithelial Barriers and Cellular and Humoral Immune Responses. *Br J Nutr* **2007**, *98*, S29–S35, doi:10.1017/S0007114507832971.
39. Chiu, S.-K.; Tsai, K.-W.; Wu, C.-C.; Zheng, C.-M.; Yang, C.-H.; Hu, W.-C.; Hou, Y.-C.; Lu, K.-C.; Chao, Y.-C. Putative Role of Vitamin D for COVID-19 Vaccination. *IJMS* **2021**, *22*, 8988, doi:10.3390/ijms22168988.
40. Velikova, T.; Fabbri, A.; Infante, M. The Role of Vitamin D as a Potential Adjuvant for COVID-19 Vaccines. *Eur Rev Med Pharmacol Sci* **2021**, *25*, 5323–5327.
41. Lai, Y.-J.; Chang, H.-S.; Yang, Y.-P.; Lin, T.-W.; Lai, W.-Y.; Lin, Y.-Y.; Chang, C.-C. The Role of Micronutrient and Immunomodulation Effect in the Vaccine Era of COVID-19. *J Chin Med Assoc* **2021**, *84*, 821–826, doi:10.1097/JCMA.0000000000000587.
42. Lee, M.-D.; Lin, C.-H.; Lei, W.-T.; Chang, H.-Y.; Lee, H.-C.; Yeung, C.-Y.; Chiu, N.-C.; Chi, H.; Liu, J.-M.; Hsu, R.-J.; et al. Does Vitamin D Deficiency Affect the Immunogenic Responses to Influenza Vaccination? A Systematic Review and Meta-Analysis. *Nutrients* **2018**, *10*, 409, doi:10.3390/nu10040409.
43. Broome, C.S.; McArdle, F.; Kyle, J.A.; Andrews, F.; Lowe, N.M.; Hart, C.A.; Arthur, J.R.; Jackson, M.J. An Increase in Selenium Intake Improves Immune Function and Poliovirus Handling in Adults with Marginal Selenium Status. *Am J Clin Nutr* **2004**, *80*, 154–162, doi:10.1093/ajcn/80.1.154.
44. Stoffel, N.U.; Uyoga, M.A.; Mutuku, F.M.; Frost, J.N.; Mwasi, E.; Paganini, D.; van der Klis, F.R.M.; Malhotra, I.J.; LaBeaud, A.D.; Ricci, C.; et al. Iron Deficiency Anemia at Time of Vaccination Predicts Decreased Vaccine Response and Iron Supplementation at Time of Vaccination Increases Humoral Vaccine Response: A Birth Cohort Study and a Randomized Trial Follow-Up Study in Kenyan Infants. *Front. Immunol.* **2020**, *11*, 1313, doi:10.3389/fimmu.2020.01313.
45. Chillon, T.S.; Demircan, K.; Heller, R.A.; Hirschbil-Bremer, I.M.; Diegmann, J.; Bachmann, M.; Moghaddam, A.; Schomburg, L. Relationship between Vitamin D Status and Antibody Response to COVID-19 mRNA Vaccination in Healthy Adults. *Biomedicines* **2021**, *9*, 1714, doi:10.3390/biomedicines9111714.
46. Parthymou, A.; Habeos, E.E.; Habeos, G.I.; Deligakis, A.; Livieratos, E.; Marangos, M.; Chartoumpekis, D.V. Sars-Cov-2 Antibody Titer 3 Months Post-Vaccination Is Affected by Age, Gender, Smoking and Vitamin D; medRxiv preprint doi: <https://doi.org/10.1101/2021.09.01.21262913>.
47. Jolliffe, D.A.; Faustini, S.E.; Holt, H.; Perdek, N.; Maltby, S.; Talaei, M.; Greenig, M.; Vivaldi, G.; Tydeman, F.; Symons, J.; et al. Determinants of Antibody Responses to Two Doses of ChAdOx1 NCoV-19 or BNT162b2 and a

- Subsequent Booster Dose of BNT162b2 or mRNA-1273: Population-Based Cohort Study (COVIDENCE UK); medRxiv preprint doi: <https://doi.org/10.1101/2022.02.14.22270930>. 433
434
48. Fulop, T.; Witkowski, J.M.; Pawelec, G.; Alan, C.; Larbi, A. On the Immunological Theory of Aging. In *Aging: Facts and Theories. Interdiscipl Top Gerontol.*; Robert, L., Fulop, T., Eds.; KARGER: Basel, Switzerland, 2014; Volume 39, pp. 163–176. 435
436
437
49. Meydani, S.N. Vitamin E Supplementation and in Vivo Immune Response in Healthy Elderly Subjects. A Randomized Controlled Trial. *JAMA* **1997**, *277*, 1380–1386, doi:10.1001/jama.277.17.1380. 438
439
50. De la Fuente, M.; Hernanz, A.; Guayerbas, N.; Victor, V.M.; Arnalich, F. Vitamin E Ingestion Improves Several Immune Functions in Elderly Men and Women. *Free Radic Res* **2008**, *42*, 272–280, doi:10.1080/10715760801898838. 440
441
51. Girodon, F. Impact of Trace Elements and Vitamin Supplementation on Immunity and Infections in Institutionalized Elderly Patients: A Randomized Controlled Trial. *Arch Intern Med* **1999**, *159*, 748, doi:10.1001/archinte.159.7.748. 442
443
444
52. Chambers, E.S.; Vukmanovic-Stejcic, M.; Turner, C.T.; Shih, B.B.; Trahair, H.; Pollara, G.; Tsaliki, E.; Rustin, M.; Freeman, T.C.; Mabbott, N.A.; et al. Vitamin D3 Replacement Enhances Antigen-Specific Immunity in Older Adults. *Immunother Adv* **2021**, *1*, Itaa008, doi:10.1093/immadv/Itaa008. 445
446
447
53. Pereira, B.; Xu, X.-N.; Akbar, A.N. Targeting Inflammation and Immunosenescence to Improve Vaccine Responses in the Elderly. *Front. Immunol.* **2020**, *11*, 583019, doi:10.3389/fimmu.2020.583019. 448
449
54. Reddi, K.; Henderson, B.; Meghji, S.; Wilson, M.; Poole, S.; Hopper, C.; Harris, M.; Hodges, S.J. Interleukin 6 Production by Lipopolysaccharide-Stimulated Human Fibroblasts Is Potently Inhibited by Naphthoquinone (Vitamin K) Compounds. *Cytokine* **1995**, *7*, 287–290, doi:10.1006/cyto.1995.0034. 450
451
452
55. Ohsaki, Y.; Shirakawa, H.; Miura, A.; Giriwono, P.E.; Sato, S.; Ohashi, A.; Iribe, M.; Goto, T.; Komai, M. Vitamin K Suppresses the Lipopolysaccharide-Induced Expression of Inflammatory Cytokines in Cultured Macrophage-like Cells via the Inhibition of the Activation of Nuclear Factor KB through the Repression of IKK α/β Phosphorylation. *J Nutr Biochem* **2010**, *21*, 1120–1126, doi:10.1016/j.jnutbio.2009.09.011. 453
454
455
456
56. Lakkireddy, M.; Gadiga, S.G.; Malathi, R.D.; Karra, M.L.; Raju, I.S.S.V.P.M.; Ragini; Chinapaka, S.; Baba, K.S.S.S.; Kandakarla, M. Impact of Daily High Dose Oral Vitamin D Therapy on the Inflammatory Markers in Patients with COVID 19 Disease. *Sci Rep* **2021**, *11*, 10641, doi:10.1038/s41598-021-90189-4. 457
458
459
57. Eckard, A.R.; O’Riordan, M.A.; Rosebush, J.C.; Lee, S.T.; Habib, J.G.; Ruff, J.H.; Labbato, D.; Daniels, J.E.; Uribe-Leitz, M.; Tangpricha, V.; et al. Vitamin D Supplementation Decreases Immune Activation and Exhaustion in HIV-1-Infected Youth. *Antivir Ther* **2017**, *23*, 315–324, doi:10.3851/IMP3199. 460
461
462
58. Janus, S.E.; Durieux, J.C.; Hajjari, J.; Carneiro, H.; McComsey, G.A. Inflammation Mediated Vitamin K and Vitamin D Effects on Vascular Calcifications in People with HIV on Active Antiretroviral Therapy. *AIDS* doi: 10.1097/QAD.0000000000003149. 463
464
465
59. Provinciali, M.; Montenovo, A.; Stefano, G.D.; Colombo, M.; Daghetta, L.; Cairati, M.; Veroni, C.; Cassino, R.; Torre, F.D.; Fabris, N. Effect of Zinc or Zinc plus Arginine Supplementation on Antibody Titre and Lymphocyte Subsets after Influenza Vaccination in Elderly Subjects: A Randomized Controlled Trial. *Age Ageing* **1998**, *27*, 715–722, doi:10.1093/ageing/27.6.715. 466
467
468
469
60. Crum-Cianflone, N.F.; Won, S.; Lee, R.; Lalani, T.; Ganesan, A.; Burgess, T.; Agan, B.K. Vitamin D Levels and Influenza Vaccine Immunogenicity among HIV-Infected and HIV-Uninfected Adults. *Vaccine* **2016**, *34*, 5040–5046, doi:10.1016/j.vaccine.2016.06.019. 470
471
472

-
61. Sundaram, M.E.; Talbot, H.K.; Zhu, Y.; Griffin, M.R.; Spencer, S.; Shay, D.K.; Coleman, L.A. Vitamin D Is Not Associated with Serologic Response to Influenza Vaccine in Adults over 50 Years Old. *Vaccine* **2013**, *31*, 2057–2061, doi:10.1016/j.vaccine.2013.02.028. 473–475
 62. Sundaram, M.E.; Meydani, S.N.; Vandermause, M.; Shay, D.K.; Coleman, L.A. Vitamin E, Vitamin A, and Zinc Status Are Not Related to Serologic Response to Influenza Vaccine in Older Adults: An Observational Prospective Cohort Study. *Nutr Res* **2014**, *34*, 149–154, doi:10.1016/j.nutres.2013.12.004. 476–478
 63. Food and Agriculture Organization of the United Nations. *Europe and Central Asia Regional Overview of Food Insecurity 2016: The Food Insecurity Transition.*; FAO: Budapest, Hungary, 2017; ISBN 978-92-5-109659-8. 479–480
 64. Bailey, R.L.; West Jr., K.P.; Black, R.E. The Epidemiology of Global Micronutrient Deficiencies. *Ann Nutr Metab* **2015**, *66*, 22–33, doi:10.1159/000371618. 481–482
 65. World Health Organization; Centers for Disease Control and Prevention (U.S.). *Worldwide Prevalence of Anaemia 1993–2005: WHO Global Database of Anaemia*; World Health Organization: Geneva, Switzerland, 2008; ISBN 978-92-4-159665-7. 483–485
 66. World Health Organization. *The World Health Report 2002: Reducing Risks, Promoting Healthy Life*, WHO, Geneva, Switzerland, 2002. 486–487
 67. Hilger, J.; Friedel, A.; Herr, R.; Rausch, T.; Roos, F.; Wahl, D.A.; Pierroz, D.D.; Weber, P.; Hoffmann, K. A Systematic Review of Vitamin D Status in Populations Worldwide. *Br J Nutr* **2014**, *111*, 23–45, doi:10.1017/S0007114513001840. 488–490
 68. Peter, S.; Friedel, A.; Roos, F.F.; Wyss, A.; Eggersdorfer, M.; Hoffmann, K.; Weber, P. A Systematic Review of Global Alpha-Tocopherol Status as Assessed by Nutritional Intake Levels and Blood Serum Concentrations. *Int J Vitam Nutr Res*. **85**, 261–281. 491–493
 69. Rowe, S.; Carr, A.C. Global Vitamin C Status and Prevalence of Deficiency: A Cause for Concern? *Nutrients* **2020**, *12*, 2008, doi:10.3390/nu12072008. 494–495
 70. Vural, Z.; Avery, A.; Kalogiros, D.I.; Coneyworth, L.J.; Welham, S.J.M. Trace Mineral Intake and Deficiencies in Older Adults Living in the Community and Institutions: A Systematic Review. *Nutrients* **2020**, *12*, 1072, doi:10.3390/nu12041072. 496–498
 71. Infante, M.; Ricordi, C.; Baidal, D.A.; Alejandro, R.; Lanzoni, G.; Sears, B.; Caprio, M.; Fabbri, A. VITAL Study: An Incomplete Picture? *Eur Rev Med Pharmacol Sci*. **23**, 3142–3147. 499–500
 72. Borsche, L.; Glauner, B.; von Mendel, J. COVID-19 Mortality Risk Correlates Inversely with Vitamin D3 Status, and a Mortality Rate Close to Zero Could Theoretically Be Achieved at 50 ng/mL 25(OH)D3: Results of a Systematic Review and Meta-Analysis. *Nutrients* **2021**, *13*, 3596, doi:10.3390/nu13103596. 501–503
 73. U.S. Centers for Disease Control and Prevention CDC Seasonal Flu Vaccine Effectiveness Studies. Available online: <https://www.cdc.gov/flu/vaccines-work/effectiveness-studies.htm> (accessed on 1 March 2022). 504–506