

A NOTE ON TORSION LENGTH AND TORSION SUBGROUPS

IAN J. LEARY AND ASHOT MINASYAN

ABSTRACT. Answering Questions 19.23 and 19.24 from the Kourovka notebook we construct polycyclic groups with arbitrary torsion lengths and give examples of finitely presented groups whose quotients by their torsion subgroups are not finitely presented.

1. INTRODUCTION

Given a group G , the *torsion subgroup*, $\text{Tor}(G)$, is the normal subgroup of G generated by all elements of finite order. The torsion length of G is defined as follows. Let $G_0 = G$ and $G_{n+1} = G_n/\text{Tor}(G_n)$, for $n \geq 0$. The *torsion length* of G , $\text{TorLen}(G)$, is the smallest $n \geq 0$ such that G_n is torsion-free. If no such n exists then we define $\text{TorLen}(G) = \omega$. Torsion length was defined and studied by Chiodo and Vyas in [3], and in [4] the same authors constructed finitely presented groups of arbitrary torsion length.

Chiodo and Vyas asked in [6, Questions 19.23 and 19.24] whether there exist finitely presented soluble groups of torsion length greater than 2 and whether there exists a finitely presented group G such that $G/\text{Tor}(G)$ is not finitely presented. Versions of these questions also appeared in [3, Subsection 3.3] and [4, Question 1]. In this note we answer both questions affirmatively.

Acknowledgements. The authors thank the anonymous referee for suggesting changes which led to the concepts introduced in Subsections 2.1 and 2.2.

2. POLYCYCLIC GROUPS OF ARBITRARILY LARGE TORSION LENGTH

2.1. Neutral homomorphisms and automorphisms of semidirect products. In this subsection we develop a few tools that will be used in the main construction.

Definition 2.1. Let K be a group and let $\beta : K \rightarrow K$ be an endomorphism.

- We define the map $\Delta_\beta : K \rightarrow K$ by $\Delta_\beta(k) = k^{-1}\beta(k)$, for all $k \in K$.
- We denote by $\mathcal{N}(\beta) \triangleleft K$ the normal closure of the subset $\Delta_\beta(K) = \{k^{-1}\beta(k) \mid k \in K\}$.
- If C is another group and $\phi : K \rightarrow C$ is a homomorphism, we say that ϕ is β -neutral if $\phi \circ \beta = \phi$, i.e., if $\phi(\beta(k)) = \phi(k)$ for all $k \in K$.

The map Δ_β can be thought of as a (non-abelian) 1-cocycle from K to K , where K acts on itself by conjugation (cf. [8, Section I.5.1]).

Lemma 2.2. Suppose that K is a group generated by a set X and $\beta \in \text{End}(K)$.

- (i) If $\phi : K \rightarrow C$ is a group homomorphism then ϕ is β -neutral if and only if $\mathcal{N}(\beta) \subseteq \ker \phi$ if and only if $\Delta_\beta(X) \subseteq \ker \phi$.
- (ii) $\mathcal{N}(\beta)$ is the normal closure in K of the subset $\Delta_\beta(X) = \{x^{-1}\beta(x) \mid x \in X\}$.

Proof. Claim (i) follows from the observation that for any $k \in K$, $\phi(\beta(k)) = \phi(k)$ is equivalent to $\phi(k^{-1}\beta(k)) = 1$ and the fact that two homomorphisms from G to C are equal if and only if they agree on X .

1991 Mathematics Subject Classification. 20F99, 20F16.

Key words and phrases. Torsion subgroups, torsion length, polycyclic groups.

For claim (ii), let $C = K/N$, where N is the normal closure of $\Delta_\beta(X)$ in K , and let $\phi : K \rightarrow C$ be the natural epimorphism. Then, according to claim (i), ϕ is β -neutral and $\mathcal{N}(\beta) \subseteq \ker \phi = N$. Clearly $N \subseteq \mathcal{N}(\beta)$, by definition, hence $\mathcal{N}(\beta) = N$. \square

Definition 2.3. Let K, H be two groups, let $\beta \in \text{Aut}(K)$ be an automorphism of K and let $\phi : K \rightarrow \text{Aut}(H)$ be a homomorphism. Denote by A the semidirect product $H \rtimes_\phi K$. We define the *extension of β to A* as the map $\beta_A : A \rightarrow A$, given by

$$\beta_A((h, k)) = (h, \beta(k)), \text{ for all } h \in H \text{ and all } k \in K.$$

In this paper we regard the semidirect product $A = H \rtimes_\phi K$ as the set $H \times K$, with the multiplication $(h_1, k_1)(h_2, k_2) = (h_1\phi(k_1)(h_2), k_1k_2)$, for all $(h_1, k_1), (h_2, k_2) \in A$.

Lemma 2.4. *Using the notation of Definition 2.3, suppose that ϕ is β -neutral. Then the following properties hold:*

- (a) *the map β_A is an automorphism of A which has the same order as β ;*
- (b) *$\mathcal{N}(\beta_A) = (1, \mathcal{N}(\beta))$ in A ;*
- (c) *the quotient $A/\mathcal{N}(\beta_A)$ is naturally isomorphic to the semidirect product $H \rtimes_{\bar{\phi}} (K/\mathcal{N}(\beta))$, where $\bar{\phi} : K/\mathcal{N}(\beta) \rightarrow \text{Aut}(H)$ is the homomorphism induced by ϕ .*

Proof. (a) Evidently β_A is a bijection with the same order as β . Consider any $(h_1, k_1), (h_2, k_2) \in A$, where $h_1, h_2 \in H$ and $k_1, k_2 \in K$, and observe that

$$\beta_A((h_1, k_1)(h_2, k_2)) = \beta_A((h_1\phi(k_1)(h_2), k_1k_2)) = (h_1\phi(k_1)(h_2), \beta(k_1)\beta(k_2)).$$

On the other hand,

$$\beta_A((h_1, k_1))\beta_A((h_2, k_2)) = (h_1, \beta(k_1))(h_2, \beta(k_2)) = (h_1\phi(\beta(k_1))(h_2), \beta(k_1)\beta(k_2)).$$

Since ϕ is β -neutral, we see that $\beta_A((h_1, k_1)(h_2, k_2)) = \beta_A((h_1, k_1))\beta_A((h_2, k_2))$, hence β_A is an automorphism of A .

(b) By Lemma 2.2(i), $\mathcal{N}(\beta) \subseteq \ker \phi$, which means that the subgroup $(1, \mathcal{N}(\beta))$ centralizes the subgroup $(H, 1)$ in A . Since $\mathcal{N}(\beta) \triangleleft K$ and A is generated by $(H, 1) \cup (1, K)$, it follows that $(1, \mathcal{N}(\beta)) \triangleleft A$. Now, according to Lemma 2.2(ii), $\mathcal{N}(\beta_A)$ is the normal closure in A of the subset

$$\Delta_{\beta_A}((H, 1)) \cup \Delta_{\beta_A}((1, K)) = \{(1, 1)\} \cup (1, \Delta_\beta(K)) = (1, \Delta_\beta(K)).$$

Therefore $\mathcal{N}(\beta_A) \subseteq (1, \mathcal{N}(\beta)) \subseteq \mathcal{N}(\beta_A)$, i.e., $\mathcal{N}(\beta_A) = (1, \mathcal{N}(\beta))$.

Claim (c) is a consequence of claim (b) and the fact that $\mathcal{N}(\beta) \subseteq \ker \phi$. \square

2.2. Exact automorphisms. Throughout this subsection we assume that F is a group and $\beta \in \text{Aut}(F)$ is an automorphism of finite order $m \in \mathbb{N}$. We define the map $\Sigma_\beta : F \rightarrow F$ by

$$\Sigma_\beta(f) = f\beta(f) \dots \beta^{m-1}(f), \text{ for all } f \in F.$$

Remark 2.5. It is easy to check that the image $\text{Im } \Delta_\beta$ is always contained in the kernel $\ker \Sigma_\beta$.

The above remark naturally motivates the following terminology, which was originally proposed to us by the referee.

Definition 2.6. We will say that the automorphism $\beta \in \text{Aut}(F)$ is *exact* if $\text{Im } \Delta_\beta = \ker \Sigma_\beta$.

For example, the order 2 automorphism of \mathbb{Z}^2 , which interchanges the standard generators, is exact. More generally, if F is a direct product of n copies of a group K and β is an automorphism of F cyclically permuting the factors then β is exact. On the other hand, the automorphism of \mathbb{Z} sending 1 to -1 is not exact.

We will use exact automorphisms to control torsion in extensions by β .

Lemma 2.7. *Let G be a group with a torsion-free normal subgroup $F \triangleleft G$ and an element $x \in G$ of finite order $n \in \mathbb{N}$, such that $G = \langle F, x \rangle$. Let $\beta \in \text{Aut}(F)$ denote the automorphism induced by conjugation by x . The following are equivalent:*

- (i) β is an exact automorphism of F ;
- (ii) if an element of the form fx , for some $f \in F$, has finite order in G then it is conjugate to x in G ;
- (iii) G contains exactly one conjugacy class of cyclic subgroups of order n .

Proof. By the assumptions, β must have order $m \in \mathbb{N}$ in $\text{Aut}(F)$, where $n = ml$, for some $l \in \mathbb{N}$. Therefore, for any $f \in F$ we have

$$(1) \quad (fx)^n = f(xfx^{-1})(x^2fx^{-2}) \dots (x^{n-1}fx^{-n+1})x^n = f\beta(f)\beta^2(f) \dots \beta^{n-1}(f) = (\Sigma_\beta(f))^l.$$

Let us show that (i) implies (ii). Suppose that β is an exact automorphism of F and $fx \in G$ is an element of finite order, for some $f \in F$. Equation (1) yields that $\Sigma_\beta(f)$ has finite order in F , and since F is torsion-free we can conclude that $\Sigma_\beta(f) = 1$. Hence $f = \Delta_\beta(h)$, for some $h \in F$, by exactness. Therefore

$$fx = h^{-1}\beta(h)x = h^{-1}xhx^{-1}x = h^{-1}xh,$$

so (ii) holds.

To show that (ii) implies (i), suppose that $f \in \ker \Sigma_\beta$ in F . Then $(fx)^n = 1$ by (1), so, in view of (ii), there must exist an element $g \in G$ such that $fx = g^{-1}xg$. Now, $g = x^t h$, for some $t \in \mathbb{Z}$ and some $h \in F$, because G is generated by F and x and $F \triangleleft G$. Hence

$$fx = g^{-1}xg = h^{-1}xh = h^{-1}\beta(h)x,$$

so $f = \Delta_\beta(h)$. Thus $\beta \in \text{Aut}(F)$ is exact, as required.

We have shown that (i) is equivalent to (ii). The equivalence between (ii) and (iii) is an easy consequence of the observation that every cyclic subgroup of order n in G contains precisely one element of the form fx , for $f \in F$, which will generate this subgroup. \square

The equivalence between (i) and (iii) in the previous lemma gives the following corollary.

Corollary 2.8. *Suppose that F is a torsion-free group with an automorphism β of order $m \in \mathbb{N}$. If β is exact then so is $\beta^p \in \text{Aut}(F)$, for any integer p coprime to m .*

Proof. Let $G = F \rtimes \langle \beta \rangle$ be the natural semidirect product of F with $\langle \beta \rangle \cong C_m$. We identify F with its image in G and let $x \in G$ denote the image of β . Then $F \triangleleft G$, $G = \langle F, x \rangle$ and $xfx^{-1} = \beta(f)$, for all $f \in F$.

Take any $p \in \mathbb{Z}$, coprime to m . Since x has order m in G , x^p also has order m , so $G = \langle F, x^p \rangle$ and conjugation by x^p induces the automorphism β^p on F . If $\beta \in \text{Aut}(F)$ is exact then G has exactly one conjugacy class of cyclic subgroups of order m , by Lemma 2.7. Therefore, the same lemma shows that $\beta^p \in \text{Aut}(F)$ is also exact. \square

Remark 2.9. The results of this subsection can be expressed in terms of the first non-abelian cohomology group of the cyclic group C_n with coefficients in a group F , which we compute using the presentation 2-complex associated to $\langle \beta \mid \beta^n \rangle$ rather than the general approach described in [8, Section I.5.1]. In this setting, the kernel of Σ_β is the set of 1-cocycles, and the image of Δ_β is the 1-coboundaries. The automorphism β of F is exact if and only if the pointed set $H^1(C_n; F)$ has just one element, and Lemma 2.7 is closely related to [8, Exercise 1 in Section I.5.1,].

The final lemma in this subsection verifies that extensions of exact automorphisms to semidirect products, as discussed in Subsection 2.1, are again exact.

Lemma 2.10. *Let K, H be groups, let $\beta \in \text{Aut}(K)$ be a finite order automorphism. Suppose that $\phi : K \rightarrow \text{Aut}(H)$ is a β -neutral homomorphism and $A = H \rtimes_{\phi} K$ is the corresponding semidirect product. If H is torsion-free and β is an exact automorphism of K then its extension β_A , given by Definition 2.3, is an exact automorphism of A .*

Proof. Suppose that β has order $m \in \mathbb{N}$ in $\text{Aut}(K)$. Then $\beta_A \in \text{Aut}(A)$ and it has the same order m , by Lemma 2.4(a). Observe that for any $h \in H$ and $k \in K$ we have

$$\begin{aligned} \Sigma_{\beta_A}((h, k)) &= (h, k)(h, \beta(k)) \dots (h, \beta^{m-1}(k)) \\ (2) \quad &= (h \cdot \phi(k)(h) \cdot \phi(k\beta(k))(h) \dots \phi(k\beta(k) \dots \beta^{m-2}(k))(h), k\beta(k) \dots \beta^{m-1}(k)) \\ &= (h \cdot \phi(k)(h) \cdot \phi(k^2)(h) \dots \phi(k^{m-1})(h), \Sigma_{\beta}(k)), \end{aligned}$$

where in the last equality we used the fact that ϕ is β -neutral.

Now, assume that $\Sigma_{\beta_A}((h, k)) = (1, 1)$. Then $\Sigma_{\beta}(k) = 1$ by (2), so $k = \Delta_{\beta}(u)$, for some $u \in K$, by exactness. Lemma 2.2(i) implies that $\phi(k) = \text{Id}_H$, so (2) shows that

$$\Sigma_{\beta_A}((h, k)) = (h^m, 1) = (1, 1).$$

Thus $h^m = 1$, whence $h = 1$ as H is torsion-free. Consequently,

$$(h, k) = (1, u^{-1}\beta(u)) = (1, u)^{-1} \beta_A((1, u)) \in \Delta_{\beta_A}(A).$$

Therefore $\ker \Sigma_{\beta_A} = \Delta_{\beta_A}(A)$, as required. \square

2.3. The main construction. The building block for our examples will be the group K , given by the presentation

$$(3) \quad K = \langle a, s, b, t \mid a^s = a^{-1}, b^t = b^{-1}, [a, b] = [a, t] = [s, b] = [s, t] = 1 \rangle,$$

where we write $a^s = sas^{-1}$ and $[a, b] = aba^{-1}b^{-1}$.

Clearly $K \cong \pi \times \pi$, where $\pi = \langle a, s \mid a^s = a^{-1} \rangle$ is the fundamental group of the Klein bottle. In particular, K is torsion-free, metabelian and polycyclic of Hirsch length 4.

We define $\beta \in \text{Aut}(K)$ to be the order 2 automorphism interchanging the two copies of π , i.e., $\beta(a) = b$, $\beta(s) = t$, $\beta(b) = a$ and $\beta(t) = s$.

Let C_2 be the cyclic group of order 2, generated by an element α , and let $\phi : K \rightarrow C_2$ be the homomorphism sending a and b to α and s and t to 1. Verification of the following elementary properties is left for the reader (Lemma 2.2(ii) may be useful for claim (ii)).

Lemma 2.11. *The group K , its automorphism β and the homomorphism ϕ enjoy the following properties.*

- (i) $\phi \circ \beta = \phi$, i.e., ϕ is β -neutral.
- (ii) The quotient $K/\mathcal{N}(\beta)$ is isomorphic to $C_2 \times C_{\infty}$, where C_2 is generated by the image of a and C_{∞} is generated by the image of s .
- (iii) The automorphism β of K is exact.

We now construct auxiliary torsion-free polycyclic groups H_n as iterated semidirect products of K with itself n times. These extensions will be defined by induction on n .

Construction 2.12. Set $H_0 = \{1\}$ and let $\beta_0 \in \text{Aut}(H_0)$ be the identity automorphism; set $H_1 = K$ and let $\beta_1 \in \text{Aut}(H_1)$ be the automorphism β defined above. Now suppose that the group H_{n-1} and an order 2 automorphism $\beta_{n-1} \in \text{Aut}(H_{n-1})$ have already been constructed, for some $n \geq 2$. We define the group H_n as the semidirect product

$$H_n = H_{n-1} \rtimes_{\phi_{n-1}} K,$$

where $\phi_{n-1} : K \rightarrow \text{Aut}(H_{n-1})$ is essentially the homomorphism ϕ defined above, whose image is the cyclic subgroup $\langle \beta_{n-1} \rangle$.

By Lemma 2.11(i), ϕ_{n-1} is β -neutral, so we can use Definition 2.3 and Lemma 2.4 to define an automorphism $\beta_n \in \text{Aut}(H_n)$ as the extension β_{H_n} , of β to H_n . Thus

$$\beta_n((h, k)) = (h, \beta(k)), \text{ for all } h \in H_{n-1} \text{ and all } k \in K.$$

Proposition 2.13. *For all $n \in \mathbb{N}$, the group H_n satisfies the following properties:*

- (i) *the quotient $H_n/\mathcal{N}(\beta_n)$ is isomorphic to $(H_{n-1} \rtimes \langle \beta_{n-1} \rangle) \times C_\infty$, where $\langle \beta_{n-1} \rangle \leq \text{Aut}(H_{n-1})$ is the cyclic subgroup of order 2 generated by β_{n-1} ;*
- (ii) *β_n is an exact automorphism of H_n .*

Proof. When $n = 1$ the claims follow from Lemma 2.11. For $n \geq 2$, (i) is an easy consequence of Lemma 2.4(c) and the definition of ϕ_{n-1} , and claim (ii) follows from Lemma 2.10. \square

Theorem 2.14. *For every integer $n \geq 1$ there exists a polycyclic group G_n of torsion length n .*

Proof. For each $n \geq 0$ we define the group G_{n+1} as follows:

$$(4) \quad G_{n+1} = \langle H_n, x \mid x^2 = 1, xgx^{-1} = \beta_n(g), \text{ for all } g \in H_n \rangle \cong H_n \rtimes \langle \beta_n \rangle,$$

where H_n and $\beta_n \in \text{Aut}(H_n)$ are provided by Construction 2.12. Thus H_n is a subgroup of G_{n+1} of index 2. Note that H_n is a polycyclic group of Hirsch length $4n$, by definition, hence so is G_{n+1} .

Let us show that the torsion length of G_{n+1} is $n + 1$ by induction on n . When $n = 0$, $G_1 \cong C_2$ has torsion length 1, thus we can further assume that $n \geq 1$.

Recall that H_n is torsion-free and $\beta_n \in \text{Aut}(H_n)$ is an exact automorphism by Proposition 2.13. Therefore, according to Lemma 2.7, every finite order element of G_{n+1} must be conjugate to a power of x . Thus $\text{Tor}(G_{n+1})$ is the normal closure of x in G_{n+1} . The definition of G_{n+1} , given in (4), implies that the quotient $G_{n+1}/\text{Tor}(G_{n+1})$ is naturally isomorphic to the quotient of H_n by $\mathcal{N}(\beta_n)$. So, Proposition 2.13.(i) yields that

$$G_{n+1}/\text{Tor}(G_{n+1}) \cong (H_{n-1} \rtimes \langle \beta_{n-1} \rangle) \times C_\infty \cong G_n \times C_\infty.$$

Evidently, the torsion length of $G_n \times C_\infty$ is the same as the torsion length of G_n , which equals n , by induction. Therefore the torsion length of G_{n+1} is $n + 1$, as claimed. \square

Remark 2.15. It is not difficult to show that the groups H_n and G_n , constructed above, are, in fact, virtually abelian. A more careful analysis reveals that for every $n \geq 1$ the group G_n contains a free abelian normal subgroup F_n such that G_n/F_n is isomorphic to the direct power $(C_2 \wr C_2)^{n-1}$. It follows that, for each $n \in \mathbb{N}$, G_n is soluble of derived length 3, hence the (restricted) direct product $\times_{n=1}^\infty G_n$ is a countable soluble group of derived length 3 with infinite torsion length.

Remark 2.16. It is easy to see that the torsion length of a polycyclic group G is at most $h(G) + 1$, where $h(G)$ denotes the Hirsch length of G . In particular, the torsion length of any polycyclic group is finite.

In [3, Subsection 3.3] Chiodo and Vyas asked whether there exists a finitely generated soluble group with infinite torsion length. This interesting question remains open. As Remark 2.16 shows, such a group cannot be polycyclic.

3. FINITELY PRESENTED GROUPS G WITH $G/\text{Tor}(G)$ NOT FINITELY PRESENTED

In [4, Question 1] and [6, 19.24] it was asked whether there is a finitely presented group G so that $G/\text{Tor}(G)$ is recursively presented but not finitely presented. We give two different constructions of such a group, one of which is soluble and the other of which is virtually torsion-free.

Construction 3.1. First we construct a soluble group G . In this case one starts with Abels' group H which is a finitely presented torsion-free soluble group of 4×4 upper-triangular matrices over $\mathbb{Z}[1/p]$ for a fixed prime p , such that the centre $Z = Z(H)$ is isomorphic to the additive group $\mathbb{Z}[1/p]$ [1]. Let C be an infinite cyclic subgroup with $C \leq Z$, and let $G = H/C$. The matrix representation of H gives rise to a sequence $Z = H_1 < H_2 < H_3 < H_4 = H$ of normal subgroups of H with each H_{i+1}/H_i torsion-free abelian, and it follows that H/Z is torsion-free. Hence the subgroup $\text{Tor}(G)$ is equal to $Z/C \cong \mathbb{Z}[1/p]/\mathbb{Z}$. This subgroup of H/C is central and not finitely generated. It follows that $G/\text{Tor}(G) \cong H/Z$ is a non-finitely presented soluble group. Since Z is clearly recursively generated, H/Z is recursively presented.

The virtually torsion-free examples rely on Bestvina-Brady groups [2], so we start by recalling these groups. Let L be a finite flag simplicial complex, i.e., a clique complex. The right-angled Artin group A_L is the group with generators the vertices of L subject only to the relations that the ends of each edge of L commute. There is a homomorphism $f_L : A_L \rightarrow \mathbb{Z}$ that sends each generator to $1 \in \mathbb{Z}$, and the kernel of f is the Bestvina-Brady group BB_L . Bestvina-Brady show that BB_L is finitely generated if and only if L is connected, and finitely presented if and only if L is simply-connected [2].

Construction 3.2. Let L be a finite connected flag complex with non-trivial finite fundamental group π , and let K be the universal cover of L . The action of π on K by deck transformations induces an action on A_K and on BB_K . The group G will be the semidirect product $G = BB_K \rtimes \pi$. Let $\overline{G} = A_K \rtimes \pi$, and note that G is a normal subgroup of \overline{G} with $\overline{G}/G \cong \mathbb{Z}$. It follows that every torsion element of \overline{G} is in G . It is easy to see that $\overline{G}/\text{Tor}(\overline{G}) = A_L$, and hence $G/\text{Tor}(G) = BB_L$, a non-finitely presented subgroup of the group A_L . Thus the group G contains infinitely many conjugacy classes of subgroups isomorphic to π (see also [7, Thm. 3] for another proof of this fact). An explicit presentation for $G/\text{Tor}(G) = BB_L$ with generators the directed edges of L is given in [5].

REFERENCES

- [1] H. Abels, *An example of a finitely presented solvable group*. In *Homological group theory (Proc. Sympos. Durham, 1977)*, London Math. Soc. Lecture Notes **36**, 205–211, Cambridge Univ. Press (Cambridge) (1979).
- [2] M. Bestvina and N. Brady, *Morse theory and finiteness properties of groups*. *Invent. Math.* **129** (1997) 445–470.
- [3] M. Chiodo and R. Vyas, *A note on torsion length*. *Comm. Alg.* **43** (2015) 4825–4835.
- [4] M. Chiodo and R. Vyas, *Torsion, torsion length and finitely presented groups*. *J. Group Theory* **21** (2018) 949–971.
- [5] W. Dicks and I. J. Leary, *Presentations for subgroups of Artin groups*. *Proc. Amer. Math. Soc.* **127** (1999) 343–348.
- [6] E.I. Khukhro and V.D. Mazurov, editors, *Unsolved problems in group theory, the Kourovka notebook*. 19th edition (2018). Sobolev Institute of Mathematics, Russian Academy of Sciences, Siberian Branch, Novosibirsk.
- [7] I. J. Leary and B. E. A. Nucinkis, *Some groups of type VF*. *Invent. Math.* **151** (2003), 135–165.
- [8] J.-P. Serre, *Galois cohomology*. Translated from the French by Patrick Ion and revised by the author. *Springer-Verlag, Berlin*, 1997. x+210 pp.

CGTA, SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SOUTHAMPTON, HIGHFIELD, SOUTHAMPTON, SO17 1BJ, UNITED KINGDOM.

Email address: I.J.Leary@soton.ac.uk, aminasyan@gmail.com