


 Cite this: *CrystEngComm*, 2022, 24, 6137

 Received 26th July 2022,
Accepted 11th August 2022

DOI: 10.1039/d2ce01029a

rsc.li/crystengcomm

Self-assembly of $[\text{Sn}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2]_6$ metallocyclic $\text{Sn}(\text{II})$ hexamer stacks with CF_3 -lined channel interiors†

 Kelsey R. Cairns, Rhys P. King,  William Levason  and Gillian Reid *

$[\text{Sn}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2]_6$ (**1**) self-assembles to form a highly unusual hexagonal metallocyclic array with bridging CF_3SO_3^- groups. The discrete hexamers stack along the *c*-axis to form hydrophobic CF_3 -lined channels with an F...F distance across the channel diameter of 0.8 nm. This $\text{Sn}(\text{II})$ complex is different from its $\text{Pb}(\text{II})$ analogue, which instead forms a discrete, doubly CF_3SO_3^- -bridged dimer of formula, $[\{\text{Pb}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2\}(\mu\text{-CF}_3\text{SO}_3)_2]$ (**3**).

Main group chemistry has attracted considerable research effort in the last two decades in response to the desire for tailored precursors for p-block materials deposition, metal-free catalysts, and porous coordination polymers, in addition to exploring new classes of compounds and reaction chemistry.¹

While many of the reported p-block complexes are derived from main group halide precursors with secondary bonding interactions to neutral group 15 and 16 donor ligands, replacing the halide with more weakly coordinating (less Lewis basic) anions, such as CF_3SO_3^- , and the bulky fluorine tetra-arylborates and fluorine aluminates, *etc.*, can provide an entry to highly Lewis acidic p-block cations that are of interest in organic/organometallic chemistry,^{2–5} while delocalised fluorine anions including CF_3SO_3^- , SiF_6^{2-} , PF_6^- , *etc.*, can be used to lower the dimensionality of porous coordination polymers and metal–organic frameworks (MOFs).^{6–9} Weakly coordinating anions, in which the negative charge is delocalised and diffuse, including triflate, fluorine tetra-arylborates and

fluorine aluminates, amongst others, have attracted considerable research effort in the field of organometallic chemistry.^{2–5} They aid solubility in low polarity solvents and can facilitate stabilisation of highly reactive species such as the univalent $[\text{Ga}(\text{PPh}_3)_3]^+$,¹⁰ and $[\text{Ag}(\text{P}_4)_2]^+$ cations,¹¹ as well as unusual homoleptic phosphine¹² or thioether complexes with alkali metal cations.¹³

The CF_3SO_3^- group has been used extensively as a weakly coordinating anion in work developing group 14 chemistry and cations thereof. Notable examples include the stabilisation of ‘naked’ $\text{Ge}(\text{II})$ dications encapsulated within 2.2.2-cryptand,¹⁴ $\text{Ge}(\text{II})$ dications bearing N-heterocyclic carbenes,¹⁵ and pnictines,¹⁶ as well as tetravalent group 14 fluoride species with soft pnictine ligands.¹⁷ We have also observed that the CF_3SO_3^- anion can engage in a range of different secondary interactions that play an important role in determining the speciation and structures adopted. For example, in $[\text{In}(\text{OTf})_2(\text{Ph}_3\text{PO})_4]_4$ – $[\text{In}\{(\text{OH})_2(\text{OTf})_4\}(\text{Ph}_3\text{PO})_2]_4$, which forms a weakly associated 24-membered pseudo-macrocyclic ring *via* H-bonding between the coordinated water molecules and the CF_3SO_3^- anions.¹⁸ In view of these observations, we became interested to investigate the coordination chemistry of the divalent group 14 ions (Ge, Sn, Pb) with the hard and sterically compact OPMe_3 ligand.

Reaction of $\text{Sn}(\text{CF}_3\text{SO}_3)_2$ with three mol equiv. of OPMe_3 in a weakly coordinating solvent such as CH_2Cl_2 or MeCN readily leads to the formation of a colourless crystalline solid, $[\text{Sn}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2]_6$, in high yield. Spectroscopic data for this species are presented in Fig. S3–S5 (ESI†) and discussed below. However, to probe the role of the CF_3SO_3^- , single crystals suitable for X-ray analysis were grown either by diffusion of hexane into a CH_2Cl_2 solution or by vapour diffusion of Et_2O into an MeCN or MeNO_2 solution with several crystals grown from different batches and solvents all producing the same crystalline form.

The material crystallises in the trigonal space group ($P\bar{3}$) and the structure determination shows that it self-assembles into discrete hexamers, $[\text{Sn}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2]_6$ (**1**) (Fig. 1(a))

School of Chemistry, University of Southampton, Southampton SO17 1BJ, UK.
E-mail: G.Reid@soton.ac.uk

† Electronic supplementary information (ESI) available: Full experimental, spectroscopic and microanalytical data for the new complexes, together with the X-ray crystallographic parameters and simulated and experimental powder X-ray patterns for the $\text{Sn}(\text{II})$ hexamer. Experimental methods and characterisation data are provided in the ESI, along with the crystallographic parameters (Table S1), alternative views of the structure of $[\text{Sn}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2]_6$ (Fig. S1), simulated and experimental PXRD patterns for $[\text{Sn}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2]_6$ (Fig. S2). Cif files for the crystal structures are available from the Cambridge Crystallographic Data Centre with CCDC numbers 2169639 ($[\text{Sn}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2]_6$, **1**) and 2169640 ($[\{\text{Pb}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2\}(\mu\text{-CF}_3\text{SO}_3)_2]$, **3**). For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d2ce01029a>






Fig. 1 (a) View of the hexameric assembly adopted by $[\text{Sn}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2]_6$ (**1**) via the bridging CF_3SO_3^- groups; (b) the coordination environment at $\text{Sn}(\text{II})$ with the atom labelling scheme. Ellipsoids are drawn at the 50% probability level and H atoms are omitted for clarity. Selected bond lengths (\AA): $\text{Sn1}-\text{O1} = 2.1557(11)$, $\text{Sn1}-\text{O2} = 2.1670(10)$, $\text{Sn1}-\text{O3} = 2.0943(11)$, $\text{Sn1}\cdots\text{O5} = 2.8578(12)$ (bridging), $\text{Sn1}\cdots\text{O6} = 3.0277(13)$ (bridging), $\text{Sn1}\cdots\text{O7} = 3.0822(12)$ (terminal); selected bond angles ($^\circ$): $\text{O1}-\text{Sn1}-\text{O2} = 82.15(4)$, $\text{O1}-\text{Sn1}-\text{O3} = 85.97(4)$, $\text{O2}-\text{Sn1}-\text{O3} = 87.49(4)$; (c) space-filling diagram of a discrete hexamer – top view; (d) space-filling diagram of a discrete hexamer – edge view.

comprising six pseudo-octahedrally coordinated $\text{Sn}(\text{II})$ centres with three mutually *fac* OPMe_3 ligands ($d(\text{Sn}-\text{OPMe}_3) = 2.094(1), 2.156(2), 2.167(1) \text{ \AA}$) and three weakly bound, CF_3SO_3^- groups (Fig. 1(b)). Two of the triflates bridge between adjacent Sn centres to form the cyclic arrangement ($d(\text{Sn}\cdots\text{OO}_2\text{SCF}_3) = 2.858(1), 3.028(2) \text{ \AA}$), while the third is bonded terminally ($d(\text{Sn}\cdots\text{OO}_2\text{SCF}_3) = 3.082(1) \text{ \AA}$).

Notably, the CF_3 groups from the six bridging CF_3SO_3^- anions all point towards the central void of the metallocyclic hexamer, with the $\text{F}\cdots\text{F}$ distance across the void diameter being $8.019(2) \text{ \AA}$ (ESI† Fig. S1). These hexamers stack along the crystallographic *c*-axis, resulting in a structure with hydrophobic CF_3 -lined nanochannels. There are no significant interactions between the hexamers. The structure determination showed the void to contain ~ 1 disordered hexane molecule per hexamer unit. The crystal has a void volume of 6.1% determined by PLATON.¹⁹ Brunauer–Emmett–Teller (BET) surface area analysis performed on compound (**1**) after heating the powdered samples *in vacuo* at 80°C overnight revealed no appreciable N_2 sorption.

Although the cyclic hexamer is very unusual, it is readily reproduced from different product batches and *via* crystallisation from different solvent systems. There is no evidence for other polymorphs and powder X-ray diffraction from the bulk product gives a good phase match to the pattern simulated from the single crystal structure (ESI† Fig. S2).

The corresponding $\text{Ge}(\text{II})$ and $\text{Pb}(\text{II})$ complexes, $\text{M}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2$, were also prepared; the former from GeCl_2 (dioxane), $\text{Me}_3\text{SiO}_3\text{SCF}_3$ and OPMe_3 in a 1:2:3 molar ratio, and the latter directly from $\text{Pb}(\text{CF}_3\text{SO}_3)_2$ and 3 mol equiv. of OPMe_3 . The $\text{Ge}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2$ (**2**) is a low-melting, waxy solid, therefore it was not possible to obtain diffraction data for this compound. Its spectroscopic and microanalytical data are consistent with the formulation shown (Fig. S4†), although it is not possible to confirm from the spectra whether the CF_3SO_3^- groups are coordinated. In the case of $\text{Pb}(\text{II})$, a white powdered solid was isolated and assigned as $[\{\text{Pb}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2\}_2(\mu\text{-CF}_3\text{SO}_3)_2]$ (**3**) based on its X-ray crystal structure (below) and elemental analysis, with solution NMR spectroscopic data also consistent with this (Fig. S5†). Despite the stoichiometry of the product being the same as for the $\text{Sn}(\text{II})$ analogue, the X-ray crystal determination reveals that it adopts a very different structure from the $\text{Sn}(\text{II})$ hexamer.

In this case the $\text{Pb}(\text{II})$ complex is a discrete dimer with two CF_3SO_3^- bridges (Fig. 2(a)), and with distorted octahedral coordination at each Pb centre *via* three OPMe_3 ligands ($d(\text{Pb}-\text{OPMe}_3) = 2.3218(17), 2.3295(17), 2.3628(18) \text{ \AA}$), and three longer O-bound CF_3SO_3^- groups (two μ^2 -bridging and one terminal) located around 2.8–3.0 \AA from the Pb atoms. Fig. 2(b) shows the packing arrangement in the *bc*-plane, but illustrating that the triflate CF_3 groups decorate the upper and lower surfaces at the interfaces between the layers.

Solution ^1H , $^{13}\text{C}\{^1\text{H}\}$ and $^{31}\text{P}\{^1\text{H}\}$ NMR spectra (Fig. S3–S5†) for the complexes dissolved in CD_2Cl_2 (for **1** and **2**) or CD_3CN (for **3**) each show only one OPMe_3 environment, with the $^{31}\text{P}\{^1\text{H}\}$ chemical shifts showing a marginal low frequency






Fig. 2 (a) View of the structure of the $[\text{Pb}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2](\mu\text{-CF}_3\text{SO}_3)_2$ dimer (**3**) with the atom labelling scheme. Selected bond lengths (Å): Pb1–O1 = 2.3628(18), Pb1–O2 = 2.3295(17), Pb1–O3 = 2.3218(17), Pb1–O4 = 2.7325(19), Pb1...O6 = 2.889(2), Pb1...O7 = 2.888(2). Selected bond angles ($^\circ$): O1–Pb1–O2 = 97.35(6), O1–Pb1–O3 = 76.97(6), O2–Pb1–O3 = 80.09(6); (b) view of the *bc* plane showing the CF_3 groups aligned on the top and bottom surfaces.

shift upon going down group 14 from (2; Ge) \rightarrow (1; Sn) \rightarrow (3; Pb). The $^{19}\text{F}\{^1\text{H}\}$ NMR shifts are sharp singlets at *ca.* -79 ppm in all three cases, consistent with ionic CF_3SO_3^- ,¹⁶ suggesting triflate dissociation in solution. The ^{119}Sn NMR resonance for (1) occurs at -793 ppm, similar to the chemical shifts reported for Sn(II) phosphines and phosphine chalcogenides with SbF_6^- anions.²⁰

The formation of the $[\text{Sn}(\text{OPMe}_3)_3(\text{CF}_3\text{SO}_3)_2]_6$ metallocycle identified in this work was very unexpected and the weakly coordinating CF_3SO_3^- linkers clearly play an important role in its assembly. Very recently Itoh and co-workers described a series of fluorinated oligoamide nanorings with diameters ranging from 0.9–1.9 nm, which undergo supramolecular polymerisation in bilayer membranes forming nanochannels with a similar dense F-atom coverage of the interior walls as observed in our hexamer assemblies. Notably, Itoh *et al.* showed that the 0.9 nm channels exhibit a water permeation flux two orders of magnitude higher than those of aquaporins and carbon nanotubes.²¹ It should be noted that the covalent oligoamide nanorings are rather more robust compared to the weakly coordinated hexameric assemblies reported in our work. Nevertheless, these results indicate that the incorporation of CF_3 groups can promote highly hydrophobic molecu-

lar and supramolecular nanochannels with fluorine-rich interior walls. There is also a growing interest in the development of both fluorinated porous materials, as well as Sn-based MOFs for a range of applications.^{4–6,22–24}

Our current work aims to understand the different behaviours observed for (1)–(3) in this work and to explore how modifying the pnictogen and/or the organic substituents influences the speciation with divalent group 14 triflates, and whether altering the denticity and architecture of the pnictine oxide increases the scope for developing other novel assemblies.

We thank the EPSRC for support through the ADEPT Programme Grant (EP/N035437/1) and EP/R513325/1. We also thank Sidrah Hussain for assistance recording the PXRD data and Alice Oakley for the BET measurements.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- For examples see: Coordination Chemistry of the s, p, and f metals, *Comprehensive Coordination Chemistry II*, ed. J. A. McCleverty and T. J. Meyer, Elsevier, 2003, vol. 3; S. Aldridge and C. Jones, Modern Main Group Chemistry (special issue), *Chem. Soc. Rev.*, 2016, **45**, 763; J. Parr, *Germanium, tin and lead*, *Comprehensive Coordination Chemistry II*, ed. J. A. McCleverty and T. J. Meyer, Elsevier, 2003, vol. 3, p. 545, and references therein.
- G. A. Lawrance, *Chem. Rev.*, 1986, **86**, 17.
- S. H. Strauss, *Chem. Rev.*, 1993, **93**, 927.
- I. Krossing, *Angew. Chem., Int. Ed.*, 2004, **43**, 2066.
- T. A. Engesser, M. R. Lichtenhaler, M. Schleep and I. Krossing, *Chem. Soc. Rev.*, 2016, **45**, 789.
- S. Noro and T. Nakamura, *NPG Asia Mater.*, 2017, **9**, e433.
- A. Kondo, H. Noguchi, L. Carlucci, D. M. Proserpio, G. Ciani, H. Kajiro, T. Ohba, H. Kanoh and K. Kaneko, *J. Am. Chem. Soc.*, 2007, **129**, 12362.
- S. Noro, Y. Hijikata, M. Inukai, T. Fukushima, S. Horike, M. Higuchi, S. Kitagawa, T. Akutagawa and T. Nakamura, *Inorg. Chem.*, 2013, **52**, 280.
- A. Kondo, A. Chinen, H. Kajiro, T. Nakagawa, K. Kato, M. Takata, Y. Hattori, F. Okino, T. Ohba, K. Kaneko and H. Kanoh, *Chem. – Eur. J.*, 2009, **15**, 7549.
- J. M. Slattery, A. Higelin, T. Bayer and I. Krossing, *Angew. Chem., Int. Ed.*, 2010, **49**, 3228.
- I. Krossing, *J. Am. Chem. Soc.*, 2001, **123**, 4603.
- M. Carravetta, M. Concistre, W. Levason, G. Reid and W. Zhang, *Chem. Commun.*, 2015, **51**, 9555.
- M. J. D. Champion, J. M. Dyke, W. Levason, M. E. Light, D. Pugh, H. Bhakhoa, L. Rhyman, P. Ramasami and G. Reid, *Inorg. Chem.*, 2015, **54**, 2497.
- P. A. Rugar, V. N. Staroverov and K. M. Baines, *Science*, 2008, **322**, 1360.
- P. A. Rugar, V. N. Staroverov, P. J. Ragogna and K. M. Baines, *J. Am. Chem. Soc.*, 2007, **129**, 15138.



- 16 R. P. King, V. K. Greenacre, W. Levason, J. M. Dyke and G. Reid, *Inorg. Chem.*, 2021, **60**, 12100.
- 17 R. P. King, M. S. Woodward, G. McRobbie, J. Grigg, W. Levason and G. Reid, *Dalton Trans.*, 2021, **50**, 14400; R. P. King, W. Levason and G. Reid, *Dalton Trans.*, 2021, **50**, 17751.
- 18 K. C. Cairns, W. Levason, G. Reid and W. Zhang, *Polyhedron*, 2021, **207**, 115367; K. C. Cairns, W. Levason, G. Reid and W. Zhang, *Polyhedron*, 2021, **210**, 115529.
- 19 A. L. Spek, *Acta Crystallogr., Sect. D: Biol. Crystallogr.*, 2009, **65**, 148.
- 20 P. A. W. Dean, *Can. J. Chem.*, 1983, **61**, 1795; P. A. W. Dean, D. D. Phillips and L. Polensek, *Can. J. Chem.*, 1981, **59**, 50; P. A. W. Dean, *Can. J. Chem.*, 1981, **59**, 50.
- 21 Y. Itoh, S. Chem, R. Hirahara, T. Konda, T. Aoki, T. Ueda, I. Shimada, J. J. Cannon, C. Shao, J. Shiomi, K. V. Tabata, H. Noji, K. Sato and T. Aida, *Science*, 2022, **376**, 738–743.
- 22 B. Y. Mingbo, S. Renliang, H. Wei, Q. Rongxin and S. He, *Colloids Surf., A*, 2021, **616**, 126322; P. M. Bhatt, Y. Belmabkhout, A. Cadiou, K. Adil, O. Shekhah, A. Shkurenko, L. J. Barbour and M. Eddaoudi, *J. Am. Chem. Soc.*, 2016, **138**, 9301; M. R. Tchalala, P. M. Bhatt, K. N. Chappanda, S. R. Tavares, K. Adil, Y. Belmabkhout, A. Shkurenko, A. Cadiou, N. Heymans, G. De Weireld, G. Maurin, K. N. Salama and M. Eddaoudi, *Nat. Commun.*, 2019, **10**, 1328.
- 23 Y. Kamakura, S. Fujisawa, K. Takahashi, H. Toshima, Y. Nakatani, H. Yoshikawa, A. Saeki, K. Ogasawara and D. Tanaka, *Inorg. Chem.*, 2021, **60**, 12691.
- 24 K. Liu, C. Li, L. Yan, M. Fan, Y. Wu, X. Meng and T. Ma, *J. Mater. Chem. A*, 2021, **9**, 27234; X. Zhou, Y. Yu, J. Yang, H. Wang, M. Jia and J. Tang, *ChemElectroChem*, 2019, **6**, 205; X. Jiang, M. Shao, K. Li, L. Ding and M. Zeng, *J. Electroanal. Chem.*, 2022, **912**, 116268.

