

Completing RHINO

Pasquale Di Bari and Adam Murphy

*Physics and Astronomy, University of Southampton,
Southampton, SO17 1BJ, U.K.*

October 21, 2022

Abstract

The right-handed (RH) Higgs-induced neutrino mixing (RHINO) model explains neutrino masses and origin of matter in the universe within a unified picture. The mixing, effectively described by a dimension five operator, is responsible both for the production of dark neutrinos, converting a small fraction of seesaw neutrinos acting as source, and for their decays. We show that including the production of source neutrinos from Higgs portal interactions, their abundance can thermalise prior to the onset of source-dark neutrino oscillations, resulting into an enhanced production of dark neutrinos that thus can play the role of decaying dark matter (DM) for a much higher seesaw scale. This can be above the sphaleron freeze-out temperature and as high as ~ 100 TeV, so that strong thermal resonant leptogenesis for the generation of the matter-antimatter asymmetry is viable. We obtain a ~ 1 TeV–1 PeV allowed dark neutrino mass range. Intriguingly, their decays can also explain a neutrino flux excess at $\mathcal{O}(100$ TeV) energies recently confirmed by the IceCube collaboration analysing 7.5yr HESE data. Our results also point to an effective scale for Higgs portal interactions nicely identifiable with the grandunified scale and many orders of magnitude below the effective scale for the mixing. We explain this hierarchy in a UV-complete model with a very heavy fermion as mediator: the first scale corresponds to the fundamental scale of new physics, while the second is much higher because of a very small coupling identifiable with a symmetry breaking parameter. Therefore, RHINO realises a simple unified model of neutrino masses and origin of matter in the universe currently under scrutiny at neutrino telescopes and potentially embeddable within a grandunified model.

1 Introduction

The origin of matter in the universe is a key issue in modern physics [1]. Its solution would guide us along that path to new physics whose existence is motivated both to a theoretical and a phenomenological level. A strong phenomenological motivation is represented by the necessity to extend the standard model (SM) in order to incorporate neutrino masses and mixing. It is then reasonable to investigate whether extensions of the SM able to explain neutrino masses and mixing can also provide a solution to the problem of the origin of matter in the universe. The minimal type-I seesaw mechanism [2], augmenting the SM particle content with the introduction of RH neutrinos, provides an elegant and minimal way to understand both the lightness of the ordinary neutrinos and the observed large mixing angles in the leptonic mixing matrix. Moreover, it can be easily embedded within many realistic models beyond the SM [3] and it leads to leptogenesis as a built-in scenario of baryogenesis for the explanation of the matter-antimatter asymmetry of the universe [4]. It is also well known that a keV seesaw RH neutrino can be efficiently produced via active-sterile neutrino mixing and play the role of (warm) DM [5]. A unified picture, combining a keV lightest seesaw neutrino DM produced from active-sterile neutrino mixing with successful leptogenesis from sterile-sterile neutrino mixing [6] and satisfying low energy neutrino data is also potentially viable [7]. However, constraints from X-ray observations and large scale structure N-body simulations require the presence of a large lepton asymmetry for an efficient resonant production [8], introducing an additional issue. In principle, such a large lepton asymmetry can be produced by the decays of one of the two heaviest seesaw neutrinos, without additional non-minimal ingredients beyond the type-I seesaw Lagrangian [9], but this requires a very strong degeneracy of the two heavier RH neutrino masses. Interestingly, a 7 keV lightest seesaw neutrino resonantly produced and playing the role of DM can also address the 3.5 keV X-ray anomaly [10]. Recently, numerical solutions fulfilling all requirements and the 3.5 keV anomaly have been shown to exist [11]. However, the relative degeneracy of the two heavier RH neutrino masses has to be very fine-tuned, at the level of 10^{-16} . In any case the XRISM satellite [12] should soon either confirm the 3.5 keV anomaly or place stringent constraints on warm DM decays, ruling out this minimal scenario for the origin of matter in the universe where new physics resides just in the type-I seesaw Lagrangian.

An alternative approach, where a RH neutrino plays the role of cold rather than warm DM, is based on adding a non renormalizable operator to the type-I seesaw Lagrangian [13]. In this case a dark neutrino, with suppressed Yukawa couplings, would be produced by its Higgs-induced mixing with a seesaw neutrino via the dimension five Anisimov

operator [13, 14]

$$\mathcal{L}_{\text{DS}}^{\Lambda} = \frac{\lambda_{\text{DS}}}{\Lambda} \Phi^{\dagger} \Phi \overline{N_{\text{D}}^c} N_{\text{S}}, \quad (1)$$

where Λ is the scale of new physics. This mixing would be both responsible for the production of the dark neutrinos and for their decays [13, 15]. Decays would be indeed induced at zero temperature by the effective small Yukawa couplings generated by the mixing [16]. In this way there is a tension between the two conditions to be imposed for the dark neutrino to be a DM candidate: reproducing the observed DM abundance and being sufficiently long-lived to evade constraints from neutrino telescopes [17]. This tension produces constraints on the neutrino mixing effective scale $\tilde{\Lambda}_{\text{DS}} \equiv \Lambda/\lambda_{\text{DS}}$ and on the mass of the dark neutrino [16], making the scenario highly predictive.

The production of dark neutrinos proceeds through their Higgs-induced mixing with the source neutrinos. The origin of the mixing is a mismatch between the Majorana mass eigenbasis and the basis where the new Higgs-induced interactions are diagonal. It has a strong temperature dependence encoded in the thermal self-energy that, in the case of neutrino mixing, can be described in terms of the effective potential or, alternatively, in terms of thermal masses. This strong temperature dependence is the key for a solution of the tension between an efficient production and a sufficiently long-lived dark neutrino, since the mixing can be much stronger in the very early universe than today, when dark neutrinos decay.

Initially, calculations of the dark neutrino abundance were done using a Landau-Zener approximation [13, 15]. However, it has been shown that, in general, this fails to provide a correct description and numerical calculations within a density matrix formalism show that the production is, in general, much less efficient [16].¹ If one starts from a vanishing source neutrino abundance, a stringent upper bound is obtained on the mass of the source neutrino mass that needs to be lighter than the W gauge boson mass. In this way the four-body decay rate gets suppressed and the dark neutrino can be the DM particle only for masses $M_{\text{D}} \gtrsim 10^7 \text{ GeV}$. However, in such a setup, strong thermal leptogenesis from (non-relativistic) decays is not viable, since the matter-antimatter asymmetry should be resonantly produced from the decays of the seesaw neutrinos with masses about twice the sphaleron freeze-out temperature $T_{\text{RH}} \sim 132 \text{ GeV}$ [18]. The asymmetry could be still generated from the mixing of the two seesaw neutrinos with GeV masses [6, 7] or also from decays [19, 20]. Therefore, compared to the νMSM , one would still have low scale leptogenesis with GeV seesaw neutrinos but the DM would be very heavy and, therefore,

¹The validity of the Landau-Zener approximation requires extremely quasi-degenerate dark-source neutrino masses.

cold rather than warm.

However, in low scale leptogenesis with GeV neutrinos the final asymmetry is produced in the relativistic regime and it would be sensitive not only to an accurate description of many different processes but also to the initial conditions. Since the reheat temperature needs to be higher than $\sim 10^9$ GeV for an efficient dark neutrino production, in this case a large pre-existing asymmetry is a rather natural outcome in many different mechanisms. Therefore, having to impose that this is negligible is certainly an unattractive feature of the scenario. In this paper we show that a rather simple and elegant solution to this problem exists without spoiling minimality.

The main point is that the Anisimov operator is actually only one term of a more general effective Lagrangian that can be written as [13] ($I, J = D, S$)

$$\mathcal{L}_A = \sum_{I,J} \frac{\lambda'_{IJ}}{\Lambda} \Phi^\dagger \Phi \overline{N_I^c} N_J, \quad (2)$$

or, explicitly, as the sum of three terms

$$\mathcal{L}_A = \frac{\lambda_{DS}}{\Lambda} \Phi^\dagger \Phi \overline{N_D^c} N_S + \frac{\lambda_{SS}}{\Lambda} \Phi^\dagger \Phi \overline{N_S^c} N_S + \frac{\lambda_{DD}}{\Lambda} \Phi^\dagger \Phi \overline{N_D^c} N_D, \quad (3)$$

having properly redefined $\lambda_{DS} = \lambda'_{DS} + \lambda'_{SD}$.² In this way also (non-renormalisable) Higgs portal interactions terms are included.³ Higgs portal interactions were assumed not to play a role in the production of dark neutrinos and they were neglected in previous papers [13, 15, 16]. Here we show that actually these interactions can influence the production of dark neutrinos in a very significant way and help not only to reconcile a solution of the DM conundrum with strong thermal (i.e., independent of the initial conditions) leptogenesis but also to allow dark neutrino to be a decaying DM for masses $M_D \sim 100$ TeV, currently hinted by measurements of high energy neutrino flux at neutrino telescopes. Notice that in Eq. (3) there is also a term describing Higgs portal interactions for dark neutrinos that might compete with the mixing term in producing the dark neutrinos. The same mixing term can also be responsible for a mixed production of dark neutrinos and source neutrinos from Higgs scatterings ($\phi + \phi^\dagger \rightarrow N_D + N_S$). However, in this paper we assume that the production of dark neutrinos from Higgs scatterings can be neglected. In the end of the paper we will discuss a model where this assumption is naturally justified. Notice, however, that direct dark neutrino production from Higgs scatterings might play a role

²This is because one simply has $\overline{N_S^c} N_D = \overline{N_D^c} N_S$.

³Renormalisable Higgs portal interactions to scalars were first considered in [21], while non-renormalisable 5-dim Higgs portal interactions to RH neutrinos in [13].

in other contexts, for example in the thermalisation of a dark sector prior to a phase transition, as in the scenario discussed in [22].

In this paper we show that including Higgs portal interactions for the source neutrinos, the allowed region in the space of parameters enlarges at higher seesaw scales, where traditional leptogenesis from decays in the non-relativistic regime can work independently of the initial conditions (strong thermal leptogenesis). This is because they can partly or fully thermalise the source neutrinos prior to the mixing, i.e., at higher temperatures than Yukawa interactions. This makes possible to reproduce the observed DM abundance for higher values of the effective RHINO scale $\tilde{\Lambda}_{\text{DS}} \equiv \Lambda/\lambda_{\text{DS}}$ and, consequently, to satisfy the constraint from four body decays also for values of the seesaw scale above the W gauge boson mass. We show that in this way the seesaw scale can be as high as ~ 100 TeV. This is still well below the leptogenesis lower bound $M_I \gtrsim 10^9$ GeV holding for hierarchical seesaw neutrinos [23] so that a resonant production [24, 25] and/or some combined tuning in the seesaw formula [26, 27, 28] are necessary to enhance the (total and/or flavoured) CP asymmetries. Moreover, such higher seesaw scales are attractive also because they can be more easily embedded within realistic models of flavour [3]. Moreover, for such higher seesaw scale, the allowed dark neutrino mass is in the range ~ 1 TeV–1 PeV, thus making possible to address the hint of a ~ 100 TeV excess in high energy neutrino data.

The paper is structured as follows. In Section 2 we briefly review the RHINO model and how the dark neutrino abundance can be calculated within a density matrix formalism. In Section 3 we include Higgs portal interactions for the source neutrinos showing how the set of kinetic equations gets modified and discuss some benchmark cases clearly illustrating their effect. Here we also show how the critical value of the effective scale for thermalisation of source neutrinos is nicely obtained to be coinciding with the grandunified scale. In Section 4 we derive the allowed regions in the dark neutrino lifetime-mass plane, for various values of the seesaw scale, reheat temperature and effective scale for the source neutrino Higgs portal interactions. In Section 5 we discuss a UV-complete model that can consistently incorporate all interactions described by the Anisimov operator. This is able to explain the values of the three effective scales as stemming from just one fundamental scale that can be nicely identified with the grandunified scale, something that should be regarded as a successful outcome of the model. Finally, in Section 6 we draw our conclusions.

2 The RHINO model

Let us now briefly review the RHINO model and how the dark neutrino abundance can be calculated within a density matrix formalism just using the Anisimov operator in Eq. (1). The SM field content is augmented by the introduction of three RH neutrinos. However, in addition to the usual seesaw Lagrangian with neutrino Yukawa couplings and Majorana mass terms, one also has a RHINO term ($I = J = 1, 2, 3; \alpha = e, \mu, \tau$)

$$-\mathcal{L}_{Y+M+\Lambda}^{\nu} = \overline{L}_{\alpha} h_{\alpha I} N_I \tilde{\Phi} + \frac{1}{2} \overline{N}_I^c M_I N_I + \sum_{I \neq J} \frac{\lambda_{IJ}}{\Lambda} \Phi^{\dagger} \Phi \overline{N}_I^c N_J + \text{h.c.}, \quad (4)$$

where the L_{α} 's are the three lepton doublets, Φ is the Higgs doublet and $\tilde{\Phi} \equiv i\sigma_2 \Phi^*$ is its dual. Notice here we are not (yet) including Higgs portal interactions corresponding to terms $I = J$. The neutrino Yukawa matrix h is written in a basis where both charged lepton mass matrix and Majorana mass matrix M_{IJ} are diagonal so that the N_I 's are the three Majorana mass eigenstates. One of the three RH neutrinos is assumed to have vanishing Yukawa couplings and, therefore, it is fully decoupled from the seesaw formula.⁴ Moreover, we assume that the Higgs-induced mixing between the two seesaw neutrinos is negligible, we will comment on this assumption. In our discussion we identify the dark neutrino N_D with the heaviest RH neutrino N_3 , so that its mass $M_D = M_3$. This corresponds to assume neutrino Yukawa matrix of the form

$$h = \begin{pmatrix} h_{e1} & h_{e2} & 0 \\ h_{\mu 1} & h_{\mu 2} & 0 \\ h_{\tau 1} & h_{\tau 2} & 0 \end{pmatrix}. \quad (5)$$

This matrix can be diagonalised by a bi-unitary transformation of the left-handed and RH neutrino fields, i.e.,

$$h = V_L^{\dagger} D_h U_R, \quad (6)$$

where $D_h \equiv \text{diag}(h_C, h_B, h_A)$ with $h_A \leq h_B \leq h_C$. Necessarily, starting from a Yukawa matrix of the form (5), one has $h_A = 0$. The RH neutrino mixing matrix U_R is of the form

$$U_R(\varepsilon) = \begin{pmatrix} \cos \varepsilon & \sin \varepsilon & 0 \\ -\sin \varepsilon & \cos \varepsilon & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (7)$$

⁴Therefore, at this stage, we should actually refer to N_D as a heavy neutral lepton. However, small Yukawa couplings are triggered by the Anisimov operator, as we are going to discuss soon.

where ε is a complex angle that generates a mixing between the seesaw neutrinos N_1 and N_2 . This mixing is necessary if one wants to have non-vanishing CP asymmetries in N_1 and N_2 -decays in order to generate a baryon asymmetry via leptogenesis. In our case, the most conservative choice, maximising the dark neutrino lifetime, is to take $\varepsilon \simeq 0$ so that approximately $U_R \simeq I$. Notice that even such an infinitesimal deviation of U_R from the identity can yield successful leptogenesis [29]. Within this setup, ordinary neutrino masses and mixing are generated by a minimal two RH neutrino type I seesaw mechanism, where the two seesaw neutrinos have to be identified with N_1 and N_2 and the lightest neutrino mass vanishes.

In order to maximise the dark neutrino lifetime, we can also assume the dark neutrino mixing only with the seesaw neutrino with smaller Yukawa coupling h_B . This plays the role of source neutrino, denoted by N_S , and it can be either the lightest RH neutrino N_1 or the next-to-lightest N_2 . For definiteness, we can choose $N_S = N_2$ so that $M_S = M_2$. For this reason in the following we will refer to a two-neutrino mixing formalism. In order to minimise the Yukawa coupling of the source neutrino and maximise the lifetime of DM, we can also assume normal ordering for the light neutrino masses. In this way one simply has $h_S = h_B = \sqrt{m_{\text{sol}} M_S}/v$. Of course, notice that with this choice one also has $h_C = \sqrt{m_{\text{atm}} M_I}/v$, where either $I = 1$, if $N_S = N_2$, or $I = 2$, if $N_S = N_1$. However, in order to have successful leptogenesis at a scale much below 10^9 GeV, necessarily $M_1 \simeq M_2$, so whether the source RH neutrino corresponds to the lightest or to the next-to-lightest seesaw neutrino does not have any implication in our discussion. In conclusion, the Lagrangian in Eq. (4) simplifies to ($I = 1, 2, 3, J = 1, 2$):

$$-\mathcal{L}_{Y+M+\tilde{\Lambda}_{\text{DS}}}^\nu = \overline{L}_\alpha h_{\alpha J} N_J \tilde{\Phi} + \frac{1}{2} \overline{N}_I^c M_I N_I + \frac{1}{\tilde{\Lambda}_{\text{DS}}} \Phi^\dagger \Phi \overline{N}_D^c N_S + \text{h.c.} . \quad (8)$$

After electroweak spontaneous symmetry breaking, the Higgs acquires a vev v that generates a neutrino Dirac mass matrix

$$m_D = v h = \begin{pmatrix} m_{D e1} & m_{D e2} & 0 \\ m_{D \mu 1} & m_{D \mu 2} & 0 \\ m_{D \tau 1} & m_{D \tau 2} & 0 \end{pmatrix} \quad (9)$$

from the usual Yukawa term but also a small off-diagonal correction to the Majorana mass term $\delta M_{\text{DS}}^\Lambda = v^2/\tilde{\Lambda}_{\text{DS}}$ generated by the Anisimov operator. Therefore, the full neutrino mass Lagrangian is now given by

$$-\mathcal{L}_{D+M+\delta M_{\text{DS}}^\Lambda}^\nu = \overline{\nu}_{L\alpha} m_{D\alpha J} N_J + \frac{1}{2} \overline{N}_I^c M_I N_I + \frac{1}{2} \overline{N}_D^c \delta M_{\text{DS}}^\Lambda N_S + \text{h.c.} . \quad (10)$$

In this way, as an effect of the Anisimov operator, the mass eigenstates change from N_D, N_S to

$$N_{D0}^\Lambda = N_{DM} \cos \theta_{\Lambda 0} - N_S \sin \theta_{\Lambda 0} \quad (11)$$

$$N_{S0}^\Lambda = N_{DM} \sin \theta_{\Lambda 0} + N_S \cos \theta_{\Lambda 0}, \quad (12)$$

where⁵

$$\theta_{\Lambda 0} = \frac{2v^2/\tilde{\Lambda}_{DS}}{M_D(1 - M_S/M_D)} \quad (13)$$

is the effective mixing angle at zero temperature. In this way the RH neutrino mixing matrix from the new mass eigenstate basis to the flavour basis can be written as

$$U_R^\Lambda(\theta_\Lambda^0) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta_\Lambda^0 & \sin \theta_\Lambda^0 \\ 0 & -\sin \theta_\Lambda^0 & \cos \theta_\Lambda^0 \end{pmatrix} \quad (14)$$

and in the new basis the Dirac mass matrix becomes

$$m_D^\Lambda = m_D U_R^\Lambda(\theta_\Lambda^0) = \begin{pmatrix} m_{D e1} & m_{D e2} \cos \theta_\Lambda^0 & m_{D e3} \sin \theta_\Lambda^0 \\ m_{D \mu 1} & m_{D \mu 2} \cos \theta_\Lambda^0 & m_{D \mu 3} \sin \theta_\Lambda^0 \\ m_{D \tau 1} & m_{D \tau 2} \cos \theta_\Lambda^0 & m_{D \tau 3} \sin \theta_\Lambda^0 \end{pmatrix}. \quad (15)$$

This shows that now the mass eigenstates created by the field N_D^Λ , those that have to be identified with the DM particles, will not be rigorously stable but decay with a lifetime [16]

$$\tau_D \simeq (\Gamma_{D \rightarrow A + \ell_S} + \Gamma_{D \rightarrow 3A + \ell_S})^{-1}, \quad (16)$$

where $\Gamma_{D \rightarrow A + \ell_S}$ is the two body decay rate given by

$$\Gamma_{D \rightarrow A + \ell_S} = \frac{h_S^2}{\pi} \left(\frac{v^2}{\tilde{\Lambda}} \right)^2 \frac{M_D}{(M_D - M_S)^2} \quad (17)$$

and $\Gamma_{D \rightarrow 3A + \ell_S}$ is the four body decay rate given, in the narrow width approximation, by

$$\Gamma_{D \rightarrow 3A + \ell_S} = \frac{\Gamma_S}{15 \cdot 2^{11} \cdot \pi^4} \frac{M_D}{M_S} \left(\frac{M_D}{\tilde{\Lambda}_{DS}} \right)^2, \quad (18)$$

where $\Gamma_S = h_S^2 M_S / (4\pi)$. The most stringent lower bound on the lifetime comes from the IceCube collaboration that, analysing 7.5 yr HESE data, has recently found approximately $\tau_D \gtrsim 10^{27} - 10^{28}$ s in the energy range 60 TeV–10 PeV [30], extending previous results [31].⁶

⁵This expression holds for $\theta_{\Lambda 0} \ll 1$. This is always verified unless one considers a case of extreme degeneracy, in which case the angle becomes maximal. Our results are not at all affected by such an approximation.

⁶At the moment, this should be regarded as an indicative approximate lower bound. We will later on discuss more detailed lower bounds that in fact depends on the DM mass and on the decay channel.




Figure 1: Self-energy diagrams from Yukawa interactions (panel (a)) and from Anisimov interactions (panels (b) and (c)).

At finite temperatures, the Yukawa and Anisimov interactions in Eq. (1) both contribute to the self-energies of the RH neutrinos affecting their propagation and mixing. These are diagrammatically shown in panels (a) and (b) of Fig. 1.

The effect can be described in terms of the effective potentials that, in the Yukawa basis, are given respectively by [32] ($I, J = D, S$)

$$V_{IJ}^Y = \frac{T^2}{8 E_J} h_J^2 \delta_{IJ}, \quad (19)$$

and

$$V_{IJ}^\Lambda \simeq \frac{T^2}{12 \Lambda} \lambda_{IJ}. \quad (20)$$

The contribution from the Anisimov interactions is not diagonal in general in the Yukawa basis and this misalignment generates a mixing that strongly depends on temperature. The evolution of the dark and source neutrino eigenstates will then be described by the Hamiltonian

$$\mathcal{H} = \begin{pmatrix} E_{\text{DM}} & \frac{T^2}{12 \Lambda_{\text{DS}}} \\ \frac{T^2}{12 \Lambda_{\text{DS}}} & E_{\text{S}} + \frac{T^2}{8 E_{\text{S}}} h_{\text{S}}^2 \end{pmatrix}. \quad (21)$$

Subtracting a term proportional to the identity, the effective mixing Hamiltonian is then given by

$$\Delta \mathcal{H} \simeq \begin{pmatrix} -\frac{\Delta M^2}{4p} - \frac{T^2}{16p} h_{\text{S}}^2 & \frac{T^2}{12 \Lambda_{\text{DS}}} \\ \frac{T^2}{12 \Lambda_{\text{DS}}} & \frac{\Delta M^2}{4p} + \frac{T^2}{16p} h_{\text{S}}^2 \end{pmatrix}, \quad (22)$$

where we used the ultra-relativistic approximation⁷ and defined $\Delta M^2 \equiv M_S^2 - M_D^2$.

The Yukawa interactions produce a source neutrino abundance that we denote by N_{N_S} and normalise it in a way that in ultrarelativistic thermal equilibrium $N_{N_S}^{\text{eq}}(z_S \ll 1) = 1$, where $z_S \equiv M_S/T = z M_S/M_D$ and $z \equiv M_D/T$. The production of source neutrinos is then described by the simple rate equation

$$\frac{dN_{N_S}}{dz} = -(D + S)(N_{N_S} - N_{N_S}^{\text{eq}}), \quad (23)$$

where $D \equiv \Gamma_D/(H z)$, $S \equiv \Gamma_S/(H z)$ and Γ_D and Γ_S are the source neutrino total decay and the $\Delta L = 1$ scattering rates respectively. Finally, $H = H(z)$ is the expansion rate given by

$$H(z) = \sqrt{\frac{8\pi^3 g_R}{90}} \frac{M_D^2}{M_{\text{P}}} \frac{1}{z^2}, \quad (24)$$

where for the value of the number of ultrarelativistic degrees of freedom we can simply take the SM value, so that $g_R = g_R^{\text{SM}} = 106.75$. The off-diagonal term in the Hamiltonian will then mix source and dark neutrinos. Adopting a monochromatic approximation, the momentum can be replaced by its average value $p \simeq 3T$ and the effective mixing Hamiltonian in the flavour basis becomes

$$\Delta \mathcal{H} \simeq \frac{\Delta M^2}{12T} \begin{pmatrix} -1 - v_S^Y & \sin 2\theta_\Lambda \\ \sin 2\theta_\Lambda & 1 + v_S^Y \end{pmatrix}. \quad (25)$$

Here we have also introduced the dimensionless effective potential $v_S^Y \equiv T^2 h_S^2/(4\Delta M^2)$ and the effective mixing angle $\sin 2\theta_\Lambda(T) \equiv T^3/(\tilde{\Lambda}_{\text{DS}} \Delta M^2)$, parameterising the misalignment between the Yukawa and the Higgs-induced interactions.

The production of dark neutrinos can be described by the density matrix equation [16]

$$\frac{d\mathcal{N}}{dz} = -\frac{i}{H(z)z} [\Delta \mathcal{H}, \mathcal{N}] - \begin{pmatrix} 0 & \frac{1}{2}(D + S)\mathcal{N}_{\text{DS}} \\ \frac{1}{2}(D + S)\mathcal{N}_{\text{SD}} & (D + S)(N_{N_S} - N_{N_S}^{\text{eq}}) \end{pmatrix}, \quad (26)$$

where the diagonal elements give the abundances of dark neutrinos, $N_{N_D} = \mathcal{N}_{\text{DD}}$, and source neutrinos, $N_{N_S} = \mathcal{N}_{\text{SS}}$. Notice how decays and scatterings also contribute to decoherence effects, damping the density matrix off-diagonal terms. Initially, for $z = z_{\text{in}}$, the density matrix is simply given by:

$$\mathcal{N}(z_{\text{in}}) = N_{N_S}(z_{\text{in}}) \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}. \quad (27)$$

⁷This assumes that the dark neutrino production occurs in the ultra-relativistic regime.

Expressing the matrices in the Pauli matrix basis, the density matrix equation can be recast in a vectorial notation. The effective Hamiltonian can be written as

$$\Delta\mathcal{H} = \frac{1}{2} \vec{V} \cdot \vec{\sigma}, \quad (28)$$

where the *effective potential vector* \vec{V} is defined as

$$\vec{V} \equiv \frac{\Delta M^2}{6T} (\sin 2\theta_\Lambda, 0, -1 - v_S^Y). \quad (29)$$

The abundance normalised density matrix is analogously recast, introducing the quantity P_0 and the polarisation vector \vec{P} , as

$$\mathcal{N} = \frac{1}{2} P_0 (1 + \vec{P} \cdot \vec{\sigma}), \quad (30)$$

in a way that

$$N_{N_D} = \frac{1}{2} P_0 (1 + P_z), \quad (31)$$

$$N_{N_S} = \frac{1}{2} P_0 (1 - P_z), \quad (32)$$

$$N_{N_D} + N_{N_S} = P_0. \quad (33)$$

Inserting Eqs. (28) and (30) into the density matrix equation (26), one obtains a set of equations for P_0 and \vec{P} :

$$\frac{d\vec{P}}{dz} = \vec{V} \times \vec{P} - \left[\frac{1}{2}(D + S) + \frac{d \ln P_0}{dz} \right] \vec{P}_T - (1 + P_z) \frac{d \ln P_0}{dz} \hat{z}, \quad (34)$$

$$\frac{dP_0}{dz} = -(D + S) (N_{N_S} - N_{N_S}^{\text{eq}}), \quad (35)$$

where we defined $\vec{P}_T \equiv P_x \hat{x} + P_y \hat{y}$ and $\vec{V} \equiv \vec{V}/(Hz)$.

In Fig. 2 we show the evolution of N_{N_D} and N_{N_S} for the benchmark values $M_S = 300 \text{ GeV}$, $M_D = 220 \text{ TeV}$, $\tau_D = 3.48 \times 10^{28} \text{ s}$ and assuming an initial vanishing N_S abundance. These particular values for M_D and τ_D correspond to best fit values found in a likelihood statistical analysis of 6 year High Energy Starting Events (HESE) IceCube where the presence of a neutrino flux contribution from neutrinophilic DM decays in addition to an astrophysical component with fixed spectral index $\gamma = 2.2$ is favoured at more than $\sim 3\sigma$ compared to the null hypothesis where there is no decaying DM [33].⁸ The horizontal line denotes $N_{N_D}^{\text{f,obs}}$, the final value of N_{N_D} that reproduces the measured

⁸Another analysis of 6 year HESE data where, differently from [33], the spectral index is let to vary and DM decays via one decay channel, also finds that the addition of a component from decaying DM improves the fit. The best fit is obtained for DM decaying into W bosons, a mass $M_{DM} \sim 400 \text{ TeV}$ and for a value of the spectral index $\gamma \simeq 2.3$ but with a lower statistical significance $\sim 2\sigma$ [34].




Figure 2: Evolution of the source and dark neutrino abundances for different values of the initial temperature and for the indicated benchmark values of M_D , M_S and τ_D . Top panel: initial thermal N_{N_S} -abundance. Bottom panel: initial vanishing N_{N_S} -abundance.

DM energy density at the present time $\Omega_{\text{DM}} h^2 \simeq 0.119$. For a generic mass M_D , one finds [16]

$$N_{N_D}^{f,\text{obs}} \simeq 1.1 \times 10^{-7} \frac{\text{GeV}}{M_D}. \quad (36)$$

One can notice how the final dark neutrino abundance falls many orders of magnitude below $N_{N_D}^{f,\text{obs}}$. The value $M_S = 300 \text{ GeV}$ corresponds to a minimum possible value to have successful leptogenesis from decays independently of the initial conditions, so that one would conclude that it is not possible to have dark neutrinos as DM and strong thermal leptogenesis in a unified picture. In the next section we will see how the introduction of Higgs portal interactions for the source neutrinos drastically changes this conclusion.

3 Including Higgs portal interactions for N_S

Let us now consider the effect of introducing Higgs portal interactions for the source neutrinos so that the Lagrangian in Eq. (8) becomes

$$-\mathcal{L}_{Y+M+\Lambda}^\nu = \overline{L}_\alpha h_{\alpha J} N_J \tilde{\Phi} + \frac{1}{2} \overline{N}_I^c M_I N_I + \frac{1}{\tilde{\Lambda}_{DS}} \Phi^\dagger \Phi \overline{N}_D^c N_S + \frac{1}{\tilde{\Lambda}_{SS}} \Phi^\dagger \Phi \overline{N}_S^c N_S + \text{h.c.}, \quad (37)$$

where we introduced the effective scale for the Higgs portal interactions $\tilde{\Lambda}_{SS} \equiv \Lambda/\lambda_{SS}$.

At zero temperature, after electroweak spontaneous symmetry breaking, they will yield a contribution $\delta M_S^\Lambda = (v^2/\tilde{\Lambda}_{SS})$ to the source neutrino mass that, however, can be safely neglected, anticipating that we will obtain $\tilde{\Lambda}_{SS} \gtrsim 10^8$ GeV. At finite temperatures they will give a contribution to the self-energy (see panel (c) in Fig. 1) and, therefore, to the effective potential, given by

$$V_{SS}^\Lambda = \frac{T^2}{12 \tilde{\Lambda}_{SS}}. \quad (38)$$

This is in addition to the term from the Yukawa couplings in Eq. (38). In this way the effective Hamiltonian in Eq. (22) would now become

$$\Delta \mathcal{H} \simeq \begin{pmatrix} -\frac{\Delta M^2}{4p} - \frac{T^2}{16p} h_S^2 - \frac{T^2}{24 \tilde{\Lambda}_{SS}} & \frac{T^2}{12 \tilde{\Lambda}_{DS}} \\ \frac{T^2}{12 \tilde{\Lambda}_{DS}} & \frac{\Delta M^2}{4p} + \frac{T^2}{16p} h_S^2 + \frac{T^2}{24 \tilde{\Lambda}_{SS}} \end{pmatrix}. \quad (39)$$

This additional term can be comparable or even larger than the term from the Yukawa couplings depending whether $T/\tilde{\Lambda}_{SS} \sim h_S^2$ or higher. However, in the relevant range of temperatures where dark neutrinos are produced the effect should be limited, at least for large values of $\tilde{\Lambda}_{SS} \gg 10^{10}$ GeV. This is because the dark neutrino production occurs at temperatures much below the resonance [16]. This conclusion is not changed by this additional term and, therefore, like also the effective potential from Yukawa couplings, one can expect that it should not considerably affect the final abundance of dark neutrinos. In any case it represents a subdominant effect compared to the contribution of Higgs scatterings to the production of source neutrinos, the main focus of this paper. For this reason we will neglect it here but it will be worth to explore its impact in a future paper.

The most important effect of Higgs portal interactions is their contribution to the source neutrino production coming from Higgs scatterings $\phi \phi^\dagger \rightarrow N_S N_S$ (diagrammatically shown in the panel (a) of Fig. 3). In this way Eq. (35) gets generalised into

$$\frac{dP_0}{dz} \simeq \frac{dN_{N_S}}{dz} = -(D + S) (N_{N_S} - N_{N_S}^{\text{eq}}) - A (N_{N_S}^2 - N_{N_S}^{\text{eq}2}), \quad (40)$$

where we defined

$$A \equiv \frac{\langle \sigma_{\phi \phi^\dagger \rightarrow N_S N_S^c} v_{\text{rel}} \rangle}{H(z) z R^3(z)}, \quad (41)$$




Figure 3: Higgs portal interactions: source neutrino production (a), dark neutrino production (b) and mixed production (c).

with $\langle \sigma_{\phi\phi^\dagger \rightarrow N_S N_S^c} v_{\text{rel}} \rangle$ the thermal averaged cross section. Here $R^3(z)$ is the portion of comoving volume where abundances are calculated, essentially a normalisation factor. With our choice, $N_{N_S}^{\text{eq}}(z_S \ll 1) = 1$, one has

$$R^3(z) = \frac{4}{3} \frac{\pi^2}{g_{N_S} \zeta(3)} \frac{z^3}{M_D^3}. \quad (42)$$

In the regime $M_S \ll T$ we are interested in, the thermal averaged cross section is simply given by [36]

$$\langle \sigma_{\phi\phi^\dagger \rightarrow N_S N_S} v_{\text{rel}} \rangle \Big|_{M_S \ll T} = \frac{\tilde{\Lambda}_{SS}^{-2}}{4\pi}. \quad (43)$$

Combining all together, we obtain:

$$A(z) = \frac{A_1}{z^2}, \quad \text{with } A_1 \equiv A(z=1) = \frac{3}{16} \frac{\zeta(3)}{\pi^3} g_{N_S} \sqrt{\frac{90}{8\pi^3 g_R}} \frac{M_D M_P}{\tilde{\Lambda}_{SS}^2}. \quad (44)$$

It is also convenient to give a numerical expression for A_1 ,

$$A_1 \simeq 1.0 \times 10^{-11} \left(\frac{M_D}{100 \text{ TeV}} \right) \left(\frac{10^{16} \text{ GeV}}{\tilde{\Lambda}_{SS}} \right)^2, \quad (45)$$

where we used the SM value $g_R = 106.75$ for the number of ultrarelativistic degrees of freedom and $g_{N_S} = 2$. It is simple to find an approximate solution for $N_{N_S}(z)$, valid in

the regime $N_{N_S} \ll N_{N_S}^{\text{eq}}$ and for $z \ll 1$, so that we can take $N_{N_S}^{\text{eq}} \simeq 1$. Assuming that the production from Higgs portal interactions dominates and taking as initial condition $N_{N_S}(z_{\text{in}}) = 0$, one immediately finds

$$N_{N_S}(z) \simeq A_1 \left(\frac{1}{z_{\text{in}}} - \frac{1}{z} \right), \quad (46)$$

implying asymptotically, but still for $z \ll 1$,

$$N_{N_S}(z_{\text{in}} \ll z \ll 1) \simeq \frac{A_1}{z_{\text{in}}} \simeq 1.0 \times \left(\frac{T_{\text{in}}}{10^{16} \text{ GeV}} \right) \left(\frac{10^{16} \text{ GeV}}{\tilde{\Lambda}_{\text{SS}}} \right)^2, \quad (47)$$

where the numerical expression has been obtained from Eq. (45) and where notice that the dependence on M_D has cancelled out. This expression of course is valid only for $N_{N_S}(z_{\text{in}} \ll z \ll 1) \ll 1$, so that it can be also read as a condition on T_{in} and $\tilde{\Lambda}_{\text{SS}}$ for the thermalisation of the source neutrinos prior to the onset of source-dark neutrino oscillations. In this case it can be also simply recast as $\tilde{\Lambda}_{\text{SS}}^2/T_{\text{in}} = 10^{16} \text{ GeV}$. Notice that, for the validity of the effective theory, one has to impose $T_{\text{in}} \lesssim \tilde{\Lambda}_{\text{SS}}$.

As initial temperature we can assume $T_{\text{in}} = T_{\text{RH}}$, where T_{RH} is the reheat temperature below which one can assume a radiation dominated regime.⁹ The result we obtained shows that for $\tilde{\Lambda}_{\text{SS}} \leq 10^{16} \text{ GeV}$ one can always obtain a full thermalisation of the source neutrino abundance by increasing T_{RH} . In particular, in the limit case $\tilde{\Lambda}_{\text{SS}} = 10^{16} \text{ GeV}$, the thermalisation can be obtained for the maximum value allowed by cosmological observations $T_{\text{RH}} = 10^{16} \text{ GeV}$.

In Fig. 4 we show, for the same values of M_S , M_D and τ_D as in Fig. 2, the evolution of N_{N_S} and N_{N_D} when Higgs portal interactions for N_S are taken into account. In the upper panel we fixed $\tilde{\Lambda}_{\text{SS}} = 10^{16} \text{ GeV}$ and we show solutions for the indicated values of T_{RH} . One can clearly see how the source neutrino abundance thermalises for the maximum value $T_{\text{RH}} = 10^{16} \text{ GeV}$, while it is suppressed linearly for lower values. Of course it should be noticed how, for $T_{\text{RH}} = 10^{16} \text{ GeV}$, one obtains a DM abundance that is greater than the observed value, clearly indicating that one can always find choices of the parameters reproducing the observed value.

In the lower panel we have conservatively imposed $T_{\text{RH}} = \tilde{\Lambda}_{\text{SS}}$ and we show the evolution of the source and dark neutrino abundances for the indicated values of $\tilde{\Lambda}_{\text{SS}} \leq 10^{16} \text{ GeV}$. Notice how in this case the thermalisation condition is always respected. Therefore, one obtains solutions that are clearly very close to those shown in the upper

⁹We do not include a production between $T_{\text{max}} > T_{\text{RH}}$ and T_{RH} [36] since this would just very slightly contribute to relax the final constraints.




Figure 4: Effect of source neutrino Higgs portal interactions on the source and dark neutrino abundances.

panel of Fig. 2 with the difference that now the thermalisation is not just assumed but obtained as the result of the Higgs portal interactions for the source neutrino.

Finally, let us highlight again that we are neglecting a possible production, pure or mixed, of dark neutrinos directly from Higgs portal interactions (shown diagrammatically in the panel (b) and panel (c) of Fig. 3). This is equivalent to say that their corresponding effective scales $\tilde{\Lambda}_{DD} \equiv \lambda_{DD}/\Lambda$ and $\tilde{\Lambda}_{DS}$, are sufficiently large that the associated production is negligible compared to the contribution from neutrino mixing. It is simple to derive a lower bound on $\tilde{\Lambda}_{DD}$ and $\tilde{\Lambda}_{DS}$ imposing that the dark neutrino relic abundance produced from Higgs portal interactions is negligible compared to the observed abundance in Eq. (36). This abundance can be calculated in the same way as the source neutrino abundance and therefore it will be given by Eq. (47) simply replacing $\tilde{\Lambda}_{SS}$ with $\tilde{\Lambda}_{DD}$ or $\tilde{\Lambda}_{DS}$. One can

then easily derive the conditions

$$\tilde{\Lambda}_{\text{DD}}, \tilde{\Lambda}_{\text{DS}} \gg 10^{22} \text{ GeV} \sqrt{\frac{T_{\text{RH}}}{10^{16} \text{ GeV}} \frac{M_{\text{D}}}{\text{PeV}}}. \quad (48)$$

We will discuss in Section 5 a model where these two conditions are naturally satisfied.

4 Allowed regions in the dark neutrino lifetime-mass plane and experimental constraints

We determined the allowed regions in the dark neutrino lifetime-mass plane and we show the results in Figs. 5–8. In the first case, as in Fig. 1, we fix $\tilde{\Lambda}_{\text{SS}} = 10^{16} \text{ GeV}$ and show the allowed regions for different values of the reheat temperature, with $T_{\text{RH}} \leq \tilde{\Lambda}_{\text{SS}}$, as indicated. The different panels are for $M_{\text{S}} = 300 \text{ GeV}$ (top) and 1 TeV (bottom) in Fig. 5 and for $M_{\text{S}} = 10 \text{ TeV}$ (top) and 100 TeV (bottom) in Fig. 6. The allowed regions are obtained imposing $N_{N_{\text{D}}} \geq N_{\text{D}}^{\text{f,obs}}$. On the borders one has exactly $N_{N_{\text{D}}} = N_{\text{D}}^{\text{f,obs}}$, while any point inside would correspond to an overabundance but this can be lowered simply lowering T_{RH} and/or increasing M_{S} .

In each panel the shadowed region for $\tau_{\text{D}} \leq 10^{28} \text{ s}$ is indicatively the region that is currently tested at neutrino telescopes such as IceCube and gamma ray telescopes such as the Fermi Gamma-Ray telescope, placing lower bounds on the lifetime. An accurate lower bound in fact depends on the mass of the DM particle and on the specific primary decay channel. It also depends on a description of the astrophysical contribution that in this case plays the role of a background. A positive signal should show up as an excess with respect to this astrophysical background. We show some of the lower bounds on the lifetime of a decaying DM recently obtained by the IceCube collaboration at 90% C.L. [30]. The thin solid line indicates the lower bound on the DM mass in the range $160 \text{ TeV} - 20 \text{ PeV}$ obtained from 7.5 yr High-Energy Starting Event (HESE) data in the energy range 60 TeV to 10 PeV for the decay channel $\text{DM} \rightarrow \text{Higgs} + \nu$. We also show, with a thick solid line, the same lower bound but for the decay channel $\text{DM} \rightarrow b + \bar{b}$. At lower masses we also indicate, with a dashed line, a lower bound obtained from 2 yr cascade events for the decay channel $\text{DM} \rightarrow \mu \bar{\mu}$ and at even lower masses, with a dotted line, the lower bound from the Fermi gamma-ray telescope for the channel $\text{DM} \rightarrow \nu \bar{\nu}$ also reported in [30]. Of course our case is different from the considered decay channels since the dark neutrino can decay both into Higgs boson and gauge bosons plus neutrino or charged leptons. All shown lower bounds should then be regarded as indicative and a dedicated analysis would be needed. Interestingly, the IceCube collaboration also confirms the presence of




Figure 5: Allowed regions in the lifetime versus mass plane for a fixed value $\tilde{\Lambda}_{\text{SS}} = 10^{16}$ GeV, for the indicated values of T_{RH} and for $M_{\text{S}} = 300$ GeV (upper panel), 1 TeV (bottom panel).

an excess at $\mathcal{O}(100 \text{ TeV})$ energies compared to an astrophysical component. The decaying DM hypothesis improves the data fit with a 2.5σ statistical significance and the best fit is found for the decay channel $\text{DM} \rightarrow b\bar{b}$ with $M_{\text{DM}} = 289 \text{ TeV}$ and $\tau_{\text{DM}} = 2.8 \times 10^{27} \text{ s}$. This best fit point is denoted by a star in the panels and it should also be regarded as indicative. It would be of course interesting to see whether a dedicated analysis within the RHINO model can also address the excess and with which statistical significance.

In Figs. 7 and 8 we show the allowed regions respectively for the indicated values of $\tilde{\Lambda}_{\text{SS}} = T_{\text{RH}}$ and again for $M_{\text{S}} = 300$ GeV (top) and 1 TeV (bottom) in Fig. 7 and for $M_{\text{S}} = 10$ TeV (top) and 100 TeV (bottom) in Fig. 8.

One can see that, as far as $\tilde{\Lambda}_{\text{SS}} = T_{\text{RH}} \gg 10^{10} \text{ GeV}$, the allowed regions get only slightly reduced when $\tilde{\Lambda}_{\text{SS}}$ decreases. On the other hand, for $\tilde{\Lambda}_{\text{SS}} = T_{\text{RH}} \lesssim 10^{10} \text{ GeV}$,




Figure 6: Allowed regions in the lifetime versus mass plane for a fixed value $\tilde{\Lambda}_{SS} = 10^{16}$ GeV, for the indicated values of T_{RH} and for $M_S = 10$ TeV (upper panel), 100 TeV (bottom panel).

the reduction becomes quite significant and for $\tilde{\Lambda}_{SS} = T_{RH} \lesssim 10^8$ GeV there is basically no allowed region, showing that $T_{RH} \gtrsim 10^8$ GeV can be regarded as a conservative lower bound for T_{RH} , of course under the assumption that dark neutrinos play the role of DM. This result is a consequence of what we have already noticed discussing the dark neutrino abundance evolution shown in the bottom panel of Fig. 4: since the onset of neutrino oscillations, and consequently the dark neutrino production, occurs approximately at $T \sim 10^9$ GeV, as far as the reheat temperature is much greater than this value there is no much variation when T_{RH} is increased, but for lower values the final dark neutrino abundance gets strongly suppressed. Notice that such a lower bound on the reheat temperature would be compatible with the well known upper bound $T_{RH} \lesssim 10^{10}$ GeV from the gravitino problem in gravity mediated supersymmetric models [35]. The same lower bound also




Figure 7: Allowed regions in the lifetime versus mass plane for the indicated values of $\tilde{\Lambda}_{\text{SS}} = T_{\text{RH}}$ for $M_S = 300 \text{ GeV}$ (top) and $M_S = 1 \text{ TeV}$.

holds on the effective scale $\tilde{\Lambda}_{\text{SS}} \gtrsim 10^8 \text{ GeV}$. This lower bound confirms the validity of having neglected the Majorana mass term originating from the Higgs portal operator for the source neutrino in Eq. (3), since one has $|\delta M_S^\Delta| \lesssim v^2 / \tilde{\Lambda}_{\text{SS}} \lesssim 0.1 \text{ MeV}$.

It should also be noticed that for $\tilde{\Lambda}_{\text{SS}} = T_{\text{RH}} = 10^{16} \text{ GeV}$, the value that maximises the dark neutrino final abundance, there are allowed regions only for $M_S \lesssim 100 \text{ TeV}$. This should be regarded as an upper bound of the model on the seesaw scale.¹⁰

¹⁰However, notice that our analysis assumes $M_D \geq M_S$. It should be understood whether there can be solutions also in the case $M_S > M_{\text{DM}}$, this interesting possibility requires a dedicated study and will be explored elsewhere.




Figure 8: As in Fig. 7 but for $M_{\text{S}} = 10 \text{ TeV}$ (top) and $M_{\text{S}} = 100 \text{ TeV}$ (bottom).

5 UV-completing RHINO

Let us now finally discuss two possible UV-complete RHINO models that were already qualitatively sketched in the conclusions of [13]. In the first case, the mediator in the Anisimov operators in Eq. (1) is a heavy scalar H with vanishing vev. In the second case, the mediator is a heavy fermion F .¹¹

5.1 Heavy scalar H as mediator

Let us consider an extension of the seesaw Lagrangian where a heavy real scalar field H (with vanishing vev) is introduced and it couples to the RH neutrinos with Yukawa

¹¹Of course there could be more than one heavy fermion, the generalisation is straightforward.




Figure 9: Feynman diagrams with a heavy scalar H as mediator and $I, J = D, S$. Integrating out H , they lead to the Feynman diagrams in panel (b) of Fig. 2 and (a), (b) and (c) of Fig. 3.

couplings y_{IJ} and to the standard Higgs field with a trilinear coupling μ :¹²

$$\mathcal{L}_H = \frac{1}{2} \partial_\mu H \partial^\mu H - \frac{1}{2} M_H^2 H^2 - \sum_{I,J} \lambda_{IJ} H \overline{N}_I^c N_J - \mu H \phi^\dagger \phi. \quad (49)$$

At scales much below M_H we can integrate out H , obtaining the effective Lagrangian

$$\mathcal{L}_H^{\text{eff}} = \frac{1}{2} \sum_{I,J,K,L} \frac{\lambda_{IJ} \lambda_{KL}}{M_H^2} (\overline{N}_I^c N_J) (\overline{N}_K^c N_L) + \frac{1}{2} \frac{\mu^2}{M_H^2} (\phi^\dagger \phi)^2 + \sum_{I,J} \frac{\mu \lambda_{IJ}}{M_H^2} \Phi^\dagger \Phi \overline{N}_I^c N_J. \quad (50)$$

One can clearly recognise the Anisimov operators in Eq. (2) where the effective scales can be identified with $\tilde{\Lambda}_{IJ} = \Lambda / \lambda_{IJ}$, where $\Lambda = M_H^2 / \mu$. Diagrammatically, the self-energy diagram in the panel (b) of Fig. 1 and the scattering diagram in panel (a) of Fig. 3 are obtained by the diagrams in Fig. 9, panel (a) and panel (b), respectively. The appealing feature of this model is that one can get a trans-Planckian value for the effective scale $\tilde{\Lambda}_{DS} \sim 10^{23}$ GeV even for $\lambda_{IJ} \sim 1$, simply choosing $\mu \ll M_{\text{GUT}}$, for example, $M_H \sim M_{\text{GUT}} \sim 10^{16}$ GeV and $\mu \sim 10^9$ GeV. However, the problem of this setup is that in this case one cannot also reproduce the effective scale $\tilde{\Lambda}_{SS} \sim 10^{16}$ GeV for the source neutrino Higgs portal interactions. In that respect, one should arbitrarily assume $\Lambda \sim 10^{16}$ GeV, for example for $\mu = M_H \sim M_{\text{GUT}} \sim 10^{16}$ GeV, $\lambda_{DS} \sim 10^{-7}$ and $\lambda_{SS} \ll 10^{-7}$ in order for $\tilde{\Lambda}_{DD}$ to satisfy the condition in Eq. (48).

However, there is a much simpler model where one can nicely understand the values of the effective scales that, as we have seen, would be able to address the DM problem

¹²This model was also discussed in [15, 36].




Figure 10: Feynman diagrams with a heavy fermion F as mediator and $I, J = D, S$. Integrating out F , they lead to the Feynman diagrams in panel (b) of Fig. 2 and in panels (a) and (b) of Fig. 4.

compatibly with successful strong thermal leptogenesis and experimental constraints from neutrino telescopes.

5.2 Heavy fermion F as mediator

Let us this time extend the seesaw Lagrangian introducing an heavy fermion doublet F with Yukawa couplings y_I to RH neutrinos, explicitly:

$$\mathcal{L}_F = \bar{F} (i \not{\partial} - M_F) F - \sum_I y_I (\bar{F} \phi N_I + \bar{N}_I \phi^\dagger F). \quad (51)$$

At scales much below M_F one can integrate out F obtaining the effective Lagrangian

$$-\mathcal{L}_F^{\text{eff}} = \sum_{I,J} \frac{y_I y_J}{M_F} \bar{N}_I N_J \phi^\dagger \phi, \quad (52)$$

where the RH side coincides with the Anisimov operators with the simple identification $\Lambda = M_F$ and $\lambda'_{IJ} = y_I y_J$. The three Anisimov operators in Eq. (3), Higgs-induced neutrino mixing, source neutrino Higgs portal interactions and dark neutrino Higgs portal interactions, can then be regarded as the low energy effective operators generated by the three diagrams in Fig. 10, respectively.

If we take $M_F \sim M_{\text{GUT}}$, $y_S \sim 1$ and $y_D \sim 10^{-7}$, one can immediately reproduce the values $\tilde{\Lambda}_{\text{SS}} \sim 10^{16}$ GeV, $\tilde{\Lambda}_{\text{DS}} \sim 10^{23}$ GeV and $\tilde{\Lambda}_{\text{DD}} \sim 10^{30}$ GeV. As we have seen, these

are the correct values to reproduce the observed DM abundance from Higgs-induced RH neutrino mixing, with source neutrino Higgs portal interactions able to thermalise the source neutrino abundance prior to the onset of the oscillations and with a suppressed contribution to dark neutrino production that we have indeed neglected. These values are much less arbitrary than the choice in the previous model, since the three couplings λ_{IJ} are the product of just two Yukawa couplings and it is non trivial that the third is obtained automatically and satisfying correctly the condition in Eq. (48). Moreover, they can be well understood imposing a \mathbb{Z}_2 symmetry under which all particles are even, except the dark neutrino that is odd. In this way the small Yukawa coupling $y_D \sim 10^{-7}$ could be regarded as a small symmetry breaking parameter connecting the visible sector to the dark sector.

6 Conclusions

We have seen how, including Higgs portal interactions for the source neutrino, the RHINO model can fully express its potential, providing a model for a decaying heavy DM particle in the range 1 TeV–1 PeV, compatible with strong thermal resonant leptogenesis and testable at neutrino telescopes. In this way RHINO can be regarded as quite a minimal model of the origin of matter and neutrino masses. Since the leptogenesis scale can be higher than the sphaleron freeze-out temperature, the final matter-antimatter asymmetry is independent of the initial conditions. Notice, moreover, that the dark neutrino abundance is independent of a possible external contribution to the production of the source neutrino abundance, since this is anyway thermalised by Higgs portal interactions. On the other hand, of course, it is not independent of a possible additional direct production of the dark neutrinos from some external mechanism, that, therefore, has to be assumed to be negligible or in any case sub-dominant. In this respect the main competitive mechanism is a possible gravitational production. However, typically, this is non negligible only for even heavier particles, for example in the case of WIMPzillas [36, 37]. Therefore, RHINOs and WIMPzillas seem to be successful candidates of DM particles in different mass ranges. In addition, based just on cosmological considerations, we obtained as an extra attractive feature of the model that the natural fundamental scale for the effective interactions, responsible within RHINO for the production and the decay of the DM, is the grandunified scale. We have also seen how our results point to a simple UV-complete model where the mediator of the Anisimov interactions is a heavy fermion. It should be appreciated how this UV-complete model simultaneously yields correctly the three effective scales in the Anisimov operators when the fundamental scale is identified with

the grandunified scale (or close to it). Our results also show that the RHINO model can nicely address the current hint from 7.5yr HESE IceCube data for a 100 TeV excess in the high energy neutrino flux on the top of an astrophysical component with spectral index $\gamma \simeq 2.2$. If an explanation in terms of a decaying DM is correct, then gradually this excess should exhibit anisotropies tracking current DM distribution, since the signal would be simply proportional to the DM density. We believe that the RHINO model would then be the leading candidate to explain such an excess, since it is the only model that genuinely *pre-dicts* such a signal [13], since the same physics is responsible both for DM production and its decays. As suggested in [15], further experimental tests could rely on the flavour composition of primary neutrinos. It would be of course also interesting to explore the potential of future planned 100 TeV colliders and possible links with flavour anomalies.

Acknowledgments

PDB acknowledges financial support from the STFC Consolidated Grant ST/T000775/1. AM is supported by a DiscNET/NGCM scholarship. PDB wishes to thank the organisers of the workshop on *Neutrino Theories* held at the Institute for theoretical physics at UAM, Madrid, 16 May - 17 June and the organisers of the workshop on the *Standard Model and beyond* held at the Corfu Summer Institute, 28 August- 8 September 2022, where part of this work was carried out. During these workshops he could benefit from stimulating talks and discussions with Sasha Belyaev, Thomas Hambye, Steve King, Rocky Kolb, Gino Isidori, Nick Mavromatos, Apostolos Pilaftsis, Subir Sarkar, Mikhail Shaposhnikov, Alexey Smirnov and Jim Talbert.

References

- [1] For a recent review and references see P. Di Bari, *On the origin of matter in the Universe*, Prog. Part. Nucl. Phys. **122** (2022), 103913 [arXiv:2107.13750 [hep-ph]].
- [2] P. Minkowski, $\mu \rightarrow e\gamma$ *At A Rate Of One Out Of 1-Billion Muon Decays?*, Phys. Lett. B **67** (1977) 421; T. Yanagida, *Horizontal gauge symmetry and masses of neutrinos*, Conf. Proc. C **7902131** (1979) 95. Proceedings of the Workshop on Unified Theory and Baryon Number of the Universe, eds. O. Sawada and A. Sugamoto (KEK, 1979) p.95; P. Ramond, Invited talk given at Conference: C79-02-25 (Feb 1979) p.265-280, CALT-68-709, *The Family Group in Grand Unified Theories*, hep-ph/9809459; M. Gell-Mann, P. Ramond and R. Slansky, in Supergravity, eds. P. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979) Conf.Proc.

- C790927 p.315, PRINT-80-0576; R. Barbieri, D. V. Nanopoulos, G. Morchio and F. Strocchi, *Neutrino Masses in Grand Unified Theories*, Phys. Lett. **90B** (1980) 91. R. N. Mohapatra and G. Senjanovic, *Neutrino Mass and Spontaneous Parity Nonconservation*, Phys. Rev. Lett. **44** (1980) 912.
- [3] S. F. King, *Models of Neutrino Mass, Mixing and CP Violation*, J. Phys. G **42** (2015), 123001 [arXiv:1510.02091 [hep-ph]].
- [4] M. Fukugita and T. Yanagida, *Baryogenesis Without Grand Unification*, Phys. Lett. B **174** (1986) 45.
- [5] S. Dodelson and L. M. Widrow, *Sterile-neutrinos as dark matter*, Phys. Rev. Lett. **72** (1994), 17-20 [arXiv:hep-ph/9303287 [hep-ph]].
- [6] E. K. Akhmedov, V. A. Rubakov and A. Y. Smirnov, *Baryogenesis via neutrino oscillations*, Phys. Rev. Lett. **81** (1998), 1359-1362 [arXiv:hep-ph/9803255 [hep-ph]].
- [7] T. Asaka and M. Shaposhnikov, *The ν MSM, dark matter and baryon asymmetry of the universe*, Phys. Lett. B **620** (2005), 17-26 [arXiv:hep-ph/0505013 [hep-ph]].
- [8] X. D. Shi and G. M. Fuller, *A New dark matter candidate: Nonthermal sterile neutrinos*, Phys. Rev. Lett. **82** (1999), 2832-2835 [arXiv:astro-ph/9810076 [astro-ph]].
- [9] L. Canetti, M. Drewes and M. Shaposhnikov, *Sterile Neutrinos as the Origin of Dark and Baryonic Matter*, Phys. Rev. Lett. **110** (2013) no.6, 061801 [arXiv:1204.3902 [hep-ph]].
- [10] K. N. Abazajian, *Resonantly Produced 7 keV Sterile Neutrino Dark Matter Models and the Properties of Milky Way Satellites*, Phys. Rev. Lett. **112** (2014) no.16, 161303 [arXiv:1403.0954 [astro-ph.CO]].
- [11] J. Ghiglieri and M. Laine, *Sterile neutrino dark matter via coinciding resonances*, JCAP **07** (2020), 012 [arXiv:2004.10766 [hep-ph]].
- [12] [XRISM Science Team], *Science with the X-ray Imaging and Spectroscopy Mission (XRISM)*, [arXiv:2003.04962 [astro-ph.HE]].
- [13] A. Anisimov and P. Di Bari, *Cold Dark Matter from heavy Right-Handed neutrino mixing*, Phys. Rev. D **80** (2009), 073017 [arXiv:0812.5085 [hep-ph]].
- [14] A. Anisimov, *Majorana Dark Matter*, [arXiv:hep-ph/0612024 [hep-ph]].

- [15] P. Di Bari, P. O. Ludl and S. Palomares-Ruiz, *Unifying leptogenesis, dark matter and high-energy neutrinos with right-handed neutrino mixing via Higgs portal*, JCAP **11** (2016), 044 [arXiv:1606.06238 [hep-ph]].
- [16] P. Di Bari, K. Farrag, R. Samanta and Y. L. Zhou, *Density matrix calculation of the dark matter abundance in the Higgs induced right-handed neutrino mixing model*, JCAP **10** (2020), 029 [arXiv:1908.00521 [hep-ph]].
- [17] P. Gondolo, G. Gelmini and S. Sarkar, *Cosmic neutrinos from unstable relic particles*, Nucl. Phys. B **392** (1993), 111-136 [arXiv:hep-ph/9209236 [hep-ph]].
- [18] M. D’Onofrio, K. Rummukainen and A. Tranberg, *Sphaleron Rate in the Minimal Standard Model*, Phys. Rev. Lett. **113** (2014) no.14, 141602 [arXiv:1404.3565 [hep-ph]].
- [19] M. Drewes, Y. Georis and J. Klarić, *Mapping the Viable Parameter Space for Testable Leptogenesis*, Phys. Rev. Lett. **128** (2022) no.5, 051801 [arXiv:2106.16226 [hep-ph]].
- [20] A. Granelli, K. Moffat and S. T. Petcov, *Flavoured resonant leptogenesis at sub-TeV scales*, Nucl. Phys. B **973** (2021), 115597 [arXiv:2009.03166 [hep-ph]].
- [21] B. Patt and F. Wilczek, *Higgs-field portal into hidden sectors*, [arXiv:hep-ph/0605188 [hep-ph]].
- [22] P. Di Bari, D. Marfatia and Y. L. Zhou, *Gravitational waves from first-order phase transitions in Majoron models of neutrino mass*, JHEP **10** (2021), 193 [arXiv:2106.00025 [hep-ph]].
- [23] S. Davidson and A. Ibarra, *A Lower bound on the right-handed neutrino mass from leptogenesis*, Phys. Lett. B **535** (2002), 25-32 [arXiv:hep-ph/0202239 [hep-ph]].
- [24] A. Pilaftsis, *CP violation and baryogenesis due to heavy Majorana neutrinos*, Phys. Rev. D **56** (1997), 5431-5451 [arXiv:hep-ph/9707235 [hep-ph]].
- [25] A. Pilaftsis and T. E. J. Underwood, *Resonant leptogenesis*, Nucl. Phys. B **692** (2004), 303-345 [arXiv:hep-ph/0309342 [hep-ph]].
- [26] S. Blanchet and P. Di Bari, *New aspects of leptogenesis bounds*, Nucl. Phys. B **807** (2009), 155-187 [arXiv:0807.0743 [hep-ph]].
- [27] S. Antusch, S. Blanchet, M. Blennow and E. Fernandez-Martinez, *Non-unitary Leptonic Mixing and Leptogenesis*, JHEP **01** (2010), 017 [arXiv:0910.5957 [hep-ph]].

- [28] K. Moffat, S. Pascoli, S. T. Petcov and J. Turner, *Leptogenesis from Low Energy CP Violation*, JHEP **03** (2019), 034 [arXiv:1809.08251 [hep-ph]].
- [29] W. Buchmuller, P. Di Bari and M. Plumacher, *Cosmic microwave background, matter - antimatter asymmetry and neutrino masses*, Nucl. Phys. B **643** (2002), 367-390 [erratum: Nucl. Phys. B **793** (2008), 362] [arXiv:hep-ph/0205349 [hep-ph]].
- [30] R. Abbasi *et al.* [IceCube], *Searches for Connections between Dark Matter and High-Energy Neutrinos with IceCube*, [arXiv:2205.12950 [hep-ex]].
- [31] M. G. Aartsen *et al.* [IceCube], *Search for neutrinos from decaying dark matter with IceCube*, Eur. Phys. J. C **78** (2018) no.10, 831 [arXiv:1804.03848 [astro-ph.HE]].
- [32] H. A. Weldon, *Covariant Calculations at Finite Temperature: The Relativistic Plasma*, Phys. Rev. D **26** (1982) 1394. H. A. Weldon, *Effective Fermion Masses of Order gT in High Temperature Gauge Theories with Exact Chiral Invariance*, Phys. Rev. D **26** (1982) 2789. H. A. Weldon, *Simple Rules for Discontinuities in Finite Temperature Field Theory*, Phys. Rev. D **28** (1983) 2007.
- [33] M. Chianese, G. Miele, S. Morisi and E. Peinado, JCAP **12** (2018), 016 doi:10.1088/1475-7516/2018/12/016 [arXiv:1808.02486 [hep-ph]].
- [34] A. Bhattacharya, A. Esmaili, S. Palomares-Ruiz and I. Sarcevic, *Update on decaying and annihilating heavy dark matter with the 6-year IceCube HESE data*, JCAP **05** (2019), 051 [arXiv:1903.12623 [hep-ph]].
- [35] M. Y. Khlopov and A. D. Linde, Phys. Lett. B **138** (1984) 265; J. R. Ellis, J. E. Kim and D. V. Nanopoulos, Phys. Lett. B **145** (1984) 181; K. Kohri, T. Moroi and A. Yotsuyanagi, Phys. Rev. D **73** (2006) 123511; M. Kawasaki, K. Kohri, T. Moroi and A. Yotsuyanagi, *Big-Bang Nucleosynthesis and Gravitino*, Phys. Rev. D **78** (2008) 065011 [arXiv:0804.3745 [hep-ph]].
- [36] E. W. Kolb and A. J. Long, *Superheavy dark matter through Higgs portal operators*, Phys. Rev. D **96** (2017) no.10, 103540 [arXiv:1708.04293 [astro-ph.CO]].
- [37] D. J. H. Chung, E. W. Kolb and A. Riotto, Phys. Rev. Lett. **81** (1998), 4048-4051 doi:10.1103/PhysRevLett.81.4048 [arXiv:hep-ph/9805473 [hep-ph]].