











## SHORT REPORT

# Biallelic variants in *CEP164* cause a motile ciliopathy-like syndrome

Laura A. Devlin<sup>1</sup>  | Janice Coles<sup>2</sup> | Claire L. Jackson<sup>2</sup> | Miguel Barroso-Gil<sup>1</sup>  | Ben Green<sup>3</sup> | Woolf T. Walker<sup>2</sup> | N. Simon Thomas<sup>4,5</sup>  | James Thompson<sup>2</sup>  | Simon A. Rock<sup>6</sup>  | Ruxandra Neatu<sup>1</sup> | Laura Powell<sup>1</sup> | Elisa Molinari<sup>1</sup> | Genomics England Research Consortium | Ian J. Wilson<sup>7</sup> | Heather J. Cordell<sup>8</sup>  | Eric Olinger<sup>1</sup>  | Colin G. Miles<sup>1</sup> | John A. Sayer<sup>1,9,10</sup>  | Gabrielle Wheway<sup>4</sup>  | Jane S. Lucas<sup>2</sup> 

<sup>1</sup>Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK

<sup>2</sup>Primary Ciliary Dyskinesia Centre, NIHR Southampton Biomedical Research Centre, University of Southampton Faculty of Medicine and University Hospital Southampton NHS Foundation Trust, Southampton, UK

<sup>3</sup>Department of Respiratory Medicine, University Hospitals NHS Trust, Portsmouth, UK

<sup>4</sup>Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, UK

<sup>5</sup>Wessex Regional Genetics Laboratory, Salisbury NSF Foundation Trust, Salisbury District Hospital, Salisbury, UK

<sup>6</sup>North East Innovation Lab, The Newcastle upon Tyne Hospitals NHS Foundation Trust, The Biosphere, Newcastle upon Tyne, UK

<sup>7</sup>Biosciences Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK

<sup>8</sup>Population Health Sciences Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK

<sup>9</sup>Renal Services Centre, The Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK

<sup>10</sup>National Institute for Health Research Newcastle Biomedical Research Centre, Newcastle upon Tyne, UK

## Correspondence

John A. Sayer, Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Central Parkway, Newcastle upon Tyne, NE1 3BZ, UK.  
Email: [john.sayer@newcastle.ac.uk](mailto:john.sayer@newcastle.ac.uk)

Jane S. Lucas, Primary Ciliary Dyskinesia Centre, NIHR Southampton Biomedical Research Centre, University of Southampton Faculty of Medicine and University Hospital Southampton NHS Foundation Trust, Southampton, SO16 6YD, UK.  
Email: [jlucas1@soton.ac.uk](mailto:jlucas1@soton.ac.uk)

## Funding information

AAIR Charity, Grant/Award Number: 1129698; Kidney Research UK, Grant/Award Numbers: Paed\_RP\_001\_20180925, ST\_001\_20171120; MRC DiMeN DTP; National Institute for Health Research, Grant/Award Number: RfPB

## Abstract

Ciliopathies may be classed as primary or motile depending on the underlying ciliary defect and are usually considered distinct clinical entities. Primary ciliopathies are associated with multisystem syndromes typically affecting the brain, kidney, and eye, as well as other organ systems such as the liver, skeleton, auditory system, and metabolism. Motile ciliopathies are a heterogeneous group of disorders with defects in specialised motile ciliated tissues found within the lung, brain, and reproductive system, and are associated with primary ciliary dyskinesia, bronchiectasis, infertility and rarely hydrocephalus. Primary and motile cilia share defined core ultra-structures with an overlapping proteome, and human disease phenotypes can reflect both primary and motile ciliopathies. *CEP164* encodes a centrosomal distal appendage protein vital for primary ciliogenesis. Human *CEP164* mutations are typically described in patients with nephronophthisis-related primary ciliopathies but have also been implicated in

Laura A. Devlin, Janice Coles, Claire L. Jackson are joint first authors.

John A. Sayer and Jane S. Lucas joint corresponding authors.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Clinical Genetics* published by John Wiley & Sons Ltd.

PB-PG-1215-20014 200470; NHS England; Northern Counties Kidney Research Fund, Grant/Award Number: 2019/01; Swiss National Science Foundation, Grant/Award Number: P2ZHP3\_195181 & P500PB\_206851

motile ciliary dysfunction. Here we describe a patient with an atypical motile ciliopathy phenotype and biallelic *CEP164* variants. This work provides further evidence that *CEP164* mutations can contribute to both primary and motile ciliopathy syndromes, supporting their functional and clinical overlap, and informs the investigation and management of *CEP164* ciliopathy patients.

#### KEYWORDS

*CEP164*, ciliopathy, genetics, mutation, primary ciliary dyskinesia

## 1 | INTRODUCTION

The primary cilium is a microtubule-based cellular appendage, which functions as a vital signalling hub with key roles in organogenesis, growth, tissue function and regeneration. It is a non-motile organelle present on nearly every mammalian cell, consisting of a core basal body (BB), distal and sub-distal appendages, the central ciliary axoneme, and transition zone.<sup>1</sup> Primary ciliopathies are defined as a group of inherited multi-organ diseases sharing a critical defect in the biogenesis, functioning or maintenance of the primary cilium. They display a wide spectrum of overlapping clinical manifestations such as cystic kidney disease, retinal degeneration, neurological disorders, skeletal defects, and developmental defects.

Another subtype of cilia are motile cilia, which are located as groups of organelles and are required to move fluid or debris across the cell or move the cell within fluid.<sup>2</sup> They are found on specialised tissues including the respiratory tract, oviduct/fallopian tract, and ventricles of the central nervous system; the sperm flagellum is also a specialised motile cilium, as well as the embryonic nodal cilia. Motile cilia share a similar ultrastructure to primary cilia but typically have a central inner microtubule pair with accessory structures that form the motility apparatus. Abnormalities in the motile cilium typically lead to primary ciliary dyskinesia (PCD) or reduced generation of multiple motile cilia. Patients often present with defective mucociliary clearance and airway disease, including a daily wet cough from infancy, bronchiectasis, serous otitis media and rhino sinusitis. Aberrations in motile cilia can also cause male and female subfertility/infertility (common), *situs inversus* (50%), and hydrocephalus (rare).<sup>3</sup>

Centrosomal protein 164 kDa (*CEP164*) (OMIM: 614848) is a centriolar protein, localised to the distal appendages of the ciliary BB.<sup>4</sup> *CEP164* is essential for ciliary vesicle recruitment, correct BB docking and axonemal extension.<sup>4,5</sup> Recent studies have indicated that *CEP164* might also be involved in motile ciliogenesis, specifically in vesicle recruitment.<sup>6</sup> Biallelic *CEP164* variants have been identified in patients with nephronophthisis (NPHP)-related ciliopathies (NPHP-RC), a primary ciliopathy disease spectrum which can be accompanied by retinal, neurological, skeletal, and developmental abnormalities (Table S1).<sup>7–11</sup> There are a few patients recorded with *CEP164*-ciliopathy disease lacking a renal phenotype. More recently, *CEP164* variants have been associated with motile-ciliary phenotypes.<sup>12</sup>

Here we describe a patient with bronchiectasis in whom we identified biallelic *CEP164* mutations as the likely cause of an atypical motile ciliary phenotype. This finding has important consequences for

the investigation, management, and prognosis of patients with *CEP164* mutations, and for the diagnostic work up of those with PCD.

## 2 | MATERIALS AND METHODS


Genetic, clinical, and in vitro investigations were carried out on a patient diagnosed with non-cystic fibrosis (CF) bronchiectasis. Detailed methodology and ethical approval are presented within supporting information.

## 3 | RESULTS

### 3.1 | A patient with bronchiectasis and motile ciliary defects is explained by biallelic *CEP164* variants

The Genomics England 100,000 Genomes Project database is a rich data source for the study of ciliopathies. Filtering for biallelic or compound heterozygous *CEP164* variants in the rare disease cohort, we identified an adult patient recruited under “non-CF bronchiectasis” in whom we identified compound heterozygous stop gain variants in *CEP164* (Figure 1A, Figure S1). The first variant, (NM\_014956.5 c.1726C > T; NP\_055771.4:p.R576\*, NM\_001271933.2:c.1735C > T, NP\_001258862.1:p.R579\*) was a known *CEP164*-ciliopathy rare allele (Table 1, Table S1), with a gnomAD allele frequency of 0.000008<sup>7,12</sup> (Table 1). The second *CEP164* variant identified in this patient (NM\_014956.5 c.4228C > T; Q1410\*), was a stop gain in the penultimate exon, not been described previously in *CEP164*-ciliopathy patients, with an allele frequency in gnomAD of 0.0008 (Table 1). Each parent of the individual was heterozygous for one *CEP164* variant, confirming segregation of these alleles. No tier 1 or 2 pathogenic variants were found within the patient’s whole genome, or deleterious biallelic/compound heterozygous variants in bronchiectasis-associated genes *CFTR*, *SCNN1B*, *SCNN1A*, *SCNN1G*, 42 OMIM primary ciliary dyskinesia genes or 301 gold standard Syscilia genes (Tables S2 and S3), and the patient was considered unsolved by Genomics England. There were heterozygous variants found in 5 ciliary genes (*DNAH6*, *MYO15A*, *DNAH2*, *BBS9*, *RTTN*), however none of these genes had a second pathogenic variant (Table S4).

The patient, in his 50’s, of Caucasian origin, underwent deep phenotypic analysis as part of his diagnostic work-up for non-CF bronchiectasis, in particular PCD. The patient had a long-standing history of



**FIGURE 1** Non-CF bronchiectasis patient with *CEP164* compound heterozygous variants, motile ciliary defects and loss of *CEP164* from the ciliary brush border. (A) Pedigree diagram showing the single affected male patient (arrow) with non-CF bronchiectasis. Compound heterozygous stop gain *CEP164* variants, segregating from each parent were identified. The patient had phenotypes shown. RTI, respiratory tract infection. (B) Immunofluorescence labelling of NB and ALI cultured epithelial cells, showing expression of the centriolar marker gamma tubulin (red), *CEP164* (green) and DAPI nuclear counterstain. In the healthy controls, *CEP164* colocalises with gamma tubulin, marking the centriolar region. The patient's NB and ALI-cultured cells show loss of *CEP164* in the centriolar region (arrows), but some aggregate *CEP164* staining remains in the cell cytoplasm. (C) Still images from high-speed video microscopy showing the patient's NB epithelium which had uncoordinated cilia with reduced ciliary beat amplitude, and lack of mucociliary clearance. There is variable cilia length, with some long cilia (arrows). (D) TEM of patient's ALI cells show that motile cilia have a mostly normal microtubule structure. (E) Analysis of TEM images of motile cilia show that only a few cilia have microtubule defects, mainly disarranged microtubules (MT), but the ultrastructure is largely normal. The 108 cilia were counted. MT, microtubule; MTD, microtubular defect; IDA, inner dynein arm defect; ODA, outer dynein arm defect.

recurrent lower respiratory tract infections and asthma, with a lifelong wet cough. The patient also reported recurrent ear infections, had tinnitus and wears hearing aids. The patient was also overweight, with a BMI of 28. There was no evidence of kidney or retinal defects following a renal ultrasound and ophthalmologist review, and the patient was fertile, having fathered a son. The patient had some difficulty reading and mild learning difficulties, but formal cognitive testing was not completed.

The patient was diagnosed with probable PCD. The diagnostic findings included low-equivocal nasal nitric oxide (164 nL/min, Normal >250 nL/min, PCD usually <77 nL/min).<sup>13</sup> Analysis of the

patient's ciliated nasal epithelium following nasal brushings (NB) and air-liquid-interface (ALI) culture showed a complete loss of *CEP164* protein at the cilia brush border (Figure 1B), consistent with the pathogenicity of the identified *CEP164* alleles. *CEP164* intracellular aggregates were found in the NB and ALI-cultured cells, indicating mislocalisation of the *CEP164* protein. High-speed video microscopy analysis of the NB sample showed the motile cilia had a dyskinetic-uncoordinated ciliary beat (accurate ciliary beat frequencies was indeterminate), whilst after ALI-culture the ciliary function appeared to have improved beat coordination with a ciliary beat frequency of 14.5 Hz (within normal 'local' range, at 37°C), but an abnormal

**TABLE 1** Compound heterozygous *CEP164* variants identified in a non-CF bronchiectasis patient within The Genomics England 100,000 Genomes Project

| Diagnosis                             | Sex  | GRCh37                  | GRCh38       | Transcript                        | Coding change | Protein change | Exon  | Zygoty | GnomAD AF |
|---------------------------------------|------|-------------------------|--------------|-----------------------------------|---------------|----------------|-------|--------|-----------|
| Non-CF bronchiectasis (Candidate PCD) | Male | 11:117257920 (maternal) | 11:117387204 | <i>CEP164</i> : ENST00000278935.8 | c.1726C > T   | p.R576*        | 15/33 | Het    | 7.9 E-06  |
|                                       |      | 11:117282575 (paternal) | 11:117411859 |                                   | c.4228C > T   | p.Q1410*       | 32/33 | Het    | 0.000815  |

'staggered beat' was present. Elongated cilia were observed in both the NB and the ALI-cultured epithelial cells; most pronounced in NB samples (Figure 1C) (Movie S1, S2, S3). Transmission electron microscopy (TEM) following ALI culture showed the motile cilia had a largely normal ultrastructure (76.9%), with only few cilia with microtubule or dynein defects (23.1%) (Figure 1D, E).

## 4 | DISCUSSION

In this study we utilised the Genomics England 100,000 Genomes Project to identify and deep phenotype a non-CF bronchiectasis patient with compound heterozygous variants in *CEP164*. One variant (NM\_014956.5 c.1726C > T; R576\*) was a known allele, previously found homozygous in *CEP164* patients with NPHP-RC and motile ciliary defects.<sup>7,12</sup> The second variant (c.4228C > T; Q1410\*) is a stop gain within the penultimate exon, not previously described in *CEP164*-ciliopathy patients, however mutations in the final exon of *CEP164* have been previously identified.<sup>7</sup> Although this second variant is annotated with 'conflicting interpretation of pathogenicity' in ClinVar and has a relatively high gnomAD frequency (0.0008, no homozygotes recorded), the patient's NB and ALI-cultured epithelial cells demonstrated loss of *CEP164* at the base of the motile cilia brush border, supporting the pathogenicity of the identified *CEP164* variants. *CEP164* aggregates were found in the cytoplasm of the patient's cells, suggesting that *CEP164* may be mislocalised, potentially due to the loss of a functioning C-terminus which normally localises *CEP164* to the centriole.<sup>4</sup> The patient's NB and ALI-cultured epithelial cells demonstrated defects in ciliary movement but ultrastructure remained largely normal. A loss of cilia was not apparent, but cilia length was occasionally elongated. We suggest that *CEP164* may have roles within motile cilia aside from initial ciliary formation. This is the first *CEP164* patient described with a largely exclusive motile ciliary disease. The patient also showed a high BMI, which may further link *CEP164* variants to an obesity phenotype, as seen in ciliopathies such as Bardet-Biedl syndrome.<sup>8</sup>

The findings from our patient suggest that pathogenic variants in *CEP164* can cause both primary and motile ciliopathies, correlating with known *CEP164* expression patterns in the human and mouse.<sup>14</sup> Indeed, there is growing evidence for 'primary cilia genes' to have overlapping clinical manifestations with motile ciliopathies. Some patients with pathogenic variants in *INVS* have NPHP, *situs inversus totalis*, and motile cilia dysgenesis.<sup>15</sup> Also, mutations in *RPGR* and *OFD1* are found in some patients with PCD, often associated with elongated airway cilia.<sup>16,17</sup> Patients with polycystic kidney disease also have an increased risk of bronchiectasis, and often show PCD phenotypes.<sup>18</sup> Our patient with mutations in *CEP164* was previously found in a cohort analysis of bronchiectasis patients, whereby monogenic ciliopathy gene variants were found in 12% of the cohort.<sup>19</sup> With an overlapping ultrastructure, and evidence pointing towards a partially shared proteome, the role of typically 'primary cilia' genes in motile ciliopathies is now important. Extending genetic panels for patients with motile ciliary phenotypes such as bronchiectasis would allow more patients with these phenotypes to be diagnosed.

In conclusion, variants in *CEP164* now need to be correlated with both primary ciliopathy and motile ciliopathy phenotypes. Patients with *CEP164* genetic variants must be fully assessed and investigated for all relevant phenotypes to allow optimal and personalised management.

## AUTHOR CONTRIBUTIONS

Gabrielle Wheway, John A. Sayer, Colin G. Miles and Jane S. Lucas conceived the study with input from Laura A. Devlin. Gabrielle Wheway, Laura A. Devlin and Miguel Barroso-Gil carried out genomic research analysis, and N. Simon Thomas carried out clinical variant analysis. Ian J. Wilson, Eric Olinger, Heather J. Cordell and Ruxandra Neatu assisted with variant interpretation. Simon A. Rock, Laura Powell and Elisa Molinari assisted in drafting and revising the manuscript. Ben Green and Jane S. Lucas clinically phenotyped the patient. Janice Coles, Claire L. Jackson, James Thompson, Woolf T. Walker, and Jane S. Lucas collected and analysed the airway data, which was interpreted with Laura A. Devlin. Laura A. Devlin wrote the initial manuscript.

## ACKNOWLEDGEMENT

The authors thank the family for involvement in the study. See supporting information for full acknowledgements.

## CONFLICT OF INTEREST

The authors declare no competing interests.

## DATA AVAILABILITY STATEMENT

Data availability statement within manuscript: "Data availability The identified variants are available within the Genomics England 100,000 Genomes project rare disease arm."

## ORCID

Laura A. Devlin  <https://orcid.org/0000-0003-4251-9903>

Miguel Barroso-Gil  <https://orcid.org/0000-0001-8245-1929>

N. Simon Thomas  <https://orcid.org/0000-0002-1494-2467>

James Thompson  <https://orcid.org/0000-0002-9285-1317>

Simon A. Rock  <https://orcid.org/0000-0001-8531-1534>

Heather J. Cordell  <https://orcid.org/0000-0002-1879-5572>

Eric Olinger  <https://orcid.org/0000-0003-1178-7980>

John A. Sayer  <https://orcid.org/0000-0003-1881-3782>

Gabrielle Wheway  <https://orcid.org/0000-0002-0494-0783>

Jane S. Lucas  <https://orcid.org/0000-0001-8701-9975>

## REFERENCES

- Reiter JF, Leroux MR. Genes and molecular pathways underpinning ciliopathies. *Nat Rev Mol Cell Biol.* 2017;18(9):533-547.
- Mitchison HM, Valente EM. Motile and non-motile cilia in human pathology: from function to phenotypes. *J Pathol.* 2017;241(2):294-309.
- Lucas JS, Davis SD, Omran H, Shoemark A. Primary ciliary dyskinesia in the genomics age. *Lancet Respir Med.* 2020;8(2):202-216.
- Schmidt KN, Kuhns S, Neuner A, Hub B, Zentgraf H, Pereira G. Cep164 mediates vesicular docking to the mother centriole during early steps of ciliogenesis. *J Cell Biol.* 2012;199(7):1083-1101.
- Rosa ES, Binó L, Johnson CM, et al. Molecular mechanisms underlying the role of the centriolar CEP164-TTBK2 complex in ciliopathies. *Structure.* 2022;30(1):114-128 e9.
- Siller SS, Sharma H, Li S, et al. Conditional knockout mice for the distal appendage protein CEP164 reveal its essential roles in airway multiciliated cell differentiation. *PLoS Genet.* 2017;13(12):e1007128.
- Chaki M, Airik R, Ghosh AK, et al. Exome capture reveals ZNF423 and CEP164 mutations, linking renal ciliopathies to DNA damage response signaling. *Cell.* 2012;150(3):533-548.
- Maria M, Lamers IJC, Schmidts M, et al. Genetic and clinical characterization of Pakistani families with Bardet-Biedl syndrome extends the genetic and phenotypic spectrum. *Sci Rep.* 2016;6:34764.
- Fujimaru T., Genetic Background and Clinicopathologic Features of Adult-Onset Nephronophthisis 2021.
- Strong A, Simone L, Krentz A, et al. Expanding the genetic landscape of oral-facial-digital syndrome with two novel genes. *Am J Med Genet A.* 2021;185(8):2409-2416.
- Vilboux T, Doherty DA, Glass IA, et al. Molecular genetic findings and clinical correlations in 100 patients with Joubert syndrome and related disorders prospectively evaluated at a single center. *Genet Med.* 2017;19(8):875-882.
- Shamseldin HE, Shaheen R, Ewida N, et al. The morbid genome of ciliopathies: an update. *Genet Med.* 2020;22(6):1051-1060.
- Shapiro AJ, Dell SD, Gaston B, et al. Nasal nitric oxide measurement in primary ciliary dyskinesia. A technical paper on standardized testing protocols. *Ann Am Thorac Soc.* 2020;17(2):e1-e12.
- Devlin LA, Ramsbottom SA, Overman LM, et al. Embryonic and foetal expression patterns of the ciliopathy gene CEP164. *PLoS One.* 2020;15(1):e0221914.
- Moalem S, Keating S, Shannon P, et al. Broadening the ciliopathy spectrum: motile cilia dyskinesia, and nephronophthisis associated with a previously unreported homozygous mutation in the INVS/NPHP2 gene. *Am J Med Genet A.* 2013;161A(7):1792-1796.
- Moore A, Escudier E, Roger G, et al. RPGR is mutated in patients with a complex X linked phenotype combining primary ciliary dyskinesia and retinitis pigmentosa. *J Med Genet.* 2006;43(4):326-333.
- Bukowy-Bieryllo Z, Rabiasz A, Dabrowski M, et al. Truncating mutations in exons 20 and 21 of OFD1 can cause primary ciliary dyskinesia without associated syndromic symptoms. *J Med Genet.* 2019;56(11):769-777.
- Moua T, Zand L, Hartman RP, et al. Radiologic and clinical bronchiectasis associated with autosomal dominant polycystic kidney disease. *PLoS One.* 2014;9(4):e93674.
- Shoemark A, Griffin H, Wheway G, et al. Genome sequencing reveals underdiagnosis of primary ciliary dyskinesia in bronchiectasis. *Eur Respir J.* 2022;2200176. [Epub ahead of print].

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Devlin LA, Coles J, Jackson CL, et al. Biallelic variants in *CEP164* cause a motile ciliopathy-like syndrome. *Clinical Genetics.* 2022;1-5. doi:10.1111/cge.14251